

ABELIAN VARIETIES ARE DE RHAM $K(\pi, 1)$

VÕ QUỐC BAO AND QUANG-KHAI NGUYEN

ABSTRACT. Motivated by the work of Esnault-Hai, one has the notion of *de Rham $K(\pi, 1)$ schemes*, defined as follows. Given a smooth proper geometrically connected scheme X over a field k of characteristic 0 and a base point $x \in X(k)$, one can define its differential fundamental group $\pi^{\text{diff}}(X/k)$, which comes from the Tannakian duality of the category of coherent integrable connections on X . Using the formalism of δ -functor, one can define natural morphisms between the group-scheme cohomology of $\pi^{\text{diff}}(X/k)$ and the de Rham cohomology of X . One says that X with $x \in X(k)$ is *de Rham $K(\pi, 1)$* if such morphisms are all isomorphisms. In this article, we prove that abelian varieties in characteristic 0 are *de Rham $K(\pi, 1)$* . In the second part of the article, we study the group-scheme cohomology of the abelianization of the differential fundamental group of a smooth proper geometrically connected scheme via its Albanese variety.

CONTENTS

1. Introduction	1
2. De Rham $K(\pi, 1)$	5
3. Differential fundamental group	10
4. Abelian varieties are de Rham $K(\pi, 1)$	11
5. Cohomology of the abelianization of the differential fundamental group	14
References	17

1. INTRODUCTION

Given a topological space X , the relation between the group cohomology of the topological fundamental group and sheaf cohomology was first studied by H. Cartan and J. Leray

2020 *Mathematics Subject Classification.* 14F40, 14F43, 14L15, 18G15, 18G40, 18M25.

Key words and phrases. de Rham cohomology; group scheme cohomology; Tannakian duality.

The work of Võ Quốc Bao is supported by the Vingroup Innovation Foundation under grant number VINIF.2024.TS.003 and by the IMU Breakout Graduate Fellowship Program. Quang-Khai Nguyen would like to thank the Vietnam Institute of Mathematics and Labex MILYON for financial support during his master's studies, and the European Union's MSCA–Horizon Europe programme (grant agreement No. 101126554) for funding his doctoral studies.

(see [CE56, Chapter XVI.8]). In some cases, these cohomologies are naturally isomorphic, and when this phenomenon happens, we say that X is $K(\pi, 1)$. A similar story has also been observed for other fundamental groups and cohomologies. For example, étale $K(\pi, 1)$ -spaces are schemes for which a comparison theorem holds between the group cohomology of the étale fundamental group and étale cohomology (see [Gro73, Ach15, Ach17]). In his ground-breaking work [Sch13], P. Scholze showed that a connected affinoid rigid-analytic variety is a pro-étale $K(\pi, 1)$ -space. In [EH06], H. Esnault and P. H. Hai studied the comparison between the cohomology of the differential fundamental group and de Rham cohomology, which marked the beginning of the investigation into a de Rham analog of $K(\pi, 1)$ spaces. In [BHT25], the authors showed that a smooth projective curve of genus greater than 0 is de Rham $K(\pi, 1)$. In this work, we aim to continue the study in *loc. cit* by showing that abelian varieties are de Rham $K(\pi, 1)$.

The notion of the differential fundamental group is first introduced in [Del89] and developed further in [EH06], which can be defined as follows. Let X be a smooth and geometrically connected algebraic variety over a field k of characteristic 0, and let $x \in X(k)$ be a rational point. The *differential fundamental group* $\pi^{\text{diff}}(X/k)$ of X with base point x is an affine group scheme over k , defined via Tannakian duality. Consider the category $\text{MIC}^{\text{coh}}(X)$ of \mathcal{O}_X -coherent sheaves equipped with k -linear integrable connections. This category is a k -linear abelian rigid tensor category equipped with the fiber functor at x

$$x^* : \text{MIC}^{\text{coh}}(X) \longrightarrow \text{Vec}_k, \quad (\mathcal{V}, \nabla) \mapsto \mathcal{V}_x,$$

which is faithfully exact, hence a neutral Tannakian category. By the Tannakian duality (see [DM82, Theorem 2.11]), this yields an affine group scheme $\pi^{\text{diff}}(X/k)$ over k such that the functor x^* induces an equivalence of tensor categories:

$$(1) \quad x^* : \text{MIC}^{\text{coh}}(X) \xrightarrow{\cong} \text{Rep}^f(\pi^{\text{diff}}(X/k)),$$

where $\text{Rep}^f(\pi^{\text{diff}}(X/k))$ denotes the category of finite-dimensional k -linear representations of $\pi^{\text{diff}}(X/k)$. This equivalence extends to an equivalence between ind-categories (i.e., between the category of inductive direct limits of vector bundles with connection and the category of all representations of $\pi^{\text{diff}}(X/k)$):

$$(2) \quad x^* : \text{MIC}^{\text{ind}}(X) \xrightarrow{\cong} \text{Rep}(\pi^{\text{diff}}(X/k)).$$

Let $H_{\text{dR}}^\bullet(X, (\mathcal{V}, \nabla))$ denote the de Rham cohomology of the scheme X with coefficients in an integrable connection (\mathcal{V}, ∇) (as defined in [Kat70]), and set $V := x^*(\mathcal{V}, \nabla)$. Let $H^\bullet(\pi^{\text{diff}}(X/k), V)$ denote the group-scheme cohomology of the affine group scheme $\pi^{\text{diff}}(X/k)$ with coefficients in the k -linear representation V of $\pi^{\text{diff}}(X/k)$, see [DG70, I] or [Jan03]. The equivalence (2) gives us the following morphism

$$(3) \quad \delta_{X/k}^i(V) : H^i(\pi^{\text{diff}}(X/k), V) \longrightarrow H_{\text{dR}}^i(X/k, (\mathcal{V}, \nabla))$$

for every $i \geq 0$. In light of [BHT25], we propose the following definition of *de Rham $K(\pi, 1)$* schemes for such data given above, which can be seen as an analogue of the notion of $K(\pi, 1)$ spaces.

Definition 1.1. A smooth geometrically connected scheme X together with a point $x \in X(k)$ is said to be *de Rham $K(\pi, 1)$* if $\delta_{X/k}(V)$ is an isomorphism for all $i \geq 0$ and for every finite-dimensional $\pi^{\text{diff}}(X/k)$ -module V .

Achinger suggested in his comments [Ach23] that abelian varieties should be de Rham $K(\pi, 1)$. Our first result confirms Achinger's expectation.

Theorem A. (*Theorem 4.3*) *Every abelian variety of dimension g over a field of characteristic 0 is de Rham $K(\pi, 1)$.*

Remark 1.2. For an abelian variety X , its differential fundamental group, up to isomorphism, does not depend on the choice of the base point, thanks to the translation. Thus, from now on, we will always consider the base point of an abelian variety to be its identity 0_X .

The main difficulty in proving Theorem A is that one can not use the Poincaré duality as in the proof of [BHT25, Theorem 4.1] for curves. Instead, we employ a transcendental method. We first prove our result when $k = \mathbb{C}$ by translating the problem into one in algebraic topology and verifying that it holds in the topological setting. In the general case, we can find a finitely generated subfield k' of k for which our given data can be defined by X' over k' . At this stage, we can compare the differential fundamental groups of X , X' and $X' \otimes_{k'} \mathbb{C}$, and compare their corresponding cohomologies by using the Lyndon-Hochschild-Serre spectral sequence. The key idea is to exploit the fact that the differential fundamental groups of abelian varieties are commutative, from which we can decompose them into direct sums of unipotent parts and reductive ones. To conclude, we apply the GAGA principle and use several base change properties of the group-scheme cohomology.

Remark 1.3. The transcendental method has also been used to study the homotopy sequence of the differential fundamental groups [Zha13].

Since de Rham cohomologies of abelian varieties are finite-dimensional, we obtain the following.

Corollary B (Corollary 4.4). *For an abelian variety of dimension g over a field of characteristic 0, the group-scheme cohomology of its differential fundamental group is always finite-dimensional and vanishes in degrees greater than $2g$. Moreover, the Euler characteristic of the group-scheme cohomology of its differential fundamental group is well-defined and equal to zero.*

Note, however, that in general we do not know whether the group-scheme cohomology of the differential fundamental group of a smooth proper geometrically connected scheme is finite-dimensional or not.

Our second study in this article involves a relation between the group-scheme cohomology of the differential fundamental group of a smooth proper geometrically connected scheme and that of its associated Albanese variety.

Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and $x \in X(k)$. Let $f : X \rightarrow \text{Alb}_X$ be the Albanese morphism from a smooth projective scheme to its Albanese variety such that f maps x to the identity element 0_{Alb_X} . Then we have the following functor:

$$f^* : \text{MIC}^{\text{coh}}(\text{Alb}_X/k) \rightarrow \text{MIC}^{\text{coh}}(X/k).$$

The pullback functor f^* is compatible with fiber functors x^* and $0_{\text{Alb}_X}^*$, hence it induces a morphism of differential fundamental groups

$$f_* : \pi^{\text{diff}}(X/k) \rightarrow \pi^{\text{diff}}(\text{Alb}_X/k),$$

see [DM82, Corollary 2.9]. Since the differential fundamental group $\pi^{\text{diff}}(\text{Alb}_X)$ is commutative by Lemma 3.2 below, the morphism f_* factor as follows:

$$\begin{array}{ccc} \pi^{\text{diff}}(X) & \longrightarrow & \pi^{\text{diff}}(\text{Alb}_X), \\ \downarrow & \nearrow & \\ \pi^{\text{diff}}(X)^{\text{ab}} & & \end{array}$$

where $\pi^{\text{diff}}(X)^{\text{ab}}$ is the largest abelian quotient (see the argument above Corollary 22 in [dS07]).

Using the projection formula, one sees that the functor f^* is fully faithful. Since the natural morphism

$$\pi^{\text{diff}}(X) \longrightarrow \pi^{\text{diff}}(X)^{\text{ab}}$$

is faithfully flat, it follows that the following functor, induced by $\pi^{\text{diff}}(X/k)^{\text{ab}} \longrightarrow \pi^{\text{diff}}(\text{Alb}_X/k)$,

$$\text{Rep}^f(\pi^{\text{diff}}(X/k)^{\text{ab}}) \longrightarrow \text{Rep}^f(\text{Alb}_X/k)$$

is fully faithful. This fully faithful functor extends to a fully faithful functor between ind-categories:

$$\text{Rep}(\pi^{\text{diff}}(X/k)^{\text{ab}}) \longrightarrow \text{Rep}(\text{Alb}_X/k).$$

Consequently,

$$V^{\pi^{\text{diff}}(\text{Alb}_X/k)} = V^{\pi^{\text{diff}}(X/k)^{\text{ab}}}$$

for any $V \in \text{Obj}(\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k)))$. Using universal δ -functor (see [Har77, page 205]) we obtain the following morphism for every $i \geq 0$:

$$(4) \quad \eta^i(V) : H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \longrightarrow H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V),$$

where V is an object of $\text{Rep}^f(\pi^{\text{diff}}(X/k)^{\text{ab}})$.

Theorem C (Theorem 5.4). *Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and a base point $x \in X(k)$. Let f be the Albanese morphism mapping x to 0_{Alb_X} , the identity. Then the morphism*

$$\eta^i(V) : H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \longrightarrow H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

is an isomorphism for all $i \geq 0$ and all $V \in \text{Obj}(\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k)))$.

Our idea to prove Theorem C comes from the complex picture, in which one can see that $\pi^{\text{diff}}(\text{Alb}_X/k)$ and $\pi^{\text{diff}}(X/k)^{\text{ab}}$ have isomorphic unipotent radicals. For general k , this observation can be proved by using the fact that the first de Rham cohomology of X and Alb_X are isomorphic, whose proof itself is transcendental. We then conclude by using the Lyndon-Hochschild-Serre spectral sequence.

Since the group-scheme cohomologies of the differential fundamental group of an Albanese variety are finite-dimensional, we have the following result involving the Euler characteristic of $\pi^{\text{diff}}(X/k)^{\text{ab}}$.

Corollary D (Corollary 5.5). *Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and a base point $x \in X(k)$. Then the group-scheme cohomology*

$$H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

is finite-dimensional for any $i \geq 0$ and any $V \in \text{Obj}(\text{Rep}^f(\pi^{\text{diff}}(X/k)^{\text{ab}}))$. Moreover, the Euler characteristic of the group-scheme cohomology of $\pi^{\text{diff}}(X/k)^{\text{ab}}$ is well-defined and equal to zero.

Using Theorem C, one can obtain similar isomorphisms between group cohomologies of the groups $\pi_1^{\text{top}}(X, x)^{\text{ab}}$ and $\pi_1^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X})$.

Corollary E. (Corollary 5.6) *Let X be a smooth projective connected complex variety, and x be a base point. Let f be the Albanese morphism sending x to 0_{Alb_X} . Then f induces isomorphisms*

$$H^i(\pi_1^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X}), V) \cong H^i(\pi_1^{\text{top}}(X, x)^{\text{ab}}, V)$$

for all $i \geq 0$ and all $V \in \text{Obj}(\text{Rep}^f(\pi_1^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X})))$.

The article is organized as follows. Section 2 is devoted to providing general properties of de Rham $K(\pi, 1)$ schemes and its relation to the pro-algebraic completion and to $K(\pi, 1)$ spaces (in the usual topological sense). In Section 3, we provide some properties of the differential fundamental groups that will be useful later on. Section 4 is devoted to prove Theorem A and Corollary B. In section 5, we prove Theorem C, Corollary D, and Corollary E.

1.1. Notations and Conventions.

- (1) k is a field of characteristic 0.
- (2) For a smooth connected scheme X over k , let $\text{MIC}(X/k)$ denote the category of quasi-coherent \mathcal{O}_X -sheaves equipped with k -linear integrable connections.
- (3) We denote $\text{MIC}^{\text{coh}}(X/k)$ the full subcategory of $\text{MIC}(X/k)$ consisting of coherent \mathcal{O}_X -sheaves equipped with k -linear integrable connections.
- (4) We denote $\pi_1^{\text{top}}(X, x)$ (resp. $\pi^{\text{diff}}(X/k, x)$) the topological fundamental group (resp. the differential fundamental group). If the base point x is clear from the context, we use $\pi^{\text{diff}}(X/k)$ to simplify the notation.
- (5) When comparing the group-scheme cohomology of the fundamental group with de Rham cohomology, we always assume that V is the representation corresponding to the integrable connection (\mathcal{V}, ∇) via the fiber functor.
- (6) We always use the subscript to denote the result of base change (of schemes, morphisms, linear maps).

2. DE RHAM $K(\pi, 1)$

The aim of this section is to review the notions of algebraically good groups, $K(\pi, 1)$ spaces, de Rham $K(\pi, 1)$ schemes and their relations.

2.1. Algebraically good group. We recall the notion of the *proalgebraic completion* of an abstract group (see [BLMM02], [Den23]).

Definition 2.1. Let Γ be an abstract group. The *proalgebraic completion* of Γ over a field k , denoted Γ^{alg} , is a proalgebraic group equipped with a morphism of groups

$$\iota: \Gamma \longrightarrow \Gamma^{\text{alg}}(k)$$

satisfying the following universal property: for any representation $\theta: \Gamma \rightarrow \text{GL}(V)$ on a finite-dimensional k -vector space V , there is a unique representation $\theta^{\text{alg}}: \Gamma^{\text{alg}} \rightarrow \text{GL}(V)$ such that

the diagram

$$\begin{array}{ccc} \Gamma & \xrightarrow{\iota} & \Gamma^{\text{alg}}(k) \\ & \searrow \theta & \downarrow \theta^{\text{alg}} \\ & & \text{GL}(V) \end{array}$$

commutes.

Let $\text{Rep}^f(\Gamma)$ be the category of k -linear finite-dimensional representations of Γ and $\text{Rep}^f(\Gamma^{\text{alg}})$ the category of k -linear finite-dimensional representation of the affine group scheme Γ^{alg} . According to the definition of pro-algebraic completion, the inclusion functor

$$(5) \quad \text{Rep}^f(\Gamma^{\text{alg}}) \longrightarrow \text{Rep}^f(\Gamma)$$

yields an equivalence of categories.

Example 2.2. *The Riemann-Hilbert correspondence, established by Deligne [Del70], provides an equivalence of Tannakian categories*

$$\text{Rep}^f\left(\pi_1^{\text{top}}(X^{\text{an}}, x)\right) \cong \text{MIC}^{\text{coh}}(X/\mathbb{C}),$$

for any complex smooth projective variety X . In other words, the differential fundamental group $\pi^{\text{diff}}(X/\mathbb{C})$ is the pro-algebraic completion over \mathbb{C} of the topological fundamental group $\pi_1^{\text{top}}(X^{\text{an}}, x)$.

The equivalence (5) extends to an equivalence between the ind-category $\text{Rep}^{\text{ind}}(\Gamma^{\text{alg}})$ and the ind-category $\text{Rep}^{\text{ind}}(\Gamma^{\text{alg}})$. Since every k -linear representation of an affine group scheme is a direct limit of k -linear finite representations (see [Wat79]), we have the following inclusion functor:

$$(6) \quad \text{Rep}(\Gamma^{\text{alg}}) \xrightarrow{\cong} \text{Rep}^{\text{ind}}(\Gamma^{\text{alg}}) \hookrightarrow \text{Rep}(\Gamma).$$

For every V a representation of Γ^{alg} , let $H^{\bullet}(\Gamma^{\text{alg}}, V)$ denote the group-scheme cohomology of the affine group scheme Γ^{alg} with coefficients in V (see [[DG70], I] or [Jan03]). Let $H^{\bullet}(\Gamma, V)$ denote the group cohomology of the abstract group Γ with coefficients in V (see [Bro94]). The equivalence (6) induces, for all $i \geq 0$, natural morphisms

$$\iota^i : H^i(\Gamma^{\text{alg}}, V) \longrightarrow H^i(\Gamma, V).$$

In general, the maps δ^i are not isomorphisms, and we are interested in the class of groups Γ for which δ^i is an isomorphism.

Definition 2.3. Let Γ be an abstract group, and let V be a finite-dimensional k -linear representation of Γ^{alg} . Then Γ is said to be *algebraically good* (over k) if for every $i \geq 0$, the canonical map

$$\iota^i : H^i(\Gamma^{\text{alg}}, V) \xrightarrow{\cong} H^i(\Gamma, V)$$

is an isomorphism.

Remark 2.4. We find that it is more convenient for us to define the definition of an algebraically good group in this way. The original one is defined differently, see [KPT09], which is equivalent to Definition 2.3, cf. [KPT09, Lemma 4.11].

Example 2.5. *The following groups are algebraically good over any field k of characteristic 0.*

- (1) *Finite groups*
- (2) *Free groups of finite type*
- (3) *Abelian groups of finite type* ([KPT09, Proposition 4.13]).

2.2. A review of $K(\pi, 1)$ spaces. We recall in this section the notion of $K(\pi, 1)$ spaces, as presented in [Ach15, Section 2.1.1]. Let X be a path-connected, semi-locally simply connected topological space, and we fix a base point $x \in X$. Classical covering space theory identifies its fundamental group $\pi_1^{\text{top}}(X, x)$ as the automorphism group of the fiber functor that sends a covering $f : X' \rightarrow X$ to its fiber $f^{-1}(x)$. This induces an equivalence between the category of covering spaces of X and the category of right $\pi_1^{\text{top}}(X, x)$ -sets. Thus \tilde{X} is a principal $\pi_1^{\text{top}}(X, x)$ -bundle over X . The functor $X \mapsto$ (isomorphism classes of principal $\pi_1^{\text{top}}(X, x)$ bundles on X) is represented by a classifying space $B\pi_1^{\text{top}}(X, x)$ with a universal principal $\pi_1^{\text{top}}(X, x)$ -bundle $E\pi_1^{\text{top}}(X, x) \rightarrow B\pi_1^{\text{top}}(X, x)$, where $E\pi_1^{\text{top}}(X, x)$ is the total space of $\pi_1^{\text{top}}(X, x)$. Hence, there exists a map of topological spaces

$$\rho : X \rightarrow B\pi_1^{\text{top}}(X, x)$$

such that the universal covering $\tilde{X} \rightarrow X$ is the pullback of the universal principal $\pi_1^{\text{top}}(X, x)$ -bundle $E\pi_1^{\text{top}}(X, x) \rightarrow B\pi_1^{\text{top}}(X, x)$.

Let $\text{Loc}(X)$ be a category of local systems on X . There is a functor

$$\rho^* : \pi_1^{\text{top}}(X, x) \rightarrow \text{Sheaves}(X)$$

which is fully faithful and its essential image is $\text{Loc}(X)$ (see [Del70, Corollary 1.4]). Since

$$(-)^{\pi_1^{\text{top}}(X, x)} = \Gamma(X, \rho^*(-)),$$

using universal δ -functor (see [Har77, page 205]) we obtain the following morphism for every $i \geq 0$:

$$\rho^i : H^i(\pi_1^{\text{top}}(X, x), M) \rightarrow H^i(X, \rho^* M),$$

where M is a $\pi_1^{\text{top}}(X, x)$ -module. In general, the maps ρ^i are not isomorphisms; and we are interested in the class of topological spaces X for which ρ^i is an isomorphism.

Proposition 2.6. ([Ach15, Proposition 2.1.1]) *The following properties are equivalent:*

- (a) $\pi_i^{\text{top}}(X) = 0$ for all $i > 1$;
- (b) *the universal cover \tilde{X} is weakly contractible*;
- (c) *for every locally constant sheaf \mathcal{F} on X , the pullback maps $H^i(X, \mathcal{F}) \rightarrow H^i(\tilde{X}, p^* \mathcal{F})$ are zero for all $i > 0$;*
- (d) *for every locally constant sheaf \mathcal{F} on X and any class $\omega \in H^i(X, \mathcal{F})$, $i > 0$, there exists a covering $f : X' \rightarrow X$ such that $f^* \omega = 0$ in $H^i(X', f^* \mathcal{F})$;*
- (e) *the natural transformation $\text{id} \rightarrow R\rho_* \rho^*$ is an isomorphism*;
- (f) *the maps ρ^i are isomorphisms for any $\pi_1^{\text{top}}(X)$ -module M and for all $i > 0$;*
- (g) *the map $\rho : X \rightarrow B\pi_1^{\text{top}}(X, x)$ is a weak homotopy equivalence.*

Motivated by Proposition 2.6, one defines:

Definition 2.7. ([Ach15, Definition 2.1.2]) A topological space X is called a $K(\pi, 1)$ space if it satisfies one of the equivalent conditions in Proposition 2.6.

Example 2.8. For all $n \geq 1$, the complex tori $(\mathbb{S}^1)^n$ are $K(\pi, 1)$ spaces since their universal covers are \mathbb{C}^n , which are contractible.

2.3. De Rham $K(\pi, 1)$ schemes. Let X be a smooth geometrically connected over a field k of characteristic 0. Assume that X has a rational point $x \in X(k)$. In this section, we will recap the notion of de Rham $K(\pi, 1)$ schemes.

We start by constructing the map (3). Let V be an object of $\text{Rep}(\pi^{\text{diff}}(X/k))$. The invariant subspace of V is defined to be (k being equipped with the trivial action):

$$V^{\pi^{\text{diff}}(X/k)} := \text{Hom}_{\pi^{\text{diff}}(X/k)}(k, V).$$

Since the fixed point functor $V \mapsto V^{\pi^{\text{diff}}(X/k)}$ is left exact, one can define the group-scheme cohomology $H^i(\pi^{\text{diff}}(X/k), V)$ as the i -th right derived functor of this functor. Since $\text{Rep}(\pi^{\text{diff}}(X/k))$ has enough injectives, we get isomorphisms of right derived functors, as stated in [Jan87, 4.2]:

$$(7) \quad H^i(\pi^{\text{diff}}(X/k), -) \cong \text{Ext}_{\text{Rep}(\pi^{\text{diff}}(X/k))}^i(k, -).$$

For a sheaf with an integrable connection (\mathcal{V}, ∇) on X/k , the *sheaf of horizontal sections* is defined as

$$(8) \quad \mathcal{V}^\nabla := \text{Ker}(\nabla : \mathcal{V} \rightarrow \Omega_{X/k}^1 \otimes \mathcal{V}).$$

This is a sheaf of k -vector spaces. The *0-th de Rham cohomology* of \mathcal{V} is defined as

$$H_{\text{dR}}^0(X/k, (\mathcal{V}, \nabla)) := f_*(\mathcal{V}^\nabla).$$

The vector space $H_{\text{dR}}^0(X/k, (\mathcal{V}, \nabla))$ can be identified with the hom-set of connections in $\text{MIC}(X/k)$

$$\{\varphi : (\mathcal{O}_X, d) \longrightarrow (\mathcal{M}, \nabla)\}.$$

Since $\text{MIC}(X/k)$ is equivalent to the category of left modules on the sheaf of differential operators $\mathcal{D}_{X/k}$, it has enough injectives (cf. [Kat70]). Thus, we can define the *higher de Rham cohomologies* $H_{\text{dR}}^i(X/k, -)$ to be the derived functors of the functor

$$H_{\text{dR}}^0(X/k, -) : \text{MIC}(X/k) \longrightarrow \text{Vec}_k.$$

Recall that the differential fundamental group $\pi^{\text{diff}}(X)$ is defined via the Tannakian duality (1). It follows that

$$(-)^{\pi^{\text{diff}}(X/k)} = \text{Hom}_{\text{Rep}^f(\pi^{\text{diff}}(X/k))}(k, -) = \text{Hom}_{\text{MIC}(X/k)}(\mathcal{O}_X, -) = H_{\text{dR}}^0(X/k, -),$$

and by using the universal δ -functor, we obtain the following morphisms:

$$(9) \quad \delta_{X/k}^i(V) : H^i(\pi^{\text{diff}}(X/k), V) \longrightarrow H_{\text{dR}}^i(X/k, (\mathcal{V}, \nabla)) \text{ for all } i \geq 0,$$

where V is a $\pi^{\text{diff}}(X/k)$ -module associated with the connection (\mathcal{V}, ∇) .

Remark 2.9. The equivalence of Tannakian categories (1) yields that

$$\delta_{X/k}^1(V) : H^1(\pi^{\text{diff}}(X/k), V) \longrightarrow H_{\text{dR}}^1(X/k, (\mathcal{V}, \nabla))$$

is an isomorphism for all V . However, higher morphisms $\delta^i, i \geq 2$ might fail to be isomorphic, for example, when $X = \mathbb{P}^1$.

When X with the base point $x \in X(K)$ satisfy $\delta_{X/k}^i(-)$ is an isomorphism for all $i \geq 0$, we say that X is de Rham $K(\pi, 1)$, see Definition 1.1.

2.4. A relationship between algebraically good groups, $K(\pi, 1)$ spaces, and de Rham $K(\pi, 1)$ schemes. When $k = \mathbb{C}$, we can relate these notions mentioned above as follows.

Proposition 2.10. *Let X be a smooth projective connected scheme over \mathbb{C} , X^{an} the associated analytic space, and $x \in X(\mathbb{C})$. Consider the following three conditions.*

- (1) X is de Rham $K(\pi, 1)$.
- (2) X^{an} is $K(\pi, 1)$.
- (3) The fundamental group $\pi_1^{\text{top}}(X^{\text{an}}, x)$ is algebraically good.

Then (1)&(2) \Rightarrow (3) and (2)&(3) \Rightarrow (1).

Proof. According to Deligne's Riemann-Hilbert correspondence [Del70, Theorem 2.17], the functor

$$\begin{aligned} \text{MIC}^{\text{coh}}(X^{\text{an}}/\mathbb{C}) &\longrightarrow \text{Loc}(X^{\text{an}}) \\ (\mathcal{V}^{\text{an}}, \nabla^{\text{an}}) &\mapsto \tilde{V} := (\mathcal{V}^{\text{an}})^{\nabla} \end{aligned}$$

is an equivalence between these two categories. Since $H_{\text{dR}}^0(X/\mathbb{C}, -) = H_{\text{dR}}^0(X^{\text{an}}/\mathbb{C}, -) = \Gamma(X^{\text{an}}, -)$, using the universal δ -functor, we obtain the following morphism:

$$\psi^i : H_{\text{dR}}^i(X/\mathbb{C}, (\mathcal{V}, \nabla)) \longrightarrow H^i(X^{\text{an}}, \tilde{V}),$$

where (\mathcal{V}, ∇) is an integrable connection on X^{an} .

Let (\mathcal{V}, ∇) be an object in $\text{MIC}^{\text{coh}}(X/\mathbb{C})$. For every $i \geq 0$, the morphism $\delta_{X/k}^i(V)$ (see (9)) fits into the following commutative diagram

$$\begin{array}{ccc} H^i(\pi^{\text{diff}}(X/\mathbb{C}), V) & \xrightarrow{\delta_{X/k}^i(V)} & H_{\text{dR}}^i(X/\mathbb{C}, (\mathcal{V}, (\mathcal{V}, \nabla))) \\ \downarrow \iota^i & & \downarrow \psi^i \\ H^i(\pi_1^{\text{top}}(X^{\text{an}}, x), V) & \xrightarrow{\rho^i} & H^i(X^{\text{an}}, \tilde{V}) \end{array}$$

since all the morphisms in the above diagram are constructed by the universal δ -functor "formalism". Deligne's comparison theorem [Del70, Theorem 6.12] implies that the morphism ψ^i is an isomorphism, and the result follows since ι^i is an isomorphism for all $i \geq 0$ by Example 2.2. \square

Remark 2.11. Proposition 2.10 can be seen as a de Rham version of [Ach15, Proposition 2.1.15].

Corollary 2.12. *Every abelian variety of dimension g over \mathbb{C} is a de Rham $K(\pi, 1)$ scheme.*

Proof. Recall that every abelian variety over \mathbb{C} is a complex torus, which is a $K(\pi, 1)$ space by Example 2.8. In addition, their fundamental groups are abelian groups of finite type, hence algebraically good by Example 2.5. We then conclude by applying Proposition 2.10. \square

3. DIFFERENTIAL FUNDAMENTAL GROUP

To prove Theorem A, we need to reduce the problem to the case over \mathbb{C} . In order to do it, we need to understand how the differential fundamental groups behave under base change.

3.1. Differential fundamental groups and base change. Let X be a smooth geometrically connected over a field k of characteristic 0 and let $x \in X(k)$. Let L be a field extension of k , and $X_L := X \otimes_k L$. We have the following Cartesian diagram

$$\begin{array}{ccc} X_L & \xrightarrow{g} & X \\ \downarrow & & \downarrow \\ \text{Spec}(L) & \longrightarrow & \text{Spec}(k). \end{array}$$

Let $\mathcal{C}(X_L)$ be the full subcategory of $\text{MIC}^{\text{coh}}(X_L/L)$ whose objects, after being pushed forward along the map $X_L \xrightarrow{g} X$, belongs to the category $\text{Ind}(\text{MIC}^{\text{coh}}(X/k))$. This $\mathcal{C}(X_L)$ is a Tannakian subcategory and its Tannakian group is precisely $\pi^{\text{diff}}(X/k)_L := \pi^{\text{diff}}(X/k) \times_k L$ thanks to [Del89, 10.38 and 10.41] and we have a map $\pi^{\text{diff}}(X_L/L) \rightarrow \pi^{\text{diff}}(X/k)_L$ induced by the inclusion of $\mathcal{C}(X_L)$ in $\text{MIC}^{\text{coh}}(X_L/L)$.

Lemma 3.1. *The map $\pi^{\text{diff}}(X_L/L) \rightarrow \pi^{\text{diff}}(X/k)_L$ is faithfully flat.*

Proof. Since $\mathcal{C}(X_L)$ is a full subcategory of $\text{MIC}^{\text{coh}}(X/L)$ and is stable under taking subquotients, the result follows from the Tannakian criterion for faithfully flat morphisms in [DM82]. \square

It follows that the fixed points $(-)^{\pi^{\text{diff}}(X/k)_L}$ and $(-)^{\pi^{\text{diff}}(X_L/L)}$ coincide. The universal δ -functor formalism yields that for every $\pi^{\text{diff}}(X/k)_L$ -module V_L , we have natural maps

$$\pi_L^i(V_L) : H^i(\pi^{\text{diff}}(X/k)_L, V_L) \rightarrow H^i(\pi^{\text{diff}}(X_L/L), V_L), i \geq 0.$$

3.2. Differential fundamental groups of abelian varieties. The aim of this section is to show that the map π_L^i constructed previously is an isomorphism when X is an abelian variety.

Lemma 3.2. *Let X be an abelian variety over a field k of characteristic 0. Then the differential fundamental group $\pi^{\text{diff}}(X/k)$ is commutative.*

Proof. Recall that we always consider the identity of an abelian variety as a base point. We will use an Eckmann-Hilton argument, see [EH62, Theorem 5.4.2]. By the Künneth theorem for

$\pi^{\text{diff}}(X/k)$ ([Del89, Corollary 10.47]), the two projections from $X \times X$ to its factors induces an isomorphism

$$\pi^{\text{diff}}(X \times X/k) \cong \pi^{\text{diff}}(X/k) \times \pi^{\text{diff}}(X/k).$$

Now, if we denote by m the multiplication $X \times X \rightarrow X$ on X , then it induces a morphism

$$\pi^{\text{diff}}(X/k) \times \pi^{\text{diff}}(X/k) \xrightarrow{m_*} \pi^{\text{diff}}(X/k),$$

thus $\pi^{\text{diff}}(X/k)$ is endowed with a structure of a commutative affine group scheme as wanted. \square

Remark 3.3. For a commutative pro-algebraic group G , one can always decompose it as a direct product of its pro-unipotent part and its pro-reductive part:

$$G = G^{\text{uni}} \times G^{\text{red}}.$$

Here, G^{uni} is the pro-unipotent radical of G .

Proposition 3.4. *Let X be an abelian variety of dimension g over a field k of characteristic 0. Let W be an object in $\text{Rep}_f(\pi^{\text{diff}}(X/k)_L)$. Then the map*

$$\pi_L^i(W) : H^i(\pi^{\text{diff}}(X/k)_L, W) \rightarrow H^i(\pi^{\text{diff}}(X_L/L), W)$$

is an isomorphism for all $i \geq 0$.

Proof. Since X is an abelian variety, the groups $\pi^{\text{diff}}(X_L/L)$ and $\pi^{\text{diff}}(X/k)_L$ are commutative by Lemma 3.2. Thus, we have decompositions

$$\pi^{\text{diff}}(X_L/L) = \pi^{\text{uni}}(X_L/L) \times \pi^{\text{red}}(X_L/L) \text{ and } \pi^{\text{diff}}(X/k) = \pi^{\text{uni}}(X/k) \times \pi^{\text{red}}(X/k)$$

into pro-unipotent parts and pro-reductive ones. The faithfully flat map $\pi^{\text{diff}}(X_L/L) \rightarrow \pi^{\text{diff}}(X/k)_L$ in Lemma 3.1 induces the faithfully flat map

$$\pi^{\text{uni}}(X_L/L) \rightarrow \pi^{\text{uni}}(X/k)_L.$$

Since $H^1(\pi^{\text{uni}}(X/k)_L, L) = H^1(\pi^{\text{uni}}(X_L/L), L)$, it follows that

$$\pi^{\text{uni}}(X/k)_L = \pi^{\text{uni}}(X_L/L) \cong (\mathbb{G}_a)^{2g},$$

by means of [LM82, (1.2) & (1.16)].

Using the Lyndon-Hochschild-Serre spectral sequence ([Jan03, Proposition 6.6]), we can compute the cohomology of the differential fundamental group in terms of the cohomology of its pro-unipotent part. We observe that the unipotent parts of $\pi^{\text{diff}}(X/k)_L$ and $\pi^{\text{diff}}(X_L/L)$ contain the same number of copies of \mathbb{G}_a . Therefore, the morphism π_L^i is an isomorphism for every $i \geq 0$. \square

4. ABELIAN VARIETIES ARE DE RHAM $K(\pi, 1)$

In this section, we prove Theorem A. Recall that we want to prove that every abelian variety (equipped with its identity element, see Remark 1.2) is de Rham $K(\pi, 1)$. We want to reduce the general case to the complex case by first finding a model of $X, x, (\mathcal{V}, \nabla)$ over some field k' embedded in both k and \mathbb{C} , then we use the structure of the differential fundamental groups of abelian varieties to compare their group-scheme cohomology. We then can conclude by invoking the complex case Corollary 2.12.

4.1. A construction. We start with a construction that aims to find our desired model over some finite generated field k' over \mathbb{Q} .

Let X be an abelian variety over a field k of characteristic 0 equipped with a point $x \in X(k)$. There is a finite generated field k' over \mathbb{Q} on which all these objects $X, k, (\mathcal{V}, \nabla)$ and the morphisms f, x are defined. In other words, we have an abelian variety X'/k' that fits into the following Cartesian diagram

$$\begin{array}{ccc} X = X' \otimes_{k'} k & \longrightarrow & X' \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & \text{Spec}(k'). \end{array}$$

Next, we fix an embedding $k' \hookrightarrow \mathbb{C}$, so we have the following a cartesian diagram

$$\begin{array}{ccc} X'' = X' \otimes_{k'} \mathbb{C} & \longrightarrow & X' \\ \downarrow & & \downarrow \\ \text{Spec}(\mathbb{C}) & \longrightarrow & \text{Spec}(k'). \end{array}$$

Here, X'' is an abelian variety over \mathbb{C} .

4.2. Proof of Theorem A. We keep the same notations as in Section 3.2 and Section 4.1.

Corollary 4.1. *The maps*

$$\pi^{\text{diff}}(X/k) \rightarrow \pi^{\text{diff}}(X'/k')_k$$

and

$$\pi^{\text{diff}}(X''/\mathbb{C}) \rightarrow \pi^{\text{diff}}(X'/k')_{\mathbb{C}}$$

are faithfully flat. The maps

$$\pi_{\mathbb{C}}^i(-) : H^i(\pi^{\text{diff}}(X'/k')_{\mathbb{C}}, -) \rightarrow H^i(\pi^{\text{diff}}(X''/\mathbb{C}), -)$$

and

$$\pi_k^i(-) : H^i(\pi^{\text{diff}}(X'/k')_k, -) \rightarrow H^i(\pi^{\text{diff}}(X/k), -)$$

are also isomorphisms for all $i \geq 0$.

Proof. It follows directly from Lemma 3.1 and Proposition 3.4. □

Lemma 4.2. *The following map*

$$\delta_{X'/k'}^i(V) : H^i(\pi^{\text{diff}}(X'/k'), V) \longrightarrow H_{\text{dR}}^i(X'/k', (\mathcal{V}, \nabla))$$

is an isomorphism for all $i \geq 0$ and $(\mathcal{V}, \nabla) \in \text{Obj}(\text{MIC}^{\text{coh}}(X'/k'))$.

Proof. Applying the flat base change theorem for de Rham cohomology ([ABC20, 23.5]) and for group-scheme cohomology ([Jan03, Proposition 4.13]), we obtain the following commutative

diagram

$$\begin{array}{ccc}
 H^i(\pi^{\text{diff}}(X'/k')_{\mathbb{C}}, V_{\mathbb{C}}) & \xrightarrow{\cong} & H^i(\pi^{\text{diff}}(X'/k'), V) \otimes \mathbb{C} \\
 \pi_{\mathbb{C}}^i(V_{\mathbb{C}}) \downarrow & & \downarrow \delta_{X'/k'}^i(V) \otimes \mathbb{C} \\
 H^i(\pi^{\text{diff}}(X''/\mathbb{C}), V_{\mathbb{C}}) & & \\
 \delta_{X''/\mathbb{C}}^i(V_{\mathbb{C}}) \downarrow & & \\
 H_{\text{dR}}^i(X''/\mathbb{C}, (\mathcal{V}_{\mathbb{C}}, \nabla_{\mathbb{C}})) & \xrightarrow{\cong} & H_{\text{dR}}^i(X'/k', (\mathcal{V}, \nabla)) \otimes \mathbb{C}.
 \end{array}$$

Here, $V_{\mathbb{C}}$ (resp. $\delta^i(X'/k')_{\mathbb{C}}$) is the base change to \mathbb{C} of V (resp. $\delta^i(X'/k')$). Corollary 2.12 yields that $\delta^i(X''/\mathbb{C})$ is an isomorphism. It follows that $\delta^i(X'/k')_{\mathbb{C}}$ is an isomorphism. Thus $\delta^i(X'/k')$ is also an isomorphism as wanted. \square

Theorem 4.3. *Let X be an abelian variety of dimension g over a field k of characteristic 0. Let (\mathcal{V}, ∇) be an object in $\text{MIC}^{\text{coh}}(X/k)$. Set $V := 0_X^*(\mathcal{V}, \nabla)$. Then*

$$(10) \quad \delta_{X/k}^i(V) : H^i(\pi^{\text{diff}}(X/k), V) \longrightarrow H_{\text{dR}}^i(X/k, (\mathcal{V}, \nabla))$$

is an isomorphism for every $i \geq 0$, that is, every abelian variety is de Rham $K(\pi, 1)$.

Proof. Let (\mathcal{V}, ∇) be an object of $\text{MIC}^{\text{coh}}(X/k)$. By the above construction, there exists an integrable connection (\mathcal{V}', ∇') on X'/k' such that (\mathcal{V}', ∇') is the pull back of (\mathcal{V}, ∇) via the morphism $X' \rightarrow X$. We set $V' := 0_X^*(\mathcal{V}', \nabla')$, then $V = V' \otimes_{k'} k$. Again, applying the flat base change theorem for de Rham cohomology ([ABC20, 23.5]) and for group cohomology ([Jan03, Proposition 4.13]), we obtain the following commutative diagram

$$\begin{array}{ccc}
 H^i(\pi^{\text{diff}}(X/k), V) & \xrightarrow{\delta^i(X/k)} & H_{\text{dR}}^i(X/k, (\mathcal{V}, \nabla)) \\
 \pi_k^i(V') \uparrow & & \uparrow \cong \\
 H^i(\pi^{\text{diff}}(X'/k'), V') \otimes k & \xrightarrow{\delta^i(X'/k')_k} & H_{\text{dR}}^i(X'/k', (\mathcal{V}', \nabla')) \otimes k.
 \end{array}$$

Corollary 4.1 yields that the map π_k^i is an isomorphism. Lemma 4.2 implies that $\delta^i(X'/k')_k$ is an isomorphism. We deduce that the map $\delta^i(X/k)$ is also an isomorphism as desired. \square

4.3. The Euler characteristic of $\pi^{\text{diff}}(X/k)$. Let X be an abelian variety of dimension g . Thanks to Theorem 4.3, the group-scheme cohomology

$$H^i(\pi^{\text{diff}}(X/k), V)$$

is finite-dimensional and vanishes when $i > 2g$. We may define the Euler characteristic of the group-scheme cohomology of the differential fundamental group of an abelian variety by

$$\chi_{\pi^{\text{diff}}}(V) := \sum_{i=0}^{2g} (-1)^i \dim_k H^i(\pi^{\text{diff}}(X), V),$$

where V is an object in $\text{Rep}^f(\pi^{\text{diff}}(X/k))$.

Corollary 4.4. *Let X be an abelian variety of dimension g over a field k of characteristic 0. Let $(\mathcal{V}, \nabla) \in \text{Obj}(\text{MIC}^{\text{coh}}(X/k))$. Denote $V := 0_X^*(\mathcal{V}, \nabla)$. For $i \geq 0$, then*

$$H^i(\pi^{\text{diff}}(X/k), V)$$

is finite-dimensional and vanishes when $i > 2g$. Moreover,

$$\chi_{\pi^{\text{diff}}}(V) = 0.$$

Proof. It suffices to prove that $\chi_{\pi^{\text{diff}}}(V) = 0$. Using flat base change for de Rham cohomology (see [ABC20, 23.5]) we may assume that the base field is \mathbb{C} . According to [Del70, Theorem 6.2, Chapter II], we have

$$H_{\text{dR}}^i(X/\mathbb{C}, (\mathcal{V}, \nabla)) \cong H^i(X^{\text{an}}, \tilde{V}),$$

and, combined with [Dim04, Proposition 2.5.4], this implies

$$\chi_{\text{dR}}(\mathcal{V}, \nabla) = n \chi_{\text{dR}}(\mathcal{O}_X, d).$$

Since $\chi_{\text{dR}}(\mathcal{O}_X, d) = 0$ and applying Theorem 4.3 once again, the result follows. \square

5. COHOMOLOGY OF THE ABELIZATION OF THE DIFFERENTIAL FUNDAMENTAL GROUP

In this section, we study the group scheme cohomology of $\pi^{\text{diff}}(X)^{\text{ab}}$. To do that, we consider the Abenese morphism $f : X \rightarrow \text{Alb}_X$. The morphism f induces, for each $i \geq 0$, a map

$$\eta^i : H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \rightarrow H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V),$$

where V is an object in $(\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X)))$ (see (4)). We aim to prove Theorem C saying that the group-scheme cohomology of the differential fundamental group of the Albanese variety coincides with the group-scheme cohomology of $\pi^{\text{diff}}(X)^{\text{ab}}$.

We start with a remark on the construction of the largest abelian quotient of a pro-algebraic group in [dS07, Page 708]. For a pro-algebraic group $G = \varprojlim_{\alpha} G_{\alpha}$ with $\mathcal{O}(G_{\alpha}) \subseteq \mathcal{O}(G)$ of finite type, we can form G^{ab} as the limit $\varprojlim_{\alpha} G_{\alpha}^{\text{ab}}$ which has the universal property: any homomorphism $G \rightarrow A$ with A is commutative group scheme factor through G^{ab} . The group-scheme G^{ab} is called the *abelianization* of G .

5.1. An observation over \mathbb{C} . Our proof is motivated by the following observation when the base field is \mathbb{C} .

Proposition 5.1. *Let X be a smooth connected projective variety over a field \mathbb{C} and $x \in X(\mathbb{C})$. Let f be the Albanese morphism mapping the base point x to 0_{Alb_X} . Then the induced morphism*

$$\pi^{\text{diff}}(X/\mathbb{C})^{\text{ab}} \xrightarrow{f_*} \pi^{\text{diff}}(\text{Alb}_X/\mathbb{C})$$

is faithfully flat, whose kernel is the finite constant group scheme $(\text{Tor}(H_1(X^{\text{an}}, \mathbb{Z})))^{\text{alg}}$ over \mathbb{C} . As a consequence, we have

$$H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \cong H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

for all $i \geq 0$ and all $V \in \text{Obj}(\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k)))$.

Proof. By the Hurewicz theorem we have

$$\pi_1^{\text{top}}(X^{\text{an}}, x)^{\text{ab}} = H_1(X^{\text{an}}, \mathbb{Z}) \text{ and } \pi_1^{\text{top}}(\text{Alb}_{X^{\text{an}}}, 0_{\text{Alb}_X}) \cong H_1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z})$$

(recall that $\text{Alb}_{X^{\text{an}}}$ is an abelian variety, cf. [BL04, Corollary 11.11.7], whence $\pi_1^{\text{top}}(\text{Alb}_{X^{\text{an}}}, 0_{\text{Alb}_X})$ is commutative). In addition, the map $H^1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z}) \rightarrow H^1(X, \mathbb{Z})$ is an isomorphism. By the universal coefficient theorem, we have

$$H^1(X^{\text{an}}, \mathbb{Z}) \cong \text{Hom}(H_1(X^{\text{an}}, \mathbb{Z}), \mathbb{Z})$$

and

$$H^1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z}) \cong \text{Hom}(H_1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z}), \mathbb{Z}).$$

Since $\text{Alb}_{X^{\text{an}}}$ is a complex torus, $H_1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z})$ has no torsion. We deduce that $H_1(X^{\text{an}}, \mathbb{Z}) \rightarrow H_1(\text{Alb}_{X^{\text{an}}}, \mathbb{Z})$, hence $\pi_1^{\text{top}}(X^{\text{an}}, x) \rightarrow \pi_1^{\text{top}}(\text{Alb}_{X^{\text{an}}}, 0_{\text{Alb}_X})$, is surjective. Moreover, its kernel is the torsion part $\text{Tor}(H_1(X^{\text{an}}, \mathbb{Z}))$, which is a finite commutative (abstract) group.

Next, by Example 2.2, we know that $\pi^{\text{diff}}(X, x)$ is the pro-algebraic completion over \mathbb{C} of $\pi_1^{\text{top}}(X^{\text{an}}, x)$. We note that the analytification of Alb_X is $\text{Alb}_{X^{\text{an}}}$ by Chow's theorem [BL04, Corollary A.4]. Since taking pro-algebraic completion is right exact and commutes with taking abelianization [BLMM02, Remark 1: (a), (c)], the map

$$\pi^{\text{diff}}(X, x)^{\text{ab}} \xrightarrow{f_*} \pi^{\text{diff}}(\text{Alb}_X, 0_{\text{Alb}_X})$$

is also faithfully flat as desired. Next, since $\text{Tor}(H_1(X, \mathbb{Z}))$ is finite and commutative, it is observable in $H_1(X^{\text{an}}, \mathbb{Z})$. It follows from [BLMM02, Remark 1: (b)] that $(\text{Tor}(H_1(X, \mathbb{Z})))^{\text{alg}}$ is the kernel of

$$\pi^{\text{diff}}(X, x)^{\text{ab}} \xrightarrow{f_*} \pi^{\text{diff}}(\text{Alb}_X, 0_{\text{Alb}_X}),$$

which is a finite commutative group scheme over \mathbb{C} .

Finally, since $\pi^{\text{diff}}(\text{Alb}_X)$ and $\pi^{\text{diff}}(X)^{\text{ab}}$ are commutative, we have decompositions into pro-unipotent parts and pro-reductive parts

$$\pi^{\text{diff}}(\text{Alb}_X) = \pi^{\text{diff,uni}}(\text{Alb}_X) \times \pi^{\text{diff,red}}(\text{Alb}_X) \text{ and } \pi^{\text{diff}}(X)^{\text{ab}} = \pi^{\text{diff,uni}}(X)^{\text{ab}} \times \pi^{\text{diff,red}}(X)^{\text{ab}}.$$

We deduce that $\pi^{\text{diff,uni}}(\text{Alb}_X) \cong \pi^{\text{diff,uni}}(X)^{\text{ab}}$. As in the proof of Proposition 3.4, we use the Lyndon-Hochschild-Serre spectral sequence ([Jan03, Proposition 6.6]) to compute the cohomologies $H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V)$ and $H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$ in terms of their pro-unipotent parts. It follows that η^i is an isomorphism for all $i \geq 0$. \square

Remark 5.2. The faithfully flat morphism $\pi^{\text{diff}}(X/\mathbb{C})^{\text{ab}} \xrightarrow{f_*} \pi^{\text{diff}}(\text{Alb}_X/\mathbb{C})$ is inspired by the work of [Lan12], [BdS17], [D'A23].

5.2. A proof of Theorem C. For the general case of k , we provide an algebraic proof of the isomorphism $H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \cong H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$ in this section.

Lemma 5.3. *Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and a base point $x \in X(k)$. Let f be the Albanese morphism mapping x to 0_{Alb_X} , the identity. Then we have an isomorphism*

$$H^1_{\text{dR}}(\text{Alb}_X, \mathcal{O}_{\text{Alb}_X}) \cong H^1_{\text{dR}}(X/k, \mathcal{O}_X).$$

Proof. Using flat base change for de Rham cohomology (see [ABC20, 23.5]), we may assume that the base field is \mathbb{C} . In this case, we have the Hodge decompositions

$$H_{dR}^1(X, \mathcal{O}_X) \cong H^1(X, \mathcal{O}_X) \oplus H^0(X, \Omega_X^1)$$

and

$$H_{dR}^1(\text{Alb}_X, \mathcal{O}_{\text{Alb}_X}) \cong H^1(X, \mathcal{O}_{\text{Alb}_X}) \oplus H^0(X, \Omega_{\text{Alb}_X}^1).$$

By Igusa's theorem [Isu55], we have an isomorphism $H^0(X, \Omega_{\text{Alb}_X}^1) \cong H^0(X, \Omega_X^1)$. By the Hodge symmetry, we also have an isomorphism $H^1(X, \mathcal{O}_{\text{Alb}_X}) \cong H^1(X, \mathcal{O}_X)$. The desired result follows. \square

Theorem 5.4. *Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and a base point $x \in X(k)$. Let f be the Albanese morphism mapping x to 0_{Alb_X} . Let V be an object of $\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k))$. Then the induced map*

$$\eta^i(V) : H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \longrightarrow H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

is an isomorphism for every $i \geq 0$.

Proof. Recall from remark 2.9 that we have isomorphisms

$$H^1(\pi^{\text{diff}}(\text{Alb}_X/k), k) \cong H_{dR}^1(\text{Alb}_X, \mathcal{O}_{\text{Alb}_X})$$

and

$$H^1(\pi^{\text{diff}}(X/k), k) \cong H_{dR}^1(X/k, \mathcal{O}_X).$$

Combining with Lemma 5.3, we have

$$(11) \quad H^1(\pi^{\text{diff}}(\text{Alb}_X/k), k) \cong H^1(\pi^{\text{diff}}(X/k), k).$$

From the fully faithful functor $\text{Rep}^f(\pi^{\text{diff}}(X/k)^{\text{ab}}) \rightarrow \text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k))$ we deduce that

$$\eta^1(V) : H^1(\pi^{\text{diff}}(\text{Alb}_X/k), V) \rightarrow H^1(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

is injective for all $V \in \text{Obj}(\text{Rep}^f(\text{Alb}_X/k))$. In particular, we have an injection

$$(12) \quad \eta^1(k) : H^1(\pi^{\text{diff}}(\text{Alb}_X/k), k) \hookrightarrow H^1(\pi^{\text{diff}}(X/k)^{\text{ab}}, k).$$

From the faithfully flat morphism $\pi^{\text{diff}}(X/k) \rightarrow \pi^{\text{diff}}(X/k)^{\text{ab}}$, we apply the five-term exact sequence to the trivial $\pi^{\text{diff}}(X/k)^{\text{ab}}$ -representation k to obtain an injection

$$(13) \quad H^1(\pi^{\text{diff}}(X/k)^{\text{ab}}, k) \hookrightarrow H^1(\pi^{\text{diff}}(X/k), k).$$

We combine (11), (12) and (13) to deduce that

$$H^1(\pi^{\text{diff}}(\text{Alb}_X/k), k) \cong H^1(\pi^{\text{diff}}(X/k)^{\text{ab}}, k).$$

Once again, by [LM82, (1.2) & (1.16)], we obtain that the unipotent radicals $\pi^{\text{diff}}(X/k)^{\text{ab}, \text{uni}}$ and $\pi^{\text{diff}, \text{uni}}(\text{Alb}_X/k)$ are isomorphic. Applying the Lyndon-Hochschild-Serre spectral sequence again yields that $H^i(\pi^{\text{diff}}(\text{Alb}_X/k), V) \cong H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$ for all $i \geq 0$ and $V \in \text{Obj}(\text{Rep}^f(\pi^{\text{diff}}(\text{Alb}_X/k)))$ as wanted. \square

Corollary 5.5. *Let X be a smooth proper geometrically connected scheme over a field k of characteristic 0 and a base point $x \in X(k)$. Then the group-scheme cohomology*

$$H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V)$$

is finite-dimensional for any $i \geq 0$ and any $V \in \text{Obj}(\pi^{\text{diff}}(X/k)^{\text{ab}})$. Moreover, the Euler characteristic of the group-scheme cohomology of $\pi^{\text{diff}}(X/k)^{\text{ab}}$ is well-defined and equal to zero.

Proof. We first prove the finiteness of group scheme cohomology of $\pi^{\text{diff}}(X/k)^{\text{ab}}$. According to Theorem 5.4, we have

$$H^1(\pi^{\text{diff}}(\text{Alb}_X/k), k) \cong H^1(\pi^{\text{diff}}(X/k)^{\text{ab}}, k).$$

This implies that the pro-unipotent part of $\pi^{\text{diff}}(X/k)^{\text{ab}}$ is $(\mathbb{G}_a)^{2g}$ where g is dimension of Albanese variety. Thanks to the Lyndon-Hochschild-Serre spectral sequence, we obtain

$$H^i(\pi^{\text{diff}}(X/k)^{\text{ab}}, V) \cong H^i((\mathbb{G}_a)^{2g}, V).$$

Using Corollary B, the desired result follows since there exists an abelian variety of dimension g for every $g \geq 0$. \square

5.3. Cohomology of the abelianization of the topological fundamental group. In this section, we prove Corollary E.

Corollary 5.6. *Let X be a smooth connected projective complex variety, and x be a base point. Let f be the Albanese morphism mapping x to 0_{Alb_X} . Let V be an object in $\text{Rep}^f(\pi^{\text{top}}(\text{Alb}, 0_{\text{Alb}_X}))$. Then the induced map*

$$\eta^i(V) : H^i(\pi^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X}), V) \longrightarrow H^i(\pi^{\text{top}}(X, x)^{\text{ab}}, V)$$

is an isomorphism for every $i \geq 0$.

Proof. In view of Theorem A, Theorem C and Proposition 2.10, it suffices to show that the groups $\pi^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X})$ and $\pi^{\text{top}}(X, x)^{\text{ab}}$ are algebraically good. Since X is a finite CW complex, $\pi^{\text{top}}(X, x)$ is finitely presented. Therefore, $\pi^{\text{top}}(X, x)^{\text{ab}}$ is abelian of finite type, hence algebraically good by Example 2.5. Since Alb_X is an abelian variety, $\pi^{\text{top}}(\text{Alb}_X, 0_{\text{Alb}_X})$ is also algebraically good. The corollary is proved. \square

Acknowledgments

We thank Francesco Baldassarri, Michel Brion, Hélène Esnault, Phung Ho Hai, Dao Van Thinh, and Tran Phan Quoc Bao for insightful discussions.

REFERENCES

- [ABC20] Y. André, F. Baldassarri, and M. Cailotto. *De Rham Cohomology of Differential Modules on Algebraic Varieties*, volume 189 of *Progress in Mathematics*. Birkhäuser, 2 edition, 2020.
- [Ach15] P. Achinger. *$K(\pi, 1)$ Spaces in Algebraic Geometry*. PhD thesis, University of California, Berkeley, 2015. Ph.D. dissertation.
- [Ach17] P. Achinger. Wild ramification and $K(\pi, 1)$ spaces. *Inventiones Mathematicae*, 210(2):453–503, 2017.
- [Ach23] Piotr Achinger. Is anything known about de rham $K(\pi, 1)$ s?, 2023. Comments on MathOverflow, version: 2023-06-04.

- [BdS17] I. Biswas and J.P. dos Santos. Abelianization of the F -divided fundamental group scheme. *Proc. Indian Acad. Sci. Math. Sci.*, 127(2), 2017.
- [BHT25] V.Q. Bao, P. H. Hai, and D. V. Thinh. Cohomology of the differential fundamental group of algebraic curves. *Bulletin des Sciences mathématiques*, 203, 2025.
- [BL04] C. Birkenhake and H. Lange. *Complex Abelian Varieties*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2004.
- [BLMM02] H. Bass, A. Lubotzky, A. R. Magid, and S. Mozes. The proalgebraic completion of rigid groups. In *Proceedings of the Conference on Geometric and Combinatorial Group Theory*, 95:19–58, 2002.
- [Bro94] K. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
- [CE56] H. Cartan and S. Eilenberg. *Homological Algebra*. Princeton University Press, 1956.
- [D'A23] Marco D'Addezio. Slopes of f -isocrystals over abelian varieties. *Documenta Mathematica*, 28(1):1–9, 2023.
- [Del70] P. Deligne. *Équations différentielles à points singuliers réguliers*, volume 163 of *Lecture Notes in Mathematics*. Springer, 1970.
- [Del89] P. Deligne. *Le Groupe Fondamental de la Droit Projective Moins Trois Points: in Galois Group over \mathbb{Q}* . Springer-Verlag, 1989.
- [Den23] C. Deninger. A pro-algebraic fundamental group for topological spaces. *Proceedings of the Steklov Institute of Mathematics*, 320(1):62–90, 2023.
- [DG70] M. Demazure and A. Grothendieck. *Schémas en groupes I: Propriétés générales des schémas en groupes*, volume 151 of *Lecture notes in Math.* Springer-Verlag, 1970.
- [Dim04] A. Dimca. *Sheaves in Topology*. Springer, 2004.
- [DM82] P. Deligne and J. Milne. *Tannakian Categories*, volume 900 of *Lecture Notes in Mathematics*. Springer-Verlag, 1982.
- [dS07] J.P. dos Santos. Fundamental group schemes for stratified sheaves. *Journal of Algebra*, 317, 2007.
- [EH62] B. Eckmann and P. J. Hilton. Group-like structures in general categories. i. multiplications and comultiplications. *Mathematische Annalen*, 145, 1962.
- [EH06] H. Esnault and P.H. Hai. The Gauss-Manin connection and Tannaka duality. *Int. Math. Res. Not.*, pages 1–35, 2006.
- [Gro73] Alexander Grothendieck. *Théorie des topos et cohomologie étale des schémas. Tome 3*, volume 305 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin–New York, 1973. Séminaire de Géométrie Algébrique du Bois Marie 1963–1964 (SGA 4).
- [Har77] Robin Hartshorne. *Algebraic Geometry*, volume 52 of *Graduate Texts in Mathematics*. Springer, 1977.
- [Isu55] Jun-ichi Isuga. A fundamental inequality in the theory of picard varieties^{*}. *Proceedings of the National Academy of Sciences*, 41(5):317–320, 1955.
- [Jan87] J.C. Jantzen. *Representations of Algebraic Groups*, volume 131 of *Pure Appl. Math.* Academic Press, Inc., Boston, MA, 1987.
- [Jan03] J.C. Jantzen. *Representations of Algebraic Groups*, volume 107 of *Mathematical Surveys and Monographs*. American Mathematical Society, second edition, 2003.
- [Kat70] N. Katz. Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin. *Inst. Hautes Études Sci. Publ. Math.*, 39:175–232, 1970.
- [KPT09] L. Katzarkov, T. Pantev, and B. Toën. Algebraic and topological aspects of the schematization functor. *Composito Mathematica*, 145(3), 2009.
- [Lan12] A. Langer. On the S -fundamental group scheme. II. *J. Inst. Math. Jussieu*, 11(4), 2012.
- [LM82] A. Lubotzky and A. Magid. Cohomology of unipotent and prounipotent groups. *Journal of Algebra*, 74:76–95, 1982.
- [Sch13] Peter Scholze. p -adic Hodge theory for rigid-analytic varieties. *Forum of Mathematics, Pi*, 1, 2013.
- [Wat79] W. C. Waterhouse. *Introduction to affine group schemes*, volume 66 of *Graduate Texts in Mathematics*. Springer-Verlag, New York - Berlin, 1979.
- [Zha13] Lei Zhang. The homotopy sequence of the algebraic fundamental group. *International Mathematics Research Notices*, 2014(22):6155–6174, 07 2013.

INSTITUTE OF MATHEMATICS, VIETNAM ACADEMY OF SCIENCE AND TECHNOLOGY

Email address, QK Nguyen: nguyen@math.univ-lyon1.fr

CAMILLE JORDAN INSTITUTE, CLAUDE BERNARD UNIVERSITY LYON 1