

ENERGY DECAY OF A VISCOELASTIC WAVE EQUATION WITH VARIABLE EXPONENT LOGARITHMIC NONLINEARITY AND WEAK DAMPING

QINGQING PENG¹ AND YIKAN LIU^{2,*}

ABSTRACT. In this paper, we investigate the energy decay of the solution to a viscoelastic wave equation with variable exponents logarithmic nonlinearity and weak damping in a bounded domain. We establish an explicit general decay result under mild conditions on the relaxation function g . Furthermore, under the general assumption $g'(t) \leq -\zeta(t)G(g(t))$ with some suitably given ζ and G , we derive a refined decay estimate improving existing results. In particular, uniform exponential and polynomial decay rates are obtained under a further special situation $g'(t) \leq -\xi(t)g^q(t)$ with $1 \leq q < 2$, extending earlier studies that were restricted to the case $1 \leq q < \frac{3}{2}$.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ ($n \geq 3$) be a bounded domain with a smooth boundary $\partial\Omega$. This article is concerned with the initial-boundary value problem for a nonlinear viscoelastic wave equation

$$\begin{cases} u_{tt} - \Delta u + \int_0^t g(t-s) \Delta u(s) \, ds + u_t = \alpha |u|^{p(x)-2} u \log |u|, & (x, t) \in \Omega \times (0, \infty), \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x), & x \in \Omega, \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, \infty), \end{cases} \quad (1.1)$$

where $\alpha > 0$ is a constant. Here we assume that the exponent p in the nonlinear term satisfies

$$p_1 := \operatorname{ess\,inf}_{x \in \Omega} p(x) > 2, \quad p_2 := \operatorname{ess\,sup}_{x \in \Omega} p(x) < \frac{2(n-1)}{n-2}. \quad (1.2)$$

Moreover, p is further assumed to be log-Hölder continuous, i.e., there exist a constant $A > 0$ such that for a.e. $x, y \in \Omega$ satisfying $|x - y| < 1$, there holds

$$|p(x) - p(y)| \leq -\frac{A}{\log |x - y|}. \quad (1.3)$$

Wave equations with logarithmic nonlinearity have attracted significant attention in recent years, and numerous studies have explored the dynamics of problem (1.1) in the absence of the memory effect (i.e., when $g = 0$). The logarithmic wave equation was first introduced by Bialynicki-Birula and Mycielski in [5, 6], who demonstrated the existence of stable and localized solutions in one

Key words and phrases. Viscoelastic wave equation, variable exponent logarithmic nonlinear, weakly damping, exponential and polynomial decay.

¹School of Mathematics and Statistics & Hubei Key Laboratory of Engineering Modeling and Scientific Computing, Huazhong University of Science and Technology, Wuhan 430074, China.

² Department of Mathematics, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan.

*Corresponding author. E-mail: liu.yikan.8z@kyoto-u.ac.jp.

spatial dimension. Subsequently, Cazenave and Haraux [7] established the well-posedness of the Cauchy problem in three spatial dimensions. Bartkowiak and Górkak [3] studied classical and weak solutions to the one-dimensional Cauchy problem, while Górkak [11] proved the global existence of weak solutions for initial data $(u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega)$ using compactness arguments. More recently, Di [8] applied the potential well method to establish global existence for the wave equation with logarithmic nonlinearity and derived exponential or polynomial decay by constructing an appropriate Lyapunov functional. Moreover, blow-up in the unstable set was also established. For further results on stability and blow-up of systems with logarithmic nonlinearities, we refer the reader to [12, 18, 24–26] and the references therein.

On the other hand, under certain assumptions on the kernel function g in the memory term, several decay and blow-up results have been established in the literature. In [14], the authors studied the stability of the initial-boundary value problem for a quasilinear viscoelastic equation. They obtained polynomial decay under mild conditions on g , and further investigated both polynomial and exponential decay under a more general condition on g . In [4], energy decay results for the viscoelastic problem were derived by constructing an appropriate Lyapunov functional under suitable assumptions on g . Additionally, the authors in [16] considered blow-up phenomena for the viscoelastic wave problem, while those in [22] analyzed the blow-up of solutions for a viscoelastic problem with variable exponents. Later, Liao [17] discussed energy decay rates for solutions to a viscoelastic wave equation with variable exponents and weak damping. Further related works can be found e.g. in [2, 13, 21, 23, 27].

Inspired by the aforementioned studies, in this article we investigate the stability and blow-up behavior of problem (1.1). This manuscript contains three main contributions. First, in proving the energy decay, we neither require the initial energy to be smaller than the depth of the potential well d , nor restrict the analysis to a stable set as required in previous works involving logarithmic nonlinearity, which relaxes the key structural assumptions. Second, the conditions imposed on the kernel g are more general than those considered in earlier studies on systems with variable exponents. Third, in Theorem 3.5, our result holds for $1 \leq q < 2$, whereas existing literature only covered the case of $1 \leq q < \frac{3}{2}$. Indeed, our findings extend and generalize previous results, particularly as those in [2, 14, 21, 25].

The organization of this paper is as follows. In Section 2, we establish the global existence of solutions and present several auxiliary lemmas. Then Section 3 and 4 are devoted to the stability analysis of problem (1.1), achieved through the construction of an appropriate Lyapunov functional and the application of multiplier methods.

2. PRELIMINARIES AND SOME LEMMAS

In this section, we introduce some notations, basic definitions, essential lemmas, and function spaces that will be used in stating and proving our main results.

Throughout, by $\|\cdot\|_k$ we denote the norm of the Lebesgue space $L^k(\Omega)$ for $1 \leq k \leq \infty$, and by (\cdot, \cdot) the inner product of $L^2(\Omega)$. In order to study problem (1.1), we start with recalling the Orlicz-Sobolev-type Banach spaces $L^{p(x)}(\Omega)$ defined as (see [9, 10])

$$L^{p(x)}(\Omega) := \left\{ f : \text{a measurable real-valued function in } \Omega \mid \int_{\Omega} |f|^p dx < \infty \right\},$$

where $p \in L^{\infty}(\Omega)$ satisfies (1.2). The norm of $L^{p(x)}(\Omega)$ is given by

$$\|f\|_{p(x)} := \inf \left\{ \lambda > 0 \mid \int_{\Omega} \left| \frac{f}{\lambda} \right|^p dx \leq 1 \right\},$$

and it is readily seen that

$$\min \left\{ \|f\|_{p(x)}^{p_1}, \|f\|_{p(x)}^{p_2} \right\} \leq \int_{\Omega} |f|^p \, dx \leq \max \left\{ \|f\|_{p(x)}^{p_1}, \|f\|_{p(x)}^{p_2} \right\}. \quad (2.1)$$

We collect several basic facts about the function spaces in the following lemmas.

Lemma 2.1 (see [1]). *Let k be a constant satisfying $2 \leq k \leq 2_* := \frac{2n}{n-2}$ with $n \geq 3$. Then there exists an optimal constant depending only on k such that*

$$\|v\|_k^k \leq B_k \|\nabla v\|_2^k, \quad \forall v \in H_0^1(\Omega).$$

Lemma 2.2 (see [9, 10]). *Let $p, q \in C(\bar{\Omega})$ satisfy $1 < p \leq q$ on $\bar{\Omega}$. Then the embedding $L^{q(x)}(\Omega) \hookrightarrow L^{p(x)}(\Omega)$ is continuous and its operator norm does not exceed $|\Omega| + 1$.*

Lemma 2.3 (see [15]). *Let p satisfy (1.2)–(1.3). Then the embedding $H_0^1(\Omega) \hookrightarrow L^{p(x)}(\Omega)$ is continuous and compact.*

Next, we fix the basic assumption on the kernel function g in the memory term.

(A1) The function $g \in C^1([0, \infty); (0, \infty))$ is non-increasing and satisfies

$$g(0) > 0, \quad \ell := 1 - \int_0^{\infty} g(s) \, ds > 0.$$

In the sequel, we denote

$$I(t) := \int_t^{\infty} g(s) \, ds, \quad K_{\delta}(s) := -\frac{g'(s)}{g(s)} + \delta, \quad M(\delta) := \int_0^{\infty} \frac{g(s)}{K_{\delta}(s)} \, ds, \quad (2.2)$$

where $\delta \in (0, 1)$ is a constant. To state our results, we define the energy functional associated with problem (1.1) as

$$\begin{aligned} E(t) := & \frac{1}{2} \|u_t(t)\|_2^2 + \frac{1}{2} \left(1 - \int_0^t g(s) \, ds \right) \|\nabla u(t)\|_2^2 + \frac{1}{2} (g \circ \nabla u)(t) \\ & - \alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} \, dx + \alpha \int_{\Omega} \frac{|u(t)|^p}{p^2} \, dx, \end{aligned} \quad (2.3)$$

where

$$(g \circ \nabla u)(t) := \int_0^t g(t-s) \|\nabla u(t) - \nabla u(s)\|_2^2 \, ds.$$

By directly differentiating (2.3) and using (1.1), it is straightforward to verify that

$$\begin{aligned} E'(t) = & \frac{1}{2} (g' \circ \nabla u)(t) - \frac{1}{2} g(t) \|\nabla u(t)\|_2^2 - \|u_t(t)\|_2^2 \\ \leq & \frac{1}{2} (g' \circ \nabla u)(t) - \|u_t(t)\|_2^2 \leq 0. \end{aligned} \quad (2.4)$$

Lemma 2.4. *Let assumption (A1) hold, p satisfy (1.2)–(1.3) and $\mu > 0$ be a constant satisfying $p_2 + \mu < 2_*$. Let $B_{p_2+\mu}$ be the optimal constant in Lemma 2.1 and define*

$$B := B_{p_2+\mu} \ell^{-\frac{p_2+\mu}{2}}, \quad R(\lambda) := \frac{1}{2} \lambda^2 - \frac{\alpha B}{\epsilon \mu p_1} \lambda^{p_2+\mu}, \quad \lambda(t) := \{\ell \|\nabla u(t)\|_2^2 + (g \circ \nabla u)(t)\}^{\frac{1}{2}}.$$

Then there holds $E(t) \geq R(\lambda(t))$.

Proof. Splitting the domain Ω into

$$\Omega_1 := \{v \in H_0^1(\Omega) \mid |v| < 1\}, \quad \Omega_2 := \{v \in H_0^1(\Omega) \mid |v| \geq 1\}, \quad (2.5)$$

we apply the Sobolev embedding theorem and Lemma 2.1 with $k = p_2 + \mu$ to deduce

$$\begin{aligned} \int_{\Omega} \frac{|v|^p \log |v|}{p} dx &= \left(\int_{\Omega_1} + \int_{\Omega_2} \right) \frac{|v|^p \log |v|}{p} dx \leq \frac{1}{\epsilon \mu p_1} \int_{\Omega_2} |v|^{p_2} |v|^\mu dx \\ &\leq \frac{1}{\epsilon \mu p_1} \int_{\Omega} |v|^{p_2 + \mu} dx \leq \frac{B_{p_2 + \mu}}{\epsilon \mu p_1} \|\nabla v\|_2^{p_2 + \mu}. \end{aligned} \quad (2.6)$$

Here we used the inequality $x^{-\mu} \log x < (\epsilon \mu)^{-1}$ for $x \geq 1$, where $\mu > 0$ is the constant stated in the theorem. Then according to the definition (2.3) of $E(t)$, we substitute $v = u(t)$ in (2.6) to estimate

$$\begin{aligned} E(t) &\geq \frac{\ell}{2} \|\nabla u(t)\|_2^2 + \frac{1}{2} (g \circ \nabla u)(t) - \frac{\alpha B_{p_2 + \mu}}{\epsilon \mu p_1} \|\nabla u(t)\|_2^{p_2 + \mu} + \frac{\alpha}{p_2^2} \int_{\Omega} |u(t)|^p dx \\ &\geq \frac{1}{2} \{\ell \|\nabla u(t)\|_2^2 + (g \circ \nabla u)(t)\} - \frac{\alpha B}{\epsilon \mu p_1} \{\ell \|\nabla u(t)\|_2^2 + (g \circ \nabla u)(t)\}^{\frac{p_2 + \mu}{2}} \\ &= \frac{1}{2} \lambda(t)^2 - \frac{\alpha B}{\epsilon \mu p_1} \lambda(t)^{p_2 + \mu} = R(\lambda(t)), \end{aligned}$$

by the definitions of B , $\lambda(t)$ and $R(\lambda)$. \square

It is easily verified that $R(\lambda)$ attains its maximum at

$$\lambda_1 := \left(\frac{\epsilon \mu p_1}{\alpha (p_2 + \mu) B} \right)^{\frac{1}{p_2 + \mu - 2}} > 0$$

with the corresponding maximum

$$E_1 := R(\lambda_1) = \left(\frac{1}{2} - \frac{1}{p_2 + \mu} \right) \lambda_1^2 > 0.$$

Lemma 2.5 (see [17, Lemma 3.4]). *Let assumption (A1) hold and p satisfies (1.2)–(1.3). Let u be a solution to problem (1.1) whose initial data satisfies*

$$E(0) < E_1, \quad \lambda(0) < \lambda_1.$$

Then there exists a constant $\lambda_2 \in (0, \lambda_1)$ such that

$$\lambda(t) = \{\ell \|\nabla u(t)\|_2^2 + (g \circ \nabla u)(t)\}^{\frac{1}{2}} \leq \lambda_2, \quad \forall t \in [0, T_{\max}),$$

where T_{\max} is the maximal existence time.

Next, we invoke the existence result for the solution to (1.1).

Proposition 2.6 (Local existence). *Let $(u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega)$ be given. Let assumption (A1) hold and p satisfy (1.2)–(1.3). Then there exists $T > 0$ such that problem (1.1) admits a unique local weak solution u on $[0, T]$.*

Combining the Faedo-Galerkin method with the proof for logarithmic nonlinearity in [8] and that for nonlinear damping and a memory term in [22], we can show the local existence stated in Proposition 2.6.

We introduce an auxiliary energy

$$\mathbb{E}(t) := E(t) + \alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} dx. \quad (2.7)$$

We prepare additional estimates for $E(t)$ and $\mathbb{E}(t)$.

Lemma 2.7. *Under the same assumptions of Lemma 2.5, there hold*

$$\alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} dx \leq \tilde{C} E(t) \leq \tilde{C} E(0), \quad (2.8)$$

$$\mathbb{E}(t) \leq (1 + \tilde{C}) E(t) \leq (1 + \tilde{C}) E(0) \quad (2.9)$$

for any $t \in [0, T_{\max})$, where

$$\tilde{C} := \frac{\frac{2\alpha\lambda_2^{p_2+\mu-2}B}{\epsilon\mu p_1}}{1 - \frac{2\alpha\lambda_2^{p_2+\mu-2}B}{\epsilon\mu p_1}}.$$

Proof. Since the second halves of (2.8) and (2.9) follow immediately from the monotonicity of $E(t)$, it suffices to verify the respective first halves.

We employ (2.6) and Lemma 2.5 to deduce

$$\begin{aligned} \alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} dx &\leq \frac{\alpha B_{p_2+\mu} \|\nabla u(t)\|_2^{p_2+\mu}}{\epsilon\mu p_1} \leq \frac{\alpha B}{\epsilon\mu p_1} \left(\ell^{\frac{1}{2}} \|\nabla u(t)\|_2 \right)^{p_2+\mu-2} \ell \|\nabla u(t)\|_2^2 \\ &\leq \frac{2\alpha\lambda_2^{p_2+\mu-2}B}{\epsilon\mu p_1} \left(E(t) + \alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} dx \right), \end{aligned}$$

which implies the first half of (2.8) by a simple rearrangement. Then the first half of (2.9) is a direct consequence of (2.8) and (2.7). \square

Remark 2.8. Clearly, (2.9) indicates that $\mathbb{E}(t)$ is uniformly bounded for all $t \in [0, T_{\max})$, implying the global existence of the solution, i.e., $T_{\max} = \infty$. At the same time, we also have $0 \leq E(t) \leq E(0)$ for all $t \in [0, \infty)$.

3. ENERGY DECAY RESULTS

This section is devoted to the statements of our main results regarding the energy decay rates of global solutions to problem (1.1). The first result is as follows.

Theorem 3.1. *Under the same assumptions of Lemma 2.5, further let α be sufficiently small. then there exists a constant $C > 0$ such that the energy $E(t)$ of the solution to (1.1) satisfies the following polynomial decay estimates*

$$\int_0^{\infty} E(t) dt \leq CE(0), \quad E(t) \leq CE(0)(1+t)^{-1}, \quad t \geq 0. \quad (3.1)$$

Next, suppose that $g(t)$ satisfies the following additional assumption.

(A2) Let $G \in C^1([0, \infty); [0, \infty))$ be either linear or strictly increasing, strictly convex and of C^2 -class on $[0, r]$ ($r \leq g(0)$) satisfying $G(0) = G'(0) = 0$, and $\zeta \in C^1([0, \infty); (0, \infty))$ be non-increasing. The function $g(t)$ satisfies the ordinary differential inequality

$$g'(t) \leq -\zeta(t)G(g(t)), \quad \forall t > 0.$$

Remark 3.2 (see [20]). If G is a strictly increasing and strictly convex C^2 function on $[0, r]$ satisfying $G(0) = G'(0) = 0$, then it admits an extension \bar{G} sharing the same monotonicity, convexity and

regularity on $[0, \infty)$. For instance, if $G(r) = a$, $G'(r) = b$ and $G''(r) = c$, then we can construct \bar{G} for $t > r$ as

$$\bar{G}(t) = \frac{C}{2}t^2 + (b - Cr)t + \left(a + \frac{C}{2}r^2 - br\right).$$

Meanwhile, in this case the convexity of G and $G(0) = 0$ yield

$$G(\theta t) \leq \theta G(t), \quad \forall \theta \in [0, 1], \quad \forall t \in (0, r].$$

With assumption (A2), we have the following result.

Theorem 3.3. *Under the same assumptions of Theorem 3.1, further let assumption (A2) hold. Then there exist constants $k_1, k_2, t_1 > 0$ such that the energy of problem (1.1) satisfies*

$$E(t) \leq k_2 G_1^{-1} \left(k_1 \int_{t_1}^t \zeta(s) \, ds \right) \quad (3.2)$$

for all $t \geq t_1$, where $G_1(t) := \int_t^r \frac{1}{s G'(s)} \, ds$ is strictly decreasing and convex on $[0, r]$ and satisfies $\lim_{t \rightarrow 0} G_1(t) = \infty$.

For better understanding, we provide an example to illustrate the result above.

Example 3.4. Let $g(t) = a \exp(-t^p)$, where $0 < p < 1$ and $a > 0$ is sufficiently small so that g satisfies assumption (A1). Then assumption (A2) is fulfilled with

$$\zeta(t) = 1, \quad G(t) = \frac{p^t}{(\ln \frac{a}{t})^{\frac{1}{p}-1}}.$$

Then direct calculation yields

$$G'(t) = \frac{(1-p) + p \ln \frac{a}{t}}{(\ln \frac{a}{t})^{\frac{1}{p}}}, \quad G''(t) = \frac{(1-p)(\ln \frac{a}{t} + \frac{1}{p})}{(\ln \frac{a}{t})^{\frac{1}{p+1}}}.$$

Thus, G satisfies assumption (A2) on $[0, r]$ for any $0 < r < a$. Further, we calculate

$$\begin{aligned} G_1(t) &= \int_t^r \frac{1}{s G'(s)} \, ds = \int_t^r \frac{(\ln \frac{a}{s})^{\frac{1}{p}}}{s(1-p + p \ln \frac{a}{s})} \, ds = \int_{\ln \frac{a}{r}}^{\ln \frac{a}{t}} \frac{\sigma^{\frac{1}{p}}}{1-p + p\sigma} \, d\sigma \\ &= \frac{1}{p} \int_{\ln \frac{a}{r}}^{\ln \frac{a}{t}} \sigma^{\frac{1}{p}-1} \left(\frac{\sigma}{\frac{1-p}{p} + \sigma} \right) \, d\sigma \leq \frac{1}{p} \int_{\ln \frac{a}{r}}^{\ln \frac{a}{t}} \sigma^{\frac{1}{p}-1} \, d\sigma \leq \left(\ln \frac{a}{t} \right)^{\frac{1}{p}}. \end{aligned}$$

Consequently, (3.2) implies $E(t) \leq k \exp(-kt^p)$.

Finally, we add an alternative assumption on $g(t)$ as follows.

(A3) Let $\xi \in C^1([0, \infty); (0, \infty))$ be non-increasing such that $\int_0^\infty \xi(s) \, ds = \infty$, and $1 \leq q < 2$. The function $g(t)$ satisfies the ordinary differential inequality

$$g'(t) \leq -\xi(t)g(t)^q, \quad \forall t > 0.$$

Replacing assumption (A2) with (A3), we obtain the following result.

Theorem 3.5. *Under the same assumptions of Theorem 3.1, further let assumption (A3) hold. Then there exist constants $K > 0$ and $K' > 0$, such that the energy of problem (1.1) satisfies*

$$E(t) \leq \begin{cases} E(0) \exp \left(1 - K \int_0^t \xi(s) \, ds \right), & q = 1, \\ E(0) \left(\frac{q}{1 + K'(q-1) \int_0^t \xi(s) \, ds} \right)^{\frac{1}{q-1}}, & 1 < q < 2. \end{cases} \quad (3.3)$$

Theorem 3.5 generalizes the results in [2, 21, 25], where $g'(t)$ was assumed to satisfy condition (A3) with $\leq p < \frac{3}{2}$, while our result holds for $1 \leq q < 2$. Moreover, the decay rates

$$E(t) \leq \begin{cases} K \exp \left(1 - \lambda \int_{t_0}^t \xi(s) \, ds \right), & p = 1, \\ K \left(\frac{1}{1 + \int_{t_0}^t \xi^{2p-1}(s) \, ds} \right)^{\frac{1}{2p-2}}, & 1 < p < \frac{3}{2} \end{cases}$$

obtained in [21] are improved under this broader framework.

4. PROOF OF ENERGY DECAY RESULTS

Throughout this section, by $C > 0$ we denote generic constants with may change line to line.

4.1. Proof of Theorem 3.1. To proceed, we shall first introduce several auxiliary functions and prepare several lemmas accordingly.

Recall the functions $I(t)$ and $M(\delta)$ defined in (2.2) and introduce

$$I_1(t) := \int_{\Omega} \int_0^t I(t-s) |\nabla u(s)|^2 \, ds \, dx, \quad I_2(t) := M(\delta)(\delta I_1(t) + E(t)), \quad \delta > 0.$$

Lemma 4.1 (see [14, Lemma 2.1]). *For $t \geq 0$, the functions $I_1(t), I_2(t)$ defined above satisfy*

$$\begin{aligned} \frac{d}{dt} I_1(t) &\leq -\frac{1}{2}(g \circ \nabla u)(t) + 2\|\nabla u(t)\|_2^2, \\ \frac{d}{dt} I_2(t) &\leq -\frac{M(\delta)}{2} \int_{\Omega} \int_0^t K_{\delta}(t-s) g(t-s) |\nabla u(t) - \nabla u(s)|^2 \, ds \, dx + 2\delta M(\delta) \|\nabla u(t)\|_2^2. \end{aligned} \quad (4.1)$$

Moreover, we have

$$\delta M(\delta) \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad (4.2)$$

Next, we further introduce

$$I_3(t) := \int_{\Omega} u_t(t) u(t) \, dx, \quad I_4(t) := - \int_{\Omega} u_t(t) \int_0^t g(t-s) (u(t) - u(s)) \, ds \, dx.$$

The following two lemmas give estimates for the derivatives of $I_3(t)$ and $I_4(t)$.

Lemma 4.2. *Let $(u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega)$ be given. Under the same assumptions of Lemma 2.5, the function $I_3(t)$ defined above satisfies*

$$\begin{aligned} \frac{d}{dt} I_3(t) &\leq -\frac{\ell}{2} \|\nabla u(t)\|_2^2 + \left(1 + \frac{B_2}{\ell} \right) \|u_t(t)\|_2^2 + \frac{1}{\ell} \int_{\Omega} \left| \int_0^t g(t-s) (\nabla u(s) - \nabla u(t)) \, ds \right|^2 \, dx \\ &\quad + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx. \end{aligned} \quad (4.3)$$

Proof. By direct differentiation and the governing equation in (1.1), we obtain

$$\begin{aligned} \frac{d}{dt} I_3(t) &\leq -\ell \|\nabla u(t)\|_2^2 + \|u_t(t)\|_2^2 + \int_{\Omega} \nabla u(t) \cdot \int_0^t g(t-s)(\nabla u(s) - \nabla u(t)) ds dx \\ &\quad - \int_{\Omega} u_t(t)u(t) dx + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| dx. \end{aligned} \quad (4.4)$$

For the third and fourth terms on the right-hand side of (4.4), we apply Cauchy's inequality to estimate

$$\begin{aligned} &\int_{\Omega} \nabla u(t) \cdot \int_0^t g(t-s)(\nabla u(s) - \nabla u(t)) ds dx \\ &\leq \frac{\ell}{4} \|\nabla u(t)\|_2^2 + \frac{1}{\ell} \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(s) - \nabla u(t)) ds \right|^2 dx, \end{aligned}$$

and

$$\begin{aligned} - \int_{\Omega} u_t(t)u(t) dx &\leq \int_{\Omega} |u_t(t)u(t)| dx \leq \int_{\Omega} \left(\frac{1}{4\delta} |u_t(t)|^2 + \delta |u(t)|^2 \right) dx \\ &\leq \frac{1}{4\delta} \|u_t(t)\|_2^2 + \delta B_2 \|\nabla u(t)\|_2^2, \end{aligned}$$

where we used Lemma 2.1 with $k = 2$ and $\delta > 0$ is a constant. Substituting the above two inequalities into (4.4), we arrive at the desired estimate (4.3) by simply choosing $\delta = \frac{\ell}{4B_2}$. \square

Lemma 4.3. *Let $(u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega)$ be given. Under the same assumptions of Lemma 2.5, the function $I_4(t)$ defined above satisfies*

$$\begin{aligned} \frac{d}{dt} I_4(t) &\leq \delta(1 + \alpha \xi_1) \|\nabla u(t)\|_2^2 + c(\delta) \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) ds \right|^2 dx \\ &\quad + \left(\frac{1}{4\delta} + \delta - \int_0^t g(s) ds \right) \|u_t(t)\|_2^2 + \frac{g(0)B_2}{4\delta} (-g' \circ \nabla u)(t), \end{aligned} \quad (4.5)$$

where $\delta > 0$ is a constant to be determined later, $c(\delta) > 0$ denotes a generic constant depending on δ such that $\lim_{\delta \rightarrow 0} c(\delta) = \infty$ and

$$\xi_1 := \frac{B_2}{e(p_1 - 2)} + \frac{B_2(p_2 - 1 + \mu)}{e\mu} \left(\frac{2(1 + \tilde{C})E(0)}{\ell} \right)^{p_2 - 2 + \mu}. \quad (4.6)$$

Proof. Directly differentiating $I_4(t)$ and employing the governing equation of (1.1) yield

$$\begin{aligned} \frac{d}{dt} I_4(t) &= - \int_{\Omega} u_{tt} \int_0^t g(t-s)(u(t) - u(s)) ds dx \\ &\quad - \int_{\Omega} u_t(t) \int_0^t g'(t-s)(u(t) - u(s)) ds dx - \int_0^t g(s) ds \|u_t(t)\|_2^2 \\ &=: \sum_{i=1}^6 I_2^i(t), \end{aligned}$$

where

$$I_2^1(t) := \int_{\Omega} \left(1 - \int_0^t g(s) ds \right) \nabla u(t) \cdot \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) ds dx,$$

$$\begin{aligned}
I_2^2(t) &:= \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx, \\
I_2^3(t) &:= \int_{\Omega} u_t(t) \left(\int_0^t g(t-s)(u(t) - u(s)) \, ds \right) \, dx, \\
I_2^4(t) &:= -\alpha \int_{\Omega} |u(t)|^{p-2} u(t) \log |u(t)| \left(\int_0^t g(t-s)(u(t) - u(s)) \, ds \right) \, dx, \\
I_2^5(t) &:= -\int_{\Omega} u_t(t) \int_0^t g'(t-s)(u(t) - u(s)) \, ds \, dx, \\
I_2^6(t) &:= -\int_0^t g(s) \, ds \|u_t(t)\|_2^2.
\end{aligned}$$

Now we estimate the each of the 6 terms above. For $I_2^1(t)$, we apply Cauchy's inequality with $\delta > 0$ to deduce

$$I_2^1(t) \leq \delta \|\nabla u(t)\|_2^2 + \frac{1}{4\delta} \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx. \quad (4.7)$$

Similarly, we estimate $I_2^3(t)$ as

$$\begin{aligned}
I_2^3(t) &\leq \delta \int_{\Omega} \left(\int_0^t g(t-s)(u(t) - u(s)) \, ds \right)^2 \, dx + \frac{1}{4\delta} \|u_t(t)\|_2^2 \\
&\leq B_2 \delta \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx + \frac{1}{4\delta} \|u_t(t)\|_2^2,
\end{aligned} \quad (4.8)$$

where we used the Poincaré inequality. In a same manner, we first estimate $I_2^5(t)$ as

$$I_2^5(t) \leq \delta \|u_t(t)\|_2^2 + \frac{1}{4\delta} \int_{\Omega} \left| \int_0^t g'(t-s)|u(t) - u(s)| \, ds \right|^2 \, dx. \quad (4.9)$$

By assumption (A1) on g , we apply the Cauchy-Schwarz inequality to treat

$$\begin{aligned}
\left| \int_0^t g'(t-s)|u(t) - u(s)| \, ds \right|^2 &\leq \int_0^t (-g'(t-s)) \, ds \int_0^t (-g'(t-s))|u(t) - u(s)|^2 \, ds \\
&\leq g(0) \int_0^t (-g'(t-s))|u(t) - u(s)|^2 \, ds.
\end{aligned}$$

Then we employ the Poincaré inequality and recall the definition of $(g \circ \nabla u)(t)$ to treat the second term on the right-hand side of (4.9) as

$$\int_{\Omega} \left| \int_0^t g'(t-s)|u(t) - u(s)| \, ds \right|^2 \, dx \leq g(0) \int_0^t (-g'(t-s))\|u(t) - u(s)\|_2^2 \, dx \leq g(0)B_2(-g' \circ \nabla u)(t),$$

indicating

$$I_2^5(t) \leq \delta \|u_t(t)\|_2^2 + \frac{g(0)B_2}{4\delta} (-g' \circ \nabla u)(t). \quad (4.10)$$

Next, we deal with $I_2^4(t)$. For $v \in H_0^1(\Omega)$, we split Ω as that in (2.5) and utilize the assumption (1.2) on p to obtain

$$||v|^{p-2}v \log|v|| \leq \begin{cases} ||v|^{p_1-2}v \log|v|| & \text{in } \Omega_1, \\ ||v|^{p_2-2}v \log|v|| & \text{in } \Omega_2. \end{cases}$$

For Ω_1 , we use the inequality $|x^{p_1-2} \log x| < \frac{1}{e(p_1-2)}$ for $0 < x < 1$ to bound

$$||v|^{p-2} v \log |v|| \leq \frac{|v|}{e(p_1-2)} \quad \text{in } \Omega_1.$$

For Ω_2 , we take advantage of the inequality $|x^{-\mu} \log x| < \frac{1}{e\mu}$ for $x > 1$, where $\mu > 0$ is sufficiently small such that $2(p_2 - 1 + \mu) < \frac{2n}{n-2}$. Then we can bound

$$||v|^{p-2} v \log |v|| \leq \frac{|v|^{p_2-1+\mu}}{e\mu} \quad \text{in } \Omega_2.$$

Combining the above two cases, we obtain

$$||v|^{p-2} v \log |v|| \leq \frac{|v|}{e(p_1-2)} + \frac{|v|^{p_2-1+\mu}}{e\mu} \quad \text{in } \Omega.$$

Therefore, we apply the above estimate with $v = u(t)$ to $I_2^4(t)$ and employ again Cauchy's inequality with $\delta > 0$ to deduce

$$\begin{aligned} I_2^4(t) &\leq \int_{\Omega} \left(\frac{|u(t)|}{e(p_1-2)} + \frac{|u(t)|^{p_2-1+\mu}}{e\mu} \right) \left| \int_0^t g(t-s)(u(t) - u(s)) \, ds \right| \, dx \\ &\leq \delta \int_{\Omega} \left(\frac{|u(t)|^2}{e(p_1-2)} + \frac{|u(t)|^{2(p_2-1+\mu)}}{e\mu} \right) \, dx \\ &\quad + \frac{1}{4\delta} \left(\frac{1}{e(p_1-2)} + \frac{1}{e\mu} \right) \int_{\Omega} \left| \int_0^t g(t-s)(u(t) - u(s)) \, ds \right|^2 \, dx \\ &\leq \delta \left(\frac{\|u(t)\|_2^2}{e(p_1-2)} + \frac{\|u(t)\|_2^{2(p_2-1+\mu)}}{e\mu} \right) + \frac{C}{4\delta} \int_{\Omega} \left| \int_0^t g(t-s)(u(t) - u(s)) \, ds \right|^2 \, dx. \end{aligned}$$

Finally, we apply Lemma 2.1 with $k = 2$ and $k = 2(p_2 - 1 + \mu)$ to dominate

$$\begin{aligned} I_2^4(t) &\leq \delta \left(\frac{B_2}{e(p_1-2)} \|\nabla u(t)\|_2^2 + \frac{B_2(p_2-1+\mu)}{e\mu} \|\nabla u(t)\|_2^{2(p_2-1+\mu)} \right) \\ &\quad + \frac{C}{4\delta} \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx. \end{aligned}$$

For $\|\nabla u(t)\|_2^{2(p_2-1+\mu)}$, we further employ the same argument as that for Lemma 2.7 to estimate

$$\|\nabla u(t)\|_2^{2(p_2-1+\mu)} = \|\nabla u(t)\|_2^{2(p_2-2+\mu)} \|\nabla u(t)\|_2^2 \leq \left(\frac{2(1 + \tilde{C})E(0)}{\ell} \right)^{p_2-2+\mu} \|\nabla u(t)\|_2^2.$$

Then we conclude

$$I_2^4(t) \leq \delta \xi_1 \|\nabla u(t)\|_2^2 + \frac{C}{4\delta} \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx, \quad (4.11)$$

where ξ_1 was defined in (4.6).

Combining all the estimates (4.7), (4.8), (4.10) and (4.11), we eventually arrived at (4.5). \square

Now, we are well prepared to proceed to the proof of Theorem 3.1. We start with introducing an auxiliary function

$$L(t) := N_1 E(t) + I_3(t) + N_2 I_4(t),$$

where $N_1, N_2 > 0$ are constants to be determined later. By Hölder's and Cauchy's inequalities, it is easy to verify that $L(t) \sim E(t)$, i.e., $L(t)$ is equivalent to $E(t)$ in the sense of energy comparison.

Fix $t_1 > 0$ and set $g_1 := \int_0^{t_1} g(s) \, ds > 0$. Combining (2.4), (4.3) and (4.5) with $\delta = \frac{\ell}{4N_2(1+\alpha\xi_1)}$, we estimate $L'(t)$ for all $t \geq t_1$ that

$$\begin{aligned} L'(t) &\leq -\frac{\ell}{4} \|\nabla u(t)\|_2^2 - \left\{ N_1 - 1 - \frac{B_2}{\ell} - N_2 \left(\frac{1}{4\delta} + \delta - g_1 \right) \right\} \|u_t(t)\|_2^2 \\ &\quad + \left(\frac{N_1}{2} - \frac{g(0)B_2N_2}{4\delta} \right) (g' \circ \nabla u)(t) + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx \\ &\quad + \left(\frac{1}{\ell} + c(\delta)N_2 \right) \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx. \end{aligned}$$

At this point, we can choose sufficiently large $N_1, N_2 > 0$ such that

$$N_3 := N_1 - 1 - \frac{B_2}{\ell} - N_2 \left(\frac{1}{4\delta} + \delta - g_1 \right) > 0, \quad \frac{N_1}{2} - \frac{g(0)B_2N_2}{4\delta} > 0.$$

Then the negativity of $(g' \circ \nabla u)(t)$ indicates

$$\begin{aligned} L'(t) &\leq -\frac{\ell}{4} \|\nabla u(t)\|_2^2 - N_3 \|u_t(t)\|_2^2 + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx \\ &\quad + N_4 \int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx, \end{aligned} \tag{4.12}$$

where $N_4 := \frac{1}{\ell} + c(\delta)N_2$. Recalling the definition of $M(\delta)$ in (2.2), we further bound

$$\begin{aligned} &\int_{\Omega} \left| \int_0^t g(t-s)(\nabla u(t) - \nabla u(s)) \, ds \right|^2 \, dx \\ &\leq \int_{\Omega} \int_0^t \frac{g(s)}{K_{\delta}(s)} \, ds \int_0^t K_{\delta}(t-s)g(t-s)|\nabla u(t) - \nabla u(s)|^2 \, ds \, dx \\ &\leq M(\delta) \int_{\Omega} \int_0^t K_{\delta}(t-s)g(t-s)|\nabla u(t) - \nabla u(s)|^2 \, ds \, dx, \end{aligned}$$

which, together with (4.12), indicates

$$\begin{aligned} L'(t) &\leq -\frac{\ell}{4} \|\nabla u(t)\|_2^2 - N_3 \|u_t(t)\|_2^2 + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx \\ &\quad + N_4 M(\delta) \int_{\Omega} \int_0^t K_{\delta}(t-s)g(t-s)|\nabla u(t) - \nabla u(s)|^2 \, ds \, dx. \end{aligned} \tag{4.13}$$

Next, we further introduce

$$J(t) := L(t) + \frac{\ell}{32} I_1(t) + 2N_4 I_2(t).$$

Then it follows from (4.1) and (4.13) that

$$\begin{aligned} J'(t) &\leq - \left(\frac{\ell}{4} - \frac{\ell}{16} - 2N_4 \delta M(\delta) \right) \|\nabla u(t)\|_2^2 - N_3 \|u_t(t)\|_2^2 - \frac{\ell}{64} (g \circ \nabla u)(t) \\ &\quad + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx. \end{aligned}$$

Owing to the convergence (4.2), we can choose sufficiently small $\delta > 0$ such that $2N_4\delta M(\delta) = \frac{\ell}{16}$. Therefore, in view of the original energy $E(t)$, we obtain

$$\begin{aligned} J'(t) &\leq -\frac{\ell}{8}\|\nabla u(t)\|_2^2 - N_3\|u_t(t)\|_2^2 - \frac{\ell}{64}(g \circ \nabla u)(t) + \alpha \int_{\Omega} |u(t)|^p \log |u(t)| \, dx \\ &\leq -\varepsilon E(t) - \left(\frac{\ell}{8} - \frac{\varepsilon}{2}\right)\|\nabla u(t)\|_2^2 - \left(N_3 - \frac{\varepsilon}{2}\right)\|u_t(t)\|_2^2 - \left(\frac{\ell}{64} - \frac{\varepsilon}{2}\right)(g \circ \nabla u)(t) \\ &\quad + \alpha \int_{\Omega} \left(1 - \frac{\varepsilon}{p}\right)|u(t)|^p \log |u(t)| \, dx + \frac{\alpha\varepsilon}{p_1^2} \int_{\Omega} |u(t)|^p \, dx, \end{aligned} \quad (4.14)$$

where $\varepsilon \in (0, p_1)$ is a constant. Employing the same domain splitting argument as that for (2.6), we estimate

$$\begin{aligned} \int_{\Omega} \left(1 - \frac{\varepsilon}{p}\right)|u(t)|^p \log |u(t)| \, dx &= \left(\int_{\Omega_1} + \int_{\Omega_2}\right) \left(1 - \frac{\varepsilon}{p}\right)|u(t)|^p \log |u(t)| \, dx \\ &\leq \frac{1}{\varepsilon\mu} \int_{\Omega_2} \left(1 - \frac{\varepsilon}{p}\right)|u(t)|^{p_2}|u(t)|^{\mu} \, dx \leq \frac{1}{\varepsilon\mu} \int_{\Omega} |u(t)|^{p_2+\mu} \, dx \\ &\leq \frac{B_{p_2+\mu}}{\varepsilon\mu} \|\nabla u(t)\|_2^{p_2+\mu} \leq \xi_2 \|\nabla u(t)\|_2^2, \end{aligned} \quad (4.15)$$

where

$$\xi_2 := \frac{B_{p_2+\mu}}{\varepsilon\mu} \left(\frac{2(1+\tilde{C})E(0)}{\ell}\right)^{\frac{p_2+\mu-2}{2}}$$

and $\mu > 0$ is a sufficiently small constant such that $p_2 + \mu < 2_* = \frac{2n}{n-2}$.

On the other hand, according to Lemma 2.3, there exists a constant $B_{p(x)} > 0$ such that

$$\|v\|_{p(x)} \leq B_{p(x)} \|\nabla v\|_2, \quad \forall v \in H_0^1(\Omega).$$

Hence, making use of (2.1), we deal with the last term in (4.14) in a similar manner as above as

$$\begin{aligned} \int_{\Omega} |u(t)|^p \, dx &\leq \max \left\{ \|u(t)\|_{p(x)}^{p_1}, \|u(t)\|_{p(x)}^{p_2} \right\} \leq \max \left\{ B_{p(x)}^{p_1} \|\nabla u(t)\|_2^{p_1}, B_{p(x)}^{p_2} \|\nabla u(t)\|_2^{p_2} \right\} \\ &\leq \xi_3 \|\nabla u(t)\|_2^2, \end{aligned} \quad (4.16)$$

where

$$\xi_3 := \max \left\{ B_{p(x)}^{p_1} \left(\frac{2(1+\tilde{C})E(0)}{\ell}\right)^{\frac{p_1-2}{2}}, B_{p(x)}^{p_2} \left(\frac{2(1+\tilde{C})E(0)}{\ell}\right)^{\frac{p_2-2}{2}} \right\}.$$

Plugging (4.15) and (4.16) into (4.14), we arrive at

$$J'(t) \leq -\varepsilon E(t) - \left(\frac{\ell}{8} - \frac{\varepsilon}{2} - \alpha\xi_2 - \frac{\alpha\varepsilon\xi_3}{p_1^2}\right) \|\nabla u(t)\|_2^2 - \left(N_3 - \frac{\varepsilon}{2}\right) \|u_t(t)\|_2^2 - \left(\frac{\ell}{64} - \frac{\varepsilon}{2}\right) (g \circ \nabla u)(t).$$

Choosing ε and α sufficiently small such that

$$\frac{\ell}{8} - \frac{\varepsilon}{2} - \alpha\xi_2 - \frac{\alpha\varepsilon\xi_3}{p_1^2} > 0, \quad N_3 - \frac{\varepsilon}{2} > 0, \quad \frac{\ell}{64} - \frac{\varepsilon}{2} > 0,$$

we eventually conclude

$$J'(t) \leq -\varepsilon E(t).$$

Since $J(t) \geq 0$ for all $t \geq 0$ and $J(t_0) \leq CE(0)$, we integrate the above inequality over $[t_0, \tau)$ for any $\tau > t_0$ to obtain

$$\int_{t_0}^{\tau} E(t) dt \leq CE(0).$$

Passing $\tau \rightarrow \infty$ yields the first estimate in (3.1), which, together with $E'(t) \leq 0$, implies the second estimate in (3.1). The proof of Theorem 3.1 is completed.

4.2. Proof of Theorem 3.3. The proof of Theorem 3.3 relies on the following key lemma.

Lemma 4.4 (see [23, Equation (44)]). *Let assumptions (A1)–(A2) be satisfied. Then for all $t > 0$, there holds*

$$(g \circ \nabla u)(t) \leq \frac{1}{c} \overline{G}^{-1} \left(\frac{c(-g' \circ \nabla u)(t)}{\zeta(t)} \right),$$

where $c \in (0, 1)$ is a sufficiently small constant, G and ζ are defined in (A2), and \overline{G} is a strictly increasing and strictly convex extension of G .

Now we can proceed to the proof of Theorem 3.3. By the Cauchy-Schwarz inequality and assumption (A1), first we obtain

$$\begin{aligned} \int_{\Omega} \left| \int_0^t g(t-s) (\nabla u(t) - \nabla u(s)) ds \right|^2 dx &\leq \int_{\Omega} \int_0^t g(s) ds \int_0^t g(t-s) |\nabla u(t) - \nabla u(s)|^2 ds dx \\ &\leq (1-\ell)(g \circ \nabla u)(t), \end{aligned}$$

which, together with (4.13)–(4.16), indicates

$$L'(t) \leq -\varepsilon E(t) + C(g \circ \nabla u)(t).$$

According to Lemma 4.4, we deduce

$$L'(t) \leq -\varepsilon E(t) + C \overline{G}^{-1} \left(\frac{c(-g' \circ \nabla u)(t)}{\zeta(t)} \right).$$

Next, defining

$$F(t) := \overline{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) L(t)$$

with a constant $\varepsilon_1 \in (0, r)$, we immediately see $F(t) \sim E(t)$. Noting that $\overline{G}''(t) \geq 0$ and $E'(t) \leq 0$, we obtain

$$\begin{aligned} F'(t) &= \overline{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) L'(t) + \varepsilon_1 \frac{E'(t)}{E(0)} \overline{G}'' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) L(t) \\ &\leq -\varepsilon E(t) \overline{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C \overline{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) \overline{G}^{-1} \left(\frac{c(-g' \circ \nabla u)(t)}{\zeta(t)} \right). \end{aligned} \quad (4.17)$$

Let \overline{G}^* be the convex conjugate of \overline{G} in the sense of Young, that is,

$$\overline{G}^*(s) := s(\overline{G}')^{-1}(s) - (\overline{G} \circ (\overline{G}')^{-1})(s), \quad s \in (0, \overline{G}'(r)]. \quad (4.18)$$

Then it is known that \overline{G}^* satisfies the generalized Young inequality

$$AB \leq \overline{G}^*(A) + \overline{G}(B), \quad \forall A \in (0, \overline{G}'(r)], \forall B \in (0, r].$$

Thus, taking

$$A = \overline{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right), \quad B = \overline{G}^{-1} \left(\frac{c(-g' \circ \nabla u)(t)}{\zeta(t)} \right)$$

in the above inequality and using (4.18), we can further estimate (4.17) as

$$\begin{aligned} F'(t) &\leq -\varepsilon E(t) \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C \bar{G}^* \left(\bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) \right) + C \frac{(-g' \circ \nabla u)(t)}{\zeta(t)} \\ &\leq -\varepsilon E(t) \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C \varepsilon_1 \frac{E(t)}{E(0)} \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C \frac{(-g' \circ \nabla u)(t)}{\zeta(t)}. \end{aligned}$$

Multiplying both sides of the above inequality by $\zeta(t)$ and using the facts that

$$(-g' \circ \nabla u)(t) \leq -CE'(t), \quad \varepsilon_1 \frac{E(t)}{E(0)} < r, \quad \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) = G' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right),$$

we derive

$$\begin{aligned} \zeta(t) F'(t) &\leq -\varepsilon \zeta(t) E(t) \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C \varepsilon_1 \zeta(t) \frac{E(t)}{E(0)} \bar{G}' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) + C(-g' \circ \nabla u)(t) \\ &\leq -(\varepsilon E(0) - C \varepsilon_1) \zeta(t) \frac{E(t)}{E(0)} G' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) - CE'(t). \end{aligned} \quad (4.19)$$

Next, we further introduce $F_1(t) := \zeta(t) F(t) + CE(t)$. Then it is readily seen that $F_1(t) \sim E(t)$, i.e., there exist constants $\beta_1, \beta_2 > 0$ such that

$$\beta_1 F_1(t) \leq E(t) \leq \beta_2 F_1(t). \quad (4.20)$$

Choosing $\varepsilon_1 > 0$ sufficiently small such that $\beta_3 := \varepsilon E(0) - C \varepsilon_1 > 0$, we differentiate $F_1(t)$ and utilize (4.19) to deduce

$$F_1'(t) \leq -\beta_3 \zeta(t) \frac{E(t)}{E(0)} G' \left(\varepsilon_1 \frac{E(t)}{E(0)} \right) = -\beta_3 \zeta(t) G_2 \left(\frac{E(t)}{E(0)} \right), \quad (4.21)$$

where we put $G_2(t) := t G'(\varepsilon_1 t)$. Since $G_2'(t) = G'(\varepsilon_1 t) + \varepsilon_1 t G''(\varepsilon_1 t)$, it follows from the strict monotonicity and the strict convexity of G on $(0, r]$ that $G_2(t) > 0$ and $G_2'(t) > 0$ for all $t \in (0, 1]$. Therefore, further defining $F_2(t) := \frac{\beta_1 F_1(t)}{E(0)}$, we see $F_2(t) \sim E(t)$ and the combination of (4.20) and (4.21) implies

$$F_2'(t) \leq -k_1 \zeta(t) G_2 \left(\frac{E(t)}{E(0)} \right) \leq -k_1 \zeta(t) G_2 \left(\frac{\beta_1 F_1(t)}{E(0)} \right) = -k_1 \zeta(t) G_2(F_2(t)),$$

where $k_1 := \frac{\beta_1 \beta_3}{E(0)}$. Hence, integrating both sides over $(t_1, t]$ yields

$$\int_{\varepsilon_1 F_2(t)}^{\varepsilon_1 F_2(t_1)} \frac{1}{\tau G'(\tau)} d\tau = \int_{t_1}^t -\frac{F_2'(s)}{G_2(F_2(s))} ds \geq k_1 \int_{t_1}^t \zeta(s) ds,$$

where we performed a change of variables with $\tau = F_2(s)$. Finally, we conclude

$$F_2(t) \leq \frac{1}{\varepsilon_1} G_1^{-1} \left(k_1 \int_{t_1}^t \zeta(s) ds \right),$$

where $G_1(t) := \int_t^r \frac{1}{s G'(s)} ds$ and we used the fact that G_1 is strictly decreasing on $(0, r]$. This eventually lead us to (3.2) in view of the equivalence $F_2(t) \sim E(t)$, which finalized the proof of Theorem 3.3.

4.3. Proof of Theorem 3.5. Similarly as the previous subsection, we first prepare a key lemma for the proof of Theorem 3.5.

Lemma 4.5 (see [19]). *Let $E : [0, \infty) \rightarrow (0, \infty)$ be a non-increasing function and $\phi : [0, \infty) \rightarrow [0, \infty)$ be a strictly increasing C^1 function such that*

$$\phi(0) = 0, \quad \lim_{t \rightarrow \infty} \phi(t) = \infty.$$

Suppose that there exist constants $\sigma \geq 0$ and $\omega > 0$ such that

$$\int_t^\infty \phi'(s) E(s)^{1+\sigma} ds \leq \frac{1}{\omega} E(0)^\sigma E(t).$$

Then the following decay estimates hold for $E(t)$.

- (1) *If $\sigma = 0$, then $E(t) \leq E(0) e^{1-\omega\phi(t)}$ for all $t \geq 0$.*
- (2) *If $\sigma > 0$, then $E(t) \leq E(0) \left(\frac{1+\sigma}{1+\omega\sigma\phi(t)}\right)^{\frac{1}{\sigma}}$ for all $t \geq 0$.*

We now proceed to the proof of Theorem 3.5 based on the above lemma.

Fixing some $T > 0$, we multiply the governing equation in (1.1) by $\xi(t)E(t)^{q-1}u(t)$ and integrate over $\Omega \times (\tau, T)$ with some $\tau \in (0, T)$ to derive

$$\begin{aligned} & \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega u_{tt}(t) u(t) dx dt + \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega u_t(t) u(t) dx dt \\ & + \int_\tau^T \xi(t) E(t)^{q-1} \|\nabla u(t)\|_2^2 dt - \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega \nabla u(t) \cdot \int_0^t g(t-s) \nabla u(s) ds dx dt \\ & = \alpha \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega |u(t)|^p \log |u(t)| dx dt, \end{aligned} \quad (4.22)$$

where it is straightforward to see

$$\begin{aligned} & \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega \nabla u(t) \cdot \int_0^t g(t-s) \nabla u(s) ds dx dt \\ & = \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega \int_0^t g(t-s) (\nabla u(s) - \nabla u(t)) \cdot \nabla u(t) ds dx dt \\ & + \int_\tau^T \xi(t) E(t)^{q-1} \int_0^t g(s) ds \|\nabla u(t)\|_2^2 dt. \end{aligned}$$

Then we substitute the above identity into (4.22) and rearrange to obtain

$$\begin{aligned} & \int_\tau^T \xi(t) E(t)^{q-1} \left(1 - \int_0^t g(s) ds\right) \|\nabla u(t)\|_2^2 dt \\ & = - \int_\tau^T \xi(t) E(t)^{q-1} \frac{d}{dt} \int_\Omega u_t(t) u(t) dx dt + \int_\tau^T \xi(t) E(t)^{q-1} \|u_t(t)\|_2^2 dt \\ & + \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega \int_0^t g(t-s) (\nabla u(s) - \nabla u(t)) \cdot \nabla u(t) ds dx dt \\ & - \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega u_t(t) u(t) dx dt + \alpha \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega |u(t)|^p \log |u(t)| dx dt. \end{aligned}$$

Then we turn to the definition of $E(t)$ in (2.3) and employ the above identity to calculate

$$\begin{aligned} 2 \int_{\tau}^T \xi(t) E(t)^q dt &= \int_{\tau}^T \xi(t) E(t)^{q-1} \left\{ \|u_t(t)\|_2^2 + \left(1 - \int_0^t g(s) ds\right) \|\nabla u(t)\|_2^2 + (g \circ \nabla u)(t) \right. \\ &\quad \left. - 2\alpha \int_{\Omega} \frac{|u(t)|^p \log |u(t)|}{p} dx + 2\alpha \int_{\Omega} \frac{|u(t)|^p}{p^2} dx\right\} dt \\ &= \sum_{i=1}^7 J_i(\tau), \end{aligned} \quad (4.23)$$

where

$$\begin{aligned} J_1(\tau) &:= 2 \int_{\tau}^T \xi(t) E(t)^{q-1} \|u_t(t)\|_2^2 dt, \\ J_2(\tau) &:= - \int_{\tau}^T \xi(t) E(t)^{q-1} \int_{\Omega} u_t(t) u(t) dx dt, \\ J_3(\tau) &:= - \int_{\tau}^T \xi(t) E(t)^{q-1} \frac{d}{dt} \int_{\Omega} u_t(t) u(t) dx dt, \\ J_4(\tau) &:= \int_{\tau}^T \xi(t) E(t)^{q-1} (g \circ \nabla u)(t) dt, \\ J_5(\tau) &:= \int_{\tau}^T \xi(t) E(t)^{q-1} \int_{\Omega} \int_0^t g(t-s) (\nabla u(s) - \nabla u(t)) \cdot \nabla u(t) ds dx dt, \\ J_6(\tau) &:= \alpha \int_{\tau}^T \xi(t) E(t)^{q-1} \int_{\Omega} \left(1 - \frac{2}{p}\right) |u(t)|^p \log |u(t)| dx dt, \\ J_7(\tau) &:= 2\alpha \int_{\tau}^T \xi(t) E(t)^{q-1} \int_{\Omega} \frac{|u(t)|^p}{p^2} dx dt. \end{aligned}$$

In the sequel, we estimate each of the 7 terms above.

Step 1 We first estimate $J_1(\tau)$ and $J_2(\tau)$. Since (2.4) implies $\|u_t(t)\|_2^2 \leq -E'(t)$, it follows immediately from the monotonicity of $\xi(t)$ and $E(t)$ that

$$\begin{aligned} J_1(\tau) &\leq -2 \int_{\tau}^T \xi(t) E(t)^{q-1} E'(t) dt \leq 2\xi(0) \left[\frac{E(t)^q}{q} \right]_{t=\tau}^{t=T} \\ &\leq \frac{2\xi(0)}{q} E(\tau)^q \leq \frac{2\xi(0) E(0)^{q-1}}{q} E(\tau). \end{aligned} \quad (4.24)$$

For $J_2(\tau)$, we apply Cauchy's inequality with $\delta \in (0, 1)$ and the above inequality to deduce

$$\begin{aligned} J_2(\tau) &\leq \int_{\tau}^T \xi(t) E(t)^{q-1} \int_{\Omega} \left(\frac{1}{4\delta} |u_t(t)|^2 + \delta |u(t)|^2 \right) dx dt = \frac{1}{8\delta} J_1(\tau) + \delta \int_{\tau}^T \xi(t) E(t)^{q-1} \|u(t)\|_2^2 dt \\ &\leq \frac{\xi(0) E(0)^{q-1}}{4\delta q} E(\tau) + \delta B_2 \int_{\tau}^T \xi(t) E(t)^{q-1} \|\nabla u(t)\|_2^2 dt, \end{aligned}$$

where again we used the Poincaré inequality. To deal with the last term above, we recall the definition (2.7) of $\mathbb{E}(t)$ and apply Lemma 2.7 to bound

$$\frac{\ell}{2} \|\nabla u(t)\|_2^2 \leq \frac{1}{2} \left(1 - \int_0^t g(s) ds \right) \|\nabla u(t)\|_2^2 \leq \mathbb{E}(t) \leq (1 + \tilde{C}) E(t). \quad (4.25)$$

Thus it reveals that

$$J_2(t) \leq \frac{\xi(0)E(0)^{q-1}}{4\delta q} E(\tau) + \frac{2\delta B_2(1+\tilde{C})}{\ell} \int_{\tau}^T \xi(t)E(t)^q dt. \quad (4.26)$$

Step 2 Next, we deal with $J_3(\tau)$. Differentiating $\xi(t)E(t)^{q-1} \int_{\Omega} u_t(t)u(t) dx$, we divide $J_3(\tau) = J_3^1(\tau) + J_3^2(\tau) + J_3^3(\tau)$ with

$$\begin{aligned} J_3^1(\tau) &:= - \int_{\tau}^T \frac{d}{dt} \left(\xi(t)E(t)^{q-1} \int_{\Omega} u_t(t)u(t) dx \right) dt, \\ J_3^2(\tau) &:= (q-1) \int_{\tau}^T \xi(t)E(t)^{q-2} E'(t) \int_{\Omega} u_t(t)u(t) dx dt \quad (q > 1), \\ J_3^3(\tau) &:= \int_{\tau}^T \xi'(t)E(t)^{q-1} \int_{\Omega} u_t(t)u(t) dx dt. \end{aligned}$$

For $J_3^1(\tau)$, again we employ the monotonicity of $\xi(t), E(t)$ and the Poincaré inequality to deduce

$$\begin{aligned} J_3^1(\tau) &= \left[\xi(t)E(t)^{q-1} \int_{\Omega} u_t(t)u(t) dx \right]_{t=T}^{t=\tau} \leq \xi(0)E(0)^{q-1} \int_{\Omega} (|u_t(\tau)u(\tau)| + |u_t(T)u(T)|) dx \\ &\leq \frac{\xi(0)E(0)^{q-1}}{2} (\|u_t(\tau)\|_2^2 + \|u(\tau)\|_2^2 + \|u_t(T)\|_2^2 + \|u(T)\|_2^2) dx \\ &\leq \frac{\xi(0)E(0)^{q-1}}{2} \{ \|u_t(\tau)\|_2^2 + \|u_t(T)\|_2^2 + B_2 (\|\nabla u(\tau)\|_2^2 + \|\nabla u(T)\|_2^2) \}. \end{aligned}$$

Similarly to the argument for (4.25), we have

$$\|u_t(t)\|_2^2 \leq 2(1+\tilde{C})E(t), \quad \|\nabla u(t)\|_2^2 \leq \frac{2(1+\tilde{C})}{\ell} E(t), \quad \forall t \geq 0,$$

and hence

$$J_3^1(\tau) \leq 2(1+\tilde{C}) \left(1 + \frac{B_2}{\ell} \right) \xi(0)E(0)^{q-1} E(\tau). \quad (4.27)$$

In an analogous manner, we estimate $J_3^2(t)$ as

$$\begin{aligned} J_3^2(\tau) &\leq -(q-1) \int_{\tau}^T \xi(t)E(t)^{q-2} E'(t) \int_{\Omega} |u_t(t)u(t)| dx dt \\ &\leq -\frac{q-1}{2} \int_{\tau}^T \xi(t)E(t)^{q-2} E'(t) (\|u_t(t)\|_2^2 + B_2 \|\nabla u(t)\|_2^2) dt \\ &\leq -(q-1)(1+\tilde{C}) \left(1 + \frac{B_2}{\ell} \right) \int_{\tau}^T \xi(t)E(t)^{q-1} E'(t) dt. \end{aligned}$$

Owing to the estimate of $\int_{\tau}^T \xi(t)E(t)^{q-1} E'(t) dt$ obtained in (4.24), we immediately get

$$\begin{aligned} J_3^2(\tau) &\leq \frac{(q-1)(1+\tilde{C})\xi(0)E(0)^{q-1}}{q} \left(1 + \frac{B_2}{\ell} \right) E(\tau) \\ &\leq (1+\tilde{C})\xi(0)E(0)^{q-1} \left(1 + \frac{B_2}{\ell} \right) E(\tau). \end{aligned} \quad (4.28)$$

For $J_3^3(\tau)$, we repeat the same argument and perform integration by parts to derive

$$\begin{aligned} J_3^3(\tau) &\leq \frac{1}{2} \int_{\tau}^T \xi'(t) E(t)^{q-1} (\|u_t(t)\|_2^2 + B_2 \|\nabla u(t)\|_2^2) dt \leq -(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell}\right) \int_{\tau}^T \xi'(t) E(t)^q dt \\ &= (1 + \tilde{C}) \left(1 + \frac{B_2}{\ell}\right) \left\{ \left[\xi(t) E(t)^q \right]_{t=T}^{t=\tau} + q \int_{\tau}^T \xi(t) E(t)^{q-1} E'(t) dt \right\} \\ &\leq (1 + \tilde{C}) \left(1 + \frac{B_2}{\ell}\right) \xi(0) E(0)^{q-1} E(\tau). \end{aligned} \quad (4.29)$$

Therefore, summing up (4.27)–(4.29) yields

$$J_3(\tau) \leq 4(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell}\right) \xi(0) E(0)^{q-1} E(\tau). \quad (4.30)$$

Step 3 To investigate $J_4(\tau)$ and $J_5(\tau)$, we shall divide the estimates into 2 cases, i.e., $q = 1$ and $1 < q < 2$.

Case 1 For $q = 1$, assumption (A3) reduces to $g'(t) \leq -\xi(t)g(t)$. Since (2.4) gives $(-g' \circ \nabla u)(t) \leq -2E'(t)$, the monotonicity of $\xi(t)$ indicates

$$\begin{aligned} J_4(\tau) &\leq \int_{\tau}^T \int_0^t \xi(t-s) g(t-s) \|\nabla u(s) - \nabla u(t)\|_2^2 ds dt \leq \int_{\tau}^T (-g' \circ \nabla u)(t) dt \\ &\leq -2 \int_{\tau}^T E'(t) dt \leq 2E(\tau). \end{aligned} \quad (4.31)$$

As for $J_5(\tau)$, we apply Cauchy's inequality with $\varepsilon > 0$ and make use of (4.25) and (4.31) to derive

$$\begin{aligned} J_5(\tau) &\leq \frac{\varepsilon}{2} \int_{\tau}^T \xi(t) \int_0^t g(t-s) \|\nabla u(t)\|_2^2 ds dt + \frac{1}{2\varepsilon} \int_{\tau}^T \int_0^t \xi(t-s) g(t-s) \|\nabla u(s) - \nabla u(t)\|_2^2 ds dt \\ &\leq \frac{\varepsilon(1 + \tilde{C})(1 - \ell)}{\ell} \int_{\tau}^T \xi(t) E(t) dt + \frac{1}{\varepsilon} E(\tau) \leq \varepsilon(1 + \tilde{C}) \int_{\tau}^T \xi(t) E(t) dt + \frac{1}{\varepsilon} E(\tau). \end{aligned} \quad (4.32)$$

Case 2 Now let us consider the case of $1 < q < 2$. Applying Hölder's inequality with q and $q^* = \frac{q}{q-1}$, we employ (4.25) and (3.1) to deduce

$$\begin{aligned} (g \circ \nabla u)(t) &= \int_0^t \|\nabla u(s) - \nabla u(t)\|_2^{\frac{2(q-1)}{q}} g(t-s) \|\nabla u(s) - \nabla u(t)\|_2^{\frac{2}{q}} ds \\ &\leq \left(\int_0^t \|\nabla u(s) - \nabla u(t)\|_2^2 ds \right)^{1-\frac{1}{q}} \left(\int_0^t g(t-s)^q \|\nabla u(s) - \nabla u(t)\|_2^2 ds \right)^{\frac{1}{q}} \\ &\leq \left(2 \int_0^t (\|\nabla u(t)\|_2^2 + \|\nabla u(s)\|_2^2) ds \right)^{1-\frac{1}{q}} (g^q \circ \nabla u)(t)^{\frac{1}{q}} \\ &\leq C \left(\int_0^t (E(t) + E(s)) ds \right)^{1-\frac{1}{q}} (g^q \circ \nabla u)(t)^{\frac{1}{q}} \leq CE(0)^{1-\frac{1}{q}} (g^q \circ \nabla u)(t)^{\frac{1}{q}}. \end{aligned}$$

Then we apply Young's inequality with $\epsilon > 0$ to estimate $J_4(\tau)$ as

$$J_4(\tau) \leq CE(0)^{1-\frac{1}{q}} \int_{\tau}^T \xi(t) E(t)^{q-1} (g^q \circ \nabla u)(t)^{\frac{1}{q}} dt$$

$$\begin{aligned}
&= CE(0)^{1-\frac{1}{q}} \int_{\tau}^T (\xi(t)E(t)^q)^{\frac{q-1}{q}} (\xi(t)(g^q \circ \nabla u)(t))^{\frac{1}{q}} dt \\
&\leq CE(0)^{1-\frac{1}{q}} \epsilon \int_{\tau}^T \xi(t)E(t)^q dt + C(\epsilon) \int_{\tau}^T \xi(t)(g^q \circ \nabla u)(t) dt.
\end{aligned}$$

For the last term above, we utilize assumption (A3) and the same argument for (4.31) to deduce

$$\begin{aligned}
\int_{\tau}^T \xi(t)(g^q \circ \nabla u)(t) dt &\leq \int_{\tau}^T \int_0^t \xi(t-s)g(t-s)^q \|\nabla u(s) - \nabla u(t)\|_2^2 ds dt \\
&\leq \int_{\tau}^T (-g' \circ \nabla u)(t) dt \leq -2 \int_{\tau}^T E'(t) dt \leq 2E(\tau),
\end{aligned}$$

and hence

$$J_4(\tau) \leq C\epsilon E(0)^{1-\frac{1}{q}} \int_{\tau}^T \xi(t)E(t)^q dt + C(\epsilon)E(\tau). \quad (4.33)$$

As for $J_5(\tau)$, we mimic the estimate in (4.32) and use (4.33) to derive

$$\begin{aligned}
J_5(\tau) &\leq \frac{\varepsilon}{2} \int_{\tau}^T \xi(t)E(t)^{q-1} \int_0^t g(t-s) \|\nabla u(t)\|_2^2 ds dt + \frac{1}{2\varepsilon} \int_{\tau}^T \xi(t)E(t)^{q-1}(g \circ \nabla u)(t) dt \\
&\leq \varepsilon(1 + \tilde{C}) \int_{\tau}^T \xi(t)E(t)^q dt + \frac{1}{2\varepsilon} J_4(\tau) \\
&\leq \varepsilon(1 + \tilde{C}) \int_{\tau}^T \xi(t)E(t)^q dt + \frac{1}{2\varepsilon} \left\{ C\epsilon E(0)^{1-\frac{1}{q}} \int_{\tau}^T \xi(t)E(t)^q dt + C(\epsilon)E(\tau) \right\} \\
&\leq \varepsilon \left(1 + \tilde{C} + CE(0)^{1-\frac{1}{q}} \right) \int_{\tau}^T \xi(t)E(t)^q + C(\varepsilon)E(\tau),
\end{aligned} \quad (4.34)$$

where we took $\epsilon = \varepsilon^2$ in the last inequality.

Step 4 Now it remains to deal with the two nonlinear terms $J_6(\tau)$ and $J_7(\tau)$. To start with, again we take advantage of the same argument in (2.5)–(2.6) to estimate

$$\begin{aligned}
\int_{\Omega} \left(1 - \frac{2}{p} \right) |u(t)|^p \log |u(t)| dx &\leq \int_{\{|u(t)| \geq 1\}} \left(1 - \frac{2}{p} \right) |u(t)|^p \log |u(t)| dx \\
&\leq \left(1 - \frac{2}{p_2} \right) \int_{\{|u(t)| \geq 1\}} |u(t)|^{p_2} \log |u(t)| dt \\
&\leq \left(1 - \frac{2}{p_2} \right) \frac{1}{\epsilon \mu} \int_{\Omega} |u(t)|^{p_2+\mu} dx \leq \frac{(p_2-2)B_{p_2+\mu}}{\epsilon \mu p_2} \|\nabla u(t)\|_2^{p_2+\mu},
\end{aligned}$$

where $\mu > 0$ satisfies $p_2 + \mu < 2_*$. Then it follows from (4.25) that

$$\begin{aligned}
J_6(\tau) &\leq \alpha \frac{(p_2-2)B_{p_2+\mu}}{\epsilon \mu p_2} \int_{\tau}^T \xi(t)E(t)^{q-1} \|\nabla u(t)\|_2^{p_2+\mu} dx \\
&\leq \alpha \frac{(p_2-2)B_{p_2+\mu}}{\epsilon \mu p_2} \left(\frac{2(1+\tilde{C})}{\ell} \right)^{\frac{p_2+\mu}{2}} \int_{\tau}^T \xi(t)E(t)^{q+\frac{p_2+\mu-2}{2}} dt \leq \alpha \xi_4 \int_{\tau}^T \xi(t)E(t)^q dt, \quad (4.35)
\end{aligned}$$

where

$$\xi_4 := \frac{(p_2 - 2)B_{p_2+\mu}}{\epsilon\mu p_2} \left(\frac{2(1 + \tilde{C})}{\ell} \right)^{\frac{p_2+\mu}{2}} E(0)^{\frac{p_2+\mu-2}{2}}.$$

Finally, for $J_7(\tau)$, we invoke the estimate (4.16) to deduce

$$\begin{aligned} J_7(\tau) &\leq \frac{2\alpha}{p_1^2} \int_\tau^T \xi(t) E(t)^{q-1} \int_\Omega |u(t)|^p dx dt \leq \frac{2\alpha\xi_3}{p_1^2} \int_\tau^T \xi(t) E(t)^{q-1} \|\nabla u(t)\|_2^2 dt \\ &\leq \frac{4\alpha(1 + \tilde{C})\xi_3}{p_1^2\ell} \int_\tau^T \xi(t) E(t)^q dt. \end{aligned} \quad (4.36)$$

At this stage, we are in a position to complete the proof of Theorem 3.5.

Case 1 For $q = 1$, we substitute (4.24), (4.26), (4.30)–(4.32), (4.35) and (4.36) into (4.23) to dominate

$$\begin{aligned} 2 \int_\tau^T \xi(t) E(t) dt &\leq \left\{ 2 + \frac{1}{\epsilon} + \left(2 + 4(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell} \right) + \frac{1}{4\delta} \right) \xi(0) \right\} E(\tau) \\ &\quad + \left\{ (1 + \tilde{C}) \left(\epsilon + \frac{2\delta B_2}{\ell} \right) + \alpha \left(\xi_4 + \frac{4(1 + \tilde{C})\xi_3}{p_1^2\ell} \right) \right\} \int_\tau^T \xi(t) E(t) dt. \end{aligned} \quad (4.37)$$

Therefore, if $\alpha > 0$ is sufficiently small such that

$$\alpha < \left(\xi_4 + \frac{4(1 + \tilde{C})\xi_3}{p_1^2\ell} \right)^{-1}, \quad (4.38)$$

then we can suitably choose sufficiently small $\epsilon, \delta > 0$ such that

$$(1 + \tilde{C}) \left(\epsilon + \frac{2\delta B_2}{\ell} \right) + \alpha \left(\xi_4 + \frac{4(1 + \tilde{C})\xi_3}{p_1^2\ell} \right) = 1.$$

Thus, we can rearrange (4.37) as

$$\int_\tau^T \xi(t) E(t) dt \leq \frac{1}{K} E(\tau), \quad K := \left\{ 2 + \frac{1}{\epsilon} + \left(2 + 4(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell} \right) + \frac{1}{4\delta} \right) \xi(0) \right\}^{-1}.$$

Since $T > 0$ was chosen arbitrarily, we passing $T \rightarrow \infty$ to obtain

$$\int_\tau^\infty \xi(t) E(t) dt \leq \frac{1}{K} E(\tau).$$

Finally, setting $\phi(t) = \int_0^t \xi(s) ds$, $\sigma = 0$ and $\omega = K$ in Lemma 4.5, we arrive at the first inequality in (3.3).

Case 2 For $1 < q < 2$, we follow the same line as above to substitute (4.24), (4.26), (4.30) and (4.33)–(4.36) into (4.23) to dominate

$$\begin{aligned} 2 \int_\tau^T \xi(t) E(t)^q dt &\leq \left\{ C(\epsilon) + \xi(0) E(0)^{q-1} \left(\frac{2}{q} + \frac{1}{4\delta q} + 4(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell} \right) \right) \right\} E(\tau) \\ &\quad + \left\{ \frac{2\delta B_2(1 + \tilde{C})}{\ell} + C\epsilon^2 E(0)^{1-\frac{1}{q}} + \epsilon \left(1 + \tilde{C} + CE(0)^{1-\frac{1}{q}} \right) \right\} \end{aligned}$$

$$+ \alpha \left(\xi_4 + \frac{4(1 + \tilde{C})\xi_3}{p_1^2 \ell} \right) \right\} \int_{\tau}^T \xi(t) E(t) dt, \quad (4.39)$$

where we set $\epsilon = \varepsilon^2$ in (4.33). Therefore, if $\alpha > 0$ satisfies the same condition (4.38) as above, we can still choose $\varepsilon, \delta > 0$ suitably such that

$$\frac{2\delta B_2(1 + \tilde{C})}{\ell} + C\varepsilon^2 E(0)^{1-\frac{1}{q}} + \varepsilon \left(1 + \tilde{C} + CE(0)^{1-\frac{1}{q}} \right) + \alpha \left(\xi_4 + \frac{4(1 + \tilde{C})\xi_3}{p_1^2 \ell} \right) = 1.$$

Then we can rearrange (4.39) as

$$\int_{\tau}^T \xi(t) E(t)^q dt \leq \frac{1}{K'} E(0)^{q-1} E(\tau),$$

where

$$K' := \left\{ C(\varepsilon) + \xi(0) E(0)^{q-1} \left(\frac{2}{q} + \frac{1}{4\delta q} + 4(1 + \tilde{C}) \left(1 + \frac{B_2}{\ell} \right) \right) \right\}^{-1}.$$

Thus, passing $T \rightarrow \infty$ yields

$$\int_{\tau}^{\infty} \xi(t) E(t)^q dt \leq \frac{1}{K'} E(0)^{q-1} E(\tau).$$

where As a result, the application of Lemma 4.5 with $\phi(t) = \int_0^t \xi(s) ds$, $\sigma = q - 1$ and $\omega = K'$ implies the second inequality in (3.3).

The proof of Theorem 3.5 is completed.

ACKNOWLEDGEMENTS

The second author is supported by JSPS KAKENHI Grant Numbers JP22K13954, JP23KK0049 and Guangdong Basic and Applied Basic Research Foundation (No. 2025A1515012248).

REFERENCES

- [1] R.A. Adams, J.F. Fournier, Sobolev Spaces (2nd Ed.), Academic Press, New York, 2003.
- [2] A.M. Al-Mahdi, M.M. Al-Gharabli, New general decay results in an infinite memory viscoelastic problem with nonlinear damping, *Bound. Value Probl.*, 2019 (2019) 140.
- [3] K. Bartkowski, P. Górká, One dimensional Klein–Gordon equation with logarithmic nonlinearities, *J. Phys. A* 41 (35) (2008) 355–201.
- [4] F. Belhannache, M.M. Algharabli, S.A. Messaoud, Asymptotic stability for a viscoelastic equation with nonlinear damping and very general type of relaxation functions, *J. Dyn. Control Syst.*, 26 (1) (2020) 45–67.
- [5] I. Bialynicki-Birula, J. Mycielski, Wave equations with logarithmic nonlinearities, *Bull. Acad. Pol. Sci. Cl 23* (4) (1975) 461–466.
- [6] I. Bialynicki-Birula, J. Mycielski, Nonlinear wave mechanics, *Ann. Physics* 100 (12) (1976) 62–93.
- [7] T. Cazenave, A. Haraux, Equations d'évolution avec nonlinéarité logarithmique, 2 (1) (1980) 21–51.
- [8] H. Di, Y. Shang, Z. Song, Initial boundary value problem for a class of strongly damped semilinear wave equations with logarithmic nonlinearity, *Nonlinear Anal. Real World Appl.* 51 (2020) 102968.
- [9] X. Fan, D. Zhao, On the spaces $L^{p(x)}(\Omega)$ and $W^{k,p(x)}(\Omega)$, *J. Math. Anal. Appl.* 263 (2) (2001) 424–446.
- [10] X.-L. Fan, Q.-H. Zhang, Existence of solutions for $p(x)$ -Laplacian Dirichlet problem, *Nonlinear Anal.* 52 (8) (2003) 1843–1852.
- [11] P. Górká, Logarithmic Klein–Gordon equation, *Acta Phys. Polon. B* 40 (2009) 59–66.
- [12] T.G. Ha, S.-H. Park, Blow-up phenomena for a viscoelastic wave equation with strong damping and logarithmic nonlinearity, *Adv. Differential Equations* 2020 (2020) 235.
- [13] N. Irikil, E. Pişkin, P. Agarwal, Global existence and decay of solutions for a system of viscoelastic wave equations of Kirchhoff type with logarithmic nonlinearity, *Math. Methods Appl. Sci.* 45 (5) (2022) 2921–3948.

- [14] K.-P. Jin, L. Jin, T.-J. Xiao, Stability of initial-boundary value problem for quasilinear viscoelastic equations, *Electron. J. Differential Equations* 2020 (80) (2020) 1–15.
- [15] D. Lars, P. Harjulehto, P. Hästö, M. Ružička, *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer, Berlin 2011.
- [16] F. Li, Q. Gao, Blow-up of solution for a nonlinear Petrovsky type equation with memory, *Appl. Math. Comput.* 274 (2016) 383–392.
- [17] M. Liao, B. Guo, X. Zhu, Energy decay rates of solutions to a viscoelastic wave equation with variable exponents and weak damping, arXiv:2011.11185v1 (2020).
- [18] M. Kafini, S. Messaoudi, Local existence and blow up of solutions to a logarithmic nonlinear wave equation with delay, *Appl. Anal.* 99 (3) (2020) 530–547.
- [19] P. Martinez, A new method to obtain decay rate estimates for dissipative systems, *ESAIM Control Optim. Calc. Var.* 4 (1999) 419–444.
- [20] M.I. Mustafa, Optimal decay rates for the viscoelastic wave equation, *Math. Methods Appl. Sci.* 41 (1) (2018) 192–204.
- [21] S.A. Messaoudi, W. Al-Khulaifi, General and optimal decay for a quasilinear viscoelastic equation, *Appl. Math. Lett.* 66 (2017) 16–22.
- [22] S.-H. Park, J.-R. Kang, Blow-up of solutions for a viscoelastic wave equation with variable exponents, *Math. Methods Appl. Sci.* 42 (6) (2019) 2083–2097.
- [23] Q. Peng, Y. Liu, Energy decay of viscoelastic equations with nonlinear damping and polynomial nonlinearity, *Appl. Anal.* 104 (18) (2025) 3497–3518.
- [24] Q. Peng, Y. Liu, Exponential stability for an infinite memory wave equation with frictional damping and logarithmic nonlinear terms, *Nonlinear Anal. Real World Appl.* 88 (2026) 104470.
- [25] Q. Peng, Z. Zhang, Stabilization and blow-up in an infinite memory wave equation with logarithmic nonlinearity and acoustic boundary conditions, *J. Syst. Sci. Complex.* 37 (4) (2024) 1368–1391.
- [26] Q. Peng, Z. Zhang, Stabilization and blow-up for a class of weakly damped Kirchhoff plate equation with logarithmic nonlinearity, *Indian J. Pure Appl. Math.* 56 (2025) 711–727.
- [27] Y. Ueda, R. Duan, S. Kawashima, Decay structure for symmetric hyperbolic systems with non-symmetric relaxation and its applications, *Arch. Rational Mech. Anal.* 205 (2012) 239–266.