

Iterating PP-packages without Choice: A Cohen symmetric seed and a localization framework

Frank Trevor Gilson

January 7, 2026

Abstract

The Partition Principle PP asserts that whenever there is a surjection $A \twoheadrightarrow B$, there is an injection $B \hookrightarrow A$. Russell conjectured in 1906 that PP is equivalent to the Axiom of Choice AC; while $AC \Rightarrow PP$ is immediate, the converse has remained open. We show that PP does not imply AC by constructing a transitive model of $ZF + DC + PP + \neg AC$.

Starting from a Cohen symmetric model \mathcal{N} of $\text{Add}(\omega, \omega_1)$ built with a countable-support symmetry filter, we fix parameters $S := A^\omega$ and $T := \mathcal{P}(S)$ and perform a class-length countable-support symmetric iteration. At successor stages we use *orbit-symmetrized packages* that split targeted surjections, yielding $PP \upharpoonright T$ and AC_{WO} , while preserving DC and ensuring that A remains non-well-orderable. A diagonal-cancellation/diagonal-lift infrastructure supplies a proper ω_1 -complete normal filter at limit stages. Finally, Ryan-Smith localization shows that under $SVC^+(T)$, PP is equivalent to $PP \upharpoonright T \wedge AC_{WO}$, so the final model satisfies PP but not AC.

Contents

1	Introduction	2
2	Preliminaries and black-box iteration technology	4
3	The Cohen symmetric seed model	5
3.1	The forcing, group action, and the countable-support filter	6
3.2	CCC and dependent choice	7
3.3	Countable supports for HS-names	10
3.4	The canonical Cohen set A and failure of choice	12

3.5	A safe SVC seed in \mathcal{N}	14
4	Reduction blueprint: global PP from localized PP	14
4.1	The localized PP-package forcing \mathbb{Q}_f	16
4.2	The AC _{WO} -package forcing \mathbb{R}_f	19
5	Iterating the packages over \mathcal{N}	21
5.1	Package symmetric systems at a stage	23
5.2	Iteration-level diagonal automorphisms	31
5.3	The modified limit filter	43
5.4	Bookkeeping: names versus interpretations	46
5.5	The symmetric iteration (black box)	48
5.6	What the iteration yields	49

1 Introduction

Whether the Partition Principle implies the Axiom of Choice is sometimes called “the oldest open problem in set theory.” The Partition Principle (PP) asserts that whenever A surjects onto B , one has $|B| \leq |A|$. Russell conjectured in 1906 that PP is equivalent to AC, but no proof was given [1]. The converse is immediate: choice provides sections for all surjections. The question of whether PP and AC are equivalent over ZF has remained open.

Main Result. We resolve this question in the negative by constructing a model of

$$\text{ZF} + \text{DC} + \text{PP} + \neg\text{AC},$$

thereby establishing that the Partition Principle does not imply the Axiom of Choice.

Theorem. *The theory $\text{ZF} + \text{DC} + \text{PP} + \neg\text{AC}$ is consistent relative to ZF.*

Strategy. The construction proceeds in two stages.

Stage 1: Seed model. We build a Cohen symmetric model \mathcal{N} over $\text{Add}(\omega, \omega_1)$ using countable supports, producing a model of $\text{ZF} + \text{DC}$ containing a non-well-orderable set $A = \{c_\alpha : \alpha < \omega_1\}$ of Cohen reals. The fixed set S witnesses $\text{SVC}(S)$, which will drive the localization argument.

Stage 2: Symmetric iteration. We perform a class-length countable-support symmetric iteration over \mathcal{N} . At each successor stage, we force sections for surjections $f : Y \rightarrow X$ with $X, Y \subseteq T := \mathcal{P}(S)$, as well as right

inverses witnessing AC_{WO} . The iteration preserves DC and the non-well-orderability of A .

Localization. The global Partition Principle follows from Ryan–Smith’s localization theorem [13], which shows that under $\text{SVC}^+(T)$, the principle PP is equivalent to the conjunction of $\text{PP} \upharpoonright T$ (the restriction to surjections between subsets of T) and AC_{WO} .

Technical innovations. Two devices are essential to making the iteration work.

(1) Orbit-symmetrized packages. A naïve attempt to force a section for a single surjection f fails because the action of $\text{Sym}(\omega_1)$ on conditions does not extend to an action on the forcing poset (automorphisms would need to map \mathbb{Q}_f to $\mathbb{Q}_{\pi \cdot f}$, a *different* poset). We resolve this by forcing with the *full orbit package*

$$\mathbb{Q}_{[f]} := \prod_{g \in [f]}^{\text{cs}} \mathbb{Q}_g,$$

the countable-support product over all g in the $\text{Sym}(\omega_1)$ -orbit of f . This ensures that every $\pi \in \text{Sym}(\omega_1)$ acts by permuting coordinates and transporting structure, yielding a genuine automorphism of $\mathbb{Q}_{[f]}$.

(2) Diagonal-cancellation symmetries. Orbit-symmetrization alone does not make the generic section s_f hereditarily symmetric: the naïve action of $\pi \in \text{Fix}(E)$ on conditions still moves the f -coordinate generic. We introduce *diagonal-cancellation automorphisms* $\hat{\pi}^D$ that act as the identity on a protected set D of coordinates (including f) while acting as π elsewhere. The stage filter includes subgroups $\Delta^\uparrow(E, \{f\})$ generated by these diagonal lifts, ensuring that the generic section is stabilized by a filter subgroup and hence belongs to the symmetric model. Crucially, diagonal lifts still act as π on the Cohen base (stage 0), so they swap Cohen reals $c_\beta \leftrightarrow c_\gamma$ for $\beta, \gamma \notin E$, preserving the $\neg\text{AC}$ witness.

Conventions. Throughout, DC denotes DC_ω , i.e., dependent choice for ω -sequences. We write $\text{PP} \upharpoonright T$ for the restriction of PP to surjections $f : Y \twoheadrightarrow X$ with both $X, Y \subseteq T$.

External tools. We rely on three external frameworks, imported as black boxes.

(1) Symmetric iteration machinery. For the countable-support symmetric iteration framework—including successor and limit constructions,

filter generation, ω_1 -completeness at limits, and the hereditarily-symmetric calculus—we cite Gilson [3]. See Remark 2.1 for the specific API items we invoke.

(2) DC preservation. For preservation of Dependent Choice in symmetric extensions, we follow Karagila [6] and Karagila–Schilhan [8], building on classical work of Pincus [10]. The criterion we use is: ω_1 -closed forcing with ω_1 -complete filter preserves DC.

(3) Localization of PP. We use Ryan–Smith’s local reflection framework [13], which provides the equivalence $\text{PP} \iff \text{PP} \upharpoonright T + \text{AC}_{\text{WO}}$ under $\text{SVC}^+(T)$. For background on SVC and Hartogs–Lindenbaum spectra, see [12, 2].

For general background on forcing and symmetric extensions we follow Jech [4] and Kunen [9].

Outline. Section 3 constructs the seed model \mathcal{N} and verifies DC, $\neg\text{AC}$, and $\text{SVC}(S)$. Section 4 recalls the Ryan–Smith localization theorem and defines the target principles $\text{PP} \upharpoonright T$ and AC_{WO} . Subsection 4.1 and Subsection 4.2 define the package forcings \mathbb{Q}_f and \mathbb{R}_f for forcing sections and right inverses. Section 5 constructs the symmetric iteration, including the orbit-symmetrization and diagonal-cancellation machinery, and assembles the final model $\mathcal{M} \models \text{ZF} + \text{DC} + \text{PP} + \neg\text{AC}$.

2 Preliminaries and black-box iteration technology

We use the general symmetric-iteration machinery from [3] (and ultimately from Karagila’s iteration framework [5]). Here we only record the “API” we will use later. We treat the following as black boxes; the precise hypotheses and proofs are available in [3].

Remark 2.1 (Iteration API to be used later). We will use, as needed, the following results from [3].

- (a) Set-length and class-length versions of symmetric iterations (cf. [3, §2.1]).
- (b) Definitions and basic properties of countable-support symmetric iterations (cf. [3, Def. 3.1]).
- (c) A DC-preservation theorem for the relevant support (cf. [3, Thm. 3.20]).
- (d) Generation and completeness/normality of limit filters in the relevant cofinality cases (cf. [3, Thm. 5.5]).

- (e) A characterization of HS names by “small support” *assuming the stage model satisfies DC* (cf. [3, Lem. 3.14]).

In our application, every stage model satisfies DC (Proposition 5.65), so the support characterization may be used freely throughout the iteration.

We emphasize that at each stage λ the iteration produces a single ambient group \mathcal{G}_λ acting by automorphisms on the full initial-segment forcing \mathbb{P}_λ (API item (b)), so the action continues to move the original Cohen coordinates; we are not iterating by repeatedly forcing over intermediate symmetric models in isolation.

Our main tool for reducing the global Partition Principle to a local instance is the "Small Violations of Choice" framework.

Definition 2.2 (Small Violations of Choice, [13]). Let S be a set.

1. We say that $\text{SVC}(S)$ holds if for every set X , there exists an ordinal η such that there is a surjection $f: S \times \eta \twoheadrightarrow X$ (equivalently, $|X| \leq^* |S \times \eta|$).
2. We say that $\text{SVC}^+(S)$ holds if for every set X , there exists an ordinal η such that there is an injection $f: X \hookrightarrow S \times \eta$ (equivalently, $|X| \leq |S \times \eta|$).

The following facts from [13] drive our localization strategy:

Fact 2.3. For any set S , $\text{SVC}(S)$ implies $\text{SVC}^+(\mathcal{P}(S))$.

Theorem 2.4 (Ryan–Smith Localization, [13]). *Assume $\text{SVC}^+(T)$. Then PP is equivalent to the conjunction of $\text{PP} \upharpoonright T$ and AC_{WO} .*

Our strategy is to construct a model satisfying $\text{SVC}(S)$ (via a Cohen seed) and then force $\text{PP} \upharpoonright \mathcal{P}(S)$ and AC_{WO} while preserving $\text{SVC}(S)$.

3 The Cohen symmetric seed model

We fix a ground model $V \models \text{ZFC}$. Let

$$\mathbb{P} = \text{Add}(\omega, \omega_1) := \text{Fn}(\omega_1 \times \omega, 2, <\omega),$$

$$\dot{c}_\alpha := \{\langle \check{n}, p \rangle : p(\alpha, n) = 1\}, c_\alpha = \dot{c}_\alpha^G$$

the standard forcing to add ω_1 -many Cohen reals, presented as finite partial functions $p : \omega_1 \times \omega \rightarrow 2$.

Lemma 3.1 (Cohen reals are pairwise distinct). *For $\alpha \neq \beta < \omega_1$, the Cohen reals c_α and c_β are distinct.*

Proof. Fix $\alpha \neq \beta < \omega_1$. Let

$$D_{\alpha,\beta} = \{p \in \text{Add}(\omega, \omega_1) : \exists n (p(\alpha, n) \neq p(\beta, n))\}.$$

Given any p , choose n such that neither (α, n) nor (β, n) is in $\text{dom}(p)$, and extend p to $q \leq p$ with $q(\alpha, n) = 0$ and $q(\beta, n) = 1$. Then $q \in D_{\alpha,\beta}$. Hence $D_{\alpha,\beta}$ is dense, so any generic G meets it and therefore $c_\alpha \neq c_\beta$. \square

3.1 The forcing, group action, and the countable-support filter

Definition 3.2 (Automorphisms of $\text{Add}(\omega, \omega_1)$). Let $\mathcal{G} := \text{Sym}(\omega_1)$ be the full permutation group of ω_1 . Each $\pi \in \mathcal{G}$ acts on \mathbb{P} by

$$(\pi p)(\pi(\alpha), n) = p(\alpha, n),$$

equivalently $(\pi p) = \{(\pi(\alpha), n, i) : (\alpha, n, i) \in p\}$. This extends (recursively) to an action on \mathbb{P} -names in the usual way.

Definition 3.3 (Countable-support filter of subgroups). For $E \subseteq \omega_1$, write

$$\text{Fix}(E) := \{\pi \in \mathcal{G} : \pi \upharpoonright E = \text{id}_E\}.$$

Let \mathcal{F} be the normal filter of subgroups of \mathcal{G} generated by the family $\{\text{Fix}(E) : E \in [\omega_1]^{\leq \omega}\}$.

A (normal) filter of subgroups is upward closed and closed under finite intersections (and conjugation).

Lemma 3.4 (Basis for the Cohen filter). *If $H \in \mathcal{F}$, then there is $E \in [\omega_1]^{\leq \omega}$ such that $\text{Fix}(E) \leq H$.*

Proof. Unwinding “ \mathcal{F} is the normal filter generated by $\{\text{Fix}(E)\}$ ”, there are $E_0, \dots, E_{k-1} \in [\omega_1]^{\leq \omega}$ and $\pi_0, \dots, \pi_{k-1} \in \mathcal{G}$ such that

$$\bigcap_{i < k} \pi_i \text{Fix}(E_i) \pi_i^{-1} \leq H.$$

But $\pi_i \text{Fix}(E_i) \pi_i^{-1} = \text{Fix}(\pi_i[E_i])$, and finite intersections of $\text{Fix}(\cdot)$'s are $\text{Fix}(\bigcup \cdot)$. Let $E := \bigcup_{i < k} \pi_i[E_i]$ (finite union of countable sets is countable). Then $\text{Fix}(E) \leq H$. \square

Corollary 3.5 (ω_1 -completeness of \mathcal{F}). \mathcal{F} is ω_1 -complete.

Proof. By Lemma 3.4 (in V) choose $E_n \in [\omega_1]^{\leq \omega}$ with $\text{Fix}(E_n) \leq H_n$. Let $E := \bigcup_{n < \omega} E_n$, which is countable in V since $V \models \text{ZFC}$. Then $\text{Fix}(E) = \bigcap_{n < \omega} \text{Fix}(E_n) \leq \bigcap_{n < \omega} H_n$, hence $\bigcap_{n < \omega} H_n \in \mathcal{F}$. \square

Definition 3.6 (Symmetric and hereditarily symmetric names). For a \mathbb{P} -name \dot{x} , define its stabilizer

$$\text{Stab}(\dot{x}) := \{\pi \in \mathcal{G} : \pi \dot{x} = \dot{x}\}.$$

We call \dot{x} *symmetric* if $\text{Stab}(\dot{x}) \in \mathcal{F}$. Let HS be the class of *hereditarily symmetric* names: $\dot{x} \in \text{HS}$ iff \dot{x} is symmetric and every name appearing in \dot{x} is in HS .

Fix $G \subseteq \mathbb{P}$ V -generic. The associated symmetric extension is

$$\mathcal{N} := \text{HS}^G = \{\dot{x}^G : \dot{x} \in \text{HS}\}.$$

3.2 CCC and dependent choice

Lemma 3.7 (CCC). $\text{Add}(\omega, \omega_1)$ is ccc. In particular, it preserves all cardinals and cofinalities.

Proof. Let $\{p_\xi : \xi < \omega_1\} \subseteq \text{Add}(\omega, \omega_1)$ be uncountable. Write $D_\xi = \text{dom}(p_\xi) \subseteq \omega \times \omega_1$, so each D_ξ is finite.

By thinning out, assume $|D_\xi| = n$ for all ξ . By the Δ -system lemma (applied to the finite sets D_ξ), thin out again to an uncountable set $I \subseteq \omega_1$ such that $\{D_\xi : \xi \in I\}$ forms a Δ -system with root R .

There are only finitely many possible values for $p_\xi \upharpoonright R$. Thin out once more so that $p_\xi \upharpoonright R = p_\eta \upharpoonright R$ for all $\xi, \eta \in I$.

Now for $\xi \neq \eta$ in I , the conditions p_ξ and p_η agree on $D_\xi \cap D_\eta = R$, hence $p_\xi \cup p_\eta$ is a condition extending both. Thus p_ξ and p_η are compatible. Therefore there is no uncountable antichain, i.e. $\text{Add}(\omega, \omega_1)$ is ccc. \square

Lemma 3.8 (ZF for symmetric extensions). Let $\langle \mathbb{P}, \mathcal{G}, \mathcal{F} \rangle$ be a symmetric system, i.e. $\mathcal{G} \leq \text{Aut}(\mathbb{P})$ and \mathcal{F} is a normal filter of subgroups of \mathcal{G} . If $G \subseteq \mathbb{P}$ is V -generic, then $\text{HS}^G = \{\dot{x}^G : \dot{x} \in \text{HS}\}$ is a transitive model of ZF and $V \subseteq \text{HS}^G \subseteq V[G]$.

Proof. This is the standard symmetric extension theorem; see, e.g., Jech chapter 15, Lemma 15.51[4]. \square

Lemma 3.9 (HS witnesses from HS-sets). *Let $\dot{A} \in \text{HS}$ and let $\varphi(v, \vec{b})$ be any formula with parameters $\vec{b} \in \text{HS}$. If $p \Vdash \exists v \in \dot{A} \varphi(v, \vec{b})$, then there exist $q \leq p$ and $\dot{a} \in \text{HS}$ such that $q \Vdash \dot{a} \in \dot{A} \wedge \varphi(\dot{a}, \vec{b})$.*

Proof. Fix $s \leq p$. Since $p \Vdash \exists v \in \dot{A} \varphi(v, \vec{b})$, there are $t \leq s$ and a name \dot{v} with $t \Vdash \dot{v} \in \dot{A} \wedge \varphi(\dot{v}, \vec{b})$. From $t \Vdash \dot{v} \in \dot{A}$, by the definition of forcing for membership there is $u \leq t$ and $\langle \dot{a}, r \rangle \in \dot{A}$ such that $u \leq r$ and $u \Vdash \dot{v} = \dot{a}$. Then $u \Vdash \varphi(\dot{a}, \vec{b})$ as well, and since $\dot{A} \in \text{HS}$ every name appearing in \dot{A} (in particular \dot{a}) is hereditarily symmetric. Thus u witnesses the conclusion. \square

Remark 3.10 (Metatheoretic countability vs. internal countability). Throughout the construction of HS names, “countable” means *countable in the ground model V* (e.g. $E \in [\omega_1]^{\leq\omega}$ is evaluated in V). This notion is used only to ensure that $\text{Fix}(E) \in \mathcal{F}$ (Definition 3.3), hence that the relevant names are symmetric. In particular, when we form supports such as $E = \bigcup_{n < \omega} E_n$ in the *metatheory*, we use that $V \models \text{ZFC}$, so a countable union of countable sets (in V) is countable (in V). We do *not* claim that these supports are countable in the symmetric model \mathcal{N} .

Theorem 3.11 (DC in the symmetric model). $\mathcal{N} \models \text{ZF} + \text{DC}$.

Proof. We verify ZF and DC separately.

ZF. By Lemma 3.8, the symmetric extension $\mathcal{N} = \text{HS}^G$ is a transitive model of ZF (containing all ordinals, as usual for symmetric extensions). We record the following symmetry lemma for the action of \mathcal{G} on \mathbb{P} for later use.

Claim 1. (Symmetry lemma) For every formula φ in the forcing language, every \mathbb{P} -condition p , every $\pi \in \mathcal{G}$, and every tuple of \mathbb{P} -names \vec{x} ,

$$p \Vdash \varphi(\vec{x}) \iff \pi p \Vdash \varphi(\pi \vec{x}).$$

Proof. By induction on the complexity of φ . The atomic cases reduce to the facts that π is an automorphism of \mathbb{P} and that the action on names is defined recursively so as to commute with membership and equality. Boolean connectives and quantifiers are then handled by the induction hypothesis in the usual way. \square

DC. Let $A, R \in \mathcal{N}$ with $A \neq \emptyset$ and $\forall x \in A \exists y \in A (xRy)$. Fix $\dot{A}, \dot{R} \in \text{HS}$ with $\dot{A}^G = A$ and $\dot{R}^G = R$. Choose $p_* \in \mathbb{P}$ such that

$$p_* \Vdash \dot{A} \neq \emptyset \wedge \forall x \in \dot{A} \exists y \in \dot{A} ((x, y) \in \dot{R}).$$

We recursively build, for each $n < \omega$,

- a countable maximal antichain $D_n \subseteq \mathbb{P}$ below p_* , and
- for each $p \in D_n$ a name $\dot{a}_{n,p} \in \text{HS}$,

so that:

- (1) $p \Vdash \dot{a}_{n,p} \in \dot{A}$ for all $p \in D_n$;
- (2) for $n \geq 1$, if $p \in D_n$ and $p \leq q \in D_{n-1}$, then $p \Vdash (\dot{a}_{n-1,q}, \dot{a}_{n,p}) \in \dot{R}$.

For $n = 0$: since $p_* \Vdash \dot{A} \neq \emptyset$, there is a maximal antichain D_0 below p_* such that for each $p \in D_0$ there is a name $\dot{a}_{0,p} \in \text{HS}$ with $p \Vdash \dot{a}_{0,p} \in \dot{A}$. (Existence of such conditions is ensured by Lemma 3.9, applied to $p_* \Vdash \dot{A} \neq \emptyset$.) Because \mathbb{P} is ccc (Lemma 3.7), we may assume D_0 is countable.

Inductively: given D_n and $\dot{a}_{n,p}$ ($p \in D_n$), fix $p \in D_n$. From $p_* \Vdash$ “ \dot{R} is total on \dot{A} ”, and $p \Vdash \dot{a}_{n,p} \in \dot{A}$, we have $p \Vdash \exists y \in \dot{A} ((\dot{a}_{n,p}, y) \in \dot{R})$. Choose a maximal antichain $D_{n+1,p}$ below p and, for each $q \in D_{n+1,p}$, choose $\dot{a}_{n+1,q} \in \text{HS}$ such that

$$q \Vdash \dot{a}_{n+1,q} \in \dot{A} \wedge (\dot{a}_{n,p}, \dot{a}_{n+1,q}) \in \dot{R}.$$

(This is justified by Lemma 3.9, applied below p to the formula $v \in \dot{A} \wedge (\dot{a}_{n,p}, v) \in \dot{R}$.)

Let $D_{n+1} := \bigcup_{p \in D_n} D_{n+1,p}$. Since D_n is an antichain, if $q \in D_{n+1,p}$ and $q' \in D_{n+1,p'}$ with $p \neq p'$, then $q \perp q'$ (otherwise p and p' would be compatible). Hence D_{n+1} is an antichain. Moreover, working in the metatheory $V \models \text{ZFC}$ (Remark 3.10), D_{n+1} is countable because it is a countable union (indexed by the countable set D_n) of countable sets $D_{n+1,p}$. Finally, D_{n+1} is a maximal antichain below p_* : given any $r \leq p_*$, pick $p \in D_n$ compatible with r (maximality of D_n), let $r' \leq r, p$, and then pick $q \in D_{n+1,p}$ compatible with r' (maximality of $D_{n+1,p}$).

Now define, for each $n < \omega$, the mixed name

$$\dot{a}_n := \{\langle \dot{a}_{n,p}, p \rangle : p \in D_n\}.$$

Then $p_* \Vdash \dot{a}_n \in \dot{A}$ for all n , and $p_* \Vdash (\dot{a}_n, \dot{a}_{n+1}) \in \dot{R}$ for all n , by the coherence of the antichains.

It remains to see that each \dot{a}_n is hereditarily symmetric and that the sequence name is hereditarily symmetric. Since each $\dot{a}_{n,p} \in \text{HS}$, its stabilizer lies in \mathcal{F} ; by definition of \mathcal{F} , for each such name choose a countable set

$E(\dot{a}_{n,p}) \subseteq \omega_1$ with $\text{Fix}(E(\dot{a}_{n,p})) \leq \text{Stab}(\dot{a}_{n,p})$. Also, for each $p \in D_n$, let $\text{supp}(p) := \{\alpha : \exists m (\alpha, m) \in \text{dom}(p)\}$, which is finite. Set

$$E_n := E(\dot{A}) \cup E(\dot{R}) \cup \bigcup_{p \in D_n} (\text{supp}(p) \cup E(\dot{a}_{n,p})),$$

a countable union of V -countable subsets of ω_1 , hence V -countable (Remark 3.10). Therefore $\text{Fix}(E_n) \in \mathcal{F}$ and $\text{Fix}(E_n) \leq \text{Stab}(\dot{a}_n)$. Then every $\pi \in \text{Fix}(E_n)$ fixes each $p \in D_n$, fixes \dot{A} and \dot{R} , and fixes each $\dot{a}_{n,p}$. Therefore $\pi \dot{a}_n = \dot{a}_n$, i.e. $\text{Fix}(E_n) \leq \text{Stab}(\dot{a}_n)$, so \dot{a}_n is symmetric; hereditariness follows since the names appearing in \dot{a}_n are among the $\dot{a}_{n,p} \in \text{HS}$. Thus $\dot{a}_n \in \text{HS}$ for all n .

Let $E := \bigcup_{n < \omega} E_n$. Then E is V -countable (Remark 3.10), hence $\text{Fix}(E) \in \mathcal{F}$, and define the sequence name

$$\dot{s} := \{\langle \check{n}, \dot{a}_n \rangle, \mathbb{1}_{\mathbb{P}} : n < \omega\}.$$

Then $\text{Fix}(E) \leq \bigcap_{n < \omega} \text{Stab}(\dot{a}_n)$, so $\text{Fix}(E) \leq \text{Stab}(\dot{s})$ since \dot{s} is built functorially from the tuple $\langle \dot{a}_n : n < \omega \rangle$, and hence $\dot{s} \in \text{HS}$. Moreover $p_* \Vdash \text{"}\dot{s}\text{ is an } \omega\text{-sequence in } \dot{A}\text{ with consecutive } \dot{R}\text{-steps"}$, so in \mathcal{N} the valuation $s = \dot{s}^G$ witnesses DC for (A, R) .

Therefore $\mathcal{N} \models \text{DC}$, and we have shown $\mathcal{N} \models \text{ZF} + \text{DC}$. \square

3.3 Countable supports for HS-names

Notation 3.12 (Restrictions to E). For $E \subseteq \omega_1$ and $p \in \text{Add}(\omega, \omega_1)$, let

$$p \upharpoonright E := p \cap (E \times \omega \times 2),$$

be viewed as a condition in $\mathbb{P}_E := \text{Add}(\omega, E)$.

For a \mathbb{P} -name τ , define recursively the restricted name $\tau \upharpoonright E$ by

$$\tau \upharpoonright E := \{\langle \sigma \upharpoonright E, p \upharpoonright E \rangle : \langle \sigma, p \rangle \in \tau\}.$$

Then $\tau \upharpoonright E$ is a \mathbb{P}_E -name (via the inclusion $\mathbb{P}_E \hookrightarrow \mathbb{P}$).

Definition 3.13 (Support). If $\dot{x} \in \text{HS}$, we say that $E \subseteq \omega_1$ is a *support* for \dot{x} if $\text{Fix}(E) \leq \text{Stab}(\dot{x})$.

Lemma 3.14 (Every HS-name has a countable support). *For every $\dot{x} \in \text{HS}$ there exists $E \in [\omega_1]^{\leq \omega}$ such that $\text{Fix}(E) \leq \text{Stab}(\dot{x})$.*

Proof. If $\dot{x} \in \text{HS}$, then in particular \dot{x} is \mathcal{F} -symmetric, so $\text{Stab}(\dot{x}) \in \mathcal{F}$. Since $\text{Stab}(\dot{x}) \in \mathcal{F}$, Lemma 3.4 yields a countable E with $\text{Fix}(E) \leq \text{Stab}(\dot{x})$. This is exactly the assertion that \dot{x} has countable support. \square

Lemma 3.15 (\mathbb{P}_E -names are hereditarily symmetric). *Let $E \in [\omega_1]^{\leq\omega}$ and $\mathbb{P}_E = \text{Add}(\omega, E) \leq \mathbb{P} = \text{Add}(\omega, \omega_1)$. If τ is a \mathbb{P}_E -name, view τ as a \mathbb{P} -name via the inclusion $\mathbb{P}_E \hookrightarrow \mathbb{P}$ (i.e. the same set-theoretic name, whose conditions all lie in $\mathbb{P}_E \subseteq \mathbb{P}$). Then $\tau \in \text{HS}$, and E is a support for τ (i.e. $\text{Fix}(E) \leq \text{Stab}(\tau)$).*

Proof. Fix $\pi \in \text{Fix}(E)$. Since π fixes E pointwise, it acts trivially on \mathbb{P}_E : for every $p \in \mathbb{P}_E$ we have $\pi p = p$. We show by induction on the rank of τ that $\pi\tau = \tau$. Indeed, if $\langle \sigma, p \rangle \in \tau$, then $p \in \mathbb{P}_E$ so $\pi p = p$, and by induction $\pi\sigma = \sigma$, hence $\langle \pi\sigma, \pi p \rangle = \langle \sigma, p \rangle \in \tau$; similarly every member of $\pi\tau$ lies in τ . Thus $\pi\tau = \tau$.

Therefore $\text{Fix}(E) \leq \text{Stab}(\tau)$. Since E is countable, $\text{Fix}(E) \in \mathcal{F}$, so τ is symmetric. The same argument applies to every subname of τ (all of whose conditions also lie in \mathbb{P}_E), so τ is hereditarily symmetric, i.e. $\tau \in \text{HS}$. \square

Lemma 3.16 (Support localization for $\text{Add}(\omega, \omega_1)$). *Let $E \subseteq \omega_1$ and let τ be a \mathbb{P} -name such that $\text{Fix}(E) \leq \text{Stab}(\tau)$. Let φ be any formula whose parameters are \mathbb{P} -names also fixed by $\text{Fix}(E)$. If $p \Vdash \varphi(\tau)$, then $p \upharpoonright E \Vdash \varphi(\tau)$.*

Proof. Suppose $p \Vdash \varphi(\tau)$ but $p \upharpoonright E \not\Vdash \varphi(\tau)$. Then there is $q \leq p \upharpoonright E$ with $q \Vdash \neg\varphi(\tau)$.

Let $F_p \subseteq \omega_1 \setminus E$ be the finite set of ordinals mentioned by p outside E (i.e. those $\alpha \notin E$ with $(\alpha, n) \in \text{dom}(p)$ for some n), and define F_q similarly. Choose a permutation $\pi \in \text{Fix}(E)$ such that $\pi[F_p]$ is disjoint from F_q and such that π fixes every element of F_q . Then πp is compatible with q (they agree on E and have disjoint requirements outside E), so let $r \leq \pi p, q$.

By the symmetry lemma (Claim 1) and $\pi \in \text{Fix}(E) \leq \text{Stab}(\tau)$, we have $\pi p \Vdash \varphi(\pi\tau) = \varphi(\tau)$, hence $r \Vdash \varphi(\tau)$. But also $r \leq q$ and $q \Vdash \neg\varphi(\tau)$, so $r \Vdash \neg\varphi(\tau)$, a contradiction. \square

Corollary 3.17 (Supported names live in the corresponding subextension). *Let $E \subseteq \omega_1$ and τ a \mathbb{P} -name with $\text{Fix}(E) \leq \text{Stab}(\tau)$. Then $\mathbb{1}_\mathbb{P} \Vdash \tau = \tau \upharpoonright E$. In particular, for any V -generic $G \subseteq \mathbb{P}$ with $G_E := G \cap \mathbb{P}_E$,*

$$\tau^G = (\tau \upharpoonright E)^{G_E} \in V[G_E].$$

Proof. We prove by induction on rank that $\mathbb{1}_\mathbb{P} \Vdash \tau = \tau \upharpoonright E$. Fix a formula of the form “ $\sigma \in \tau$ ” or “ $\sigma = \tau$ ” and use Lemma 3.16 to show that whenever p decides such a statement about τ , its restriction $p \upharpoonright E$ decides it the same way. This yields that membership in τ^G is determined by G_E and the recursive definition of $\tau \upharpoonright E$, and the induction closes. \square

3.4 The canonical Cohen set A and failure of choice

For each $\alpha < \omega_1$, define the canonical \mathbb{P} -name \dot{c}_α for the α -th Cohen real by

$$\dot{c}_\alpha := \{(\check{n}, p) : p(\alpha, n) = 1\}.$$

Let $c_\alpha := \dot{c}_\alpha^G \in 2^\omega$.

Definition 3.18 (The Cohen set).

$$A := \{c_\alpha : \alpha < \omega_1\}.$$

Lemma 3.19. $A \in \mathcal{N}$.

Proof. For each $\alpha < \omega_1$, the canonical name \dot{c}_α for the α -th Cohen real has support $\{\alpha\}$, hence is hereditarily symmetric. Let

$$\dot{A} = \{(\dot{c}_\alpha, \mathbb{1}_\mathbb{P}) : \alpha < \omega_1\}.$$

Any $\pi \in \mathcal{G}$ permutes the \dot{c}_α 's, so $\pi(\dot{A}) = \dot{A}$. Thus $\text{Stab}(\dot{A}) = \mathcal{G} \in \mathcal{F}$, and \dot{A} is symmetric. Since its members are hereditarily symmetric, $\dot{A} \in \text{HS}$, hence $A = \text{val}(\dot{A}, G) \in \mathcal{N}$. Indeed \dot{c}_α is a $\mathbb{P}_{\{\alpha\}}$ -name, hence $\dot{c}_\alpha \in \text{HS}$ by Lemma 3.15. \square

Proposition 3.20 (Non-well-orderability). *In \mathcal{N} , the set A is not well-orderable. In particular, $\mathcal{N} \models \neg\text{WO}$ and hence $\neg\text{AC}$.*

Proof. We show that in \mathcal{N} there is no injection from A into any ordinal. Since AC implies that every set is well-orderable (equivalently, injects into an ordinal), this will imply that A is not well-orderable in \mathcal{N} , hence $\mathcal{N} \models \neg\text{WO}$ and therefore $\mathcal{N} \models \neg\text{AC}$.

Suppose toward a contradiction that there are an ordinal θ and $\dot{f} \in \text{HS}$ such that \dot{f}^G is an injection $A \rightarrow \theta$ in \mathcal{N} . Choose $p_0 \in G$ and θ with

$$p_0 \Vdash \text{``}\dot{f} : \dot{A} \rightarrow \check{\theta} \text{ is an injection.''}$$

By Lemma 3.14 fix a countable support $E \in [\omega_1]^{\leq\omega}$ for \dot{f} , i.e. $\text{Fix}(E) \leq \text{Stab}(\dot{f})$.

Let $\text{supp}(p) \subseteq \omega_1$ denote the finite set of ordinals α such that $(\alpha, n) \in \text{dom}(p)$ for some $n \in \omega$. Choose $\alpha \in \omega_1 \setminus (E \cup \text{supp}(p_0))$. Since p_0 forces that $\dot{f}(\dot{c}_\alpha)$ is an ordinal $< \theta$, there are $q \leq p_0$ and $\xi < \theta$ such that

$$q \Vdash \dot{f}(\dot{c}_\alpha) = \check{\xi}.$$

Now choose $\beta \in \omega_1 \setminus (E \cup \text{supp}(q))$ with $\beta \neq \alpha$, and let $\pi \in \text{Sym}(\omega_1)$ be the transposition swapping α and β and fixing every other ordinal. Then $\pi \in \text{Fix}(E)$, so $\pi \dot{f} = \dot{f}$, and also $\pi \dot{c}_\alpha = \dot{c}_\beta$.

By the standard automorphism invariance of the forcing relation,

$$q \Vdash \dot{f}(\dot{c}_\alpha) = \check{\xi} \implies \pi q \Vdash (\pi \dot{f})(\pi \dot{c}_\alpha) = \check{\xi},$$

so using $\pi \dot{f} = \dot{f}$ and $\pi \dot{c}_\alpha = \dot{c}_\beta$ we obtain

$$\pi q \Vdash \dot{f}(\dot{c}_\beta) = \check{\xi}.$$

Because q mentions α but not β , and π fixes every ordinal in $\text{supp}(q) \setminus \{\alpha\}$, the conditions q and πq are compatible; indeed, they have disjoint requirements on the α -row versus the β -row, and agree on their common domain. Let $r := q \cup \pi q$, so $r \in \mathbb{P}$ and $r \leq q, \pi q$. Then

$$r \Vdash \dot{f}(\dot{c}_\alpha) = \check{\xi} \wedge \dot{f}(\dot{c}_\beta) = \check{\xi}.$$

Finally, strengthen r to a condition $r' \leq r$ forcing $\dot{c}_\alpha \neq \dot{c}_\beta$: choose $n \in \omega$ with $(\alpha, n), (\beta, n) \notin \text{dom}(r)$ and put

$$r' := r \cup \{((\alpha, n), 0), ((\beta, n), 1)\}.$$

Then $r' \Vdash \dot{c}_\alpha(n) \neq \dot{c}_\beta(n)$, hence $r' \Vdash \dot{c}_\alpha \neq \dot{c}_\beta$. But $r' \leq r$ still forces $\dot{f}(\dot{c}_\alpha) = \dot{f}(\dot{c}_\beta)$, contradicting the injectivity of \dot{f} on \dot{A} .

This contradiction shows that no injection $A \rightarrow \theta$ exists in \mathcal{N} for any ordinal θ . Therefore A is not well-orderable in \mathcal{N} , and consequently $\mathcal{N} \models \neg\text{WO}$ and $\mathcal{N} \models \neg\text{AC}$. \square

Remark 3.21 (Dedekind-infinity). Unlike the finite-support generalized Cohen model (which produces a Dedekind-finite A ; cf. [11, §2.4]), the present countable-support filter allows many countable subsets of A : for each countable $E \subseteq \omega_1$ in V , the name $\dot{A}_E := \{(\dot{c}_\alpha, \mathbf{1}) : \alpha \in E\}$ is hereditarily symmetric, so $A_E := \{c_\alpha : \alpha \in E\} \in \mathcal{N}$.

Lemma 3.22 (Well-orders are rigid). *If (X, \prec) is a well-order and $h : X \rightarrow X$ is a bijection such that*

$$x \prec y \iff h(x) \prec h(y) \quad (x, y \in X),$$

then $h = \text{id}_X$.

Proof. Suppose σ is an order automorphism of $(X, <)$ and $\sigma \neq \text{id}_X$. Let x_0 be the $<$ -least element of X with $\sigma(x_0) \neq x_0$.

If $\sigma(x_0) < x_0$, then $\sigma(x_0) < x_0$ implies (by minimality of x_0) that $\sigma(\sigma(x_0)) = \sigma(x_0)$. But order preservation yields $\sigma(\sigma(x_0)) < \sigma(x_0)$, a contradiction.

If $\sigma(x_0) > x_0$, then apply the previous case to σ^{-1} : since σ^{-1} is also an order automorphism and $\sigma^{-1}(x_0) < x_0$ (because $x_0 < \sigma(x_0)$), we get a contradiction.

Hence $\sigma = \text{id}_X$. □

3.5 A safe SVC seed in \mathcal{N}

Proposition 3.23 (SVC(A^ω) in the Cohen symmetric model). *Work in the Cohen symmetric model \mathcal{N} , and let $S := A^\omega$. Then $\mathcal{N} \models \text{SVC}(S)$.*

Proof. Fix a nonempty $X \in \mathcal{N}$ and choose $\dot{X} \in \text{HS}$ with $X = \dot{X}^G$.

By Lemma 3.14, fix a countable support $E \subseteq \omega_1$ for \dot{X} , i.e. $\text{Fix}(E) \leq \text{Stab}(\dot{X})$. Let $\dot{X} \upharpoonright E$ be the restricted \mathbb{P}_E -name from Notation above. By Corollary 3.17,

$$X = \dot{X}^G = (\dot{X} \upharpoonright E)^{G_E} \in V[G_E],$$

where $G_E := G \cap \mathbb{P}_E$.

Since $V[G_E] \models \text{ZFC}$, there is an ordinal η and a surjection $h : \eta \twoheadrightarrow X$ in $V[G_E]$. Let \dot{h} be a \mathbb{P}_E -name for h . Every permutation $\pi \in \text{Sym}(\omega_1)$ which fixes E pointwise acts trivially on \mathbb{P}_E and hence fixes \dot{h} , so $\dot{h} \in \text{HS}$ (indeed, it has support E). Therefore $h = \dot{h}^G \in \mathcal{N}$.

Finally, since $S = A^\omega$ is nonempty in \mathcal{N} , the map $f : S \times \eta \rightarrow X$ defined by $f(s, \xi) = h(\xi)$ is a surjection in \mathcal{N} . This proves $\mathcal{N} \models \text{SVC}(S)$. □

Definition 3.24 (Fixed seed parameters: S and T). Work in \mathcal{N} and let

$$S := A^\omega, \quad T := \mathcal{P}(S).$$

4 Reduction blueprint: global PP from localized PP

We now pin down the exact hypotheses and the parameter set to which we will localize PP.

Definition 4.1 (Localized PP). Let T be a set.

1. $\text{PP} \upharpoonright T$ is the statement that for all $X, Y \subseteq T$, if there is a surjection $f : Y \twoheadrightarrow X$, then there is an injection $i : X \hookrightarrow Y$.
2. $\text{PP}(T)$ is the statement that for every set X , if there is a surjection $f : T \twoheadrightarrow X$, then there is an injection $i : X \hookrightarrow T$.

Definition 4.2 (SVC^+). For a set T , $\text{SVC}^+(T)$ denotes the statement that for every set X there exists an ordinal η and an injection $j : X \hookrightarrow T \times \eta$.

Fact 4.3 (Ryan–Smith). For every set S , $\text{SVC}(S)$ implies $\text{SVC}^+(\mathcal{P}(S))$. [13, §2.1, Fact]

Remark 4.4. Let S, T be as in Definition 3.24. By Fact 4.3 we have $\mathcal{N} \models \text{SVC}^+(T)$, and we will keep this T fixed throughout the construction.

Theorem 4.5 (Ryan–Smith localization of PP). *Assume $\text{SVC}^+(T)$. Then*

$$\text{PP} \iff (\text{PP} \upharpoonright T \wedge \text{AC}_\text{WO}).$$

[13, Proposition 3.17] (stated there with parameter “ S ”; we apply it with $S := T$)

Remark 4.6. Ryan–Smith formulate PP as the equivalence $|X| \leq |Y| \iff |X| \leq^* |Y|$. Since $|X| \leq |Y| \Rightarrow |X| \leq^* |Y|$ holds in ZF , this is equivalent to our surjection-to-injection formulation of PP .

Proposition 4.7 (The AC_WO conjunct cannot be omitted). *There is a model $M \models \text{ZF}$ and a set $S^* \in M$ such that*

$$M \models \text{SVC}^+(S^*) \wedge \text{PP}(S^*) \wedge \neg \text{PP}.$$

In particular, global PP cannot be recovered from a small-choice hypothesis together with a single-parameter instance $\text{PP}(S^)$.*

Reference. This is witnessed by Cohen’s first model; see [13, Proposition 3.22]. \square

Proposition 4.8 (The localized $\text{PP} \upharpoonright S$ component cannot be omitted). *There is a model $M \models \text{ZF}$ and a set $S^* \in M$ such that*

$$M \models \text{AC}_\text{WO} \wedge \text{SVC}^+(S^*) \wedge \text{PP}(S^*) \wedge \neg \text{PP}.$$

Consequently, even assuming AC_WO and small-choice hypotheses, one cannot in general replace the localized requirement $\text{PP} \upharpoonright S^$ by the weaker single-parameter assertion $\text{PP}(S^*)$ and still obtain global PP .*

Reference. This is witnessed by the Feferman-style model N_{\aleph_1} of Truss; see [13, Proposition 3.23]. \square

Remark 4.9 (Why we keep both conjuncts). The preceding propositions are included only to justify (conceptually) why the localization theorem is stated with both components present: the “choice” side AC_{WO} and the “localized comparability” side $\text{PP} \upharpoonright T$. We will force *both* conjuncts $\text{PP} \upharpoonright T$ and AC_{WO} in Theorem 4.5. In Ryan–Smith’s analysis the corresponding reduction is sharp: in general one cannot drop the displayed conjuncts (nor substantially weaken the hypotheses) without losing the equivalence; see [13, Props. 3.22 and 3.23].

4.1 The localized PP-package forcing \mathbb{Q}_f

Fix a transitive model $M \models \text{ZF} + \text{DC}$ extending the Cohen seed \mathcal{N} (in practice, M will be an intermediate stage of our symmetric iteration). Work with the fixed parameter $T = \mathcal{P}(S)$ from Definition 3.24.

Definition 4.10 (Localized splitting principle). Write $\text{PP}^{\text{split}} \upharpoonright T$ for the assertion that for all $X, Y \subseteq T$ and all surjections $f : Y \twoheadrightarrow X$, there exists a right inverse $s : X \rightarrow Y$ with $f \circ s = \text{id}_X$ (necessarily injective: if $s(x_1) = s(x_2)$ then $x_1 = f(s(x_1)) = f(s(x_2)) = x_2$.)

Remark 4.11 (Why we allow *localized* splittings). The global statement “every surjection splits” (i.e. every surjection has a right inverse) is equivalent to AC (a right inverse to $\bigsqcup_{i \in I} A_i \rightarrow I$ is exactly a choice function). So we are *not* attempting to force global splittings.

Moreover, $\text{PP}^{\text{split}} \upharpoonright T$ only concerns surjections whose domain and codomain lie inside the fixed parameter $T = \mathcal{P}(S)$, and is therefore far weaker than the global splitting scheme equivalent to AC .

However, for our Ryan–Smith reduction we only need to secure localized PP over the fixed parameter $T = \mathcal{P}(S)$. For that purpose it is convenient to force the *stronger* localized principle $\text{PP}^{\text{split}} \upharpoonright T$, which implies $\text{PP} \upharpoonright T$ outright. This choice buys us a canonical, highly closed package poset at each stage.

For the choice principles on well-orderable sets, Ryan–Smith prove that assuming $\text{SVC}(S)$,

$$\text{AC}_{\text{WO}} \iff \text{AC}_{\text{WO}}(S) \iff \text{every function } g : S \rightarrow \lambda \text{ splits (for each } \lambda)$$

(see [13, Prop. 3.2]). This is the equivalence we invoke in Proposition 5.71.

(We also note that Ryan–Smith’s localization for PP uses both conjuncts: $\text{PP} \upharpoonright T$ and AC_{WO} ; see [13, Prop. 3.17] and Question 3.18 therein, and see [13, Props. 3.22 and 3.23] for counterexamples showing the conjuncts cannot in general be omitted.)

Definition 4.12 (PP-package for a fixed surjection). Let $X, Y \subseteq T$ and let $f : Y \twoheadrightarrow X$ be a surjection in M . Define \mathbb{Q}_f to be the poset whose conditions are *countable partial injective sections of f* :

$$p \in \mathbb{Q}_f \iff \begin{cases} p \text{ is a function } p : \text{dom}(p) \rightarrow Y, \\ \text{dom}(p) \subseteq X \text{ is countable,} \\ p \text{ is injective,} \\ \forall x \in \text{dom}(p) \ (f(p(x)) = x). \end{cases}$$

The order is extension: $q \leq p$ iff $q \supseteq p$.

Remark 4.13 (Size and chain condition). The forcing \mathbb{Q}_f can be large and is not expected to be ccc in general (e.g. $|\mathbb{Q}_f|$ may be comparable to $|X|^{\leq\omega} \cdot |Y|^{\leq\omega}$). This is harmless for our purposes: what we use is countable closure (hence no new reals and DC preservation), not any chain condition.

Lemma 4.14 (Basic density). *For each $x \in X$, the set*

$$D_x := \{p \in \mathbb{Q}_f : x \in \text{dom}(p)\}$$

is dense in \mathbb{Q}_f .

Proof. Fix $p \in \mathbb{Q}_f$ and $x \in X$ with $x \notin \text{dom}(p)$. Choose any $y \in Y$ with $f(y) = x$ (exists since f is surjective). We claim $y \notin \text{ran}(p)$: if $y = p(x')$ for some $x' \in \text{dom}(p)$ then $x = f(y) = f(p(x')) = x'$, contradicting $x \notin \text{dom}(p)$. Thus $q := p \cup \{(x, y)\}$ is still a partial function, still injective, and still satisfies $f(q(x)) = x$; hence $q \in \mathbb{Q}_f$ and $q \leq p$ with $x \in \text{dom}(q)$. \square

Lemma 4.15 (Countable closure). *\mathbb{Q}_f is Countably-closed (sometimes called ω_1 -closed).*

Proof. Let $\langle p_n : n < \omega \rangle$ be a decreasing sequence in \mathbb{Q}_f . Put $p := \bigcup_{n < \omega} p_n$. Since each $p_{n+1} \supseteq p_n$, this union is a function. It is injective, and for every $x \in \text{dom}(p)$ we have $f(p(x)) = x$ because this holds already in some p_n containing x .

It remains to see that $\text{dom}(p) = \bigcup_{n < \omega} \text{dom}(p_n)$ is countable in M . Here we use $M \models \text{DC}$, hence $M \models \text{AC}_\omega$, so a countable union of countable sets is countable in M . (This is the only place in the closure argument where DC is used.) Therefore $p \in \mathbb{Q}_f$, and clearly $p \leq p_n$ for all n . \square

Remark 4.16 (Closure depends on DC). The ω_1 -closure of \mathbb{Q}_f is proved inside M using DC (equivalently AC_ω), via “countable union of countable sets is countable”. In a bare ZF context this need not hold, so the closure claim should be read as relative to our standing hypothesis $M \models \text{ZF} + \text{DC}$.

Proposition 4.17 (What \mathbb{Q}_f forces). *If $G \subseteq \mathbb{Q}_f$ is M -generic, then*

$$s_G := \bigcup G$$

is a total injective function $s_G : X \rightarrow Y$ with $f \circ s_G = \text{id}_X$. In particular, $M[G] \models "X \hookrightarrow Y"$.

Proof. By Lemma 4.14, G meets D_x for every $x \in X$, so $x \in \text{dom}(s_G)$ for all x and s_G is total on X . Injectivity and $f(s_G(x)) = x$ are preserved under unions of compatible conditions, hence hold for s_G . \square

Lemma 4.18 (Countably closed forcing adds no new ω -sequences and preserves DC). *Let $M \models \text{ZF} + \text{DC}$ be transitive and let $\mathbb{P} \in M$ be countably closed in M (i.e. every decreasing ω -sequence in M has a lower bound in M). Then:*

(1) \mathbb{P} adds no new ω -sequences of ordinals.

(2) Forcing with \mathbb{P} over M preserves DC.

Proof. (1) *No new ω -sequences of ordinals.* Let $p_0 \in \mathbb{P}$ and $\dot{a} \in M$ with $p_0 \Vdash \dot{a} : \check{\omega} \rightarrow \text{Ord}$. For each $n < \omega$, let

$$D_n := \left\{ p \leq p_0 : \exists \alpha \in \text{Ord} \ (p \Vdash \dot{a}(\check{n}) = \check{\alpha}) \right\}.$$

Each D_n is dense below p_0 .

Define the set of *states*

$$\Sigma := \left\{ (n, p, \vec{\alpha}) : n < \omega, p \in \mathbb{P}, p \leq p_0, \vec{\alpha} \in \text{Ord}^n, p \Vdash \forall k < n \dot{a}(\check{k}) = \check{\alpha}(\check{k}) \right\}.$$

Let R be the relation on Σ given by

$$(n, p, \vec{\alpha}) R (n+1, q, \vec{\beta})$$

iff $q \leq p$, $\vec{\beta} \in \text{Ord}^{n+1}$ extends $\vec{\alpha}$, and $q \Vdash \dot{a}(\check{n}) = \check{\alpha}(\check{n})$. Then R is total on Σ by density of D_n .

By DC in M , there is an R -chain $\langle (n, p_n, \vec{\alpha}_n) : n < \omega \rangle \in M$ starting from some $(0, p_0, \emptyset) \in \Sigma$ (note $\text{Ord}^0 = \{\emptyset\}$). By countable closure, there is $p_\omega \in \mathbb{P}$ with $p_\omega \leq p_n$ for all n .

Let $g := \bigcup_{n < \omega} \vec{\alpha}_n$. Since the chain is an element of M , Replacement in M yields $g \in M$ and $g : \omega \rightarrow \text{Ord}$. By construction, $p_\omega \Vdash \forall n \dot{a}(\check{n}) = g(\check{n})$, hence $p_\omega \Vdash \dot{a} = \check{g}$. Thus \mathbb{P} adds no new ω -sequences of ordinals.

(2) *Preservation of DC.* Let $p_0 \in \mathbb{P}$ and $\dot{X}, \dot{R}, \dot{x}_0 \in M$ with

$$p_0 \Vdash \dot{x}_0 \in \dot{X} \wedge (\forall x \in \dot{X})(\exists y \in \dot{X}) x \dot{R} y.$$

Define the state space

$$\Gamma := \{(n, p, \dot{x}) : n < \omega, p \in \mathbb{P}, p \leq p_0, \dot{x} \in M, p \Vdash \dot{x} \in \dot{X}\}.$$

Define S on Γ by

$$(n, p, \dot{x}) S (n+1, q, \dot{y})$$

iff $q \leq p$ and $q \Vdash \dot{y} \in \dot{X} \wedge \dot{x} \dot{R} \dot{y}$. Totality of S follows from the premise forced by p_0 .

By DC in M , pick an S -chain $\langle (n, p_n, \dot{x}_n) : n < \omega \rangle \in M$ starting with $(0, p_0, \dot{x}_0) \in \Gamma$. By countable closure, let $p_\omega \leq p_n$ for all n .

Let

$$\dot{f} := \{\langle \langle \check{n}, \dot{x}_n \rangle, 1_{\mathbb{P}} \rangle : n < \omega\}.$$

Since the chain is in M , Replacement gives $\dot{f} \in M$. Then $p_\omega \Vdash \dot{f} : \check{\omega} \rightarrow \dot{X}$ and $p_\omega \Vdash \forall n \dot{f}(\check{n}) \dot{R} \dot{f}(\check{n}+1)$, which is exactly DC in the extension. \square

Corollary 4.19 (No new ω -sequences; DC preservation). \mathbb{Q}_f adds no new ω -sequences of ordinals (in particular, no new reals). Moreover, over a $\text{ZF} + \text{DC}$ ground, forcing with \mathbb{Q}_f preserves DC.

Proof. Apply Lemma 4.18 with $\mathbb{P} = \mathbb{Q}_f$. Countable closure holds by Lemma 4.15. \square

4.2 The AC_{WO} -package forcing \mathbb{R}_f

Fix a transitive model $M \models \text{ZF} + \text{DC}$ extending the Cohen seed \mathcal{N} (again, in practice M will be an intermediate stage of our iteration). Retain the fixed seed parameter S from Remark 4.4.

Definition 4.20 (The Lindenbaum bound $\aleph^*(S)$). For a set S in M , let $\aleph^*(S)^M$ denote the least ordinal κ such that there is no surjection $S \twoheadrightarrow \kappa$ in M . Equivalently, $\kappa = \sup\{\alpha : \exists \text{ a surjection } S \twoheadrightarrow \alpha\} + 1$.

Remark 4.21. In ZF the ordinal $\aleph^*(S)$ exists for every set S (it is the Lindenbaum number of S); see e.g. Jech [4, §(Hartogs/Lindenbaum)].

Remark 4.22 (Why the bound $\lambda < \aleph^*(S)$ is the correct target). Assuming $\text{SVC}(S)$, Ryan-Smith shows that AC_{WO} is equivalent to the bounded scheme $\text{AC}_{<\aleph^*(S)}$ (more precisely, $\text{AC}_{<\aleph^*(S)}(S)$ in his notation); see [13, Cor. 3.3]. For fixed λ , the corresponding “splitting” reformulations of AC_λ appear in [13, Prop. 3.2]. Thus, in the presence of $\text{SVC}(S)$ it suffices to schedule splittings only for surjections onto ordinals $\lambda < \aleph^*(S)$.

Definition 4.23 (AC_{WO} -package for a fixed surjection onto an ordinal). Let λ be an ordinal with $\lambda < \aleph^*(S)^M$, let Y be a set in M , and let $f : Y \rightarrow \lambda$ be a surjection in M . Define \mathbb{R}_f to be the poset of *countable partial right inverses* to f :

$$r \in \mathbb{R}_f \iff \begin{cases} r \text{ is a function } r : \text{dom}(r) \rightarrow Y, \\ \text{dom}(r) \subseteq \lambda \text{ is countable,} \\ \forall \xi \in \text{dom}(r) \ (f(r(\xi)) = \xi). \end{cases}$$

The order is extension: $s \leq r$ iff $s \supseteq r$.

Remark 4.24 (Total sections are automatically injective). If $s : \lambda \rightarrow Y$ satisfies $f \circ s = \text{id}_\lambda$, then s is injective: if $s(\xi_1) = s(\xi_2)$ then $\xi_1 = f(s(\xi_1)) = f(s(\xi_2)) = \xi_2$.

Remark 4.25 (No localization to T for AC_{WO}). Unlike the PP -packages \mathbb{Q}_f , which are explicitly localized to $X, Y \subseteq T = \mathcal{P}(S)$, the AC_{WO} -package must be global in nature: AC_{WO} speaks about *all* well-ordered index sets. The point of Remark 4.22 is that, assuming $\text{SVC}(S)$, it is enough to schedule splittings for surjections onto $\lambda < \aleph^*(S)$; this reduction is precisely what makes the bookkeeping manageable.

Lemma 4.26 (Basic density). *For each $\xi < \lambda$, the set*

$$D_\xi := \{r \in \mathbb{R}_f : \xi \in \text{dom}(r)\}$$

is dense in \mathbb{R}_f .

Proof. Fix $r \in \mathbb{R}_f$ and $\xi < \lambda$ with $\xi \notin \text{dom}(r)$. Choose any $y \in Y$ with $f(y) = \xi$ (exists since f is surjective). Then $s := r \cup \{(\xi, y)\}$ is still a function, still satisfies $f(s(\xi)) = \xi$, and has countable domain; hence $s \in \mathbb{R}_f$ and $s \leq r$ with $\xi \in \text{dom}(s)$. \square

Lemma 4.27 (Countable closure). *Assume $M \models \text{ZF} + \text{DC}$. Then \mathbb{R}_f is countably closed.*

Proof. Let $\langle r_n : n < \omega \rangle$ be a decreasing sequence in \mathbb{R}_f . Put $r := \bigcup_{n < \omega} r_n$. This is a function, and it satisfies $f(r(\xi)) = \xi$ for every $\xi \in \text{dom}(r)$ since this already holds in some r_n containing ξ .

Finally, $\text{dom}(r) = \bigcup_{n < \omega} \text{dom}(r_n)$ is a countable union of countable sets. Since $M \models \text{DC}$ (hence $M \models \text{AC}_\omega$), $\text{dom}(r)$ is countable in M . Therefore $r \in \mathbb{R}_f$, and clearly $r \leq r_n$ for all n . \square

Remark 4.28 (Closure depends on DC). The ω_1 -closure of \mathbb{R}_f is proved inside M using DC (equivalently AC_ω), via “countable union of countable sets is countable”. In bare ZF this need not hold.

Proposition 4.29 (What \mathbb{R}_f forces). *If $G \subseteq \mathbb{R}_f$ is M -generic, then*

$$s_G := \bigcup G$$

is a total function $s_G : \lambda \rightarrow Y$ with $f \circ s_G = \text{id}_\lambda$ (hence s_G is injective). In particular, $M[G] \models \text{“}f \text{ splits.}”$

Proof. By Lemma 4.26, G meets D_ξ for every $\xi < \lambda$, so $\xi \in \text{dom}(s_G)$ for all $\xi < \lambda$ and s_G is total on λ . The equation $f(s_G(\xi)) = \xi$ is preserved under unions of compatible conditions, so $f \circ s_G = \text{id}_\lambda$. \square

Corollary 4.30 (No new reals; DC preservation (package-level)). *\mathbb{R}_f adds no new reals (indeed, no new ω -sequences of ordinals). Moreover, forcing with \mathbb{R}_f over a ZF + DC ground preserves DC.*

Proof. Apply Lemma 4.18 with $\mathbb{P} = \mathbb{R}_f$. Countable closure holds by Lemma 4.27. \square

Remark 4.31 (Package products). When we say “finite product of package posets” we mean an arbitrary finite-support product of finitely many factors (not necessarily just two), typically consisting of one AC_{WO} -package \mathbb{R}_f together with finitely many localized PP-packages $\mathbb{Q}_{g_0}, \dots, \mathbb{Q}_{g_{n-1}}$ scheduled at the same stage.

5 Iterating the packages over \mathcal{N}

Remark 5.1 (Roadmap of Section 5). This section describes a class-length countable-support *symmetric iteration* built to force local splitting principles while preserving DC and killing AC. Here is the dependency flow.

- (1) **Coordinates and products at successor stages.** At each successor stage $\alpha + 1$ the forcing factor is a *finite product* of orbit packages, and package coordinates are tracked by the triple system of Definition 5.22. This is the bookkeeping substrate for all later “fix a coordinate” arguments.
- (2) **Diagonal infrastructure.** Subsection 5.2 defines iteration-level diagonal lifts $\hat{\pi}^D$ and the associated subgroups $\Delta_\lambda^\uparrow(E, D)$, where $D \subseteq \text{Coords}_{<\lambda}$ is *countable* and (E, D) is required to be *globally admissible* (Definition 5.26). The successor “diagonal kernel” elements (Definition 5.23 and Lemma 5.25) supply the cancellation needed to define coherent lifts and to verify group membership.
- (3) **The modified limit filter.** Subsection 5.3 defines the limit-stage filter $\tilde{\mathcal{F}}_\lambda^*$ (Definition 5.48) generated by pushforwards together with *globally admissible* Δ^\uparrow -subgroups. The key structural fact is the *core lemma* (Lemma 5.50): every $H \in \tilde{\mathcal{F}}_\lambda^*$ contains some $\Delta_\lambda^\uparrow(E, D)$ with globally admissible parameters. Conjugation is handled by Lemma 5.38 together with the admissible-hull step (Lemma 5.40), so normal-filter generation never “escapes” admissibility.
- (4) **Bookkeeping by names (not interpretations).** Subsection 5.4 fixes the class bookkeeping \mathcal{B} of codes for potential \mathbb{P}_α -names (Definition 5.59). At stage α we test the *current* interpretation in M_α and add the corresponding orbit package factor(s) only when a code currently yields a relevant surjection instance not yet split.
- (5) **Black-box symmetric iteration.** Subsection 5.5 invokes the symmetric-iteration framework to assemble the class-length iteration $\langle \mathbb{P}_\alpha, \mathcal{G}_\alpha, \tilde{\mathcal{F}}_\alpha^* : \alpha \in \text{Ord} \rangle$, with $\mathbb{P}_0 = \text{Add}(\omega, \omega_1)$ and successor stages $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha * \dot{\mathbb{S}}_\alpha$ where $\dot{\mathbb{S}}_\alpha$ is the finite product of the scheduled orbit package factors.
- (6) **What the iteration yields.** Subsection 5.6 extracts the main consequences in the final model \mathcal{M} : ZF and DC are preserved along the iteration; all surjections $Y \twoheadrightarrow X$ with $X, Y \subseteq T$ split (hence $\text{PP}^{\text{split}} \upharpoonright T$); SVC(S) persists (hence $\text{SVC}^+(T)$); AC_{wo} holds via scheduled right inverses; the Ryan–Smith localization theorem upgrades $(\text{PP} \upharpoonright T \wedge \text{AC}_{\text{wo}})$ to full PP; and finally $\neg\text{AC}$ holds because the Cohen set A is not well-orderable (Proposition 5.64) using the core lemma plus a nontrivial diagonal lift acting on A .

A reader who wants only the global theorem can skim Subsections 5.3–5.6 while treating the package/diagonal machinery as the engine ensuring the required stabilizers lie in $\tilde{\mathcal{F}}_\lambda^*$.

We now describe the symmetric iteration that forces the two conjuncts $\mathbf{PP}^{\text{split}} \upharpoonright T$ and \mathbf{AC}_{WO} over the Cohen symmetric seed model \mathcal{N} , while preserving $\neg\mathbf{AC}$ (witnessed by the non-well-orderability of A). All technical iteration infrastructure is imported via the API of Remark 2.1. The purpose of this section is to isolate the *package systems*, the bookkeeping targets, and the verification that the final symmetric model satisfies the desired principles.

Remark 5.2 (Modularity of the construction). The argument is deliberately modular: the seed model technology (Section 3), the localized \mathbf{PP} -packages (Section 4.1), the \mathbf{AC}_{WO} -packages (Section 4.2), and the symmetric-iteration “API” (Remark 2.1) are separated so that if a hypothesis later needs to be strengthened or a package definition adjusted, it can typically be done locally without rewriting unrelated parts of the paper.

5.1 Package symmetric systems at a stage

Fix a transitive $M \models \mathbf{ZF} + \mathbf{DC}$ extending \mathcal{N} (in practice $M = M_\alpha$ at an intermediate stage of the iteration). Keep the parameters S and $T = \mathcal{P}(S)$ as in Remark 4.4.

Remark 5.3 (An ambient $\text{Sym}(\omega_1)$ -action on $T = \mathcal{P}(S)$). We will often use the ambient action of $\text{Sym}(\omega_1)$ on the stage-0 seed parameter $S = A^\omega$ and on $T = \mathcal{P}(S)$ by permuting the Cohen reals $c_\alpha \in A$. Concretely, for $\pi \in \text{Sym}(\omega_1)$ we set $\pi \cdot c_\alpha := c_{\pi(\alpha)}$, extend pointwise to sequences $s \in A^\omega$, and then define $\pi \cdot X := \{\pi \cdot s : s \in X\}$ for $X \subseteq S$, i.e. $\pi \cdot X \in T$. We regard this as an action which fixes all ordinals pointwise; i.e. $\pi \cdot \xi = \xi$ for every ordinal ξ .

We furthermore extend this action to every set x with $\text{tc}(x) \subseteq V(T)$ by recursion on rank:

$$\pi \cdot x := \{\pi \cdot y : y \in x\}.$$

In particular, if $f : X \rightarrow Y$ is a function with $\text{tc}(f) \subseteq V(T)$ (e.g. $X, Y \subseteq T$), we identify f with its graph and set

$$\pi \cdot f := \{(\pi \cdot a, \pi \cdot b) : (a, b) \in f\}.$$

Lemma 5.4 (Kernel acts trivially on T). *Let α be any stage and let $k \in \ker(\pi_\alpha^0)$. Then k fixes A , $S = A^\omega$, and $T = \mathcal{P}(S)$ pointwise for the ambient*

$\text{Sym}(\omega_1)$ -action of Remark 5.3. Consequently, k fixes every set/function whose transitive closure is contained in $V(T) \cup \text{Ord}$; in particular, if $f : X \rightarrow Y$ with $X, Y \subseteq T$, or $f : S \times \eta \rightarrow \lambda$ with $\eta, \lambda \in \text{Ord}$, then $k \cdot f = f$.

Proof. Put $\sigma := \pi_\alpha^0(k) \in \text{Sym}(\omega_1)$. Since $k \in \ker(\pi_\alpha^0)$ we have $\sigma = \text{id}$. By Remark 5.3, the ambient action on A , on $S = A^\omega$, and on $T = \mathcal{P}(S)$ is induced from the action $\alpha \mapsto \sigma(\alpha)$ on indices of the Cohen reals. Therefore $\sigma = \text{id}$ implies that k fixes A , and hence also S and T , pointwise.

For the final assertion, use the recursive extension of the action to $V(T)$ from Remark 5.3 and argue by induction on rank that every x with $\text{tc}(x) \subseteq V(T) \cup \text{Ord}$ is fixed pointwise (using that k fixes all ordinals pointwise, and the recursive definition of the action on $V(T)$). In particular, if $f : X \rightarrow Y$ with $X, Y \subseteq T$, then $\text{tc}(f) \subseteq V(T) \cup \text{Ord}$ (viewing f as its graph), so $k \cdot f = f$. \square

Definition 5.5 (Orbit-symmetrized package systems). Let $M \models \text{ZF} + \text{DC}$ be a transitive stage model and let $\mathcal{G} = \text{Sym}(\omega_1)$ act on T as in Remark 5.3. Fix a surjection instance f in M .

Write $[f] := \{\pi \cdot f : \pi \in \mathcal{G}\}$ for the \mathcal{G} -orbit of f (a set in M by Replacement).

(a) **Orbit PP^{split} $\upharpoonright T$ -package.** If $f : Y \rightarrow X$ with $X, Y \subseteq T$, define

$$\mathbb{Q}_{[f]} := \prod_{g \in [f]}^{\text{cs}} \mathbb{Q}_g,$$

where \mathbb{Q}_g is the section forcing from Definition 4.12. A condition $\vec{p} \in \mathbb{Q}_{[f]}$ is a function with countable support $\text{supp}(\vec{p}) := \text{dom}(\vec{p}) \subseteq [f]$ such that $\vec{p}(g) \in \mathbb{Q}_g$ for all $g \in \text{dom}(\vec{p})$, ordered by coordinatewise extension.

(b) **Orbit AC_{WO}-package (in S -presentation).** If η is an ordinal, $\lambda < \aleph^*(S)^M$, and $f : S \times \eta \rightarrow \lambda$ is a surjection in M , define

$$\mathbb{R}_{[f]} := \prod_{g \in [f]}^{\text{cs}} \mathbb{R}_g,$$

where \mathbb{R}_g is the partial right-inverse forcing from Definition 4.23.

In both cases \mathcal{G} acts on the orbit package by permuting the factors together with the *canonical* transport induced by π on the defining parameters. More precisely, for $\pi \in \mathcal{G}$ and $g \in [f]$ let

$$A_{\pi,g} : \mathbb{S}_g \rightarrow \mathbb{S}_{\pi \cdot g}$$

be the transport-of-structure isomorphism induced by π (i.e. apply π to every ordinal and every element of T appearing in a condition, using the ambient \mathcal{G} -action on T from Remark 5.3 and the fact that \mathbb{S}_g is defined functorially from g). Then the orbit action on conditions is

$$(\pi \cdot \vec{p})(g) := A_{\pi, \pi^{-1} \cdot g}(\vec{p}(\pi^{-1} \cdot g)).$$

Lemma 5.6 (Orbit packages are ω_1 -closed and preserve DC). *Assume $M \models \text{ZF} + \text{DC}$ and let $\mathbb{S}_{[f]}$ be either $\mathbb{Q}_{[f]}$ or $\mathbb{R}_{[f]}$. Then:*

- (i) $\mathbb{S}_{[f]}$ is ω_1 -closed in M ;
- (ii) $\mathbb{S}_{[f]}$ adds no new ω -sequences of ordinals (in particular, no new reals);
- (iii) forcing with $\mathbb{S}_{[f]}$ over M preserves DC.

Proof. (i) Let $\langle \vec{p}_n : n < \omega \rangle$ be decreasing in $\mathbb{S}_{[f]}$. For each $g \in [f]$ define $q_g := \bigcup_{n < \omega} p_{n,g}$, where $p_{n,g}$ is the g -coordinate of \vec{p}_n (taking $p_{n,g} = \emptyset$ if $g \notin \text{supp}(\vec{p}_n)$). Since each factor \mathbb{Q}_g is ω_1 -closed by Lemma 4.15 (and each factor \mathbb{R}_g by Lemma 4.27), q_g is a condition in that factor. Let \vec{q} have support $\text{supp}(\vec{q}) := \bigcup_{n < \omega} \text{supp}(\vec{p}_n)$ and coordinates q_g for $g \in \text{supp}(\vec{q})$. Because $M \models \text{DC}$ (hence AC_ω), a countable union of countable sets is countable, so $\text{supp}(\vec{q})$ is countable. Thus $\vec{q} \in \mathbb{S}_{[f]}$ and $\vec{q} \leq \vec{p}_n$ for all n .

(ii) and (iii) now follow by applying Lemma 4.18 to $\mathbb{P} = \mathbb{S}_{[f]}$, using (i) for countable closure. \square

Remark 5.7 (Limit-stage ω_1 -completeness). At limit stages, ω_1 -completeness (and normality) of the limit filter is provided by the iteration machinery (API item (d), cf. [3, Thm. 5.5]).

Lemma 5.8 (Well-definedness of the orbit package actions). *In each case of Definition 5.5, if \vec{p} is a condition of the orbit package $\mathbb{S}_{[f]}$ (i.e. $\mathbb{Q}_{[f]}$ or $\mathbb{R}_{[f]}$) and $\pi \in \text{Sym}(\omega_1)$, then $\pi \cdot \vec{p}$ is again a condition of the same forcing, and the action respects the order.*

Proof. By construction, each $A_{\pi,g}$ is a poset isomorphism $\mathbb{S}_g \cong \mathbb{S}_{\pi \cdot g}$, and the transport maps satisfy the cocycle law

$$A_{\pi \circ \rho, g} = A_{\pi, \rho \cdot g} \circ A_{\rho, g}$$

for all $\pi, \rho \in \text{Sym}(\omega_1)$ and $g \in [f]$. Therefore $(\pi \circ \rho) \cdot \vec{p} = \pi \cdot (\rho \cdot \vec{p})$ and $\text{id} \cdot \vec{p} = \vec{p}$, so this is a well-defined action by poset automorphisms. Since the action only permutes coordinates and applies coordinatewise isomorphisms, it preserves the ideal of supports. \square

Lemma 5.9 (Countable support of package coordinates). *For every package coordinate d at any successor stage $\alpha + 1$ of the iteration, there exists a countable set $E_d \subseteq \omega_1$ such that $\text{Fix}(E_d) \leq \text{Stab}(d)$, where $\text{Stab}(d) = \{\pi \in \text{Sym}(\omega_1) : \pi \cdot d = d\}$ is the stabilizer of d as an element of its orbit package.*

Proof. We proceed by induction on stages. At stage 0 there are no package coordinates, so the statement is vacuous.

At successor stage $\alpha + 1$, package coordinates are elements of orbit packages $[f] = \{\pi \cdot f : \pi \in \text{Sym}(\omega_1)\}$, where f is a surjection in the symmetric model M_α .

Step 1: The seed surjection f has countable support.

Since $f \in M_\alpha$, there is a hereditarily symmetric \mathbb{P}_α -name \dot{f} with $f = \dot{f}^{G_\alpha}$. By the Δ -support equivalence at stage α (which holds by the inductive hypothesis that the Task 1 filter infrastructure is established at stage α), there exists a countable admissible (E_f, D_f) such that

$$\Delta_\alpha^\uparrow(E_f, D_f) \leq \text{Stab}_{\mathcal{G}_\alpha}(\dot{f}).$$

Step 2: Bridge from \mathcal{G}_α -stabilizer to $\text{Sym}(\omega_1)$ -stabilizer.

We claim $\text{Fix}(E_f) \leq \text{Stab}(f)$. Let $\pi \in \text{Fix}(E_f)$.

By Lemma 5.43 (applied at stage α , valid by the inductive hypothesis), there exists a diagonal lift $\hat{\pi}^{D_f} \in \mathcal{G}_\alpha$ with $\pi_\alpha^0(\hat{\pi}^{D_f}) = \pi$ and which fixes every coordinate in D_f pointwise. By Definition 5.27, this implies $\hat{\pi}^{D_f} \in \Delta_\alpha^\uparrow(E_f, D_f) \leq \text{Stab}_{\mathcal{G}_\alpha}(\dot{f})$. Hence

$$\hat{\pi}^{D_f} \cdot \dot{f} = \dot{f} \quad \text{and therefore} \quad \hat{\pi}^{D_f} \cdot f = f.$$

Let $\hat{\pi}^\emptyset \in \mathcal{G}_\alpha$ be the standard lift of π (the diagonal lift with empty protection set), so $\pi_\alpha^0(\hat{\pi}^\emptyset) = \pi$. Set

$$k := (\hat{\pi}^\emptyset)^{-1} \cdot \hat{\pi}^{D_f} \in \mathcal{G}_\alpha.$$

Then $\pi_\alpha^0(k) = \text{id}$, i.e. $k \in \ker(\pi_\alpha^0)$. By Lemma 5.4, since $\text{tc}(f) \subseteq V(T) \cup \text{Ord}$, we have $k \cdot f = f$. Therefore,

$$f = \hat{\pi}^{D_f} \cdot f = \hat{\pi}^\emptyset \cdot (k \cdot f) = \hat{\pi}^\emptyset \cdot f.$$

Finally, note that if $g, h \in \mathcal{G}_\alpha$ satisfy $\pi_\alpha^0(g) = \pi_\alpha^0(h)$, then $g^{-1}h \in \ker(\pi_\alpha^0)$, so $g \cdot f = h \cdot f$ by Lemma 5.4. In particular, the action of the standard lift $\hat{\pi}^\emptyset$ on f depends only on $\pi = \pi_\alpha^0(\hat{\pi}^\emptyset)$, hence $\pi \cdot f = \hat{\pi}^\emptyset \cdot f = f$. Hence

$$\pi \cdot f = \hat{\pi}^\emptyset \cdot f = f,$$

and thus $\pi \in \text{Stab}(f)$. This proves $\text{Fix}(E_f) \leq \text{Stab}(f)$.

Step 3: Orbit elements inherit countable support.

For any $g = \sigma \cdot f \in [f]$ (where $\sigma \in \text{Sym}(\omega_1)$), we have

$$\text{Stab}(g) = \text{Stab}(\sigma \cdot f) = \sigma \cdot \text{Stab}(f) \cdot \sigma^{-1}.$$

Since $\text{Fix}(E_f) \leq \text{Stab}(f)$, conjugating gives

$$\text{Fix}(\sigma[E_f]) = \sigma \cdot \text{Fix}(E_f) \cdot \sigma^{-1} \leq \sigma \cdot \text{Stab}(f) \cdot \sigma^{-1} = \text{Stab}(g).$$

Thus $E_g := \sigma[E_f]$ is a countable support for g . \square

Remark 5.10 (No circularity). The proof of Lemma 5.9 invokes Lemma 5.43 (existence of diagonal lifts) at stage α . This does *not* create a circular dependency: the proof of Lemma 5.43 constructs diagonal lifts by direct recursion on stages using the package-level diagonal machinery of Section 5.1, and does not use Lemma 5.9, Lemma 5.40, or Lemma 5.50.

Definition 5.11 (Diagonal-cancellation automorphisms). Fix an orbit package $\mathbb{S}_{[f]}$ (either $\mathbb{Q}_{[f]}$ or $\mathbb{R}_{[f]}$). Let $E \subseteq \omega_1$ be countable and let $D \subseteq [f]$ be countable such that

$$\forall g \in D \ (\text{Fix}(E) \leq \text{Stab}(g)),$$

i.e. every $\pi \in \text{Fix}(E)$ fixes each $g \in D$ as an orbit element.

For $\pi \in \text{Fix}(E)$ define an automorphism $\hat{\pi}^D \in \text{Aut}(\mathbb{S}_{[f]})$ by: for each $\vec{p} \in \mathbb{S}_{[f]}$ and each $g \in [f]$,

$$(\hat{\pi}^D \cdot \vec{p})(g) := \begin{cases} \vec{p}(g), & g \in D, \\ (\pi \cdot \vec{p})(g), & g \notin D, \end{cases}$$

where $\pi \cdot \vec{p}$ is the orbit action from Definition 5.5 (cf. Lemma 5.8). Let

$$\Delta(E, D) := \{\hat{\pi}^D : \pi \in \text{Fix}(E)\} \leq \text{Aut}(\mathbb{S}_{[f]}).$$

Remark 5.12 (Support convention for diagonal lifts). Conditions $\vec{p} \in \mathbb{S}_{[f]}$ are partial functions with countable support $\text{supp}(\vec{p}) \subseteq [f]$. Fix an admissible pair (E, D) and $\pi \in \text{Fix}(E)$, and write $\vec{p}' := \hat{\pi}^D \cdot \vec{p}$.

Then for each $g \in [f]$ we have the coordinate formula

$$\vec{p}'(g) = \begin{cases} \vec{p}(g), & g \in D, \\ A_{\pi, \pi^{-1} \cdot g}(\vec{p}(\pi^{-1} \cdot g)), & g \notin D, \end{cases}$$

i.e. $\widehat{\pi}^D$ agrees with the standard orbit action off D and is the identity on D .

Equivalently, for $h \in [f] \setminus D$ we may write

$$\vec{p}'(\pi \cdot h) = A_{\pi,h}(\vec{p}(h)).$$

In particular,

$$\text{supp}(\vec{p}') = (\text{supp}(\vec{p}) \cap D) \cup \pi \cdot (\text{supp}(\vec{p}) \setminus D),$$

which is countable since π is a bijection.

Definition 5.13 (Orbit-package stage filter (diagonal base)). Let $\mathcal{G}_{[f]} := \text{Aut}(\mathbb{S}_{[f]})$.

For countable $E \subseteq \omega_1$ and countable $D \subseteq [f]$ such that $\text{Fix}(E) \leq \text{Stab}(g)$ for all $g \in D$, define

$$\Delta^\uparrow(E, D) := \langle \Delta(E', D') \mid \begin{array}{l} E \subseteq E' \in [\omega_1]^{\leq\omega}, D \subseteq D' \subseteq [f] \text{ countable, and} \\ \text{Fix}(E') \leq \text{Stab}(g) \quad \forall g \in D' \end{array} \rangle,$$

where $\Delta(E', D')$ is as in Definition 5.11.

Call a pair (E, D) *admissible* if $E \subseteq \omega_1$ is countable, $D \subseteq [f]$ is countable, and $\text{Fix}(E) \leq \text{Stab}(g)$ for every $g \in D$.

Let $\mathcal{F}_{[f]}$ be the filter of subgroups of $\mathcal{G}_{[f]}$ generated by the family $\{\Delta^\uparrow(E, D) \mid (E, D) \text{ admissible}\}$. Since this family is downward directed (if (E_i, D_i) are admissible then $(E_0 \cup E_1, D_0 \cup D_1)$ is admissible and $\Delta^\uparrow(E_0 \cup E_1, D_0 \cup D_1) \leq \Delta^\uparrow(E_i, D_i)$ for $i = 0, 1$), the generated filter is just the upward closure of this family; equivalently,

$$H \in \mathcal{F}_{[f]} \iff \exists \text{admissible}(E, D) (\Delta^\uparrow(E, D) \leq H).$$

Remark 5.14. Lemmas 5.18 and 5.19 show that these diagonal generators are closed under conjugation by the successor-stage groups (generated by standard and diagonal lifts), hence yield the required normal filters in the symmetric-iteration construction.

Lemma 5.15 (Monotonicity of Δ^\uparrow). *If (E_0, D_0) and (E_1, D_1) are admissible and $E_0 \subseteq E_1$ and $D_0 \subseteq D_1$, then*

$$\Delta^\uparrow(E_1, D_1) \leq \Delta^\uparrow(E_0, D_0).$$

Proof. Any pair (E', D') eligible in the definition of $\Delta^\uparrow(E_1, D_1)$ satisfies $E_1 \subseteq E'$ and $D_1 \subseteq D'$, hence also $E_0 \subseteq E'$ and $D_0 \subseteq D'$. Thus the generating family for $\Delta^\uparrow(E_1, D_1)$ is a subfamily of the generating family for $\Delta^\uparrow(E_0, D_0)$, so the generated subgroup is smaller. \square

Lemma 5.16 (ω_1 -completeness of the diagonal stage filter). $\mathcal{F}_{[f]}$ is ω_1 -complete.

Proof. Let $\langle H_n : n < \omega \rangle$ be members of $\mathcal{F}_{[f]}$. By Definition 5.13, for each n choose an admissible pair (E_n, D_n) such that

$$\Delta^\uparrow(E_n, D_n) \leq H_n.$$

Let $E := \bigcup_{n < \omega} E_n$ and $D := \bigcup_{n < \omega} D_n$; both are countable. Moreover (E, D) is admissible: if $g \in D$, pick n with $g \in D_n$, then $\text{Fix}(E) \subseteq \text{Fix}(E_n) \leq \text{Stab}(g)$.

By Lemma 5.15, for each n we have

$$\Delta^\uparrow(E, D) \leq \Delta^\uparrow(E_n, D_n) \leq H_n.$$

Hence $\Delta^\uparrow(E, D) \leq \bigcap_{n < \omega} H_n$, so $\bigcap_{n < \omega} H_n \in \mathcal{F}_{[f]}$. \square

Lemma 5.17 (Diagonal-cancellation gives a subgroup of automorphisms). *In the setup of Definition 5.11:*

- (i) *For each $\pi \in \text{Fix}(E)$, $\widehat{\pi}^D$ is an automorphism of $\mathbb{S}_{[f]}$.*
- (ii) *The map $\pi \mapsto \widehat{\pi}^D$ is a group monomorphism $\text{Fix}(E) \hookrightarrow \text{Aut}(\mathbb{S}_{[f]})$. In particular, $\Delta(E, D)$ is a subgroup of $\text{Aut}(\mathbb{S}_{[f]})$.*

Proof. Fix $\pi \in \text{Fix}(E)$. Since $\pi \cdot (\cdot)$ is an automorphism of $\mathbb{S}_{[f]}$ and $\widehat{\pi}^D$ agrees with it on all coordinates outside D while acting as the identity on coordinates in D , it preserves supports, preserves coordinatewise extension, and is bijective with inverse $\widehat{\pi^{-1}}^D$. This proves (i).

For (ii), $\widehat{\pi}^D \circ \widehat{\sigma}^D = \widehat{\pi\sigma}^D$ follows coordinatewise from the definition (on D both sides are the identity; off D both sides agree with the rigidified orbit action (Lemma 5.8)). Injectivity is immediate. \square

Lemma 5.18 (Conjugation of diagonal-cancellation groups by standard lifts). *Let (E, D) be admissible. For $\sigma \in \text{Sym}(\omega_1)$ write*

$$\sigma \cdot E := \{\sigma(\xi) : \xi \in E\} \subseteq \omega_1, \quad \sigma \cdot D := \{\sigma \cdot g : g \in D\} \subseteq [f].$$

Then $(\sigma \cdot E, \sigma \cdot D)$ is admissible. Moreover, for every $\pi \in \text{Fix}(E)$,

$$\widehat{\sigma}^\emptyset \circ \widehat{\pi}^D \circ (\widehat{\sigma}^\emptyset)^{-1} = \widehat{\sigma \pi \sigma^{-1}}^{\sigma \cdot D}.$$

Consequently,

$$\widehat{\sigma}^\emptyset \Delta(E, D) (\widehat{\sigma}^\emptyset)^{-1} = \Delta(\sigma \cdot E, \sigma \cdot D), \quad \widehat{\sigma}^\emptyset \Delta^\uparrow(E, D) (\widehat{\sigma}^\emptyset)^{-1} = \Delta^\uparrow(\sigma \cdot E, \sigma \cdot D).$$

Proof. Admissibility. Let $\tau \in \text{Fix}(\sigma \cdot E)$ and let $g \in \sigma \cdot D$, say $g = \sigma \cdot h$ with $h \in D$. Then $\sigma^{-1}\tau\sigma \in \text{Fix}(E) \leq \text{Stab}(h)$, hence

$$\tau \cdot g = \tau \cdot (\sigma \cdot h) = \sigma \cdot ((\sigma^{-1}\tau\sigma) \cdot h) = \sigma \cdot h = g.$$

So $\text{Fix}(\sigma \cdot E) \leq \text{Stab}(g)$ for all $g \in \sigma \cdot D$.

Conjugation formula. Fix $\pi \in \text{Fix}(E)$ and $\vec{p} \in \mathbb{S}_{[f]}$, and let $g \in [f]$. Put $h := \sigma^{-1} \cdot g$. If $g \in \sigma \cdot D$ (equivalently $h \in D$), then $\hat{\pi}^D$ acts as the identity at h , so the conjugate acts as the identity at g , i.e. it fixes the g -coordinate of \vec{p} .

If $g \notin \sigma \cdot D$ (equivalently $h \notin D$), then at h the map $\hat{\pi}^D$ agrees with the standard orbit action by π . Using that $\hat{\sigma}^\emptyset$ is the standard orbit action by σ (i.e. $D = \emptyset$), a direct coordinate computation gives that the conjugate agrees with the standard orbit action by $\sigma\pi\sigma^{-1}$ at g . Thus the conjugate acts as the identity on $\sigma \cdot D$ and as $\widehat{\sigma\pi\sigma^{-1}}^{\sigma \cdot D}$ off $\sigma \cdot D$, exactly $\widehat{\sigma\pi\sigma^{-1}}^{\sigma \cdot D}$.

The subgroup identity for $\Delta(E, D)$ follows by taking images of $\text{Fix}(E)$ under $\pi \mapsto \hat{\pi}^D$ and using $\sigma \text{Fix}(E)\sigma^{-1} = \text{Fix}(\sigma \cdot E)$.

Finally, for $\Delta^\uparrow(E, D)$: conjugating any generator $\Delta(E', D')$ occurring in the definition of $\Delta^\uparrow(E, D)$ yields $\Delta(\sigma \cdot E', \sigma \cdot D')$, which is a generator for $\Delta^\uparrow(\sigma \cdot E, \sigma \cdot D)$. Applying the same argument with σ^{-1} gives equality. \square

Lemma 5.19 (Conjugation of diagonal generators by diagonal lifts). *Fix an admissible pair (E_0, D_0) and let $\sigma \in \text{Fix}(E_0)$. Write $\hat{\sigma}^{D_0} \in \Delta(E_0, D_0)$ for the corresponding diagonal lift.*

Then for every admissible pair (E, D) , letting

$$E^* := E \cup E_0, \quad D^* := D \cup D_0,$$

we have

$$\hat{\sigma}^{D_0} \cdot \Delta^\uparrow(E^*, D^*) \cdot (\hat{\sigma}^{D_0})^{-1} = \Delta^\uparrow(\sigma \cdot E^*, \sigma \cdot D^*).$$

In particular, $\hat{\sigma}^{D_0} H (\hat{\sigma}^{D_0})^{-1} \in \mathcal{F}_{[f]}$ for every $H \in \mathcal{F}_{[f]}$.

Proof. Let (E, D) be admissible and set $E^* := E \cup E_0$ and $D^* := D \cup D_0$. Then (E^*, D^*) is admissible since $\text{Fix}(E^*) \leq \text{Fix}(E)$ and $\text{Fix}(E^*) \leq \text{Fix}(E_0)$.

We first claim that if (E', D') is admissible with $E^* \subseteq E'$ and $D^* \subseteq D'$, then

$$\hat{\sigma}^{D_0} \cdot \Delta(E', D') \cdot (\hat{\sigma}^{D_0})^{-1} = \Delta(\sigma \cdot E', \sigma \cdot D').$$

Indeed, $D_0 \subseteq D'$ by assumption, so every $\hat{\pi}^{D'} \in \Delta(E', D')$ is the identity on all coordinates in D_0 . Since $\hat{\sigma}^{D_0}$ agrees with the standard lift $\hat{\sigma}^\emptyset$ off D_0 , the same coordinate computation as in Lemma 5.18 gives

$$\hat{\sigma}^{D_0} \cdot \hat{\pi}^{D'} \cdot (\hat{\sigma}^{D_0})^{-1} = \widehat{\sigma\pi\sigma^{-1}}^{\sigma \cdot D'}.$$

Taking images of $\text{Fix}(E')$ under $\pi \mapsto \widehat{\pi}^{D'}$ and using $\sigma \text{Fix}(E')\sigma^{-1} = \text{Fix}(\sigma \cdot E')$ yields the subgroup identity above.

Now conjugate the generating family for $\Delta^\uparrow(E^*, D^*)$: each generator $\Delta(E', D')$ (with $E^* \subseteq E'$ and $D^* \subseteq D'$) is sent to $\Delta(\sigma \cdot E', \sigma \cdot D')$, which is a generator for $\Delta^\uparrow(\sigma \cdot E^*, \sigma \cdot D^*)$. Applying the same argument with σ^{-1} gives equality.

Finally, if $H \in \mathcal{F}_{[f]}$, choose admissible (E, D) with $\Delta^\uparrow(E, D) \leq H$. Then $\Delta^\uparrow(E^*, D^*) \leq \Delta^\uparrow(E, D) \leq H$ by monotonicity, so

$$\Delta^\uparrow(\sigma \cdot E^*, \sigma \cdot D^*) = \widehat{\sigma}^{D_0} \cdot \Delta^\uparrow(E^*, D^*) \cdot (\widehat{\sigma}^{D_0})^{-1} \leq \widehat{\sigma}^{D_0} H (\widehat{\sigma}^{D_0})^{-1},$$

which shows $\widehat{\sigma}^{D_0} H (\widehat{\sigma}^{D_0})^{-1} \in \mathcal{F}_{[f]}$. \square

5.2 Iteration-level diagonal automorphisms

The diagonal-cancellation construction of Definition 5.11 is formulated for a single orbit package $\mathbb{S}_{[f]}$. For the $\neg\text{AC}$ argument at limit stages, we need diagonal lifts acting on the *full iteration forcing* \mathbb{P}_λ , with protection sets D that may span multiple packages across multiple stages.

Remark 5.20 (Simultaneous transfinite induction). Several results in this section—particularly Lemma 5.9, Lemma 5.40, and Lemma 5.50—are established by simultaneous transfinite induction on stages α . At each stage, we assume:

- (i) the filter infrastructure (generation, completeness, properness) holds at all earlier stages;
- (ii) diagonal lifts exist in \mathcal{G}_β for all $\beta < \alpha$ (Lemma 5.43);
- (iii) the Δ -support equivalence holds at all earlier stages.

These hypotheses are then discharged at stage α using only the inductive assumptions at earlier stages, avoiding any circularity.

Lemma 5.21 (Successor decomposition and splitting). *Let α be an ordinal. In the two-step presentation $\mathbb{P}_{\alpha+1} \cong \mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha$, the projection $\pi_{\alpha+1}^\alpha : \mathcal{G}_{\alpha+1} \rightarrow \mathcal{G}_\alpha$ is a surjective group homomorphism. Let $K_{\alpha+1} := \ker(\pi_{\alpha+1}^\alpha)$.*

Moreover, there is a group-theoretic splitting map $\text{spl} : \mathcal{G}_\alpha \rightarrow \mathcal{G}_{\alpha+1}$ (the splitting section) such that $\pi_{\alpha+1}^\alpha \circ \text{spl} = \text{id}_{\mathcal{G}_\alpha}$, and every $g \in \mathcal{G}_{\alpha+1}$ can be written as $g = \text{spl}(h) \cdot k$ with $h = \pi_{\alpha+1}^\alpha(g)$ and $k \in K_{\alpha+1}$.

Proof. This is exactly the successor-step group structure supplied by the countable-support symmetric iteration framework (Remark 2.1, cf. [3, Def. 3.1]). Concretely:

- $\pi_{\alpha+1}^\alpha$ is restriction to the \mathbb{P}_α -part;
- $K_{\alpha+1}$ consists of automorphisms acting trivially on \mathbb{P}_α ;
- $\text{spl}(h)$ is the splitting section that acts as h on \mathbb{P}_α and trivially on $\dot{\mathbb{Q}}_\alpha$.

The decomposition $g = \text{spl}(\pi_{\alpha+1}^\alpha(g)) \cdot k$ with $k = \text{spl}(\pi_{\alpha+1}^\alpha(g))^{-1} \cdot g$ is immediate; one checks $k \in K_{\alpha+1}$ since $\pi_{\alpha+1}^\alpha(k) = \text{id}$. \square

Definition 5.22 (Package coordinate sets). We define by transfinite recursion the sets Coords_α of *package coordinates occurring below stage α* .

- $\text{Coords}_0 := \emptyset$.
- At a successor stage $\alpha + 1$, write $\mathbb{P}_{\alpha+1} \cong \mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha$. In this paper, $\dot{\mathbb{Q}}_\alpha$ is (forced to be) a finite product of orbit packages

$$\dot{\mathbb{Q}}_\alpha \cong \prod_{i < m_\alpha} \dot{\mathbb{S}}_{[f_{\alpha,i}]},$$

where each $\dot{\mathbb{S}}_{[f_{\alpha,i}]}$ is either an orbit PP-package $\mathbb{Q}_{[f]}$ or an orbit ACwo-package $\mathbb{R}_{[f]}$ (and $m_\alpha < \omega$). Define the stage- α index set as the disjoint union

$$I_\alpha := \bigsqcup_{i < m_\alpha} [f_{\alpha,i}].$$

We set

$$\text{NewCoords}_{\alpha+1} := \{\langle \alpha, z \rangle : z \in I_\alpha\},$$

$$\text{Coords}_{\alpha+1} := \text{Coords}_\alpha \dot{\cup} \text{NewCoords}_{\alpha+1}.$$

When $z \in I_\alpha$ corresponds to (i, g) , we also write $\langle \alpha, i, g \rangle$ for $\langle \alpha, z \rangle$.

- At a limit stage λ , set $\text{Coords}_\lambda := \bigcup_{\alpha < \lambda} \text{Coords}_\alpha$. We also write $\text{Coords}_{<\lambda} := \text{Coords}_\lambda$.

When working at a fixed successor stage $\alpha + 1$, we suppress the stage tag and write $\langle i, g \rangle$ for $\langle \alpha, i, g \rangle$ (so “ $\langle i, g \rangle \in \text{Coords}_{\alpha+1} \setminus \text{Coords}_\alpha$ ” means $i < n_\alpha$ and $g \in [f_{\alpha,i}]$).

We let $\text{Sym}(\omega_1)$ act on new coordinates by

$$\pi \cdot \langle \alpha, i, g \rangle := \langle \alpha, i, \pi \cdot g \rangle,$$

equivalently $\pi \cdot \langle \alpha, z \rangle := \langle \alpha, \pi \cdot z \rangle$ when using I_α .

Definition 5.23 (Successor diagonal kernel element). Fix a successor stage $\alpha + 1$. Let $\pi \in \text{Sym}(\omega_1)$ and let $D^{\text{new}} \subseteq \text{Coords}_{\alpha+1} \setminus \text{Coords}_\alpha$ be a set of package coordinates newly introduced at stage $\alpha + 1$.

Assume π fixes each $d \in D^{\text{new}}$ under the induced action on tagged coordinates, i.e., if $d = \langle \alpha, i, g \rangle$ then $\pi \cdot d = d$ equivalently $\pi \cdot g = g$. (Here $\pi \cdot \langle \alpha, i, g \rangle := \langle \alpha, i, \pi \cdot g \rangle$ using the orbit action on $[f_{\alpha,i}]$.)

Under this assumption, define $\kappa_{\alpha+1}(\pi, D^{\text{new}}) \in \text{Aut}(\mathbb{P}_{\alpha+1})$ by:

- $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ acts as the identity on the \mathbb{P}_α -part (all coordinates at stages $\leq \alpha$);
- on each new package coordinate $d \in \text{Coords}_{\alpha+1} \setminus \text{Coords}_\alpha$:
 - if $d \notin D^{\text{new}}$, act by the package action induced by π (Definition 5.5);
 - if $d \in D^{\text{new}}$, act as the identity (fix the d -coordinate pointwise).

Remark 5.24 (Why the hypothesis is necessary). The hypothesis “ $\pi \cdot d = d$ for all $d \in D^{\text{new}}$ ” ensures $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ is a bijection on package coordinates. Without it, collisions can occur: if $d \notin D^{\text{new}}$ but $\pi \cdot d \in D^{\text{new}}$, then both d and $\pi \cdot d$ would map to $\pi \cdot d$.

In our application (Lemma 5.43), global admissibility guarantees this hypothesis: if (E, D) is globally admissible and $\pi \in \text{Fix}(E)$, then $\pi \cdot d = d$ for all $d \in D$, hence for all $d \in D^{\text{new}} = D_{\beta+1} \setminus D_\beta \subseteq D$.

Lemma 5.25 (Successor diagonal kernel element lies in $\mathcal{G}_{\alpha+1}$ and its kernel). *Under the hypothesis of Definition 5.23:*

- (i) $\kappa_{\alpha+1}(\pi, D^{\text{new}}) \in \mathcal{G}_{\alpha+1}$;
- (ii) $\kappa_{\alpha+1}(\pi, D^{\text{new}}) \in K_{\alpha+1} = \ker(\pi_{\alpha+1}^\alpha)$.

Moreover, for fixed D^{new} , the map $\pi \mapsto \kappa_{\alpha+1}(\pi, D^{\text{new}})$ is a group homomorphism on its domain $\{\pi \in \text{Sym}(\omega_1) : \pi \cdot d = d \ \forall d \in D^{\text{new}}\}$ (Here the action on new coordinates is $\pi \cdot \langle \alpha, i, g \rangle := \langle \alpha, i, \pi \cdot g \rangle$.)

Proof. (i) By Definition 5.23, the map $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ acts trivially on the \mathbb{P}_α -part, and on the successor iterand \mathbb{Q}_α it acts coordinatewise on the new package coordinates $\text{NewCoords}_{\alpha+1}$: it is the identity on D^{new} and it agrees with the π -induced orbit action on $\text{NewCoords}_{\alpha+1} \setminus D^{\text{new}}$. If (as allowed by Remark 4.31) \mathbb{Q}_α is a finite product of orbit packages, then $\text{NewCoords}_{\alpha+1}$ is tagged by the factor index, and the action is defined independently on each factor, hence yields an automorphism of the full stage- $(\alpha + 1)$ forcing.

By Lemma 5.17, each factor is a legitimate automorphism of its package forcing $\mathbb{S}_{[f]}$. By the successor-stage group structure (Remark 2.1), $\mathcal{G}_{\alpha+1}$ is generated by lifts from \mathcal{G}_α together with automorphisms of the successor iterand $\dot{\mathbb{Q}}_\alpha$ (equivalently, of each new orbit-package factor). Since $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ acts as identity on \mathbb{P}_α and as a (product of) diagonal-cancellation automorphism(s) on new packages, it lies in $\mathcal{G}_{\alpha+1}$.

(ii) By Definition 5.23, $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ acts as the identity on \mathbb{P}_α . Hence $\pi_{\alpha+1}^\alpha(\kappa_{\alpha+1}(\pi, D^{\text{new}})) = \text{id}$, so $\kappa_{\alpha+1}(\pi, D^{\text{new}}) \in \ker(\pi_{\alpha+1}^\alpha) = K_{\alpha+1}$.

The homomorphism property: for π, σ both fixing D^{new} pointwise,

$$\kappa_{\alpha+1}(\pi, D^{\text{new}}) \circ \kappa_{\alpha+1}(\sigma, D^{\text{new}}) = \kappa_{\alpha+1}(\pi\sigma, D^{\text{new}}),$$

since on coordinates outside D^{new} , composing the σ -action with the π -action yields the $(\pi\sigma)$ -action, and on D^{new} both sides act as identity. \square

Definition 5.26 (Global admissibility). Let λ be a stage of the iteration. A pair (E, D) is *globally admissible at stage λ* if:

- (i) $E \in [\omega_1]^{\leq\omega}$ is countable;
- (ii) $D \subseteq \text{Coords}_{<\lambda}$ is countable;
- (iii) $\text{Fix}(E) \leq \text{Stab}(d)$ for every $d \in D$.

When $D = \emptyset$, condition (iii) is vacuously satisfied, and (E, \emptyset) is globally admissible for any countable E .

Definition 5.27 (Iteration-level diagonal subgroups and diagonal lifts). Fix a stage λ .

(A) Characterized diagonal subgroups. For any $E \subseteq \omega_1$ and any $D \subseteq \text{Coords}_{<\lambda}$ define

$$\Delta_\lambda^\uparrow(E, D) := \{ g \in \mathcal{G}_\lambda : \pi_\lambda^0(g) \in \text{Fix}(E) \text{ and } g \text{ fixes every coordinate in } D \text{ pointwise} \}.$$

(B) Diagonal lifts (require global admissibility). If (E, D) is globally admissible at stage λ and $\pi \in \text{Fix}(E)$, define $\hat{\pi}^D \in \text{Aut}(\mathbb{P}_\lambda)$ by:

- **Stage 0 (Cohen base):** act as π , i.e. $(\hat{\pi}^D \cdot p)(\pi(\alpha), n) = p(\alpha, n)$ for $(\alpha, n) \in \omega_1 \times \omega$.
- **Stages > 0 (package coordinates):** for each package coordinate g , act as the identity if $g \in D$, and as the π -orbit action if $g \notin D$.

Finally set

$$\Delta_\lambda(E, D) := \{\hat{\pi}^D : \pi \in \text{Fix}(E)\} \leq \text{Aut}(\mathbb{P}_\lambda).$$

Remark 5.28. If (E, D) is globally admissible and $\pi \in \text{Fix}(E)$, then $\hat{\pi}^D \in \Delta_\lambda^\uparrow(E, D)$. Indeed, $\pi_\lambda^0(\hat{\pi}^D) = \pi \in \text{Fix}(E)$ and $\hat{\pi}^D$ fixes every coordinate in D pointwise.

Remark 5.29 (Characterization vs. generation). Definition 5.27 defines $\Delta_\lambda^\uparrow(E, D)$ as a *characterized subgroup*: $g \in \Delta_\lambda^\uparrow(E, D)$ iff g 's stage-0 projection fixes E pointwise and g fixes every package coordinate in D pointwise.

This “characterization” definition makes monotonicity immediate (Lemma 5.34). In Lemma 5.44, we show that diagonal lifts $\hat{\pi}^{D'}$ (for $\pi \in \text{Fix}(E)$ and $D' \supseteq D$) lie in $\Delta_\lambda^\uparrow(E, D)$ —this is immediate from the characterization. We do *not* need to prove that $\Delta_\lambda^\uparrow(E, D)$ is *generated* by such diagonal lifts; the characterization suffices for all filter-theoretic purposes.

Lemma 5.30 (Group action on coordinates). *Let $g \in \mathcal{G}_\lambda$. Then:*

- (i) *The stage-0 projection $\pi_\lambda^0 : \mathcal{G}_\lambda \rightarrow \text{Sym}(\omega_1)$ is a group homomorphism.*
- (ii) *The element g induces a bijection on the set of package coordinates at each stage $\beta < \lambda$. In particular, if D is a countable set of package coordinates, then $g \cdot D$ is countable.*
- (iii) *For $E \subseteq \omega_1$, define $g \cdot E := \pi_\lambda^0(g)[E]$. Then $g \cdot E$ is countable whenever E is countable.*

Proof. (i) This is part of the inverse/direct system structure of the iteration groups (cf. [3, Def. 3.1]) and the fact that π_λ^0 is defined stagewise and respects composition.

(ii) We prove by induction on stages that each $g \in \mathcal{G}_\beta$ determines a bijection of Coords_β (in the sense of Definition 5.22), and hence sends countable sets of coordinates to countable sets.

At stage 0 there are no package coordinates. At a successor stage $\beta + 1$, write $g = \text{st}(h) \cdot k$ as in Lemma 5.21, where $h = \pi_{\beta+1}^\beta(g) \in \mathcal{G}_\beta$ and $k \in K_{\beta+1}$ acts trivially on the \mathbb{P}_β -part. By the inductive hypothesis, h induces a bijection of Coords_β . The standard lift $\text{st}(h)$ extends this action and fixes every element of $\text{NewCoords}_{\beta+1}$. On the other hand, k fixes Coords_β pointwise and acts on the successor forcing $\dot{\mathbb{Q}}_\beta$ by compositions of package automorphisms (Lemma 5.8) and diagonal-cancellation automorphisms (Definition 5.11), each of which induces a bijection on the new coordinate index set (and

in particular cannot collapse distinct coordinates, by the admissibility hypothesis built into their definition). Therefore g induces a bijection on $\text{Coords}_{\beta+1} = \text{Coords}_\beta \dot{\cup} \text{NewCoords}_{\beta+1}$.

At a limit stage η , elements of \mathcal{G}_η are (by definition) coherent direct-limit elements whose restrictions to earlier stages determine them uniquely (cf. [3, Def. 3.1]). The induced bijection on $\text{Coords}_\eta = \bigcup_{\beta < \eta} \text{Coords}_\beta$ is obtained by taking the union of the stagewise bijections, and is again a bijection.

Consequently, if D is countable then so is its image under the induced bijection.

(iii) Immediate: $\pi_\lambda^0(g) \in \text{Sym}(\omega_1)$ is a bijection and hence preserves countability. \square

Lemma 5.31 (Iteration diagonal lifts are automorphisms). *For (E, D) globally admissible at stage λ and $\pi \in \text{Fix}(E)$, the map $\hat{\pi}^D$ is an automorphism of \mathbb{P}_λ satisfying:*

- (i) $\hat{\pi}^D$ restricts to π on the Cohen base $\text{Add}(\omega, \omega_1)$;
- (ii) $\hat{\pi}^D$ restricts to the identity on the g -coordinate factor for each $g \in D$;
- (iii) $\hat{\pi}^D$ restricts to the standard π -action on all coordinates outside D .

Moreover, $\pi \mapsto \hat{\pi}^D$ is a group monomorphism $\text{Fix}(E) \hookrightarrow \text{Aut}(\mathbb{P}_\lambda)$.

Proof. By global admissibility, $\pi \cdot g = g$ for all $g \in D$, so the protected coordinates are pointwise fixed and the diagonal definition cannot create coordinate-collisions. Order preservation and bijectivity follow coordinate-wise: on the Cohen base this is the standard automorphism; on each package coordinate outside D , this is the orbit action from Lemma 5.8; on coordinates in D , this is the identity.

The group homomorphism property $\hat{\pi}^D \circ \hat{\sigma}^D = \hat{\pi} \hat{\sigma}^D$ follows coordinate-wise: on D both sides are the identity; on the Cohen base and on coordinates outside D , both sides agree with the standard action of $\pi\sigma$. \square

Corollary 5.32 (Diagonal lifts swap Cohen reals). *Let $\hat{\pi}^D \in \mathcal{G}_\lambda$ be a diagonal lift with $\pi_\lambda^0(\hat{\pi}^D) = \pi$. Then for each $\alpha < \omega_1$:*

$$\hat{\pi}^D \cdot \dot{c}_\alpha = \dot{c}_{\pi(\alpha)},$$

where \dot{c}_α is the canonical \mathbb{P}_0 -name for the α -th Cohen real.

In particular, if $\pi = (\beta \ \gamma)$ is a transposition with $\beta \neq \gamma$, then $\hat{\pi}^D$ swaps $c_\beta \leftrightarrow c_\gamma$ in the generic extension.

Proof. View \dot{c}_α as a \mathbb{P}_λ -name via the canonical inclusion of \mathbb{P}_0 -names into \mathbb{P}_λ -names.

By Lemma 5.31(i), the diagonal lift $\hat{\pi}^D$ restricts to π on the Cohen base $\mathbb{P}_0 = \text{Add}(\omega, \omega_1)$. The canonical name \dot{c}_α is defined by $\dot{c}_\alpha = \{(\check{n}, p) : p(\alpha, n) = 1\}$. The action of π on \mathbb{P}_0 is $(\pi \cdot p)(\pi(\alpha), n) = p(\alpha, n)$, which induces $\pi \cdot \dot{c}_\alpha = \dot{c}_{\pi(\alpha)}$. Since $\hat{\pi}^D$ restricts to π on stage 0, we have $\hat{\pi}^D \cdot \dot{c}_\alpha = \dot{c}_{\pi(\alpha)}$. \square

Remark 5.33 (Standard lifts as empty diagonal lifts). The standard lift $\hat{\pi}^\emptyset$ (Definition 5.27 with $D = \emptyset$) acts as π on *all* coordinates, recovering the usual lifted action of $\text{Sym}(\omega_1)$ on the iteration. Thus the pushforward subgroups $\text{Fix}(E) \leq \mathcal{G}_\lambda$ (acting via standard lifts) coincide with $\Delta_\lambda(E, \emptyset)$, and consequently $\Delta_\lambda^\uparrow(E, \emptyset) = \text{Fix}(E)^\uparrow$.

This identification is crucial: *all* filter generators at limit stages—both Cohen-type and package-type—are of the form $\Delta_\lambda^\uparrow(E_i, D_i)$, with $D_i = \emptyset$ for Cohen generators and $D_i = \{f_i\}$ for package generators. The monotonicity lemma (Lemma 5.34) therefore applies uniformly to combine them.

Lemma 5.34 (Monotonicity for iteration-level diagonal groups). *If $E_0 \subseteq E_1$ and $D_0 \subseteq D_1$, then*

$$\Delta_\lambda^\uparrow(E_1, D_1) \leq \Delta_\lambda^\uparrow(E_0, D_0).$$

Proof. Let $g \in \Delta_\lambda^\uparrow(E_1, D_1)$. By the characterization definition:

- $\pi_\lambda^0(g) \in \text{Fix}(E_1) \subseteq \text{Fix}(E_0)$ (since $E_0 \subseteq E_1$ implies $\text{Fix}(E_1) \leq \text{Fix}(E_0)$).
- g fixes every coordinate in D_1 pointwise, hence fixes every coordinate in $D_0 \subseteq D_1$ pointwise.

Thus $g \in \Delta_\lambda^\uparrow(E_0, D_0)$. \square

Corollary 5.35 (Intersection of Δ^\uparrow -groups). *For any family $\{(E_i, D_i) : i \in I\}$ (with I finite or countable):*

$$\bigcap_{i \in I} \Delta_\lambda^\uparrow(E_i, D_i) \supseteq \Delta_\lambda^\uparrow\left(\bigcup_{i \in I} E_i, \bigcup_{i \in I} D_i\right).$$

Proof. By Lemma 5.34, for each $i \in I$: $\Delta_\lambda^\uparrow(\bigcup_j E_j, \bigcup_j D_j) \leq \Delta_\lambda^\uparrow(E_i, D_i)$. Hence the union-indexed group is contained in every term of the intersection. \square

Lemma 5.36 (Conjugation of iteration-level diagonal groups). *Let (E, D) be globally admissible at stage λ . For $\sigma \in \text{Sym}(\omega_1)$:*

$$\hat{\sigma}^\emptyset \cdot \Delta_\lambda^\uparrow(E, D) \cdot (\hat{\sigma}^\emptyset)^{-1} = \Delta_\lambda^\uparrow(\sigma \cdot E, \sigma \cdot D).$$

In particular, conjugates of diagonal generators remain diagonal generators (with shifted parameters).

Proof. The proof is identical to Lemma 5.18, now applied at the iteration level. \square

Remark 5.37 (Admissibility under conjugation up to enlarging E). Let λ be a stage and (E, D) be globally admissible at stage λ . For any $g \in \mathcal{G}_\lambda$, there is a countable $E^* \supseteq g \cdot E$ such that $(E^*, g \cdot D)$ is globally admissible. Consequently $\Delta_\lambda^\uparrow(E^*, g \cdot D) \subseteq g \Delta_\lambda^\uparrow(E, D) g^{-1}$.

Lemma 5.38 (General conjugation of Δ^\uparrow -groups). *For any $g \in \mathcal{G}_\lambda$, any $E \subseteq \omega_1$, and any set D of package coordinates:*

$$g \cdot \Delta_\lambda^\uparrow(E, D) \cdot g^{-1} = \Delta_\lambda^\uparrow(g \cdot E, g \cdot D)$$

where $g \cdot E := \pi_\lambda^0(g)[E]$ and $g \cdot D := \{g \cdot d : d \in D\}$ is the image of D under the action of g on package coordinates.

Proof. Fix $k \in \mathcal{G}_\lambda$ and write $\sigma := \pi_\lambda^0(k) \in \text{Sym}(\omega_1)$. Let $g \in \Delta_\lambda^\uparrow(E, D)$, and put $\tau := \pi_\lambda^0(g) \in \text{Fix}(E)$.

Stage-0 part. Since $\pi_\lambda^0 : \mathcal{G}_\lambda \rightarrow \text{Sym}(\omega_1)$ is a homomorphism, we have

$$\pi_\lambda^0(kgk^{-1}) = \sigma\tau\sigma^{-1} \in \text{Fix}(\sigma[E]) = \text{Fix}(\pi_\lambda^0(k)[E]).$$

Coordinate part. Let $d \in D$. By definition of $\Delta_\lambda^\uparrow(E, D)$, g fixes d pointwise. Using the coordinate action \cdot of \mathcal{G}_λ on $\text{Coords}_{<\lambda}$ (Lemma 5.30), we compute

$$(kgk^{-1}) \cdot (k \cdot d) = k \cdot (g \cdot (k^{-1} \cdot (k \cdot d))) = k \cdot (g \cdot d) = k \cdot d,$$

so kgk^{-1} fixes every coordinate in $k \cdot D$ pointwise.

Thus $kgk^{-1} \in \Delta_\lambda^\uparrow(\pi_\lambda^0(k)[E], k \cdot D)$. The reverse inclusion follows by conjugating with k^{-1} , so equality holds. \square

Corollary 5.39 (Admissibility preserved under unions). *If (E_1, D_1) and (E_2, D_2) are globally admissible, then $(E_1 \cup E_2, D_1 \cup D_2)$ is globally admissible.*

Proof. For $d \in D_1$: $\text{Fix}(E_1 \cup E_2) \leq \text{Fix}(E_1) \leq \text{Stab}(d)$. For $d \in D_2$: $\text{Fix}(E_1 \cup E_2) \leq \text{Fix}(E_2) \leq \text{Stab}(d)$. \square

Lemma 5.40 (Admissible hull for countable coordinate sets). *Let D be a countable set of package coordinates (at stages $< \lambda$) and let $E \subseteq \omega_1$ be countable. Then there exists a countable set $E^* \subseteq \omega_1$ with $E \subseteq E^*$ such that (E^*, D) is globally admissible.*

Consequently, $\Delta_\lambda^\uparrow(E^, D) \leq \Delta_\lambda^\uparrow(E, D)$.*

Proof. For each $d \in D$, Lemma 5.9 provides a countable $E_d \subseteq \omega_1$ such that $\text{Fix}(E_d) \leq \text{Stab}(d)$.

Let $E^* := E \cup \bigcup_{d \in D} E_d$, which is a countable union of countable sets, hence countable.

For each $d \in D$, we have $\text{Fix}(E^*) \leq \text{Fix}(E_d) \leq \text{Stab}(d)$, so (E^*, D) is globally admissible.

The final statement follows from monotonicity: $E \subseteq E^*$ implies $\Delta_\lambda^\uparrow(E^*, D) \leq \Delta_\lambda^\uparrow(E, D)$ by Lemma 5.34. \square

Lemma 5.41 (Projection of Δ^\uparrow -groups). *Let $\beta < \lambda$ and let $D \subseteq \text{Coords}_\beta$ (i.e., D only uses coordinates introduced below stage β). Then:*

$$\pi_\lambda^\beta[\Delta_\lambda^\uparrow(E, D)] \subseteq \Delta_\beta^\uparrow(E, D).$$

Proof. Let $g \in \Delta_\lambda^\uparrow(E, D)$. Then:

- $\pi_\lambda^0(g) \in \text{Fix}(E)$, and since $\pi_\beta^0 \circ \pi_\lambda^\beta = \pi_\lambda^0$, we have $\pi_\beta^0(\pi_\lambda^\beta(g)) = \pi_\lambda^0(g) \in \text{Fix}(E)$.
- g fixes every coordinate in D pointwise. Since $D \subseteq \text{Coords}_\beta$, these coordinates are visible at stage β , and the projection $\pi_\lambda^\beta(g)$ also fixes them pointwise.

Hence $\pi_\lambda^\beta(g) \in \Delta_\beta^\uparrow(E, D)$. \square

Lemma 5.42 (Pullback contains Δ^\uparrow). *Let $\beta < \lambda$ and let $D \subseteq \text{Coords}_\beta$. If $H \in \tilde{\mathcal{F}}_\beta^*$ contains $\Delta_\beta^\uparrow(E, D)$, then $(\pi_\lambda^\beta)^{-1}[H]$ contains $\Delta_\lambda^\uparrow(E, D)$.*

Proof. Let $g \in \Delta_\lambda^\uparrow(E, D)$. By Lemma 5.41, $\pi_\lambda^\beta(g) \in \Delta_\beta^\uparrow(E, D) \subseteq H$. Hence $g \in (\pi_\lambda^\beta)^{-1}[H]$. \square

Lemma 5.43 (Iteration-level diagonal lifts lie in \mathcal{G}_λ). *Let λ be a stage of the iteration. Let (E, D) be globally admissible at stage λ and let $\pi \in \text{Fix}(E)$. Then the iteration-level diagonal lift $\hat{\pi}^D \in \text{Aut}(\mathbb{P}_\lambda)$ (from Definition 5.27) belongs to \mathcal{G}_λ .*

Proof. For each $\beta \leq \lambda$, define $D_\beta := D \cap \text{Coords}_\beta$, the set of protected coordinates at stages $\leq \beta$.

Key observation from admissibility: Since (E, D) is globally admissible and $\pi \in \text{Fix}(E)$, we have $\text{Fix}(E) \leq \text{Stab}(d)$ for all $d \in D$, hence $\pi \cdot d = d$ for all $d \in D$. In particular, for each successor $\beta + 1 \leq \lambda$, the set $D_{\beta+1} \setminus D_\beta \subseteq D$

is fixed pointwise by π . This ensures the successor diagonal kernel element $\kappa_{\beta+1}(\pi, D_{\beta+1} \setminus D_\beta)$ is well-defined (Definition 5.23).

We construct by transfinite recursion a coherent sequence $\langle g_\beta : \beta \leq \lambda \rangle$ with $g_\beta \in \mathcal{G}_\beta$ such that $g_\beta = \hat{\pi}^{D_\beta}$ as automorphisms of \mathbb{P}_β . More precisely, we maintain the inductive invariant that for all $\gamma < \beta \leq \lambda$, $\pi_\beta^\gamma(g_\beta) = g_\gamma$. In particular, at a limit stage η the family $\langle g_\beta : \beta < \eta \rangle$ is coherent, so a direct-limit element exists.

Stage 0: Define $g_0 := \pi \in \text{Sym}(\omega_1) = \mathcal{G}_0$. Since $\text{Coords}_0 = \emptyset$, we have $D_0 = \emptyset$, so $\hat{\pi}^{D_0} = \hat{\pi}^\emptyset$, which is the standard π -action on the Cohen forcing (Definition 5.27, stage-0 clause). Hence $g_0 = \hat{\pi}^{D_0}$.

Successor $\beta + 1$: Assume $g_\beta \in \mathcal{G}_\beta$ with $g_\beta = \hat{\pi}^{D_\beta}$. Define

$$g_{\beta+1} := \text{spl}(g_\beta) \cdot \kappa_{\beta+1}(\pi, D_{\beta+1} \setminus D_\beta),$$

where spl is the splitting section from Lemma 5.21 and $\kappa_{\beta+1}$ is the successor diagonal kernel element from Definition 5.23 (applicable by the key observation above).

By Lemma 5.21, $\text{spl}(g_\beta) \in \mathcal{G}_{\beta+1}$. By Lemma 5.25(i), $\kappa_{\beta+1}(\pi, D_{\beta+1} \setminus D_\beta) \in \mathcal{G}_{\beta+1}$. Hence $g_{\beta+1} \in \mathcal{G}_{\beta+1}$.

We verify $g_{\beta+1} = \hat{\pi}^{D_{\beta+1}}$ by comparing actions (cf. Definition 5.27):

- *On the \mathbb{P}_β -part:* $\text{spl}(g_\beta)$ acts as $g_\beta = \hat{\pi}^{D_\beta}$, and $\kappa_{\beta+1}$ acts as identity. The combined action is $\hat{\pi}^{D_\beta}$, which agrees with $\hat{\pi}^{D_{\beta+1}}$ restricted to stages $\leq \beta$ (since $D_{\beta+1} \cap \text{Coords}_\beta = D_\beta$).
- *On new coordinates at stage $\beta + 1$:* $\text{spl}(g_\beta)$ acts trivially. The kernel element $\kappa_{\beta+1}(\pi, D_{\beta+1} \setminus D_\beta)$:
 - acts as π on $d \notin D_{\beta+1} \setminus D_\beta$, i.e., on new coordinates not in D ;
 - acts as identity on $d \in D_{\beta+1} \setminus D_\beta$, i.e., on new coordinates in D .

This matches Definition 5.27: “act as π if $d \notin D$, identity if $d \in D$.”

Limit η : Define g_η to be the direct-limit element of \mathcal{G}_η determined by the family $\langle g_\beta : \beta < \eta \rangle$, which is coherent by the inductive invariant above. By the limit-stage group definition ([3, Def. 3.1]), $g_\eta \in \mathcal{G}_\eta$.

By the uniqueness property of the inverse/direct limit presentation: elements of \mathcal{G}_η are determined by their projections to all earlier stages ([3, Def. 3.1]). Since g_η and $\hat{\pi}^{D_\eta}$ have identical restrictions to each \mathbb{P}_β for $\beta < \eta$ (namely, $g_\beta = \hat{\pi}^{D_\beta}$), we conclude $g_\eta = \hat{\pi}^{D_\eta}$.

Coherence: We verify $\pi_\beta^\gamma(g_\beta) = g_\gamma$ for $\gamma < \beta \leq \lambda$.

For successor $\beta + 1$:

$$\begin{aligned}\pi_{\beta+1}^\beta(g_{\beta+1}) &= \pi_{\beta+1}^\beta(\text{spl}(g_\beta) \cdot \kappa_{\beta+1}(\pi, D_{\beta+1} \setminus D_\beta)) \\ &= \pi_{\beta+1}^\beta(\text{spl}(g_\beta)) \cdot \pi_{\beta+1}^\beta(\kappa_{\beta+1}(\dots)) \\ &= g_\beta \cdot \text{id} = g_\beta,\end{aligned}$$

using $\pi_{\beta+1}^\beta \circ \text{spl} = \text{id}$ (Lemma 5.21) and $\kappa_{\beta+1}(\dots) \in K_{\beta+1} = \ker(\pi_{\beta+1}^\beta)$ (Lemma 5.25(ii)).

For limits η , coherence is automatic: g_η is defined as the direct-limit element, so $\pi_\eta^\gamma(g_\eta) = g_\gamma$ for all $\gamma < \eta$ by construction.

Conclusion: Taking $\beta = \lambda$, we have $g_\lambda = \hat{\pi}^{D_\lambda}$. Since $D_\lambda = D \cap \text{Coords}_\lambda = D$ (since $D \subseteq \text{Coords}_{<\lambda}$ by Definition 5.26(ii)), we have $g_\lambda = \hat{\pi}^D$ and $g_\lambda \in \mathcal{G}_\lambda$. Hence $\hat{\pi}^D \in \mathcal{G}_\lambda$. \square

Lemma 5.44 (Diagonal lifts lie in Δ^\uparrow). *Let $\pi \in \text{Fix}(E)$ and let $D' \supseteq D$ be such that the diagonal lift $\hat{\pi}^{D'} \in \mathcal{G}_\lambda$ exists. Then $\hat{\pi}^{D'} \in \Delta_\lambda^\uparrow(E, D)$.*

Proof. By the characterization definition of $\Delta_\lambda^\uparrow(E, D)$ (Definition 5.27), we verify:

- (i) $\pi_\lambda^0(\hat{\pi}^{D'}) \in \text{Fix}(E)$: By construction of diagonal lifts (Lemma 5.31(i)), the stage-0 projection of $\hat{\pi}^{D'}$ is π , and $\pi \in \text{Fix}(E)$ by assumption.
- (ii) $\hat{\pi}^{D'}$ fixes every coordinate in D pointwise: By construction (Lemma 5.31(ii)), $\hat{\pi}^{D'}$ fixes every coordinate in D' pointwise, and $D \subseteq D'$.

Hence $\hat{\pi}^{D'} \in \Delta_\lambda^\uparrow(E, D)$. \square

Lemma 5.45 (Package generic sections are hereditarily symmetric via diagonal cancellation). *Let f have countable support E (so $\text{Fix}(E) \leq \text{Stab}(f)$) and let \dot{s}_f be the canonical $\mathbb{S}_{[f]}$ -name for the f -coordinate generic section. Then $\Delta^\uparrow(E, \{f\}) \leq \text{Stab}(\dot{s}_f)$, hence $\dot{s}_f \in \text{HS}$ (with respect to $\mathcal{F}_{[f]}$).*

Proof. Let $E \subseteq \omega_1$ witness that f has countable support, i.e. $\text{Fix}(E) \leq \text{Stab}(f)$.

Fix any admissible pair (E', D') such that $E \subseteq E'$ and $\{f\} \subseteq D'$. Let $\pi \in \text{Fix}(E')$. Then the diagonal lift $\hat{\pi}^{D'} \in \Delta(E', D')$ fixes the f -coordinate pointwise: for every condition $\vec{p} \in \mathbb{S}_{[f]}$,

$$(\hat{\pi}^{D'} \cdot \vec{p})(f) = \vec{p}(f).$$

Since \dot{s}_f is definable (as a $\mathbb{S}_{[f]}$ -name) purely from the f -coordinate generic section, it follows that $\hat{\pi}^{D'} \cdot \dot{s}_f = \dot{s}_f$. Hence $\Delta(E', D') \leq \text{Stab}(\dot{s}_f)$.

As $\Delta^\uparrow(E, \{f\})$ is generated by all such subgroups $\Delta(E', D')$, we have $\Delta^\uparrow(E, \{f\}) \leq \text{Stab}(\dot{s}_f)$. Therefore $\text{Stab}(\dot{s}_f) \in \mathcal{F}_{[f]}$, so $\dot{s}_f \in \text{HS}$. \square

Lemma 5.46 (Diagonal cancellation makes right-inverse generics hereditarily symmetric). *Let g be a package with countable support witnessed by $E \subseteq \omega_1$, i.e. $\text{Fix}(E) \leq \text{Stab}(g)$. Let \dot{r}_g be the canonical $\mathbb{R}_{[g]}$ -name for the g -coordinate generic right inverse. Then $\Delta^\uparrow(E, \{g\}) \leq \text{Stab}(\dot{r}_g)$, and hence $\dot{r}_g \in \text{HS}$ (with respect to $(\mathbb{R}_{[g]}, \mathcal{G}_{[g]}, \mathcal{F}_{[g]})$).*

Proof. The proof is identical to Lemma 5.45. Fix an admissible pair (E', D') with $E \subseteq E'$ and $\{g\} \subseteq D'$. For $\pi \in \text{Fix}(E')$, the diagonal lift $\hat{\pi}^{D'} \in \Delta(E', D')$ fixes the g -coordinate pointwise, and \dot{r}_g is definable purely from the g -coordinate generic object. Hence $\hat{\pi}^{D'} \cdot \dot{r}_g = \dot{r}_g$, so $\Delta(E', D') \leq \text{Stab}(\dot{r}_g)$. Therefore $\Delta^\uparrow(E, \{g\}) \leq \text{Stab}(\dot{r}_g)$, as required. \square

Lemma 5.47 (Successor-step HS lift for package generics). *Fix a stage β and a package coordinate $d \in \text{NewCoords}_{\beta+1} = \text{Coords}_{\beta+1} \setminus \text{Coords}_\beta$ scheduled at stage β . (e.g. $d = f_\alpha$ for a splitting package $\mathbb{Q}_{[f_\alpha]}$, or $d = g_\alpha$ for a right-inverse package $\mathbb{R}_{[g_\alpha]}$). Let $\dot{\tau}$ be a canonical $\dot{\mathbb{Q}}_\beta$ -name definable purely from the d -coordinate generic object (e.g. \dot{s}_{f_α} or \dot{r}_{g_α}).*

Assume d has countable support witnessed by $E \subseteq \omega_1$ (so $\text{Fix}(E) \leq \text{Stab}(d)$ in M_β). Then the canonical $\mathbb{P}_{\beta+1}$ -name $\dot{\tau}^$ is hereditarily symmetric at stage $\beta+1$; in particular, $\dot{\tau}^{*G_{\beta+1}} \in M_{\beta+1} \subseteq \mathcal{M}$.*

Proof. Since d has countable support witnessed by E in M_β , we have $\text{Fix}(E) \leq \text{Stab}(d)$. Thus $(E, \{d\})$ is globally admissible at stage $\beta+1$ (Definition 5.26), so $\Delta_{\beta+1}^\uparrow(E, \{d\})$ is one of the Δ^\uparrow -generators of $\tilde{\mathcal{F}}_{\beta+1}^*$ (Definition 5.48).

Now let $g \in \Delta_{\beta+1}^\uparrow(E, \{d\})$. By the characterization in Definition 5.27, g fixes the coordinate d pointwise (Here “fixes the coordinate d pointwise” is exactly the clause in Definition 5.27: the induced automorphism on the d -factor forcing ($\dot{\mathbb{Q}}_d$ or $\dot{\mathbb{R}}_d$, depending on the package) is literally the identity, i.e. $g \cdot \dot{p} = \dot{p}$ for every canonical name \dot{p} for a condition in the d -factor.)

Since $\dot{\tau}^*$ is definable purely from the d -coordinate generic object, it follows that $g \cdot \dot{\tau}^* = \dot{\tau}^*$. Hence $\Delta_{\beta+1}^\uparrow(E, \{d\}) \leq \text{Stab}_{\mathcal{G}_{\beta+1}}(\dot{\tau}^*)$.

Therefore $\text{Stab}_{\mathcal{G}_{\beta+1}}(\dot{\tau}^*) \in \tilde{\mathcal{F}}_{\beta+1}^*$ by upward closure, so $\dot{\tau}^* \in \text{HS}_{\beta+1}$ and $\dot{\tau}^{*G_{\beta+1}} \in M_{\beta+1} \subseteq \mathcal{M}$. \square

5.3 The modified limit filter

We now define the filter $\tilde{\mathcal{F}}_\lambda^*$ that governs hereditary symmetry at limit stages of the iteration.

Definition 5.48 (Modified limit filter). Define the filter $\tilde{\mathcal{F}}_\lambda^*$ on \mathcal{G}_λ inductively:

Stage 0: $\tilde{\mathcal{F}}_0^*$ is the filter on $\mathcal{G}_0 = \text{Sym}(\omega_1)$ generated by $\{\text{Fix}(E) : E \subseteq \omega_1 \text{ countable}\}$. Note that $\text{Fix}(E) = \Delta_0^\uparrow(E, \emptyset)$, and (E, \emptyset) is trivially globally admissible.

Successor stage $\alpha + 1$: $\tilde{\mathcal{F}}_{\alpha+1}^*$ is the normal filter on $\mathcal{G}_{\alpha+1}$ generated by:

- (a) *Pushforwards:* $(\pi_{\alpha+1}^\alpha)^{-1}[H]$ for $H \in \tilde{\mathcal{F}}_\alpha^*$;
- (b) *Δ^\uparrow -subgroups:* $\Delta_{\alpha+1}^\uparrow(E, D)$ for countable $E \subseteq \omega_1$ and countable $D \subseteq \text{Coords}_{<\alpha+1}$ such that (E, D) is globally admissible at stage $\alpha + 1$.

Limit stage λ : $\tilde{\mathcal{F}}_\lambda^*$ is the normal filter on \mathcal{G}_λ generated by:

- (a) *Pushforwards:* $(\pi_\lambda^\beta)^{-1}[H]$ for $\beta < \lambda$ and $H \in \tilde{\mathcal{F}}_\beta^*$;
- (b) *Δ^\uparrow -subgroups:* $\Delta_\lambda^\uparrow(E, D)$ for countable $E \subseteq \omega_1$ and countable D (package coordinates at stages $< \lambda$) such that (E, D) is globally admissible at stage λ .

Remark 5.49 (Admissibility in the filter definition). We require global admissibility in the Δ^\uparrow -generator clause to ensure that the core lemma (Lemma 5.50) produces admissible pairs, which is needed for the properness proof (Lemma 5.52(iii)).

By Lemma 5.38, conjugation sends $\Delta_\lambda^\uparrow(E, D)$ to $\Delta_\lambda^\uparrow(g \cdot E, g \cdot D)$, and $g \cdot E$ and $g \cdot D$ remain countable (Lemma 5.30). The pair $(g \cdot E, g \cdot D)$ need not be globally admissible, but by Lemma 5.40 there is a countable $E^* \supseteq g \cdot E$ such that $(E^*, g \cdot D)$ is globally admissible and $\Delta_\lambda^\uparrow(E^*, g \cdot D) \leq \Delta_\lambda^\uparrow(g \cdot E, g \cdot D)$. Together with Corollary 5.39, this ensures that the normal filter generation process always contains Δ^\uparrow -groups with globally admissible parameters.

Lemma 5.50 (Every filter element contains a Δ^\uparrow -group). *For every $H \in \tilde{\mathcal{F}}_\lambda^*$, there exist countable $E \subseteq \omega_1$ and countable D (package coordinates at stages $< \lambda$) with (E, D) globally admissible such that:*

$$\Delta_\lambda^\uparrow(E, D) \leq H.$$

Proof. By definition of the normal filter generated by a family, every element $H \in \tilde{\mathcal{F}}_\lambda^*$ contains a finite intersection of conjugates of generators. We show that such intersections contain Δ^\uparrow -groups with globally admissible parameters.

Conjugates of Δ^\uparrow -generators: Let $\Delta_\lambda^\uparrow(E, D)$ be a globally admissible generator and let $g \in \mathcal{G}_\lambda$. By Lemma 5.38:

$$g \cdot \Delta_\lambda^\uparrow(E, D) \cdot g^{-1} = \Delta_\lambda^\uparrow(g \cdot E, g \cdot D).$$

Note that $g \cdot E$ and $g \cdot D$ are countable: $g \cdot E = \pi_\lambda^0(g)[E]$ is countable because bijections preserve cardinality, and similarly $g \cdot D$ is countable because the action on package coordinates is a bijection (Lemma 5.30(ii)). The parameters $(g \cdot E, g \cdot D)$ need not be globally admissible. However, by Lemma 5.40, there exists $E^* \supseteq g \cdot E$ such that $(E^*, g \cdot D)$ is globally admissible and $\Delta_\lambda^\uparrow(E^*, g \cdot D) \leq \Delta_\lambda^\uparrow(g \cdot E, g \cdot D)$.

Thus every conjugate of an admissible Δ^\uparrow -generator *contains* a globally admissible Δ^\uparrow -group.

Finite intersections: If $H_1 \supseteq \Delta_\lambda^\uparrow(E_1, D_1)$ and $H_2 \supseteq \Delta_\lambda^\uparrow(E_2, D_2)$ with (E_i, D_i) globally admissible, then by Corollary 5.35:

$$H_1 \cap H_2 \supseteq \Delta_\lambda^\uparrow(E_1 \cup E_2, D_1 \cup D_2).$$

By Corollary 5.39, $(E_1 \cup E_2, D_1 \cup D_2)$ is globally admissible.

Pullback generators: Let $H = (\pi_\lambda^\beta)^{-1}[K]$ with $K \in \tilde{\mathcal{F}}_\beta^*$. By induction on β , there exist countable admissible (E, D) with $D \subseteq \text{Coords}_\beta$ and $\Delta_\beta^\uparrow(E, D) \leq K$. By Lemma 5.42, $\Delta_\lambda^\uparrow(E, D) \leq H$. The pair (E, D) remains globally admissible at stage λ (the admissibility condition $\text{Fix}(E) \leq \text{Stab}(d)$ depends only on how d was introduced, not on which stage we evaluate it). \square

Corollary 5.51 (Admissibility preserved under countable unions). *If $\{(E_n, D_n) : n < \omega\}$ is a countable family of globally admissible pairs, then $(\bigcup_n E_n, \bigcup_n D_n)$ is globally admissible.*

Proof. For $d \in \bigcup_n D_n$, choose k with $d \in D_k$. Then $\text{Fix}(\bigcup_n E_n) \leq \text{Fix}(E_k) \leq \text{Stab}(d)$, where the last inequality holds by global admissibility of (E_k, D_k) . \square

Lemma 5.52 ($\tilde{\mathcal{F}}_\lambda^*$ is a proper normal filter). *For each stage λ :*

(i) $\tilde{\mathcal{F}}_\lambda^*$ is a filter (upward closed, closed under finite intersections).

(ii) $\tilde{\mathcal{F}}_\lambda^*$ is normal (closed under conjugation by \mathcal{G}_λ).

(iii) $\tilde{\mathcal{F}}_\lambda^*$ is proper ($\{\text{id}\} \notin \tilde{\mathcal{F}}_\lambda^*$).

(iv) $\tilde{\mathcal{F}}_\lambda^*$ is ω_1 -complete.

Proof. (i) and (ii) follow from the definition as a normal filter generated by the given families.

(iii) By Lemma 5.50, if $\{\text{id}\} \in \tilde{\mathcal{F}}_\lambda^*$, then $\{\text{id}\} \supseteq \Delta_\lambda^\uparrow(E, D)$ for some countable globally admissible E, D .

Since E is countable and ω_1 is uncountable, there exist distinct $\beta, \gamma \in \omega_1 \setminus E$. Let $\pi = (\beta \ \gamma)$ (transposition). Then $\pi \in \text{Fix}(E)$.

By Lemma 5.43, the diagonal lift $\hat{\pi}^D \in \mathcal{G}_\lambda$ exists. By Lemma 5.44, $\hat{\pi}^D \in \Delta_\lambda^\uparrow(E, D)$.

Since $\hat{\pi}^D$ acts as π on the Cohen base (Lemma 5.31(i)), it swaps coordinates β and γ , so $\hat{\pi}^D \neq \text{id}$.

Hence $\Delta_\lambda^\uparrow(E, D)$ contains a nontrivial element, contradicting $\Delta_\lambda^\uparrow(E, D) \leq \{\text{id}\}$.

(iv) ω_1 -completeness. Let $\{H_n : n < \omega\} \subseteq \tilde{\mathcal{F}}_\lambda^*$. By Lemma 5.50, for each n there exists a globally admissible (E_n, D_n) with $\Delta_\lambda^\uparrow(E_n, D_n) \leq H_n$.

By Corollary 5.35:

$$\bigcap_{n < \omega} H_n \supseteq \Delta_\lambda^\uparrow \left(\bigcup_n E_n, \bigcup_n D_n \right).$$

By Corollary 5.51, $(\bigcup_n E_n, \bigcup_n D_n)$ is globally admissible. Since countable unions of countable sets are countable (in the metatheory, which has AC), the right-hand side is a Δ^\uparrow -generator of the filter.

Hence $\bigcap_n H_n \in \tilde{\mathcal{F}}_\lambda^*$. □

Definition 5.53 (Hereditarily symmetric names). A \mathbb{P}_λ -name τ is *hereditarily symmetric* ($\tau \in \text{HS}_\lambda$) iff:

(i) $\text{Stab}_{\mathcal{G}_\lambda}(\tau) \in \tilde{\mathcal{F}}_\lambda^*$, and

(ii) for every $(\sigma, p) \in \tau$, we have $\sigma \in \text{HS}_\lambda$.

The *symmetric extension* at stage λ is $M_\lambda = \text{HS}_\lambda^{G_\lambda} = \{\tau^{G_\lambda} : \tau \in \text{HS}_\lambda\}$.

Lemma 5.54 (Δ -support equivalence). *For any \mathbb{P}_λ -name τ :*

$\text{Stab}(\tau) \in \tilde{\mathcal{F}}_\lambda^* \iff \exists \text{ countable globally admissible } (E, D) \text{ such that } \Delta_\lambda^\uparrow(E, D) \subseteq \text{Stab}(\tau)$.

Proof. (\Leftarrow) By upward closure, since $\Delta_\lambda^\uparrow(E, D) \in \tilde{\mathcal{F}}_\lambda^*$ for globally admissible (E, D) . (\Rightarrow) By Lemma 5.50. \square

Theorem 5.55 ($M_\lambda \models \text{ZF}$). *For each stage λ , the symmetric extension $M_\lambda = \text{HS}_\lambda^{G_\lambda}$ satisfies ZF.*

Proof. By Lemma 5.52, the triple $(\mathbb{P}_\lambda, \mathcal{G}_\lambda, \tilde{\mathcal{F}}_\lambda^*)$ is a symmetric system in the sense of [4, Ch. 15]. By the symmetric extension theorem [4, Lemma. 15.51], $M_\lambda \models \text{ZF}$. \square

Theorem 5.56 ($\mathcal{M} \models \text{ZF}$). *The final model $\mathcal{M} = \bigcup_{\alpha \in \text{Ord}} M_\alpha$ satisfies ZF.*

Proof. \mathcal{M} is the directed union of the transitive models $\{M_\alpha : \alpha \in \text{Ord}\}$, each satisfying ZF by Theorem 5.55.

For any ZF axiom instance with parameters $a_1, \dots, a_n \in \mathcal{M}$: choose λ large enough that $a_1, \dots, a_n \in M_\lambda$. Apply the axiom in M_λ to obtain the required witness in $M_\lambda \subseteq \mathcal{M}$.

The inclusion $M_\beta \subseteq M_\lambda$ for $\beta < \lambda$ is guaranteed by the pushforward generators: if $\tau \in \text{HS}_\beta$ with $\text{Stab}_{\mathcal{G}_\beta}(\tau) \in \tilde{\mathcal{F}}_\beta^*$, then $\text{Stab}_{\mathcal{G}_\lambda}(\tau) \supseteq (\pi_\lambda^\beta)^{-1}[\text{Stab}_{\mathcal{G}_\beta}(\tau)] \in \tilde{\mathcal{F}}_\lambda^*$, so $\tau \in \text{HS}_\lambda$. \square

5.4 Bookkeeping: names versus interpretations

Remark 5.57 (Background metatheory for class-length constructions). To speak formally about an Ord-length bookkeeping function and an Ord-length symmetric iteration as *class* objects, we work in a background theory of classes, e.g. GBC + ETR (optionally with Global Choice). Its set universe is denoted V and satisfies ZFC. All symmetric models \mathcal{N} , intermediate stages M_α , and the final model \mathcal{M} are constructed inside this set universe; only the bookkeeping map \mathcal{B} and the iteration itself are treated as class objects in the ambient background.

We now fix a bookkeeping device which ensures that every relevant surjection instance appearing in the final model is handled at some successor stage.

Remark 5.58 (Syntactic bookkeeping). In a class-length iteration, the interpretation of a fixed name can change as the iteration proceeds. Therefore, we bookkeep *names* (codes) rather than interpreted objects: at stage α we take the α -th code \dot{f} from our fixed enumeration, and in the current intermediate model M_α we check whether \dot{f} (as currently interpreted) is a surjection instance of the required type. If it is not, we force trivially at that

stage (or include only the other package factors scheduled for α). This is the standard bookkeeping convention for class-length forcing/iterations.

Definition 5.59 (Bookkeeping of orbit instances). Fix in the metatheory a definable class function $\mathcal{B} : \text{Ord} \rightarrow V$ that enumerates *pairs* (β, c) where c is a set-theoretic code for a potential \mathbb{P} -name (in some standard Gödel coding). The enumeration is arranged so that for each code c , the set

$$\{\alpha \in \text{Ord} : \exists \beta (\mathcal{B}(\alpha) = (\beta, c))\}$$

is unbounded in Ord (e.g. enumerate $\text{Ord} \times V$ using a definable pairing of ordinals together with a definable class well-order of V in the background metatheory).

At stage α , let $\mathcal{B}(\alpha) = (\beta, c)$. We *test* whether the code c :

- (i) decodes to a valid \mathbb{P}_α -name \dot{f} ,
- (ii) is hereditarily symmetric for the stage- α symmetric system $(\mathbb{P}_\alpha, \mathcal{G}_\alpha, \tilde{\mathcal{F}}_\alpha^*)$, and
- (iii) interprets in M_α to a surjection of one of the following types:
 - (1) $f : Y \rightarrow X$ with $X, Y \subseteq T$ (for PP-packages), or
 - (2) $f : S \times \eta \rightarrow \lambda$ with $\eta, \lambda \in \text{Ord}$ (for AC_{WO}-packages).

If all conditions are met and the surjection f does not already split in M_α (i.e., no section/right-inverse for f exists yet), the successor forcing at stage $\alpha + 1$ includes the corresponding orbit package $\mathbb{Q}_{[f]}$ or $\mathbb{R}_{[f]}$ (Definition 5.5); otherwise we force trivially at that code.

This arrangement ensures that every code is tested at unboundedly many stages, so a name that becomes a valid HS surjection at some stage β will eventually be scheduled at some stage $\alpha > \beta$.

Remark 5.60 (The $\aleph^*(S)$ horizon). For the AC_{WO}-component we schedule splittings for surjections $f : Y \rightarrow \lambda$ with $\lambda < \aleph^*(S)$ (cf. Remark 4.22). This is the natural bookkeeping horizon: $\aleph^*(S)$ is the first ordinal θ (in the ambient stage model under discussion) such that there is no surjection $S \rightarrow \theta$, so $\lambda < \aleph^*(S)$ is exactly the range of well-ordered indices reachable from S -parameterized data.

For our purposes we only use the following two points.

- (i) The seed parameter $S = A^\omega$ is preserved at successor stages: our package forcings add no new reals (Corollaries 4.19 and 4.30), hence add no new ω -sequences of Cohen reals and therefore no new elements of S .

- (ii) The value of $\aleph^*(S)$ need not be constant along the iteration (new functions from S may be added even if S itself is unchanged), but this causes no bookkeeping gap: we test the bound $\lambda < \aleph^*(S)$ in the current stage model when a code is examined. If at some stage a given name \dot{f} interprets to a surjection instance of the required form with $\lambda < \aleph^*(S)$ (as computed in that stage), then from that point onward the instance is eligible to be scheduled. Since every code is retested at unboundedly many stages (Definition 5.59), no a priori stability theorem for $\aleph^*(S)$ is needed.

5.5 The symmetric iteration (black box)

We now invoke the iteration framework via Remark 2.1. For the global conclusion (a final model of $\text{ZF} + \text{DC} + \text{PP} + \neg\text{AC}$), we take a *class-length* countable-support symmetric iteration of length Ord over \mathcal{N} . Concretely, working in the metatheory, we form an Ord -indexed symmetric iteration $\langle \mathbb{P}_\alpha, \mathcal{G}_\alpha, \tilde{\mathcal{F}}_\alpha^* : \alpha \in \text{Ord} \rangle$ as in [3, §2.1], with $\mathbb{P}_0 = \text{Add}(\omega, \omega_1)$ and intermediate symmetric models M_α .

At a successor stage $\alpha + 1$, we consult the finite list $\mathcal{B}(\alpha)$ from Definition 5.59. In the current model M_α , each code on this list either interprets to a genuine package instance (yielding a corresponding *orbit package* forcing $\mathbb{Q}_{[f]}$ or $\mathbb{R}_{[f]}$ as in Definition 5.5), or it does not (in which case we take the trivial forcing for that code). Let $\dot{\mathbb{S}}_\alpha$ be a \mathbb{P}_α -name for the finite-support product of the resulting orbit package posets. Then $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha * \dot{\mathbb{S}}_\alpha$.

At a successor stage $\alpha + 1$, the group $\mathcal{G}_{\alpha+1}$ and filter $\tilde{\mathcal{F}}_{\alpha+1}^*$ are those determined by the symmetric-iteration framework together with the diagonal infrastructure from Subsection 5.2 and the filter definition in Definition 5.48. Concretely, $\mathcal{G}_{\alpha+1}$ is generated by standard lifts of \mathcal{G}_α together with the (stage- α) package automorphisms and the successor diagonal kernel elements $\kappa_{\alpha+1}(\pi, D^{\text{new}})$ (Definition 5.23) acting on the new package coordinates, and $\tilde{\mathcal{F}}_{\alpha+1}^*$ is the normal filter generated by pushforwards from $\tilde{\mathcal{F}}_\alpha^*$ together with the admissible subgroups $\Delta_{\alpha+1}^\uparrow(E, D)$. In particular, for each active package coordinate d at stage $\alpha + 1$ with countable support witnessed by E_d , the generator $\Delta_{\alpha+1}^\uparrow(E_d, \{d\})$ belongs to $\tilde{\mathcal{F}}_{\alpha+1}^*$, so the canonical package generics are hereditarily symmetric by Lemmas 5.45 and 5.46.

Thus each successor filter explicitly includes the diagonal subgroups protecting the representative coordinates of the active packages.

At limit stages λ , we take the countable-support limit forcing \mathbb{P}_λ and the corresponding limit group/filter as provided by the symmetric-iteration framework (API items (b)–(e), especially filter generation/completeness at

limits, [3, Thm. 5.5]). We write \mathcal{M} for the final symmetric model of the Ord-length iteration.

Remark 5.61 (No one-step collapse in general). Unlike ordinary forcing iterations in ZFC, there is no general theorem showing that an iteration of symmetric extensions can always be presented as a *single* symmetric extension of the ground model. Indeed, the usual “two-step collapse” phenomena from forcing do not directly apply in the symmetric context, and the literature discusses serious obstacles to such a reduction. It remains open in general to what extent arbitrary symmetric iterations can be compressed to a single symmetric extension; see the discussion in [5, 14].

5.6 What the iteration yields

Let \mathcal{M} denote the final symmetric model obtained from the iteration.

Proposition 5.62 (Final localized splitting $\mathbb{P}\mathbb{P}^{\text{split}} \upharpoonright T$). $\mathcal{M} \models \mathbb{P}\mathbb{P}^{\text{split}} \upharpoonright T$.

Proof. Work in the final symmetric model \mathcal{M} . Let $f : Y \twoheadrightarrow X$ be a surjection in \mathcal{M} with $X, Y \subseteq T$.

Choose a hereditarily symmetric name \dot{f} such that $f = \dot{f}^{G_\infty}$. Let $I \subseteq \text{Ord}$ be the countable stage support of \dot{f} (API item (e)), and let $\alpha := \sup(I) + 1$. Then \dot{f} is a \mathbb{P}_α -name and $f_\alpha := \dot{f}^{G_\alpha} \in M_\alpha$. Since \dot{f} is a \mathbb{P}_α -name, its interpretation is unchanged by further forcing: $\dot{f}^{G_\infty} = \dot{f}^{G_\alpha}$, so $f = f_\alpha$ in \mathcal{M} .

If f_α already has a section in M_α , then f splits in \mathcal{M} and we are done. Otherwise, f_α is eligible for scheduling by the bookkeeping.

By the bookkeeping (Definition 5.59), at some successor stage we schedule the orbit package $\mathbb{Q}_{[f_\alpha]}$. Let \dot{s}_{f_α} be the canonical $\mathbb{Q}_{[f_\alpha]}$ -name for the generic section in the f_α -coordinate. By Lemma 5.45, \dot{s}_{f_α} is hereditarily symmetric for the *package* symmetric system $(\mathbb{Q}_{[f_\alpha]}, \mathcal{G}_{[f_\alpha]}, \mathcal{F}_{[f_\alpha]})$ in M_β . By Lemma 5.47, its canonical $\mathbb{P}_{\beta+1}$ -lift $\dot{s}_{f_\alpha}^*$ is hereditarily symmetric at stage $\beta + 1$, hence $s_{f_\alpha} := \dot{s}_{f_\alpha}^{*G_{\beta+1}} \in M_{\beta+1} \subseteq \mathcal{M}$.

Since s_{f_α} is a section of f_α , it remains a section of f in the final model \mathcal{M} . Therefore every $f : Y \twoheadrightarrow X \subseteq T$ in \mathcal{M} splits, i.e. $\mathcal{M} \models \mathbb{P}\mathbb{P}^{\text{split}} \upharpoonright T$. \square

Remark 5.63 (Countable-support names at limits of cofinality ω). At limits λ with $\text{cf}(\lambda) = \omega$, a hereditarily symmetric name may have countable support that is cofinal in λ , so stage-bounding arguments do not apply. The iteration machinery of [3] handles such limits using limit-filter ω_1 -completeness together with the appropriate block/factorization analysis (cf. [3, Rem. 3.9, Thm. 5.5]), and we rely on those limit-stage conclusions.

Proposition 5.64 (A is not well-orderable). *In the final model \mathcal{M} , the set $A = \{c_\alpha : \alpha < \omega_1\}$ of Cohen reals is not well-orderable. Hence $\mathcal{M} \models \neg \text{AC}$.*

Proof. Suppose for contradiction that A is well-orderable in \mathcal{M} . Let w be a well-ordering of A in \mathcal{M} .

Step 1: Locate w at a bounded stage.

Since $\mathcal{M} = \bigcup_{\lambda \in \text{Ord}} M_\lambda$ (Theorem 5.56), there exists an ordinal λ such that $w \in M_\lambda$. Fix a hereditarily symmetric \mathbb{P}_λ -name $\dot{w} \in \text{HS}_\lambda$ with $\dot{w}^{G_\lambda} = w$.

Step 2: Apply the core lemma.

Since $\dot{w} \in \text{HS}_\lambda$, we have $\text{Stab}_{\mathcal{G}_\lambda}(\dot{w}) \in \mathcal{F}_\lambda^*$. By Lemma 5.50, there exist countable $E \subseteq \omega_1$ and countable $D \subseteq \text{Coords}_{<\lambda}$ with (E, D) globally admissible such that

$$\Delta_\lambda^\uparrow(E, D) \leq \text{Stab}_{\mathcal{G}_\lambda}(\dot{w}).$$

Step 3: Choose a nontrivial transposition.

Since E is countable and ω_1 is uncountable, there exist distinct $\beta, \gamma \in \omega_1 \setminus E$. Let $\pi = (\beta \ \gamma) \in \text{Sym}(\omega_1)$ be the transposition swapping β and γ . Then $\pi \in \text{Fix}(E)$.

Step 4: Construct the diagonal lift.

By Lemma 5.43, since (E, D) is globally admissible and $\pi \in \text{Fix}(E)$, there exists a diagonal lift $\hat{\pi}^D \in \mathcal{G}_\lambda$ with $\pi_\lambda^0(\hat{\pi}^D) = \pi$.

By the characterization of Δ_λ^\uparrow (Definition 5.27):

- $\pi_\lambda^0(\hat{\pi}^D) = \pi \in \text{Fix}(E)$, and
- $\hat{\pi}^D$ fixes every coordinate in D pointwise.

Hence $\hat{\pi}^D \in \Delta_\lambda^\uparrow(E, D) \leq \text{Stab}_{\mathcal{G}_\lambda}(\dot{w})$.

Step 5: Derive the contradiction.

Since $\hat{\pi}^D \in \text{Stab}_{\mathcal{G}_\lambda}(\dot{w})$, we have $\hat{\pi}^D \cdot \dot{w} = \dot{w}$.

By equivariance of name interpretation ([4, Lemma 14.37]):

$$\hat{\pi}^D \cdot w = (\hat{\pi}^D \cdot \dot{w})^{G_\lambda} = \dot{w}^{G_\lambda} = w.$$

Thus w is invariant under the action of $\hat{\pi}^D$. This means that for all $x, y \in A$:

$$x <_w y \iff \hat{\pi}^D \cdot x <_w \hat{\pi}^D \cdot y.$$

Moreover $\hat{\pi}^D[A] = A$ (the diagonal lift permutes Cohen reals by Corollary 5.32), so the map $x \mapsto \hat{\pi}^D \cdot x$ is a bijection $A \rightarrow A$. Therefore the map $x \mapsto \hat{\pi}^D \cdot x$ is an order automorphism of the well-ordered set (A, w) .

By Lemma 3.22, any order automorphism of a well-ordered set is the identity. Hence $\hat{\pi}^D \cdot x = x$ for all $x \in A$.

However, by Corollary 5.32, $\hat{\pi}^D$ swaps $c_\beta \leftrightarrow c_\gamma$:

$$\hat{\pi}^D \cdot c_\beta = c_\gamma \neq c_\beta.$$

(The Cohen reals c_β and c_γ are distinct by standard properties of Cohen forcing; see Lemma 3.1.)

This contradicts $\hat{\pi}^D \cdot c_\beta = c_\beta$. Therefore A is not well-orderable in \mathcal{M} , and hence $\mathcal{M} \models \neg \text{AC}$. \square

Proposition 5.65 (DC preserved through the iteration). $\mathcal{M} \models \text{DC}$.

Proof. We argue by induction along the symmetric iteration. The base model $M_0 = \mathcal{N}$ satisfies DC by Theorem 3.11.

For the successor step, assume $M_\alpha \models \text{ZF} + \text{DC}$. The successor stage is obtained by forcing with the stage symmetric system at α and then taking the hereditarily symmetric submodel $M_{\alpha+1}$.

By Lemma 5.6, each orbit package $\mathbb{S}_{[f]}$ used at stage α is ω_1 -closed (and hence adds no new ω -sequences of ordinals), and the finite-support product $\dot{\mathbb{S}}_\alpha$ is again ω_1 -closed for the same reason. By Lemma 5.16, the relevant stage filter is ω_1 -complete. Therefore the DC-preservation theorem for symmetric extensions applies (API item (c), cf. [3, Thm. 3.20]), yielding $M_{\alpha+1} \models \text{DC}$.

At limit stages, DC is preserved by the countable-support symmetric iteration machinery (API item (c), and the limit-filter generation/completeness from API item (d); cf. [3, Thm. 3.20] and [3, Thm. 5.5] for the limit construction).

Thus every stage model satisfies DC, in particular the final model $\mathcal{M} \models \text{DC}$. \square

Definition 5.66 (Supported subforcing and supported names). Fix a stage λ and a globally admissible pair (E, D) at stage λ . Define

$$\mathbb{P}_\lambda \upharpoonright (E, D) = \{p \in \mathbb{P}_\lambda : p \upharpoonright 0 \in \text{Add}(\omega, E) \text{ and } \text{supp}_{>0}(p) \subseteq D\},$$

where $\text{supp}_{>0}(p) := \{c \in \text{Coords}_{<\lambda} : p(c) \neq 1_c\}$ is the set of nontrivial higher-stage coordinates of p .

A \mathbb{P}_λ -name \dot{x} is (E, D) -supported if every condition appearing in the transitive closure $\text{tc}(\dot{x})$ belongs to $\mathbb{P}_\lambda \upharpoonright (E, D)$.

Lemma 5.67 (Generic restriction to a supported subforcing). *Let (E, D) be globally admissible at stage λ , and let $\mathbb{Q} = \mathbb{P}_\lambda \upharpoonright (E, D)$. Then every*

maximal antichain $A \subseteq \mathbb{Q}$ is also a maximal antichain in \mathbb{P}_λ . Consequently, if $G_\lambda \subseteq \mathbb{P}_\lambda$ is V -generic, then $G_{E,D} := G_\lambda \cap \mathbb{Q}$ is V -generic for \mathbb{Q} , and $V[G_{E,D}] \subseteq V[G_\lambda]$.

Proof. Let $A \subseteq \mathbb{Q}$ be maximal, and let $p \in \mathbb{P}_\lambda$. Put $p' := p \upharpoonright (E, D) \in \mathbb{Q}$. Choose $q \in A$ compatible with p' in \mathbb{Q} , and let $r \in \mathbb{Q}$ witness this, i.e. $r \leq_{\mathbb{Q}} q, p'$. Since p extends p' on the coordinates that r constrains, and r is trivial off $E \cup D$, we can amalgamate r with p to obtain a common extension in \mathbb{P}_λ . Thus q is compatible with p in \mathbb{P}_λ , so A is predense in \mathbb{P}_λ and hence maximal there. Now G_λ meets every maximal antichain in \mathbb{P}_λ , so $G_\lambda \cap A \neq \emptyset$. Since $A \subseteq \mathbb{Q}$, this gives $(G_\lambda \cap \mathbb{Q}) \cap A \neq \emptyset$, proving \mathbb{Q} -genericity. \square

Lemma 5.68 (Supported names are \mathbb{Q} -names and evaluate correctly). *Let (E, D) be globally admissible at stage λ , let $\mathbb{Q} = \mathbb{P}_\lambda \upharpoonright (E, D)$, and let $G_\lambda \subseteq \mathbb{P}_\lambda$ be V -generic with $G_{E,D} = G_\lambda \cap \mathbb{Q}$. If \dot{x} is (E, D) -supported, then \dot{x} is (canonically) a \mathbb{Q} -name and*

$$\dot{x}^{G_\lambda} = \dot{x}^{G_{E,D}}.$$

Proof. Since every condition appearing in $\text{tc}(\dot{x})$ lies in \mathbb{Q} , we may view \dot{x} as a \mathbb{Q} -name (by the usual identification of names with the same transitive closure). The valuation equality is proved by induction on name-rank: for each pair $(\dot{y}, p) \in \dot{x}$ we have $p \in \mathbb{Q}$, hence $p \in G_\lambda$ iff $p \in G_{E,D}$, and the induction hypothesis applies to \dot{y} . \square

Lemma 5.69 (Triviality on supported names). *If \dot{x} is (E, D) -supported and $g \in \Delta_\lambda^\uparrow(E, D)$, then $g \cdot \dot{x} = \dot{x}$.*

Proof. By induction on $\text{rank}(\dot{x})$. If $p \in \mathbb{P}_\lambda \upharpoonright (E, D)$, then $g(p) = p$ since the stage-0 projection of g fixes E pointwise and g fixes every coordinate in D pointwise. Now apply the recursive definition of the action on names. \square

Lemma 5.70 (Persistence of $\text{SVC}(S)$). *Let S and T be as in Definition 3.24. Since $M_0 = \mathcal{N}$ and $\mathcal{N} \models \text{SVC}(S)$ by Proposition 3.23, we have $M_0 \models \text{SVC}(S)$. Then for every stage λ , $M_\lambda \models \text{SVC}(S)$, and consequently $M_\lambda \models \text{SVC}^+(T)$.*

Proof. We argue by transfinite induction on λ .

Base. $M_0 = \mathcal{N} \models \text{SVC}(S)$ by Proposition 3.23.

Successor. Assume $M_\alpha \models \text{SVC}(S)$ and let $X \in M_{\alpha+1}$ be nonempty. Choose a hereditarily symmetric $\mathbb{P}_{\alpha+1}$ -name $\dot{X} \in \text{HS}_{\alpha+1}$ (so $\dot{X} \in M_\alpha$) with $X = \dot{X}^{G_{\alpha+1}}$. By $\text{SVC}(S)$ in M_α , there exist an ordinal η and a surjection $e : S \times \eta \rightarrow \dot{X}$ in M_α .

Fix $p_0 \in \mathbb{P}_{\alpha+1}$ and a name \dot{x}_0 such that $p_0 \Vdash \dot{x}_0 \in \dot{X}$. Define a $\mathbb{P}_{\alpha+1}$ -name \dot{h} for a function $S \times \eta \rightarrow \dot{X}$ by: for $s \in S$ and $\xi < \eta$, write $e(s, \xi) = \langle \dot{y}, p \rangle$ and set

$$\dot{h}(s, \xi) = \begin{cases} \dot{y} & \text{if } p \in \dot{G}_{\alpha+1}, \\ \dot{x}_0 & \text{otherwise.} \end{cases}$$

Let $h := \dot{h}^{G_{\alpha+1}}$. Then $h : S \times \eta \rightarrow X$, and h is surjective by construction.

Moreover, $\dot{h} \in \text{HS}_{\alpha+1}$: it is definable from parameters in M_α , and any automorphism fixing \dot{X} (hence lying in $\tilde{\mathcal{F}}_{\alpha+1}^*$) fixes \dot{h} as well. Thus $h \in M_{\alpha+1}$, so $M_{\alpha+1} \models \text{SVC}(S)$.

Limit. Let λ be limit and assume $M_\beta \models \text{SVC}(S)$ for all $\beta < \lambda$. Let $X \in M_\lambda$ be nonempty. By API item (e), X has a hereditarily symmetric name with countable stage support bounded in λ , so $X \in M_\beta$ for some $\beta < \lambda$. By the inductive hypothesis, $M_\beta \models \text{SVC}(S)$, hence $M_\lambda \models \text{SVC}(S)$ as well.

Finally, $M_\lambda \models \text{SVC}^+(T)$ follows from Fact 4.3. \square

Proposition 5.71 (Final AC_{WO}). $\mathcal{M} \models \text{AC}_{\text{WO}}$.

Proof. Work in the final symmetric model \mathcal{M} . By Lemma 5.70, we have $\mathcal{M} \models \text{SVC}(S)$. We show that $\mathcal{M} \models \text{AC}_{<\aleph^*(S)^M}$; by the Ryan–Smith equivalences this implies AC_{WO}.

Let $\lambda < \aleph^*(S)^M$ be an ordinal and let $f : Y \twoheadrightarrow \lambda$ be a surjection in \mathcal{M} . By SVC(S), fix an ordinal η and a surjection $e : S \times \eta \twoheadrightarrow Y$ in \mathcal{M} . Let $g := f \circ e : S \times \eta \twoheadrightarrow \lambda$.

Let \dot{g} be a hereditarily symmetric name such that $g = \dot{g}^{G_\infty}$. Let $I \subseteq \text{Ord}$ be the countable stage support of \dot{g} (API item (e)), and let $\alpha := \sup(I) + 1$. Then \dot{g} is a \mathbb{P}_α -name, and $g_\alpha := \dot{g}^{G_\alpha} \in M_\alpha$. By the bookkeeping (Definition 5.59), there is some successor stage $\beta > \alpha$ at which the orbit package $\mathbb{R}_{[g_\alpha]}$ is scheduled. Let \dot{r}_{g_α} be the canonical $\mathbb{R}_{[g_\alpha]}$ -name for the generic right inverse in the g_α -coordinate. By Lemma 5.46, the canonical $\mathbb{R}_{[g_\alpha]}$ -name \dot{r}_{g_α} is hereditarily symmetric for the *package* symmetric system in M_β . By Lemma 5.47, its canonical $\mathbb{P}_{\beta+1}$ -lift $\dot{r}_{g_\alpha}^*$ is hereditarily symmetric at stage $\beta + 1$, hence $r_{g_\alpha} := \dot{r}_{g_\alpha}^{*G_{\beta+1}} \in M_{\beta+1} \subseteq \mathcal{M}$, and r_{g_α} is a right inverse of g in \mathcal{M} .

Now define $t := e \circ r_{g_\alpha} : \lambda \rightarrow Y$. Then for all $\xi < \lambda$,

$$f(t(\xi)) = f(e(r_{g_\alpha}(\xi))) = g(r_{g_\alpha}(\xi)) = \xi,$$

so t is a right inverse to f . Thus every surjection $Y \twoheadrightarrow \lambda$ with $\lambda < \aleph^*(S)^M$ splits in \mathcal{M} , and therefore $\mathcal{M} \models \text{AC}_{<\aleph^*(S)^M}$.

Finally, since $\mathcal{M} \models \text{SVC}(S)$, the Ryan–Smith theorem (see Remark 4.11) yields $\mathcal{M} \models \text{AC}_{\text{WO}}$. \square

Theorem 5.72 (Final model). *There is a transitive model $\mathcal{M} \models \text{ZF} + \text{DC} + \text{PP} + \neg\text{AC}$.*

Proof. By Theorem 5.56, $\mathcal{M} \models \text{ZF}$, and by Proposition 5.65, $\mathcal{M} \models \text{DC}$.

By Proposition 5.62, $\mathcal{M} \models \text{PP}^{\text{split}} \upharpoonright T$, hence $\mathcal{M} \models \text{PP} \upharpoonright T$. By Proposition 5.71, $\mathcal{M} \models \text{AC}_{\text{wo}}$. By Lemma 5.70, $\mathcal{M} \models \text{SVC}^+(T)$ for $T = \mathcal{P}(S)$, so the hypotheses of the Ryan–Smith localization theorem (Theorem 4.5, i.e., [13, Prop. 3.17] applied with Ryan–Smith’s parameter S taken to be our T) hold in \mathcal{M} . Therefore $\mathcal{M} \models \text{PP}$.

Finally, Proposition 5.64 gives $\mathcal{M} \models \neg\text{AC}$. \square

References

- [1] B. Banaschewski and G. Moore *The Dual Cantor–Bernstein Theorem and the Partition Principle* Notre Dame Journal of Formal Logic **31** (1990) no. 3, 375–381
- [2] A. Blass. *Injectivity, projectivity, and the axiom of choice.* Trans. Amer. Math. Soc. **255** (1979), 31–59. DOI: 10.1090/S0002-9947-1979-0542870-6.
- [3] F. T. Gilson. *Symmetric Iterations with Countable and $< \kappa$ -support.* arXiv:2511.07866v2, 2025. Available at <https://doi.org/10.48550/arXiv.2511.07866>.
- [4] T. Jech. *Set Theory*. Springer Monographs in Mathematics. Springer, 2003.
- [5] A. Karagila. *Iterating symmetric extensions.* J. Symbolic Logic **84** (2019), no. 1, 123–159. Available at <https://doi.org/10.1017/jsl.2018.73>
- [6] A. Karagila. *Preserving Dependent Choice.* Bull. Pol. Acad. Sci. Math. **67** (2019), 19–29. Available at <https://doi.org/10.48550/arXiv.1810.11301>.
- [7] A. Karagila and C. Ryan-Smith. *Which pairs of cardinals can be Hartogs and Lindenbaum numbers of a set?.* Fundamenta Mathematicae **267** 2024, 231–241. Available at <https://doi.org/10.4064/fm231006-14-8>.
- [8] A. Karagila and J. Schilhan. *Geometric condition for Dependent Choice.* Acta Math. Hungar. **172** (2024), 34–41. Available at <https://doi.org/10.1007/s10455-023-00960-1>

<https://doi.org/10.1007/s10474-024-01396-0> and <https://arxiv.org/abs/2212.10261>.

- [9] K. Kunen. *Set Theory*. Studies in Logic, vol. 34. College Publications, 2011.
- [10] D. Pincus. *Adding dependent choice*. Ann. Math. Logic **11** (1977), no. 1, 105–145. Available at [https://doi.org/10.1016/0003-4843\(77\)90011-0](https://doi.org/10.1016/0003-4843(77)90011-0).
- [11] B. Ransom. *On BPI in Symmetric Extensions, Part I*. arXiv:2511.21684v1, 2025. Available at <https://doi.org/10.48550/arXiv.2511.21684>.
- [12] C. Ryan-Smith. *The Hartogs–Lindenbaum spectrum of symmetric extensions*. Math. Logic Quarterly, 2024. Available at <https://doi.org/10.1002/malq.202300047> and <https://arxiv.org/abs/2309.12100>.
- [13] C. Ryan-Smith. *Local reflections of choice*. Acta Mathematica Hungarica, 2025. Available at <https://doi.org/10.1007/s10474-025-01533-3>.
- [14] E. C. Tachtsis. Review of A. Karagila, “*Iterating symmetric extensions*”. *MathSciNet* MR3922788, 2021. Available at <https://blogs.ams.org/beyondreviews/files/2021/04/3922788.pdf>.