

On a Smoothed Dirichlet Divisor Problem

Olivier Bordellès and Florian Daval

Abstract

Hardy showed that $\sum_{n \leq x} \tau(n) - x(\log x + 2\gamma - 1)$ is not $o(x^{1/4})$. In this article, we prove that $\sum_{n \leq x} \tau(n)(1 - \frac{x}{n}) - xP(\log x) = \frac{1}{4} + O\left(\frac{\log x}{x^{1/4}}\right)$, where P is a polynomial of degree 2. As a corollary, this estimate enables us to settle a conjecture surmised by Berkane, Bordellès, and Ramaré dealing with the positivity of an integral of the error term in the Dirichlet divisor problem. All results are entirely explicit and allow us to study the proximity between the remainder of the Dirichlet divisor problem and its logarithmic version.

1 Introduction

The multiplicative function which counts the number of divisors of n , denoted by $\tau(n)$ or sometimes $d(n)$, is a classical subject of study in number theory. By the convolution identity $\tau = \mathbf{1} \star \mathbf{1}$, we have $\sum_{n \leq x} \tau(n) = \sum_{n \leq x} \lfloor x/n \rfloor$, and therefore sums of fractional parts play a significant role in determining the order of magnitude of this sum. The ongoing conjecture is $\sum_{n \leq x} \tau(n) - x(\log x + 2\gamma - 1) = O(x^{1/4+\varepsilon})$ for all $\varepsilon > 0$, but seems to be currently out of reach. After applying the hyperbola principle, historically introduced by Dirichlet in this context, we obtain a shorter sum of fractional parts

$$\begin{aligned} \sum_{n \leq x} \tau(n) &= 2 \sum_{n \leq \sqrt{x}} \left\lfloor \frac{x}{n} \right\rfloor - (\lfloor \sqrt{x} \rfloor)^2 \\ &= 2xH(\sqrt{x}) - 2 \sum_{n \leq \sqrt{x}} \left(\left\{ \frac{x}{n} \right\} - \frac{1}{2} \right) + \lfloor \sqrt{x} \rfloor - (\lfloor \sqrt{x} \rfloor)^2 \end{aligned}$$

where H is the harmonic sum. The heart of the divisor problem lies in estimating the sum

$$\sum_{n \leq \sqrt{x}} \psi\left(\frac{x}{n}\right) \tag{1}$$

where $\psi(x) := \{x\} - \frac{1}{2}$, and Voronoï showed that (1) does not exceed $\ll x^{1/3} \log x$. The best result to date is due to Li and Yang [8] who proved that, for x sufficiently large and all $\varepsilon > 0$, we have

$$\sum_{n \leq \sqrt{x}} \psi\left(\frac{x}{n}\right) \ll x^{\theta+\varepsilon}$$

with $\theta = \frac{1646}{6881} + \frac{25\sqrt{1717}}{13762} \doteq 0.31448\dots$, slightly improving on the previous record held by Huxley in [7]. In [4], the following logarithmic version is investigated to obtain other explicit results with an application for class numbers of quartic number fields

$$\left| \sum_{n \leq x} \frac{\tau(n)}{n} - \left(\frac{1}{2} \log^2 x + 2\gamma \log x + \gamma^2 - 2\gamma_1 \right) \right| \ll \frac{1.001}{x^{1/2}} \quad (x \geq 2). \tag{2}$$

Applying the hyperbola principle again, we derive

$$x \sum_{n \leq x} \frac{\tau(n)}{n} = 2x \sum_{n \leq \sqrt{x}} \frac{H(x/n)}{n} - x(H(\sqrt{x}))^2.$$

Under the conjecture $O(x^{1/4+\varepsilon})$ aforementioned, a summation by parts gives an error term equal to $O(x^{1/4+\varepsilon})$ for $\sum \tau(n)(1 - x/n)$ and $\sum \tau(n)(x/n)$. But fractional parts also appear in the harmonic sum, since

$$H(x/n)(x/n) = (x/n) \log(x/n) + \gamma x/n - \psi(x/n) + O(n/x). \tag{3}$$

So the sum $\sum_{n \leq x} \tau(n)(1 - x/n)$ contains no fractional part in the main term. The difficult part of the divisor problem is thus eliminated in this smooth version, and we shall prove below that the remainder is unconditionally $\frac{1}{4} + O(x^{-1/4} \log x)$. This error term is analysed in a first part using the Euler-Maclaurin formula, and in a second part with exponential sums. The number $\frac{1}{4}$ can be seen as a residue since the Dirichlet series of $\tau(n)$ is $\zeta(s)^2$ and $\zeta(0)^2 = \frac{1}{4}$.

Theorem 1. *For all real numbers $x \geq 1$, define*

$$\Delta(x) := \sum_{n \leq x} \tau(n) - x(\log x + 2\gamma - 1)$$

and $\delta(x) := \sum_{n \leq x} \frac{\tau(n)}{n} - \left(\frac{1}{2} \log^2 x + 2\gamma \log x + \gamma^2 - 2\gamma_1 \right)$.

Then

$$\Delta(x) - x\delta(x) = x \int_x^\infty \frac{\Delta(u)}{u^2} du = \frac{1}{4} + r(x)$$

where

$$|r(x)| \leq \begin{cases} \frac{1}{8} + \frac{0.316}{\sqrt{x}} + \frac{1}{64x}, & \text{if } x \geq 1; \\ \frac{\log x}{x^{1/4}} + \frac{0.236}{\sqrt{x}} + \frac{1}{64x}, & \text{if } x \geq 300. \end{cases}$$

Incidentally, this result solves unconditionally [2, Conjecture 1.4] with the exponent $\frac{7}{4}$ replaced by 2 as proposed in [2, Question 8.7]. See [2, Section 8] for more background on this conjecture. It should be pointed out [14] that Mahatab and Mukhopadhyay have settled asymptotically this conjecture. See also [9, Theorem 1.1] for a related topic.

Corollary 1. *For all $x \geq 1$,*

$$\int_x^\infty \frac{\Delta(u)}{u^2} du \geq 0.$$

It should be stressed that removing the sum (1) is not sufficient to get the above result. Indeed, without taking precautions, the term $O(1)$ becomes $O(\log x)$. To obtain the term $\frac{1}{4} + o(1)$, it is necessary to have at our disposal some effective results of the Chowla-Walum conjecture. Since the orders of magnitude in the Dirichlet divisors problem are between $x^{1/4}$ and $x^{1/2}$, we can convert explicit results between $\sum_{n \leq x} \tau(n)$ and $\sum_{n \leq x} \tau(n)/n$ with almost no loss. For example, Hardy's result $\Delta(x) \neq o(x^{1/4})$ transfers directly to the logarithmic version, whereas integration by parts would not have succeeded. Let us give another example. The left-hand side of (4) below comes from [2, Lemma 7.1], so that Theorem 1 yields at once the conversion

$$|\Delta(x)| \leq 0.397 x^{1/2} \implies |\delta(x)| \leq \frac{0.397}{x^{1/2}} + \frac{0.38}{x} \quad (4)$$

valid for all $x \geq 5560$, without having to do any computer calculations. Furthermore, this improves the result (2) by a factor 2, while a summation by parts of (4) would only yield $3 \times 0.397/x^{1/2}$.

To conclude this introduction, let us have a look if we replace the Dirichlet divisor function τ by the Möbius function μ . Let $M(t) = \sum_{n \leq t} \mu(n)$ be the Mertens function, and $m(t) = \sum_{n \leq t} \mu(n)/n$ be its logarithmic version. The Dirichlet series attached to the function μ is $1/\zeta(s)$, and we have the residue $1/\zeta(0) = -2$. Nevertheless, we have in this case

$$M(x) - xm(x) = x \int_x^\infty \frac{M(u)}{u^2} du = -2 + r(x)$$

with $r(x) \neq o(x^{1/2})$ and $M(x) - xm(x) < 0$ for $x \leq 18349$. These oscillations are due to the existence of nontrivial zeros of the Riemann zeta function with real part $\frac{1}{2}$. Regarding the proximity of the two summation functions, we have the following result in [5]

$$\frac{2}{3} \leq \frac{\sup_{t \leq x} |m(t)|t}{\sup_{t \leq x} |M(t)|} \leq \frac{3}{2} \quad (x \geq 94).$$

2 Proofs

2.1 Notations

In what follows, we will use the first three Bernoulli polynomials $B_j(x)$:

$$\begin{array}{cccc} j & 1 & 2 & 3 \\ B_j(x) & x - \frac{1}{2} & x^2 - x - \frac{1}{6} & x^3 - \frac{3x^2}{2} + \frac{x}{2} \end{array}$$

and their attached periodic Bernoulli functions $B_j(\{x\})$. Notice that it is customary to denote $B_1(\{x\}) := \psi(x) = \{x\} - \frac{1}{2}$ the 1st periodic Bernoulli function. We also will use the classical exponential notation $e(x) := e^{2i\pi x}$, and $H(x) := \sum_{n \leq x} \frac{1}{n}$ will be the harmonic sum. For all $x \geq 1$, define $R_1(x)$ and $R_2(x)$ to be respectively the error terms in the following asymptotic formulas

$$H(x) = \log x + \gamma - \frac{\psi(x)}{x} + R_1(x)$$

and

$$\sum_{n \leq x} \frac{1}{n} \log \left(\frac{x}{n} \right) = \frac{1}{2} \log^2 x + \gamma \log x - \gamma_1 + R_2(x)$$

where $\gamma \doteq 0.5772\dots$ is the Euler-Mascheroni constant and $\gamma_1 \doteq -0.0728\dots$ is the 1st Riemann-Stieltjes constant.

2.2 Basic explicit estimates

2.2.1 Estimates using the Euler-Maclaurin summation formula

Lemma 1. *For all $x \geq 1$, we have*

$$|R_1(x)| \leq \frac{1}{8x^2}.$$

Proof. Applying the Euler-Maclaurin summation formula [1, p. 16] we have the integral representation

$$R_1(x) = \int_x^\infty \frac{\psi(t)}{t^2} dt.$$

Define $\psi_2(x) := \int_1^x \psi(t) dt = \frac{1}{2}(\psi(x))^2 - \frac{1}{8}$. Integrating by parts, we derive

$$x^2 R_1(x) = -\psi_2(x) + 2x^2 \int_x^\infty \frac{\psi_2(t)}{t^3} dt. \quad (5)$$

Since $-\frac{1}{8} \leq \psi_2(x) \leq 0$, we get

$$-\frac{1}{8} \leq 2x^2 \int_x^\infty \frac{\psi_2(t)}{t^3} dt \leq 0$$

which proves $x^2 |R_1(x)| \leq \frac{1}{8}$ as claimed. \square

Lemma 2. *For all $x \geq 1$, we have*

$$|R_2(x)| \leq \frac{0.132}{x^2}$$

and

$$|(R_2 - R_1)(x)| \leq \frac{0.033}{x^3}.$$

Proof. By [1, equations 10 and 11 p. 38, ex. 34 p. 41 (correction p. 113)] we have

$$R_2(x) = \int_x^\infty \frac{\psi(t)}{t^2} \left(1 + \log \left(\frac{x}{t} \right) \right) dt$$

so that $x^2 |R_2(x)| \leq \frac{1}{8}(1 + e^{-3}) < 0.132$ for all $x \geq 1$. Now, an integration by parts gives

$$(R_2 - R_1)(x) = \int_x^\infty \frac{\psi(t) \log(x/t)}{t^2} dt = -\frac{1}{2} \int_x^\infty B_2(\{t\}) g(t) dt$$

where $g(t) := (1 + 2\log(\frac{x}{t}))/t^3$. Following the book [1], we divide the integral so that the function g is monotonic with a constant sign to apply the 2nd mean value theorem. Since $\int_1^z B_2(\{t\}) dt = \frac{1}{3}B_3(\{z\})$ and $c := \sup_{z \geq 0} |B_3(\{z\})| = \frac{\sqrt{3}}{36}$, we derive $\left| \int_{z_1}^{z_2} B_2(\{t\}) dt \right| \leq \frac{2c}{3}$, and hence the 2nd mean value theorem yields

$$\begin{aligned} \left| \int_x^{e^{1/2}x} B_2(\{t\})g(t) dt \right| &\leq \frac{2c}{3} g(x) = \frac{2c}{3x^3} \\ \left| \int_{e^{1/2}x}^{e^{5/2}x} B_2(\{t\})g(t) dt \right| &\leq -\frac{2c}{3} g(e^{5/2}x) = \frac{c(1+5e^{-5})}{3x^3} \\ \left| \int_{e^{5/2}x}^{\infty} B_2(\{t\})g(t) dt \right| &\leq -\frac{2c}{3} g(e^{5/2}x) = \frac{c(1+5e^{-5})}{3x^3}. \end{aligned}$$

Therefore, we finally get

$$x^3 |(R_2 - R_1)(x)| \leq \frac{x^3}{2} \left| \int_x^{\infty} B_2(\{t\})g(t) dt \right| \leq \frac{c}{3}(2 + 5e^{-5}) \leq 0.033$$

completing the proof. \square

2.2.2 Estimates for the divisor function

The following result is [2, Lemma 3.1].

Lemma 3. *For all $x \geq 1$, we have*

$$\sum_{n \leq x} \tau(n) = x(\log x + 2\gamma - 1) - 2 \sum_{n \leq \sqrt{x}} \psi\left(\frac{x}{n}\right) + \frac{1}{4} + 2xR_1(\sqrt{x}) - (\psi(\sqrt{x}))^2.$$

We now derive a similar estimate for the logarithmic sum.

Lemma 4. *For all $x \geq 1$, we have*

$$\begin{aligned} x \sum_{n \leq x} \frac{\tau(n)}{n} &= x(\log(\sqrt{x}) + \gamma)^2 - x(R_1(\sqrt{x}))^2 + 2\sqrt{x}R_1(\sqrt{x})\psi(\sqrt{x}) - (\psi(\sqrt{x}))^2 \\ &\quad + 2x \sum_{k \leq \sqrt{x}} \frac{1}{k} \log\left(\frac{\sqrt{x}}{k}\right) - 2 \sum_{k \leq \sqrt{x}} \psi\left(\frac{x}{k}\right) + 2x \sum_{k \leq \sqrt{x}} \frac{1}{k} R_1\left(\frac{x}{k}\right). \end{aligned}$$

Proof. Let $1 \leq y \leq x$ be a parameter at our disposal. Dirichlet's hyperbola principle yields

$$\sum_{n \leq x} \frac{\tau(n)}{n} = \sum_{k \leq y} \frac{1}{k} H\left(\frac{x}{k}\right) + \sum_{k \leq x/y} \frac{1}{k} H\left(\frac{x}{k}\right) - H(y)H\left(\frac{x}{y}\right).$$

Now

$$\begin{aligned} \sum_{k \leq y} \frac{1}{k} H\left(\frac{x}{k}\right) &= \sum_{k \leq y} \frac{1}{k} \log\left(\frac{x}{k}\right) + \gamma H(y) - \frac{1}{x} \sum_{k \leq y} \psi\left(\frac{x}{k}\right) + \sum_{k \leq y} \frac{1}{k} R_1\left(\frac{x}{k}\right) \\ &= \sum_{k \leq y} \frac{1}{k} \log\left(\frac{y}{k}\right) + \left(\log\left(\frac{x}{y}\right) + \gamma\right) H(y) - \frac{1}{x} \sum_{k \leq y} \psi\left(\frac{x}{k}\right) + \sum_{k \leq y} \frac{1}{k} R_1\left(\frac{x}{k}\right), \end{aligned}$$

and similarly

$$\sum_{k \leq x/y} \frac{1}{k} H\left(\frac{x}{k}\right) = \sum_{k \leq x/y} \frac{1}{k} \log\left(\frac{x/y}{k}\right) + (\log y + \gamma) H\left(\frac{x}{y}\right) - \frac{1}{x} \sum_{k \leq x/y} \psi\left(\frac{x}{k}\right) + \sum_{k \leq x/y} \frac{1}{k} R_1\left(\frac{x}{k}\right).$$

Note that

$$H(y) \left(\log\left(\frac{x}{y}\right) + \gamma \right) + H\left(\frac{x}{y}\right) (\log y + \gamma) - H(y)H\left(\frac{x}{y}\right) = (\log y + \gamma) \left(\log\left(\frac{x}{y}\right) + \gamma \right) - \mathfrak{R}_1(y)\mathfrak{R}_1\left(\frac{x}{y}\right)$$

where $\mathfrak{R}_1(t) := R_1(t) - \frac{\psi(t)}{t}$. Hence

$$\begin{aligned} \sum_{n \leq x} \frac{\tau(n)}{n} &= (\log y + \gamma) \left(\log \left(\frac{x}{y} \right) + \gamma \right) - \mathfrak{R}_1(y) \mathfrak{R}_1 \left(\frac{x}{y} \right) \\ &\quad + \sum_{k \leq y} \frac{1}{k} \log \left(\frac{y}{k} \right) - \frac{1}{x} \sum_{k \leq y} \psi \left(\frac{x}{k} \right) + \sum_{k \leq y} \frac{1}{k} R_1 \left(\frac{x}{k} \right) \\ &\quad + \sum_{k \leq x/y} \frac{1}{k} \log \left(\frac{x/y}{k} \right) - \frac{1}{x} \sum_{k \leq x/y} \psi \left(\frac{x}{k} \right) + \sum_{k \leq x/y} \frac{1}{k} R_1 \left(\frac{x}{k} \right). \end{aligned}$$

Choosing $y = \sqrt{x}$ yields the desired result. \square

Lemma 5. *For all $x \geq 1$, we have*

$$\left| 2x \sum_{k \leq \sqrt{x}} \frac{1}{k} R_1 \left(\frac{x}{k} \right) \right| \leq 0.125 \left(1 + \frac{1}{\sqrt{x}} \right).$$

Proof. Lemma 1 implies $|R_1(x/k)| \leq 0.125 k^2/x^2$, and $\sum_{n=1}^N 2n = N^2 + N$ yields the bound $2 \sum_{k \leq \sqrt{x}} k \leq x + \sqrt{x}$. \square

The largest contribution in the $O(1)$ -term in Theorem 1 comes from the 0.125 above, and the main aim of the next section is to replace it by a $o(1)$. This will be achieved in Lemma 10 below.

2.3 Explicit estimates of generalized Chowla-Walum sums

In this section, we will use the following notation: for $x > 1$, $\alpha > 1$, $\beta \in \mathbb{Z}_{\geq 0}$ and $j \in \mathbb{Z}_{\geq 1}$, set

$$G_{\alpha, \beta, j}(x) := \sum_{n \leq x^{1/\alpha}} n^\beta B_j \left(\left\{ \frac{x}{n} \right\} \right).$$

2.3.1 Results

Proposition 1. *Let $1 < \alpha < 3$, $\beta \in \mathbb{Z}_{\geq 0}$ and $j \in \mathbb{Z}_{\geq 2}$. Set $\Gamma_j := \frac{2\eta(j)j!}{(2\pi)^j}$ where $\eta(j) = \zeta(j)$ if j is even, $\eta(j) = 1$ otherwise. Then, for all $x \geq 1$, we have*

$$|G_{\alpha, \beta, j}(x)| \leq 4\Gamma_j \left(\left(\zeta \left(j - \frac{1}{2} \right) + \frac{1}{4} \right) x^{\frac{\beta}{\alpha} - \frac{1}{2\alpha} + \frac{1}{2}} + \zeta \left(j + \frac{1}{2} \right) x^{\frac{\beta}{\alpha} + \frac{3}{2\alpha} - \frac{1}{2}} \right) \mathcal{L}_{\alpha, \beta}(x)$$

where $\mathcal{L}_{\alpha, \beta}(x) := \frac{3 - \alpha}{2\alpha(\beta + 1)} \frac{\log x}{\log 2} + 1$.

Remark 1. When $\alpha \geq 3$, the trivial bound $|G_{\alpha, \beta, j}(x)| \leq \Gamma_j x^{\frac{\beta+1}{\alpha}}$ is better than the result above. Applying to $(\alpha, \beta, j) = (2, 1, 2)$, we derive the next estimate.

Corollary 2. *For all $x \geq 300$, we have*

$$\left| \sum_{n \leq x^{1/2}} n B_2 \left(\left\{ \frac{x}{n} \right\} \right) \right| < x^{3/4} \log x.$$

2.3.2 Tools

Lemma 6 (Kusmin-Landau). *Let $N < N_1 \leq 2N$ be positive integers and $f \in C^1[N, N_1]$ such that f' is non-decreasing and that there exist $c_1 \geq 1$ and $0 < \lambda_1 < 1$ such that, for all $x \in [N, N_1]$, we have*

$$k + \lambda_1 \leq f'(x) \leq k + 1 - \lambda_1 \quad (k \in \mathbb{Z}). \quad (6)$$

Then

$$\left| \sum_{N < n \leq N_1} e(\pm f(n)) \right| \leq \frac{2}{\pi \lambda_1}.$$

Proof. See [3, Corollary 6.2]. \square

Lemma 7. Let $N < N_1 \leq 2N$ be positive integers and $f \in C^2[N, N_1]$ such that there exists a real number $\lambda_2 \in (0, \pi^{-1})$ such that, for all $x \in [N, N_1]$, we have $f''(x) \geq \lambda_2$. Assume that $f'(x) \notin \mathbb{Z}$ for all $x \in (N, N_1)$. Then

$$\left| \sum_{N < n \leq N_1} e(\pm f(n)) \right| \leq \frac{4}{\sqrt{\pi \lambda_2}}.$$

Proof. First note that, under the hypotheses of the lemma, the function f' is continuous and strictly increasing on $[N, N_1]$, and hence is one-to-one. Define $u := f'(N)$ and $v := f'(N_1)$, so that $f'([N, N_1]) = [u, v]$. Since $f'(x) \notin \mathbb{Z}$ for all $x \in (N, N_1)$, $f'([N, N_1]) \subseteq ([u], [u] + 1)$. Let $M_1, M_2 \in (N, N_1)$ be real numbers such that $f'(M_1) = u + t$ and $f'(M_2) = v - t$, where $t \in (0, 1)$ is a parameter at our disposal. Split the sum into 3 subsums:

$$\sum_{N < n \leq N_1} e(f(n)) = \sum_{N < n \leq M_1} e(f(n)) + \sum_{M_1 < n \leq M_2} e(f(n)) + \sum_{M_2 < n \leq N_1} e(f(n))$$

and we treat the 1st and 3rd sums trivially. The mean value theorem implies that there exists a real number $c \in (N, 2N)$ such that

$$\begin{aligned} \left| \sum_{N < n \leq M_1} e(f(n)) \right| &\leq \max(M_1 - N, 1) = \max\left(\frac{f'(M_1) - f'(N)}{f''(c)}, 1\right) \\ &\leq \max\left(\frac{u + t - u}{\lambda_2}, 1\right) = \max\left(\frac{t}{\lambda_2}, 1\right). \end{aligned}$$

Similarly, for some $d \in (N, 2N)$, we derive

$$\left| \sum_{M_2 < n \leq N_1} e(f(n)) \right| \leq \max\left(\frac{f'(N_1) - f'(M_2)}{f''(d)}, 1\right) \leq \max\left(\frac{t}{\lambda_2}, 1\right).$$

The 2nd sum is treated with Lemma 6. Note that f' is increasing and, for all $x \in [M_1, M_2]$, we have

$$\lfloor u \rfloor + t \leq u + t \leq f'(x) \leq v - t \leq \lfloor u \rfloor + 1 - t,$$

so that Lemma 6 applies with (6) used with $\lambda_1 = t$ and $k = \lfloor u \rfloor$, and yields

$$\left| \sum_{M_1 < n \leq M_2} e(f(n)) \right| \begin{cases} = 0, & \text{if } \lfloor M_2 \rfloor - M_1 < 1; \\ \leq \max\left(1, \frac{2}{\pi t}\right), & \text{if } \lfloor M_2 \rfloor - M_1 \geq 1. \end{cases}.$$

Therefore, we deduce that

$$\left| \sum_{N < n \leq N_1} e(\pm f(n)) \right| \leq 2 \max\left(\frac{t}{\lambda_2}, 1\right) + \max\left(1, \frac{2}{\pi t}\right)$$

and the required bound follows by choosing $t = \sqrt{\lambda_2 \pi^{-1}}$, and noticing that, with this choice of t and the condition $\lambda_2 < \pi^{-1}$, we have $t < \pi^{-1}$ and hence $\frac{2}{\pi t} > 2$. \square

We now derive an explicit version of the Van der Corput inequality.

Lemma 8 (van der Corput). Let $N < N_1 \leq 2N$ be positive integers and $f \in C^2[N, N_1]$ such that there exist two real numbers $c_2 \geq 1$ and $\lambda_2 > 0$ such that, for all $x \in [N, N_1]$, we have

$$\lambda_2 \leq f''(x) \leq c_2 \lambda_2.$$

Then

$$\left| \sum_{N < n \leq N_1} e(\pm f(n)) \right| \leq \frac{4}{\sqrt{\pi}} \left(c_2 N \lambda_2^{1/2} + 2 \lambda_2^{-1/2} \right).$$

Proof. If $\lambda_2 \geq \pi^{-1}$, then

$$\frac{4N\lambda_2^{1/2}}{\sqrt{\pi}} \geq \frac{4N}{\pi} > N \geq N_1 - N \geq \left| \sum_{N < n \leq N_1} e(f(n)) \right|$$

so that we may suppose $\lambda_2 < \pi^{-1}$. We pick the numbers u and v of the proof of Lemma 7 and set

$$[u, v] \cap \mathbb{Z} := \{m + 1, \dots, m + K\}$$

where $m \in \mathbb{Z}$ et $K \in \mathbb{Z}_{\geq 1}$. For all integers $k \in \{1, \dots, K + 1\}$, define

$$J_k := (f')^{-1} \left((m + k - 1, m + k] \cap [u, v] \right) \cap \mathbb{Z}.$$

Note that $f'(x) \notin \mathbb{Z}$ whenever $x \in \overset{\circ}{J}_k$, so that Lemma 7 implies that

$$\left| \sum_{N < n \leq N_1} e(f(n)) \right| \leq \sum_{k=1}^{K+1} \left| \sum_{n \in J_k} e(f(n)) \right| \leq \frac{4(K+1)}{\sqrt{\pi\lambda_2}}$$

and, by the mean value theorem, we have

$$K - 1 \leq v - u = f'(N_1) - f'(N) \leq c_2(N_1 - N)\lambda_2 \leq c_2 N \lambda_2$$

completing the proof. \square

Remark 2. A quite similar result has been proven in [11, Lemma 2.10], but with a different method. The coefficient of the main term of our result is slightly weaker, but we have improved that of the secondary term.

2.3.3 Proof of Proposition 1

We start by recording a well-known estimate for the periodic Bernoulli functions.

Lemma 9. Let $j \in \mathbb{Z}_{\geq 2}$ and set $\Gamma_j := \frac{2\eta(j)j!}{(2\pi)^j}$ where $\eta(j) = \zeta(j)$ if j is even, $\eta(j) = 1$ otherwise. Then

$$\sup_{x \in \mathbb{R}} |B_j(\{x\})| \leq \Gamma_j.$$

Proof. When $j = 2h$ is even, $|B_{2h}(\{x\})| \leq |B_{2h}|$ for all $x \in \mathbb{R}$ and $h \in \mathbb{Z}_{\geq 1}$, and the desired bound follows from [12, (9.6) p. 17]. When j is odd, a result due to Lehmer (see [10, Exercise 5.(e) p. 504] for instance) shows that, for all $x \in \mathbb{R}$ and $h \in \mathbb{Z}_{\geq 1}$, we have

$$|B_{2h+1}(\{x\})| \leq \frac{(2h+1)!}{2^{2h}\pi^{2h+1}}$$

and the asserted estimate follows. \square

We are now in a position to prove Proposition 1.

Proof of Proposition 1. For $N \in \mathbb{Z}_{\geq 1}$, $\beta \geq 0$ and $j \in \mathbb{Z}_{\geq 2}$, define the partial sums

$$G_{N,\beta,j} := \sum_{N < n \leq 2N} n^\beta B_j \left(\left\{ \frac{x}{n} \right\} \right).$$

Let $T \leq x^{1/\alpha}$ be a parameter to be chosen later. Write

$$G_{\alpha,\beta,j}(x) = \left(\sum_{n \leq T} + \sum_{T < n \leq x^{1/\alpha}} \right) n^\beta B_j \left(\left\{ \frac{x}{n} \right\} \right)$$

so that, setting $L(x, T) := \frac{\log(x^{1/\alpha}/T)}{\log 2} + 1$, we get

$$\begin{aligned} |G_{\alpha,\beta,j}(x)| &\leq \sup_{t \in \mathbb{R}} |B_j(\{t\})| T^{\beta+1} + \left| \sum_{T < n \leq x^{1/\alpha}} n^\beta B_j\left(\left\{\frac{x}{n}\right\}\right) \right| \\ &\leq \Gamma_j T^{\beta+1} + \max_{T < N \leq x^{1/\alpha}} |G_{N,\beta,j}| L(x, T) \end{aligned} \quad (7)$$

where we used Lemma 9 and where we splitted the interval $(T, x^{1/\alpha}]$ into dyadic intervals $(N, 2N]$, whose number does not exceed $\leq L(x, T)$. Therefore, the problem amounts to estimating the sum $G_{N,\beta,j}$.

▷ Since $j \geq 2$, we may expand the Bernoulli functions in Fourier series, the convergence being uniform: for all $t \in \mathbb{R}$

$$B_j(\{t\}) = -\frac{j!}{(2\pi i)^j} \sum_{m \neq 0} \frac{e(mt)}{m^j}.$$

▷ By partial summation, we get

$$|G_{N,\beta,j}| \leq \Gamma_j N^\beta \sum_{m \geq 1} m^{-j} \max_{N \leq N_1 \leq 2N} \left| \sum_{N < n \leq N_1} e\left(\frac{mx}{n}\right) \right|.$$

▷ Applying Lemma 8 to the exponential sum with $\lambda_2 = \frac{mx}{4N^3}$ et $c_2 = 8$, we derive

$$\begin{aligned} |G_{N,\beta,j}| &\leq \Gamma_j N^\beta \sum_{m \geq 1} m^{-j} \max_{N \leq N_1 \leq 2N} \left(8N \left(\frac{mx}{4N^3} \right)^{1/2} + 2 \left(\frac{4N^3}{mx} \right)^{1/2} \right) \\ &\leq 4\Gamma_j N^\beta \sum_{m \geq 1} m^{-j} \left(\sqrt{\frac{mx}{N}} + \frac{N^{3/2}}{\sqrt{mx}} \right) \\ &= 4\Gamma_j \left(\zeta(j - \frac{1}{2}) x^{1/2} N^{\beta-1/2} + \zeta(j + \frac{1}{2}) x^{-1/2} N^{\beta+3/2} \right). \end{aligned} \quad (8)$$

Reporting (8) in (7) then yields

$$|G_{\alpha,\beta,j}(x)| \leq \Gamma_j T^{\beta+1} + 4\Gamma_j \left(\zeta(j - \frac{1}{2}) x^{\frac{\beta}{\alpha} - \frac{1}{2\alpha} + \frac{1}{2}} + \zeta(j + \frac{1}{2}) x^{\frac{\beta}{\alpha} - \frac{1}{2\alpha} + \frac{1}{2}} \right) L(x, T).$$

Now choose $T = x^{\frac{1}{\beta+1}(\frac{\beta}{\alpha} - \frac{1}{2\alpha} + \frac{1}{2})}$. Note that the condition $1 < \alpha < 3$ ensures that $T \leq x^{1/\alpha}$. This choice of T yields the announced result. \square

2.3.4 Proof of Corollary 2

When $(\alpha, \beta, j) = (2, 1, 2)$, we have $4\Gamma_2 = \frac{2}{3}$, and thus $1.9 < 4\Gamma_2 \left(\zeta\left(\frac{3}{2}\right) + \frac{1}{4} \right) < 1.91$ and $0.89 < 4\Gamma_2 \zeta\left(\frac{5}{2}\right) < 0.9$, so that

$$|G_{2,1,2}(x)| < 2.81 x^{3/4} \left(1 + \frac{\log x}{8 \log 2} \right) < x^{3/4} \log x$$

as soon as $x \geq 300$. \square

2.4 Finalization of the proof of Theorem 1

Lemma 10. *For all $x \geq 300$, we have*

$$2x \left| \sum_{k \leq \sqrt{x}} \frac{1}{k} R_1\left(\frac{x}{k}\right) \right| \leq \frac{\log x}{x^{1/4}} + \frac{0.047}{x^{1/2}}.$$

Proof. We proceed as in Lemmas 1 and 2. By integration by parts, we have

$$R_1(t) = -\frac{B_2(\{t\})}{2t^2} + 2 \int_t^\infty \frac{B_2(\{u\})}{u^3} du := -\frac{B_2(\{t\})}{2t^2} + Q(t)$$

so that

$$2x \left| \sum_{k \leq \sqrt{x}} \frac{1}{k} R_1\left(\frac{x}{k}\right) \right| \leq \frac{1}{x} \left| \sum_{k \leq \sqrt{x}} k B_2\left(\left\{\frac{x}{k}\right\}\right) \right| + 2x \left| \sum_{k \leq \sqrt{x}} \frac{1}{k} Q\left(\frac{x}{k}\right) \right|.$$

By Corollary 2, the 1st sum is $\leq \frac{\log x}{x^{1/4}}$, and by the 2nd mean value theorem, we derive $|Q(t)| \leq \frac{4c}{3t^3}$ where $c = \frac{\sqrt{3}}{36}$, so that the 2nd sum does not exceed

$$\leq \frac{8c}{3x^2} \sum_{k \leq \sqrt{x}} k^2 = \frac{8c}{3x^2} \frac{B_3(\sqrt{x}+1) - B_3(0)}{3} =: f(x).$$

Since $x \geq 300$, we get $\sqrt{x} f(x) \leq \sqrt{300} f(300) \leq 0.047$, completing the proof. \square

Proof of Theorem 1. Set $P(X) := -\frac{1}{2}X^2 - (2\gamma - 1)X + 2(\gamma + \gamma_1) - \gamma^2 - 1$. By Lemmas 3 and 4, we get

$$\begin{aligned} r(x) &:= \sum_{n \leq x} \tau(n) - x \sum_{n \leq x} \frac{\tau(n)}{n} - xP(\log x) - \frac{1}{4} \\ &= x (R_1(\sqrt{x}))^2 - 2\sqrt{x} R_1(\sqrt{x}) \psi(\sqrt{x}) - 2x (R_2 - R_1)(\sqrt{x}) - 2x \sum_{k \leq \sqrt{x}} \frac{1}{k} R_1\left(\frac{x}{k}\right) \end{aligned}$$

and Lemma 5 applied to the last sum yields

$$|r(x)| \leq \frac{\alpha^2}{x} + \frac{\alpha}{\sqrt{x}} + \frac{2\beta}{\sqrt{x}} + \alpha \left(1 + \frac{1}{\sqrt{x}}\right) = \alpha + \frac{2(\alpha + \beta)}{\sqrt{x}} + \frac{\alpha^2}{x}$$

where $\alpha := \sup_{z \geq 1} |R_1(z)|z^2$ and $\beta := \sup_{z \geq 1} |R_2(z) - R_1(z)|z^3$. Using Lemmas 1 and 2, we get $\alpha \leq 0.125$ and $\beta \leq 0.033$, proving the 1st estimate in Theorem 1. To prove the 2nd part, we use Lemma 10 instead of Lemma 5, yielding

$$|r(x)| \leq \frac{\alpha^2}{x} + \frac{\alpha}{\sqrt{x}} + \frac{2\beta}{\sqrt{x}} + \frac{\log x}{x^{1/4}} + \frac{0.047}{x^{1/2}} = \frac{\log x}{x^{1/4}} + \frac{\alpha + 2\beta + 0.047}{\sqrt{x}} + \frac{\alpha^2}{x}$$

which provides the announced estimates. \square

Proof of Corollary 1. By summation by parts, we derive

$$\frac{\Delta(x)}{x} - \delta(x) = \int_x^\infty \frac{\Delta(u)}{u^2} du.$$

When $x \geq 7$, the first estimate of Theorem 1 is less than 0.247 and so

$$\Delta(x) - x\delta(x) \geq \frac{1}{4} - r(x) \geq 0.003.$$

To complete the proof in the case $1 \leq x < 7$, we split the interval $[1, 7)$ into subintervals of the shape $I_m := [m, m+1)$, $m = 1, \dots, 6$, so that, for all $x \in I_m$, $\Delta(x) - x\delta(x) = Q_m(x) - xP(\log x)$, where $Q_m(x) := \sum_{n \leq x} \tau(n) - x \sum_{n \leq x} \frac{\tau(n)}{n}$ is a polynomial of degree 1, and P is the polynomial of degree 2 aforementioned. For instance, if $m = 5$, $Q_5(x) = -\frac{229}{60}x + 10$ and therefore, for all $x \in [5, 6)$,

$$Q_5(x) - xP(\log x) = \frac{1}{2}x(\log x)^2 + (2\gamma - 1)x \log x + x \left(\gamma^2 - 2\gamma - 2\gamma_1 - \frac{169}{60} \right) + 10$$

which is ≥ 0 whenever $5 \leq x < 6$. There is no problem gluing the intervals together for the overall function is continuous. We use 40 decimal places for γ and γ_1 thanks to [6, Lemme 4] or [13, Exercise 7.2], which is more than sufficient to get the desired result in this case. \square

Acknowledgments

The authors would like to express their sincere gratitude to Olivier Ramaré for bringing the unpublished results of Mahatab and Mukhopadhyay to their attention and for providing them with the references [6] and [13].

References

- [1] M. BALAZARD, *Le Théorème des Nombres Premiers*, Coll. Nano, Eds. Calvage et Mounet, 2016.
- [2] D. BERKANE, O. BORDELLÈS, AND O. RAMARÉ, Explicit upper bounds for the remainder term in the divisor problem, *Math. Comput.* **81** (2012), 1025–1051.
- [3] O. BORDELLÈS, *Arithmetic Tales, Advanced Edition*, UTX, Springer, 2020.
- [4] M. CULLY-HUGILL AND T. TRUDGIAN, Two explicit divisor sums, *Ramanujan J.* **56** (2021), 1382–4090.
- [5] F. DAVAL, Explicit conversions between summatory functions of the Möbius function, *Comb. Number Theory* **14** (2025), 163–188.
- [6] S. ETTAHI AND L. SUREL, *Calcul Scientifique de Constantes Arithmétiques*, Inst. Math. Marseille, Aix-Marseille Univ., 2019.
- [7] M. N. HUXLEY, Exponential sums and lattice points III, *Proc. London Math. Soc.* **87** (2003), 591–609.
- [8] X. LI AND X. YANG, An improvement on Gauss’s circle problem and Dirichlet’s divisor problem, *arXiv e-prints*, 32 pp., 2023, <https://arxiv.org/abs/2308.14859>.
- [9] K. MAHATAB AND A. MUKHOPADHYAY, Omega theorems for the twisted divisor function, *Funct. Approximatio, Comment. Math.* **62** (2020), 171–186.
- [10] H. L. MONTGOMERY AND R. C. VAUGHAN, *Multiplicative Number Theory I. Classical Theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2007.
- [11] D. PATEL, *Explicit sub-Weyl bound for the Riemann Zeta function*, PhD Thesis. The Ohio State University, 2021.
- [12] H. RADEMACHER, *Topics in Analytic Number Theory*, Springer, 1973.
- [13] O. RAMARÉ, *Excursions in Multiplicative Number Theory*, Birkhäuser Advanced Texts Basler Lehrbücher, Birkhäuser, 2022.
- [14] O. RAMARÉ, *Personal Communication*.