

ON THE SUBTRACTIVE IDEAL STRUCTURE OF COMMUTATIVE SEMIRINGS

PUBLI SENGUPTA, AMARTYA GOSWAMI, PRONAY BISWAS, AND SUJIT KUMAR SARDAR

ABSTRACT. In the theory of commutative semirings, the lack of additive inverses creates a structural divergence between ideals and congruences that does not exist in ring theory. The aim of this article is to restore critical ideal-theoretic properties via the subtractive property. We first prove a subtractive analogue of Krull's existence theorem, guaranteeing the existence of k -prime ideals disjoint from multiplicative sets. We show that in arithmetic semirings, the distinction between k -irreducible and k -strongly irreducible ideals vanishes, a coherence that we show is preserved under localisation. We investigate the structural properties and coincidence phenomena among associated subclasses of k -ideals in Laskerian semirings, von Neumann regular semirings, unique factorisation semidomains, principal ideal semidomains, and weakly Noetherian semirings. Finally, within the framework of additively idempotent semirings, we tether subtractive ideal-theoretic structures to underlying order-theoretic constraints, thereby obtaining new characterizations of k -prime and k -semiprime ideals. In that process, we also establish that every absolutely k -prime ideal is k -prime and every k -maximal ideal is absolutely k -prime.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Basic results	5
4. On quotient semirings	8
5. Under localisation	10
6. In special semirings	13
6.1. Additively idempotent semirings	17
Acknowledgement	21
References	21

1. INTRODUCTION

Since Vandiver introduced *rings without negatives* [36, 37], the study of semirings has continued to evolve. Although semirings are often viewed as a natural generalisation of both rings and distributive lattices, they have yet to settle into a definitive position between these two theories. Throughout, the term *semiring* tacitly refers to a commutative semiring $(S, +, 0, \cdot, 1)$ with multiplicative identity 1 and an element 0, which is the additive identity and also multiplicatively *absorbing*, that is, $a \cdot 0 = 0$, for all $a \in S$. In contrast to the ring-theoretic case, ideals and congruences in semirings are no longer in bijective correspondence. For example, take the semiring of non-negative integers, following Golan [12], we denote this by $(\mathbb{N}, +, 0)$, and fix two natural numbers $a, b \in \mathbb{N} \setminus \{0, 1\}$. Let $\mathcal{D} := \{(x, y) \in \mathbb{N} \mid a \text{ divides } |x - y|, \text{ and } x > b, y > b\}$. Then $k = \mathcal{D} \cup \{(x, x) \mid x \in \mathbb{N}\}$ is a congruence on \mathbb{N} . Albeit the zeroth class $[0]_k$ is the singleton $\{(0, 0)\}$, which

2020 *Mathematics Subject Classification.* 16Y60.

Key words and phrases. semiring, strongly irreducible ideal, radical, k -ideal, subtractive ideal, additively idempotent semiring.

shows that we cannot recover the original congruence from the zeroth class. *De facto*, there are semirings which are ideal-simple but not congruence-simple, see [31, Section 2]. Therefore, in the sense of Gumm and Ursini [17], semirings are not *ideal determined*. As a generalisation of rings, semirings support a wider array of ideals (*cf.* Definition 2.2) and congruences (*cf.* Definition 2.12), including structures that are either trivial or do not arise in classical ring theory. For example, the class of k -ideals (also known as subtractive ideals) is trivial in rings, since each ideal of a ring is a k -ideal. Moreover, the class of *strong ideals* in rings is incoherent. The ideal theory of semirings has been extensively studied; see [2, 3, 7, 13–16, 22, 27, 32]. Despite its early appearance in an AMS notice [20], Henriksen’s notion of a k -ideal has proved to be a persistent and productive idea in semiring theory. For a detailed account of the historical development of semiring theory, we refer the reader to Głazek [11]. Several authors, including Katsov, Abuhlail, Nam, and Il’in, have focused on characterizing algebraic properties of semirings admitting a rich class of k -ideals; see [4, 6, 21, 25, 26]. The study of the class of k -ideals appears in the earlier works [18, 19, 24, 28, 33, 34, 38] and has been taken up again in more recent work [15]. This work is a sequel to [15]. We study the classes of k -prime, k -semiprime, k -primary, k -irreducible, and k -strongly irreducible ideals, together with their associated congruence classes, both in general semirings and across several distinguished classes of semirings. For the corresponding of ideals in rings we refer to [1, 5, 35]. The so-called “exchange principal” (*cf.* [24, Proposition 3.5] and [15, Proposition 3.11, Proposition 3.21]), namely

$$k\text{-}X\text{-ideal} \iff k\text{-ideal} + X\text{-ideal},$$

is true for k -prime, k -semiprime, k -irreducible and k -strongly irreducible ideals, which make these classes more tractable.

Let us now retrace the main results established in this paper. In Section 3, we begin by examining examples and non-examples of subtractive and strongly subtractive semirings. We show that the ideal lattice of a strongly subtractive semiring is distributive, while that of a subtractive semiring is known to be modular. Consequently, strongly subtractive semirings provide a new source of examples of arithmetic semirings (see, Definition 3.12). In Proposition 3.8, we establish a subtractive analogue of Krull’s theorem for semirings, guaranteeing the existence of k -prime ideals. Although the classes of k -prime and k -primary ideals are shown to be distinct in Remark 2.9, however, Theorem 3.9 shows that they coincide in von Neumann regular semirings. We also record a correspondence between k -strongly irreducible ideals and i_k -systems. The section concludes by proving that, within the class of arithmetic semirings, every k -irreducible ideal is in fact k -strongly irreducible, a phenomenon that does not occur in general semirings.

Section 4 investigates the structural behaviour of distinguished subclasses of k -ideals under semiring homomorphisms and analyses their structural persistence in quotient semirings. We begin by establishing that the class of k -prime ideals is invariant under k -contraction. An analogous stability result is obtained for k -irreducible ideals under a suitably restricted class of semiring homomorphisms (Proposition 4.2). In Theorem 4.6, we prove that k -strongly irreducible ideals descend to quotient semirings, and that the converse implication holds in the arithmetic setting. This result strengthens the corresponding theorem of Atani and Garfami; see [3, Theorem 2.7]. Finally, we show that the k -irreducibility of an ideal can be detected purely at the level of the quotient, namely, an ideal is k -irreducible if and only if the zero ideal of the associated quotient semiring is k -irreducible.

In Section 5, we investigate the behaviour of k -ideals and their subclasses under localisation of semirings. The central result, Theorem 5.2, extends Azizi’s correspondence to the semiring setting by establishing a bijection between k -proper strongly irreducible ideals of $T^{-1}S$ and those k -proper strongly irreducible ideals of S that are disjoint from T . We then analyse the stability of k -prime ideals under extension and contraction along semiring homomorphisms, describing the induced maps on k -spectra and their injectivity and surjectivity properties (Proposition 5.5). Finally, we examine the behaviour of k -primary and k -strongly

irreducible ideals under localisation and contraction (Propositions 5.7 and 5.8). Taken together, these results show that localisation preserves and reflects essential ideal-theoretic structure in the subtractive framework.

The Section 6, is devoted to a detailed analysis of k -ideals and their variants in several important classes of semirings, with particular emphasis on k -strongly irreducible ideals and their relationship with k -prime and k -primary ideals. The section is divided into two complementary parts.

In the first part, we study the behaviour of k -strongly irreducible ideals in Laskerian semirings, von Neumann regular semirings, unique factorisation semidomains, principal ideal semidomains, and weakly Noetherian semirings. Proposition 6.2 establishes foundational links between k -strong irreducibility, k -primeness, and k -primary decomposition, showing in particular that in Laskerian and von Neumann regular semirings these notions coincide. In the setting of unique factorisation semidomains, Lemma 6.4 and Theorem 6.5 provide characterizations of k -strongly irreducible ideals in terms of prime power decompositions and least common multiples, and show that such ideals are necessarily k -primary. Localization properties of k -strongly irreducible ideals are analysed in Corollary 6.7, yielding correspondence results for unique factorisation semidomains and von Neumann regular semirings. Further structural consequences are obtained for principal ideal semidomains, where non-zero k -prime ideals are shown to be k -maximal, and for weakly Noetherian semirings, whereas Proposition 6.12 demonstrates that every k -irreducible ideal is k -primary.

The subsection 6.1 focuses on additively idempotent semirings, *alias*, B_1 -algebras. We give a new order-theoretic perspective of k -prime and k -semiprime ideals in Theorem 6.15 and Theorem 6.16, respectively, using the natural partial order. For additively idempotent unique factorisation semidomains, Theorem 6.17 characterizes k -strongly irreducible ideals through least common multiples. We then establish that saturated ideals (see Definition 6.19) coincide precisely with k -ideals (Theorem 6.22), clarifying the relationship between ideals and excellent congruences in this setting. We conclude the paper by showing that in an additively idempotent semiring, the class of absolutely k -prime is contained in k -prime ideals and the class of k -maximal ideals is contained in absolutely k -prime ideals.

2. PRELIMINARIES

This section collects the basic definitions and results needed to keep the exposition self-contained. We adopt the standard terminology and foundational material as presented in Golan's work [12]. Unless otherwise stated, all semirings considered here are commutative with identity. A semiring homomorphism $f: S \rightarrow S'$ is a map satisfying $f(x + y) = f(x) + f(y)$, $f(xy) = f(x)f(y)$, $f(0_S) = 0_{S'}$ and $f(1_S) = 1_{S'}$.

Definition 2.1 (Page 2, [32]). Suppose S is a semiring.

- (1) A non-zero and non-unit element r of S is said to be *irreducible* if $r = r_1 r_2$, where $r_1, r_2 \in S$, then either r_1 or r_2 is a unit.
- (2) An element p of S is said to be *prime* if the principal ideal $\langle p \rangle$ is prime.
- (3) A semiring S is said to be a *semidomain* if $ab = ac$ implies $b = c$, for all $b, c \in S$ and all non-zero $a \in S$.
- (4) Let S be a semidomain. Then S is called a *unique factorisation semidomain* if the following conditions hold:
 - (a) Each irreducible element of S is a prime element of S .
 - (b) Any non-zero, non-unit element of S can be uniquely written as a product of irreducible elements of S .

We follow the definitions of *prime*, *semiprime* and *maximal* ideals as presented in Golan [12, Chapter 6, Chapter 7].

Definition 2.2 ([20], [12]). An ideal I of a semiring S is said to be a k -ideal if $x + y \in I$ and $y \in I$ implies $x \in I$. Moreover, I is called strong if $a + b \in I$ implies $a, b \in I$.

Remark 2.3. In literature k -ideals are also known as *subtractive ideals* (see [12, Chapter 6]) and *semistrong ideals* (see [10, Section 1]).

It is straightforward to verify that every strong ideal is a k -ideal, though the converse need not hold.

Definition 2.4 (Definition 2.2, [15]). Let S be a semiring I be an ideal of S . The k -closure of I is defined by

$$\mathcal{C}_k(I) := \{s \in S \mid s + x \in I \text{ for some } x \in I\}.$$

We will heavily use the following properties of k -closure operators lifted from [15, Lemma 2.3].

Lemma 2.5. In what follows, I and $\{I_\lambda\}_{\lambda \in \Lambda}$ are ideals of semiring S .

- (1) $\mathcal{C}_k(I)$ is the smallest k -ideal containing I .
- (2) $\mathcal{C}_k(\mathcal{C}_k(I)) = \mathcal{C}_k(I)$.
- (3) $\mathcal{C}_k(\bigcap_{\lambda \in \Lambda} I_\lambda) = \bigcap_{\lambda \in \Lambda} \mathcal{C}_k(I_\lambda)$.

Let us recall the definitions k -contraction and k -extension of k -ideals from [15, Definition 4.1].

Definition 2.6. Let S , and S' be two semirings and $\varphi: S \rightarrow S'$ be a semiring homomorphism.

- (1) If J is a k -ideal of S' , then the *k -contraction* of J , denoted by J^c , is defined by $\varphi^{-1}(J)$.
- (2) If I is a k -ideal of S , then the *k -extension* of I , denoted by I^e , is defined by $\mathcal{C}_k(\langle \varphi(I) \rangle)$.

Remark 2.7. By a k -contracted ideal A of S , we mean, there exists a k -ideal I of S' such that $A = I^c$. Similarly, if B is a k -extended ideal of S' , then $B = I^e$, for some k -ideal I of S .

We recall some subclasses of k -ideals that play a central role throughout this work.

Definition 2.8 (Definition 3.4, Definition 3.20, [15]). Let S be a semiring.

- (1) A k -ideal I of S is said to be *k -strongly irreducible* if $A \cap B \subseteq I$ implies $A \subseteq I$ or $B \subseteq I$, for two k -ideals A, B of S and *k -irreducible* if $A \cap B = I$ implies either $A = I$ or $B = I$.
- (2) A k -ideal I of S is said to be *k -prime* if $AB \subseteq I$ implies $A \subseteq I$ or $B \subseteq I$, for two k -ideals A, B of S .
- (3) If I is a k -ideal of S , it is said to be *k -semiprime* if $A^2 \subseteq I$ implies $A \subseteq I$.
- (4) A k -ideal I of S is called *k -primary* if $ab \in I$ and $a \notin I$ implies $b^n \in I$, for some $n \in \mathbb{N}$.

Remark 2.9. Despite their close relationship, the classes of k -prime, k -semiprime, and k -strongly irreducible ideals do not coincide in general semirings. We note two instances where this divergence occurs.

- (1) Clearly, every k -prime ideal is k -strongly irreducible and k -semiprime. The converse is not always true, like in $(\mathbb{N}, +, \cdot)$, the ideal $4\mathbb{N}$ is k -strongly irreducible but not k -prime. Moreover, the ideal $6\mathbb{N}$ in \mathbb{N} is a k -semiprime ideal, which is neither k -prime nor k -strongly irreducible. Moreover, the ideal $p^2\mathbb{N}$, where p is any prime number, is a k -primary ideal which is not k -prime.
- (2) Each k -strongly irreducible ideal is k -irreducible, whereas the converse is not true in general. Consider the semiring $\mathbb{N}[x, y]$ and the ideal $\langle x, y \rangle$. Then $\langle x, y \rangle$ is a k -ideal, and it cannot be written as an intersection of two proper k -ideals, that is, it is k -irreducible. If $I = \langle x, y^2 \rangle$ and $J = \langle x^2, y \rangle$, then $\langle x, y^2 \rangle \cap \langle x^2, y \rangle \subseteq \langle x, y \rangle$. But $\langle x, y^2 \rangle \not\subseteq \langle x, y \rangle$ and $\langle x^2, y \rangle \not\subseteq \langle x, y \rangle$.

We also note that not all prime, semiprime, and strongly irreducible ideals are k -ideals in general semirings. For instance, take the unique maximal ideal $\mathbb{Q}_+[X] \setminus \mathbb{Q}_+$ in the semiring $\mathbb{Q}_+[X]$, which is not a k -ideal.

We denote the collection of all prime ideals of a semiring S by $\text{Spec}(S)$, and the collection of all k -prime ideals by $\text{Spec}_k(S)$.

Definition 2.10 (Definition 3.12, [15]). The k -radical of a k -ideal I of a semiring S is defined as

$$\mathcal{R}_k(I) := \bigcap_{I \subseteq P} \{P \mid P \in \text{Spec}_k(S)\}.$$

Throughout the paper, we repeatedly invoke the following elementary properties of the k -radical closure (lifted from [15, Lemma 3.13]) whenever required.

Lemma 2.11. *In the following, I and J are ideals of a semiring S .*

- (1) $\mathcal{R}_k(I)$ is a k -ideal containing I .
- (2) $\mathcal{R}_k(IJ) = \mathcal{R}_k(I \cap J) = \mathcal{R}_k(I) \cap \mathcal{R}_k(J)$.

Note that the appropriate congruence theoretic counterpart of k -ideals is the k -congruences, in the sense of Han [19, Definition 3.2].

Definition 2.12. Let S be a semiring. A congruence ρ on S is said to be a k -congruence if $\rho = K_I$, for some ideal I of S .

There exists a one-one correspondence between k -ideals and k -congruences, see [19, Theorem 3.8].

Definition 2.13. A k -congruence θ is said to be *irreducible* if for any two k -congruence ρ_1, ρ_2 , and $\rho_1 \cap \rho_2 = \theta$, implies that $\rho_1 = \theta$ or $\rho_2 = \theta$.

If S is an additively idempotent semiring, a natural partial order on S is defined as follows: for $x, y \in S$,

$$x \leq y \iff x + y = y.$$

Then, we have the following from [24, Proposition 3.11].

Proposition 2.14. *Let S be an additively idempotent semiring and I be an ideal. Then I is a k -ideal if and only if for all $x \in I$ and $y \leq x$, we have $y \in I$.*

3. BASIC RESULTS

The ideal theory of a semiring has a remarkably different flavour from that of a ring. While the lattice of ideals of a ring is always modular, this property fails in general for semirings; see [12, Example 6.36]. In contrast, k -ideals retain many of the familiar properties of ring ideals, for example, the lattice of all k -ideals with a meaningful join and meet is modular; see [33, Theorem 2.4]. A semiring can be k -ideal simple, that is, the only k -ideals are the trivial ones. We give some examples of k -ideal simple semirings.

Example 3.1.

- (1) The semiring $\mathbb{Q}_+[\sqrt{2}]$ with usual addition and multiplication is k -ideal simple; see [9, Section 5.2].
In general, a semiring S satisfying the condition:

$$1 + xa = ya, \text{ for all } a \in S \setminus \{0\} \text{ and for some } x, y \in S,$$

has trivial k -ideals. For a proof, see [34, Corollary 2.8].

- (2) Consider the semiring $C^\infty(X) = \{f: X \rightarrow (0, \infty) \mid f \text{ is continuous}\} \cup \{0\}$ with pointwise addition and pointwise multiplication, where X is a topological space and the constant function 0 which is the additive identity and an absorbing element. Since $1 + \infty = \infty$ and the constant function ∞ is also multiplicatively absorbing, then ∞ belongs to every ideal, but 1 is not. Therefore, the semiring $C^\infty(X)$ is a k -ideal simple semiring.

Definition 3.2. A semiring S is said to be a k -semiring if each ideal of S is a k -ideal. Moreover, S is called *strongly k -semiring* if each ideal of S is a strong ideal.

In literature, k -semirings and strongly k -semirings are also known as *subtractive semirings* and *strongly subtractive*, see [21, 25, 26]. There are numerous examples of k -semirings; we list a few of them.

Example 3.3.

- (1) Any bounded distributive lattice $(L, \vee, \wedge, 0, 1)$ when considered as a semiring. In fact, it is a strongly k -semiring.
- (2) Product of rings and bounded distributive lattices, both considered as semirings.
- (3) The semiring of non-negative real-valued continuous functions over an F -space, see [7, Theorem 3.9]
- (4) The semiring of non-negative real-valued measurable functions over a measurable space; see [8, Corollary 3.2].

Remark 3.4. The semirings in (3) and (4) of Example 3.3 are cancellative. In contrast, there are semirings which are cancellative, but not every ideal is a k -ideal. For example, take the semiring $C^+(X)$ for any Tychonoff space X . Then with pointwise addition and multiplication, $C^+(X)$ is a cancellative semiring, but not all ideals are k -ideals; see [7, Example 3.1].

While k -ideals recover much of the behaviour of ring ideals, the next theorem shows that strong ideals align naturally with lattice ideals. We denote the collection of all ideals of a semiring S by $\text{Id}(S)$.

Theorem 3.5. *Let S be a strongly subtractive semiring. Then the lattice $(\text{Id}(S), +, \cap)$ is distributive.*

Proof. It is enough to show that $I \cap (J + K) \subseteq (I \cap J) + (I \cap K)$ for any I, J and K in $\text{Id}(S)$. Let $a \in I \cap (J + K)$. Then $a = x + y \in I$, for some $x \in J$ and $y \in K$. Since I is strong, $x \in I$ and $y \in I$. We conclude that $a \in (I \cap J) + (I \cap K)$ and hence $I \cap (J + K) \subseteq (I \cap J) + (I \cap K)$. \square

As a corollary, we get that the ideal lattice of any bounded distributive lattice is distributive. Next, we note an elementary fact, which will be used freely in what follows.

Lemma 3.6. *Every k -maximal ideal of a semiring is k -prime.*

Proof. Let S be a semiring and let M be a k -maximal ideal of S . Let $ab \in M$ and $a \notin M$, for some $a, b \in S$. Then $M \subseteq M + \langle a \rangle$, which further implies that $M \subseteq M + \langle a \rangle \subseteq \mathcal{C}_k(M + \langle a \rangle)$. Since M is a k -maximal ideal, we must have $\mathcal{C}_k(M + \langle a \rangle) = S$. Therefore $1 \in \mathcal{C}_k(M + \langle a \rangle)$. By definition, $1 + ax_1 + m_1 = ax_2 + m_2$, for some $x_1, x_2 \in S$ and $m_1, m_2 \in M$. We get $b + abx_1 + bm_1 = abx_2 + bm_2$. So, $b \in \mathcal{C}_k(M) = M$ as M is a k -ideal. Hence, M is a k -prime ideal. \square

Proposition 3.7. *Let S be a semiring and $x \in S$. Then the following are equivalent.*

- (1) $1_S \in \mathcal{C}_k(\langle x \rangle)$.
- (2) x cannot be in any k -maximal ideal of S .

Proof. Let x be an element that does not belong to any k -maximal ideal of S . Since every proper k -ideal is contained in a k -maximal ideal of S , the ideal $\mathcal{C}_k(\langle x \rangle)$ cannot be a proper k -ideal. Therefore, $\mathcal{C}_k(\langle x \rangle) = S$ and so $1_S \in \mathcal{C}_k(\langle x \rangle)$. Conversely, let $1_S \in \mathcal{C}_k(\langle x \rangle)$. Let I be a k -maximal ideal of S such that $x \in I$. Then $\langle x \rangle \subseteq I$, and hence, $\mathcal{C}_k(\langle x \rangle) \subseteq \mathcal{C}_k(I) = I$. From this, we conclude that $1_S \in I$, a contradiction. Hence $x \notin I$. \square

As in the case of rings, the correspondence between prime ideals and m -systems is well established in semiring theory; see [12, Corollary 7.11]. Golan further proves a Krull-type converse in [12, Proposition 7.12]. The following proposition provides a subtractive analogue of Krull's result for semirings.

Proposition 3.8. *Let T be a multiplicatively closed set of a semiring S and let I be a k -ideal maximal among the k -ideals disjoint from T . Then I is k -prime.*

Proof. To prove I is k -prime, we need to show that I is prime. Let $ab \in I$ such that $a, b \notin I$. Consider the k -ideal $\mathcal{C}_k(I + \langle a \rangle)$. Then $\mathcal{C}_k(I + \langle a \rangle)$ is a k -ideal properly containing I , and therefore, intersects T . So, there exists $s_1 \in \mathcal{C}_k(I + \langle a \rangle)$, which implies $s_1 + i_1 + x_1 a = i_2 + x_2 a$, for some $i_1, i_2 \in I$ and $x_1, x_2 \in S$. Then, $s_1 b + i_1 b + x_1 a b = i_2 b + x_2 a b$. This implies $s_1 b \in I$ as $i_1, i_2, a b \in I$. Similarly, $\mathcal{C}_k(I + \langle b \rangle)$ is a k -ideal properly containing I , and so, it intersects T . Therefore, there exists $s_2 \in \mathcal{C}_k(I + \langle b \rangle)$. Then, $s_2 + j_1 + y_1 b = j_2 + y_2 b$, for some $j_1, j_2 \in I$ and $y_1, y_2 \in S$. So, $s_2 a + j_1 a + y_1 b a = j_2 a + y_2 b a$, which implies $s_2 a \in I$, as $j_1, j_2, a b \in I$. We have,

$$(s_1 + i_1 + x_1 a)(s_2 + j_1 + y_1 b) = (i_2 + x_2 a)(j_2 + y_2 b).$$

Thus

$$s_1 s_2 + s_1 + s_1 j_1 + y_1 s_1 b + i_1 (s_2 + j_1 + y_1 b) + x_1 s_2 a + j_1 x_1 a + x_1 y_1 a b = i_2 (j_2 + y_2 b) + j_2 x_2 a + x_2 y_2 a b.$$

Therefore, we get $s_1 s_2 \in I$, which is a contradiction to the fact that I is disjoint from T . Hence I is k -prime. \square

Although k -primary and k -prime ideals are distinct in general (c.f. Remark 2.9), the following theorem shows that this distinction disappears for von Neumann regular semirings.

Theorem 3.9. *In a von Neumann regular semiring, the classes of k -prime and k -primary ideals coincide.*

Proof. Let S be a von Neumann regular semiring and I be a k -prime ideal of S . Then I is a k -ideal of S . Therefore, $xy \in I$ implies $x \in I$ or $y \in I$, which further implies I is k -primary. Conversely, suppose that I is a k -primary ideal of S . Let $xy \in I$. Then $x \in I$ or $y^n \in I$, for some natural number n . If $x \in I$, then I is prime. Now, let $y^n \in I$. Since S is von Neumann regular for $y \in S$, there exists $x \in S$ such that $y = y^2 x$. This implies $y^{n-2} y = y^{n-2} y^2 x$, which further implies $y^{n-1} = y^n x \in I$, as I is an ideal of S . Again, $y^{n-3} y = y^{n-3} y^2 x$ implies $y^{n-2} = y^{n-1} x \in I$. Continuing in this way, we have $y^2 \in I$. Thus, $y = y^2 x \in I$. \square

Motivated by the correspondence between strongly irreducible ideals and i -systems [12, Proposition 7.33], we introduce a subtractive variant of the notion of an i -system.

Definition 3.10. A non-empty subset A of a semiring S is an i_k -system if $a, b \in A$ implies that $\mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \cap A \neq \emptyset$.

Theorem 3.11. *Let I be a k -ideal of a semiring S . Then the following conditions are equivalent.*

- (1) I is k -strongly irreducible.
- (2) If $a, b \in S$ satisfying $\mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \subseteq I$, then $a \in I$ or $b \in I$.
- (3) $S \setminus I$ is an i_k -system.

Proof. (1) \Rightarrow (2): Clear from the definition.

(2) \Rightarrow (3): Suppose $a, b \in S \setminus I$ and if possible, let $\mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \cap (S \setminus I) = \emptyset$. This implies $\mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \subseteq I$. So, $a \in I$ or $b \in I$, from (2), which is a contradiction. Therefore, $\mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \cap (S \setminus I) \neq \emptyset$. Hence, $S \setminus I$ is an i_k -system.

(3) \Rightarrow (1): Let A, B be two k -ideals of S such that $A \cap B \subseteq I$, but $A \not\subseteq I$ and $B \not\subseteq I$. Then, there exist $x \in A \setminus I$ and $y \in B \setminus I$. By (3), as $S \setminus I$ is an i_k -system, there exists $z \in \mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) \cap (S \setminus I)$. This further implies $z \in A \cap B$ such that $z \notin I$, a contradiction. So, we get $A \subseteq I$ or $B \subseteq I$. Thus I is k -strongly irreducible. \square

As observed in Remark 2.9, k -strongly irreducible and k -irreducible ideals may differ in general. In the sequel, we show that they coincide for a special class of semirings, after introducing the necessary definition.

Definition 3.12. A semiring S is said to be an *arithmetic semiring* if $A + (B \cap C) = (A + B) \cap (A + C)$, or equivalently, $A \cap (B + C) = (A \cap B) + (A \cap C)$, for any ideals A, B, C of S . In other words, in an arithmetic semiring, the set of all ideals of S forms a distributive lattice.

Proposition 3.13. *The following are equivalent for a semiring S .*

1. S is an arithmetic semiring.
2. For any $a, b, c \in S$, $\langle a \rangle \cap (\langle b \rangle + \langle c \rangle) = (\langle a \rangle \cap \langle b \rangle) + (\langle a \rangle \cap \langle c \rangle)$

Proof. (1) \Rightarrow (2): Obvious.

(2) \Rightarrow (1): Suppose (2) holds. Let I, J , and K be any elements of $\text{Id}(S)$. Then $(I \cap J) + (I \cap K) \subseteq I \cap (J + K)$ is always true in any lattice. Let $x \in I \cap (J + K)$. Therefore $x = i$ and $x = j + k$, for some $i \in I$, $j \in J$, and $k \in K$. Then, by the hypothesis, we have

$$x \in \langle i \rangle \cap (\langle j \rangle + \langle k \rangle) = (\langle i \rangle \cap \langle j \rangle) + (\langle i \rangle \cap \langle k \rangle) \subseteq (I \cap J) + (I \cap K).$$

Hence $I \cap (J + K) \subseteq (I \cap J) + (I \cap K)$. □

From Theorem 3.5, we know that all strongly subtractive semirings are arithmetic. In the next theorem, we show that the classes of k -strongly irreducible and k -irreducible ideals coincide in arithmetic semirings.

Theorem 3.14. *In an arithmetic semiring S , a k -ideal I is k -strongly irreducible if and only if it is k -irreducible.*

Proof. We know that every k -strongly irreducible ideal is k -irreducible. To prove the converse part, let I be a k -irreducible ideal of S . Let $A \cap B \subseteq I$, where A and B are two k -ideals of S . Now, since S is an arithmetic semiring, we have $I + (A \cap B) = (I + A) \cap (I + B)$. As $A \cap B \subseteq I$, we get $I = I + (A \cap B) = (I + A) \cap (I + B)$. Then

$$\mathcal{C}_k(I + A) \cap \mathcal{C}_k(I + B) = \mathcal{C}_k((I + A) \cap (I + B)) = \mathcal{C}_k(I) = I.$$

Therefore, we have $\mathcal{C}_k(I + A) = I$ or $\mathcal{C}_k(I + B) = I$. This further implies $I = I + A$ or $I = I + B$. So, $A \subseteq I$ or $B \subseteq I$. Hence I is k -strongly irreducible. □

4. ON QUOTIENT SEMIRINGS

In this section, we investigate the behaviour of k -ideals, k -irreducible ideals, and k -strongly irreducible ideals under semiring homomorphisms and passage to quotient semirings. The following proposition shows that k -prime ideals are stable under k -contractions.

Proposition 4.1. *Let S and S' be two semirings and $\varphi: S \rightarrow S'$ be a semiring homomorphism. if J is a k -prime ideal of S' , then the k -contraction of J , J^c is a k -prime ideal of S .*

Proof. We only prove the ‘prime part’. Let $xy \in J^c = \varphi^{-1}(J)$. This implies $\varphi(xy) \in J$, which further implies $\varphi(x)\varphi(y) \in J$. Then $\varphi(x) \in J$ or $\varphi(y) \in J$. Therefore $x \in \varphi^{-1}(J) = J^c$ or $y \in \varphi^{-1}(J) = J^c$. Thus J^c is a k -prime ideal of S . □

To prove the analogous result for the class of k -strongly irreducible ideals, we need to restrict the semiring homomorphism. Note that for rings, the following conditions are equivalent for a homomorphism $\varphi: S \rightarrow S'$,

- (1) $\ker \varphi \subseteq \langle x \rangle$ for each $x \notin \ker \varphi$.
- (2) If $\varphi(a) = \varphi(b) \neq 0$, then $\langle a \rangle = \langle b \rangle$.

In fact, if φ is surjective and satisfies any one of the above conditions, then $\varphi(\langle a \rangle \cap \langle b \rangle) = \langle \varphi(a) \rangle \cap \langle \varphi(b) \rangle$.

Proposition 4.2. *Let $\varphi: S \rightarrow S'$ be a surjective semiring homomorphism that satisfies the property: if $\varphi(a) = \varphi(b) \neq 0$, then $\langle a \rangle = \langle b \rangle$. If J is a k -strongly irreducible ideal of S' , then J^c is a k -strongly irreducible ideal of S .*

Proof. Suppose $a \notin J^c$ and $b \notin J^c$. It follows that $\varphi(a), \varphi(b) \notin J$. Since J is a k -strongly irreducible ideal, there exists

$$s \in \mathcal{C}_k(\langle \varphi(a) \rangle) \cap \mathcal{C}_k(\langle \varphi(b) \rangle) = \mathcal{C}_k(\langle \varphi(a) \rangle \cap \langle \varphi(b) \rangle) = \mathcal{C}_k(\varphi(\langle a \rangle \cap \langle b \rangle))$$

such that $s \notin J$. Since φ is surjective, there exists $s' \in S$ such that $s = \varphi(s')$. Furthermore, there must exist $t_1, t_2 \in \langle a \rangle \cap \langle b \rangle$ such that $\varphi(s) + \varphi(t_1) = \varphi(t_2)$. Thus $s' \in \mathcal{C}_k(\langle a \rangle \cap \langle b \rangle)$ with $s' \notin J^c$. \square

Recall that a *congruence* on a semiring S is an equivalence relation on S that is also a subsemiring of $S \times S$. We begin by recalling from [12, Page 78] the notion of a *Bourne congruence* associated with an ideal I of S .

Definition 4.3. Let S be a semiring and I be an ideal of S . We define a relation on S given by $r_1 \sim r_2$ if and only if there exist $i_1, i_2 \in I$ such that $r_1 + i_1 = r_2 + i_2$. This defines an equivalence relation and in fact a congruence on S , and equivalence classes are denoted by $r + I$ and the collection of equivalence classes by S/I . Then S/I is a semiring with respect to the operations $(r_1 + I) + (r_2 + I) = r_1 + r_2 + I$ and $(r_1 + I)(r_2 + I) = r_1 r_2 + I$.

The following lemmas summarise the basic properties of ideals and k -ideals in quotient semirings; we mention them without proof.

Lemma 4.4. *Let S be a semiring. Then the following hold:*

- (1) *If $a \in I$, then $a + I = I$, for all ideals I of S .*
- (2) *If I is a k -ideal of S , then $c + I = I$ if and only if $c \in I$.*
- (3) *I, J are ideals of S such that $I \subseteq J$, then J/I is an ideal of S/I . In particular, if J is a k -ideal of S , then J/I is a k -ideal of S/I .*
- (4) *if $1 + I \in J/I$, for ideals I, J with $I \subseteq J$, then $S/I = J/I$.*

Lemma 4.5. *Let I be an ideal of a semiring S . If J, K, L are k -ideals containing I , then $(J/I) \cap (K/I) = L/I$ if and only if $J \cap K = L$.*

In [3, Theorem 2.7], Atani and Garfami established one direction of the correspondence for k -strongly irreducible ideals in quotient semirings. The following theorem strengthens this result by proving the converse in the setting of arithmetic semirings.

Theorem 4.6. *Let S be a semiring. I is an ideal of S and J be a k -strongly irreducible ideal of S containing I . Then J/I is a k -strongly irreducible ideal of S/I . The converse is true if S is an arithmetic semiring.*

Proof. For the first part of the proof, see [3, Theorem 2.7]. For the converse, suppose S is an arithmetic semiring and let $A \cap B \subseteq I$, where A, B are two k -ideals of S . Then $(A \cap B) + I \subseteq J + I = J$, as $I \subseteq J$. This implies $(A + I) \cap (B + I) \subseteq J$, which further implies, $((A + I)/I) \cap ((B + I)/I) \subseteq J/I$ by Lemma 4.5. As, J/I is a k -strongly irreducible ideal, $((A + I)/I) \subseteq J/I$ or $((B + I)/I) \subseteq J/I$. Let $x \in A \subseteq A + I$. Then $x + I \in ((A + I)/I) \subseteq J/I$. Therefore, $x + I = j + I$, for some $j \in J$, which implies $x + a_1 = j + a_2$ for some $a_1, a_2 \in I$. This follows, $x \in J$ as J is a k -ideal of S . Thus $A \subseteq J$. Similarly, if $((B + I)/I) \subseteq J/I$, we have $B \subseteq J$. Hence, J is k -strongly irreducible. \square

The following result is significant in the sense that it shows k -irreducibility is preserved and reflected under passage to quotients, allowing the study of k -irreducible ideals to be reduced to the zero ideal in an appropriate quotient semiring.

Theorem 4.7. *Let I be a k -ideal in a semiring S . Then I is k -irreducible if and only if the zero ideal $\langle \bar{0} \rangle = I/I$ in S/I is k -irreducible.*

Proof. Suppose, I is a k -irreducible ideal of S and $I \neq S$. Then, there exists $a \in S$ such that $a + I \neq I$. As, I is a k -ideal, the zero ideal $\langle \bar{0} \rangle$ is also a k -ideal, by Lemma 4.4. Let $J/I \cap H/I = \langle \bar{0} \rangle$. Then, by Lemma 4.5, $J \cap H = I$. As, I is k -irreducible, $J = I$ or $H = I$. Therefore, $J/I = \langle \bar{0} \rangle$, or $H/I = \langle \bar{0} \rangle$.

Conversely, let $\langle \bar{0} \rangle$ in S/I is k -irreducible. Suppose, $J \cap H = I$, where J, H are k -ideals of S . Then J/I and H/I are also k -ideals of S such that $J/I \cap H/I = \langle \bar{0} \rangle$. So, $J/I = \langle \bar{0} \rangle$ or $H/I = \langle \bar{0} \rangle$. If $J/I = \langle \bar{0} \rangle$, we have $a + I = I$, for all $a \in J$. Therefore, $a \in I$. So, $J \subseteq I$. Now, if $y \in I$, then $y + I = I$. This implies $y + I \in J/I$. So, $y + i_1 = j + i_2$, for some $j \in J$ and $i_1, i_2 \in I$. Therefore, $y \in J$ as J is a k -ideal and $I \subseteq J$. Thus $J = I$. Similarly, if $H/I = \langle \bar{0} \rangle$, then we have $H = I$. \square

5. UNDER LOCALISATION

Our aim in this section is to study the behaviour of the class of k -ideals and its subclasses under localisation. Let S be a semiring and let T be a multiplicatively closed subset of S . For each k -ideal I of the semiring $T^{-1}S$, we consider

$$(\star) \quad I^c := \{x \in S \mid x/1 \in I\} = I \cap S, \quad \text{and} \quad C := \{I^c \mid I \text{ is an ideal of } T^{-1}S\}.$$

Lemma 5.1. *Let S be a semiring and T be a multiplicatively closed subset of S . If $I \in C$, then $(T^{-1}I)^c = I$.*

Proof. Let $x \in I$. Then $x/1 \in T^{-1}I$ implies $x \in (T^{-1}I)^c$. Therefore, $I \subseteq (T^{-1}I)^c$. For the reverse inclusion, suppose $y \in (T^{-1}I)^c$. As $I \in C$, we have $I = J^c$, for some k -ideal J of $T^{-1}S$. Now $y \in (T^{-1}I)^c$ implies $\frac{y}{1} = \frac{i}{s}$, for some $i \in I$ and $s \in T$. Then $ys = i \in I$, which implies $ys \in I = J^c$. Therefore $\frac{ys}{1} \in J$. It follows that $\frac{y}{1} = (\frac{1}{s})(\frac{ys}{1}) \in J$. So, $y \in J^c = I$. Hence $(T^{-1}I)^c = I$. \square

In [1, Theorem 3.1], Azizi established a correspondence between proper strongly irreducible ideals of a localized ring and proper strongly irreducible ideals of the ring itself. The next theorem extends this correspondence to the setting of semirings by considering subtractive ideals, thereby generalizing Azizi's result beyond the classical ring-theoretic framework.

Theorem 5.2. *Let S be a semiring and T be a multiplicatively closed subset of S . Then there is a one-to-one correspondence between the k -proper strongly irreducible ideals of $T^{-1}S$ and k -proper strongly irreducible ideals of S contained in C (defined in (\star)) which do not meet T .*

Proof. Suppose I is a k -proper strongly irreducible ideal of $T^{-1}S$. Then it follows that I^c is a k -ideal of S and $I^c \cap T = \emptyset$ since I cannot contain a unit. Let $A \cap B \subseteq I^c$ for k -ideals A, B of S . Then it follows that

$$(T^{-1}A) \cap (T^{-1}B) = T^{-1}(A \cap B) \subseteq T^{-1}I^c = I.$$

Therefore, $T^{-1}A \subseteq I$ or $T^{-1}B \subseteq I$. Hence $A \subseteq (T^{-1}A)^c \subseteq I^c$ or $B \subseteq (T^{-1}B)^c \subseteq I^c$. Conversely, suppose I is a k -proper strongly irreducible ideal of S such that $I \cap T = \emptyset$. Let $A \cap B \subseteq T^{-1}I$, where A and B are k -strongly irreducible ideals of $T^{-1}S$. Then $A^c \cap B^c = (A \cap B)^c \subseteq (T^{-1}I)^c$. Note that $I = (T^{-1}I)^c$. Thus $A^c \cap B^c \subseteq I$, and hence, $A^c \subseteq I$ or $B^c \subseteq I$. We then have $A = T^{-1}(A^c) \subseteq T^{-1}I$ or $B = T^{-1}(B^c) \subseteq T^{-1}I$. \square

Motivated by Azizi's results [1, Theorem 3.4] characterizing the relationships among strongly irreducible, prime, and maximal ideals in localized rings, we extend these ideas to the semiring context by formulating their subtractive counterparts in localized semirings.

Proposition 5.3. *If S is a semiring, then the following are equivalent.*

- (1) *Every k -primary ideal of S is a k -strongly irreducible ideal.*
- (2) *For any k -prime ideal P of S , every k -primary ideal of S_P is a k -strongly irreducible ideal.*
- (3) *For any k -maximal ideal P of S , every k -primary ideal of S_P is a k -strongly irreducible ideal.*

Proof. (1) \Rightarrow (2): Let I be a k -primary ideal of S_P . Then I^c is a k -ideal. Let $xy \in I^c$. This implies $\frac{xy}{1} \in I$, which further implies $\frac{x}{1} \cdot \frac{y}{1} \in I$. It follows that, $\frac{x}{1} \in I$ or $(\frac{y}{1})^n \in I$. If $\frac{x}{1} \in I$, we have $x \in I^c$, and if $\frac{y^n}{1} = (\frac{y}{1})^n \in I$, we get $y^n \in I^c$. Now, $I^c \cap (S \setminus P) = \emptyset$, $I^c \in C$. By our assumption, I^c is a k -strongly irreducible ideal of S_P . Therefore, by Theorem 5.2, we obtain $I = T^{-1}I^c$, where $T = S \setminus P$, implying that I is a k -strongly irreducible ideal of S_P .

(2) \Rightarrow (3): Since every k -maximal ideal is k -prime, the proof is clear.

(3) \Rightarrow (1): Suppose, I is a k -primary ideal of S and let M be a k -maximal ideal of S . Then by Corollary 6.7, $I_M = T^{-1}I$, where $T = S \setminus M$, is a k -primary ideal of S_M . Then by our assumption, I_M is a k -strongly irreducible ideal of S . As I is a k -primary ideal of S , we have $(I_M)^c = I$. Thus, I is k -strongly irreducible. \square

The next lemma provides a structural blueprint of k -ideals under localisation.

Lemma 5.4. *Let T be a multiplicative closed subset of a semiring S and $f: S \rightarrow T^{-1}S$ be the natural homomorphism defined by $f(r) := \frac{r}{1}$. If I is a k -ideal of S , then $I^{ec} = \bigcup_{s \in S} (I : s)$. Hence $I^e = \langle 1 \rangle$ if and only if I meets T .*

Proof. If I is a k -ideal of S , then

$$I^e = \mathcal{C}_k(\langle f(I) \rangle) = \mathcal{C}_k(T^{-1}I) = T^{-1}I.$$

Thus

$$x \in I^{ec} = (T^{-1}I)^c \iff \frac{x}{1} \in T^{-1}I \iff \frac{x}{1} = \frac{i}{s} \iff xs = i \in I \iff x \in \bigcup_{s \in T} (I : s),$$

for some $i \in I$, $s \in T$. Next, let $I^e = \langle 1 \rangle = T^{-1}S$. This implies $1 \in I^e$. Then $1 = \frac{1}{1} \in I^{ec} = \bigcup_{s \in T} (I : s)$. Therefore $1 \cdot s' \in I$, for some $s' \in T$. So, $I \cap T \neq \emptyset$. Thus I meets T . Conversely, suppose $x \in I \cap T$. Then $\frac{x}{1} \in T^{-1}I = I^e$. So, $1 = (\frac{1}{x}) \cdot (\frac{x}{1}) \in I^e$. Therefore we get $I^e = \langle 1 \rangle$. \square

Proposition 5.5. *Let $\varphi: S \rightarrow S'$ be a semiring homomorphism and $\varphi_!: \text{Spec}_k(S') \rightarrow \text{Spec}_k(S)$ be the associated map of φ defined by $\varphi_!(Q) := Q^c$. Let P and Q respectively be k -prime ideals of S and S' . Then*

- (1) P is the k -contraction of a k -prime ideal of S' if and only if $P^{ec} = P$.
- (2) P is a k -contracted ideal if and only if $\varphi_!$ is surjective.
- (3) Q is a k -extended ideal implies that $\varphi_!$ is injective.

Proof. (1) Let $P = Q^c$, where Q is a k -prime ideal of S' . Then $P = Q^c = \varphi^{-1}(Q)$. Therefore $\varphi(P) = Q$. Obviously $P \subseteq P^{ec}$. For the reverse inclusion, let $x \in P^{ec}$. This implies $x \in \varphi^{-1}(P^e)$, which further implies $\varphi(x) \in P^e = \mathcal{C}_k(\langle \varphi(P) \rangle)$. Then $\varphi(x) + r \in \langle \varphi(P) \rangle$, for some $r \in \langle \varphi(P) \rangle = \langle Q \rangle$. Therefore $r = \sum_{i=1}^n r_i q_i$, where $r_i \in S'$ and $q_i \in Q$, for all $i = 1, 2, \dots, n$. Since $q_i \in Q$, $r_i \in S'$, we have $r = \sum_{i=1}^n r_i q_i \in Q$. So, $\varphi(x) + r \in Q$. This implies $\varphi(x) \in Q$, as Q is a k -ideal of S' . Therefore $x \in \varphi^{-1}(Q) = Q^c = P$. Thus $P^{ec} = P$. Conversely, suppose $P^{ec} = P$. Let T be the image of $S \setminus P$ in S' . Now

$$S \setminus P = S \setminus P^{ec} = S \setminus \varphi^{-1}(P^e) = \varphi^{-1}(S' \setminus P^e).$$

Therefore, P^e does not meet T . Since P is a k -ideal of S , P^e is a k -ideal of S' . Then by Lemma 5.4, $(P^e)^e$ is a proper k -ideal of $T^{-1}S'$. So, it is contained in a k -maximal ideal M of $T^{-1}S'$, i.e., $(P^e)^e$ is contained in a k -prime ideal M of $T^{-1}S'$. Let Q be the contraction of M in S' . Then $Q = M^c$. Therefore, Q is k -prime, as M is k -prime. Now,

$$P^e \subseteq (P^e)^e \subseteq M^c = Q.$$

Let $r \in Q \cap T$, if possible. Then $r \in Q = M^c$, which implies $\frac{r}{1} \in M$. This further implies $1 = (\frac{1}{r}) \cdot (\frac{r}{1}) \in M$, which is impossible. Therefore $Q \cap T = \emptyset$. Since $P^e \subseteq Q$, we have $P^{ec} \subseteq Q^c$, which gives $P \subseteq Q^c$. If there exists some $x \in Q^c$ such that $x \notin P$, then $\varphi(x) \in Q$ and $\varphi(x) \in \varphi(S \setminus P) = T$. This contradicts the fact that $Q \cap T = \emptyset$. Thus, $P = Q^c$.

(2) Let $P \in \text{Spec}_k(S)$. Suppose P is a k -contracted ideal. Therefore $P = I^c$, for some k -ideal I of S' . Now

$$P^{ec} = I^{cec} = I^c = P.$$

This implies P is the k -contraction of a k -prime ideal, say J . Then $J \in \text{Spec}_k(S')$ and $J^c = P$. Thus, $\varphi(J) = J^c = P$, that is, $\varphi_!$ is surjective. Conversely, suppose that $\varphi_!$ is surjective. Let P be a k -prime ideal of S . Then there exists $Q \in \text{Spec}_k(S')$ such that $\varphi_!(Q) = P$, which implies $Q^c = P$. Hence, P is a k -contracted ideal.

(3) Let every k -prime ideal of S' is a k -extended ideal. If possible, let there exist $Q_1, Q_2 \in \text{Spec}_k(S')$ such that $\varphi_!(Q_1) = \varphi_!(Q_2)$. This implies $Q_1^c = Q_2^c$. Now, $Q_1 = J_1^e$ and $Q_2 = J_2^e$, for some k -ideals J_1, J_2 of S . $Q_1^c = Q_2^c$ implies $Q_1^{ce} = Q_2^{ce}$, which further implies $J_1^{ece} = J_2^{ece}$. So, $J_1^e = J_2^e$. Therefore $Q_1 = Q_2$. Thus $\varphi_!$ is injective. \square

Lemma 5.6. *Let S be a semiring and T be a multiplicatively closed subset of S . Then*

- (1) I^c is a k -ideal of S ,
- (2) $T^{-1}I$ is a k -ideal of $T^{-1}S$.

Proof. (1) Let $x, x+y \in I^c$. Then $\frac{x}{1}, \frac{x+y}{1} \in I$. This implies $\frac{x}{1} + \frac{y}{1} \in I$, which further implies $\frac{y}{1} \in I$ as I is a k -ideal. Therefore $y \in I^c$.

(2) Let $\frac{x}{s_1}, \frac{x}{s_1} + \frac{y}{s_2} \in T^{-1}I$. Then $\frac{xs_2+ys_1}{s_1s_2} = \frac{i}{s}$, for some $i \in I$ and $s \in T$. This implies $xs_2s + ys_1s = is_1s_2 \in I$. Now, $\frac{x}{s_1} = \frac{i'}{s'}$, for some $i' \in I$ and $s' \in T$, which implies $xs' \in I$, and so, $xs_2ss' \in I$. Therefore, $xs_2ss' + ys_1ss' \in I$ implying $ys_1ss' \in I$, as I is a k -ideal. Then $\frac{y}{s_2} = \frac{ys_1ss'}{s_2s_1ss'} \in T^{-1}I$, i.e., $\frac{y}{s_2} \in T^{-1}I$. Hence, $T^{-1}I$ is a k -ideal of $T^{-1}S$. \square

As an extension to the results established by Atani in [2, Lemma 9] in the study of localisation of semirings, we examine the behaviour of specific classes of subtractive ideals under the operation of taking k -radicals.

Proposition 5.7. *Let I be a k -primary ideal of a semiring S with $\mathcal{R}_k(I) = P$. Then the following hold:*

- (1) IS_T is a k -primary ideal of S_T .
- (2) If $P \cap T = \emptyset$, then $IS_T \cap S = I$.
- (3) If $P \cap T = \emptyset$, then $(IS_T :_{S_T} PS_T) = (I :_S P)S_T$.
- (4) If $P \cap T = \emptyset$, and J is a k -ideal of S such that $JS_T \subseteq IS_T$, then $J \subseteq I$.

Proof. (1) If I is a k -primary ideal of S , then IS_T is a k -ideal of S , by Lemma 5.6. Let $(\frac{x}{s_1}) \cdot (\frac{y}{s_2}) \in IS_T$, where $\frac{x}{s_1}, \frac{y}{s_2} \in S_T$. Then $\frac{xy}{s_1s_2} = \frac{i}{s}$, for some $i \in I$, $s \in T$. So, $xys \in I$. Since I is a k -primary ideal, either $xs \in I$ or $y^n \in I$, for some natural number n . If $xs \in I$, then $\frac{x}{s_1} = \frac{xs}{ss_1} \in IS_T$. If $y^n \in I$, then $(\frac{y}{s_2})^n = \frac{y^n}{s_2^n} \in IS_T$.

(2) Clearly, we have $I \subseteq IS_T \cap S$. For the reverse inclusion, let $x \in IS_T \cap S$. Then $\frac{x}{1} \in IS_T$. Therefore, there exists some $i \in I$ and $s \in T$ such that $\frac{x}{1} = \frac{i}{s}$. This implies $xs = i \in I$. Since I is k -primary, either $x \in I$ or $s^n \in I$ for some natural number n . Now, $s^n \in I$ implies $s \in \mathcal{R}_k(I) = P$, which is a contradiction. So, $x \in I$.

(3) Let $\frac{x}{s} \in (I :_S P)$. Then $xP \subseteq I$. Therefore, for all $\frac{p}{s_1} \in PS_T$, we have $\frac{xp}{s_1s} \in IS_T$. This implies that $(\frac{x}{s}) \cdot (\frac{p}{s_1}) \in IS_T$, which further implies, $\frac{x}{s} PS_T \subseteq IS_T$. So, $\frac{x}{s} \in (IS_T :_{S_T} PS_T)$. We have to show $xy \in I$, for all $y \in P$. Now, for all $y \in P$, clearly $(\frac{x}{s}) \cdot (\frac{y}{1}) = \frac{xy}{s} \in IS_T$. Then, there exist $a \in I$, $b \in T$ such that $\frac{xy}{s} = \frac{a}{b}$.

Therefore $(xy)b \in I$. As I is a k -primary ideal of S , either $xy \in I$ or $b^n \in I$, for some natural number n . If $b^n \in I$, we have $b \in \mathcal{R}_k(I) = P$, a contradiction. Thus $xy \in I$. Hence the proof.

(4) Let $x \in J$. Since $\frac{x}{1} \in JS_T \subseteq IS_T$, there exist $i \in I$ and $s \in T$ such that $\frac{x}{1} = \frac{i}{s}$, which implies $xs \in I$. Therefore, I is a k -primary ideal implies $x \in I$ or $s^n \in I$. If $s^n \in I$, then we have $s \in \mathcal{R}_k(I) = P$, a contradiction to the fact that $P \cap T = \emptyset$. Therefore, $x \in I$. Hence $J \subseteq I$. \square

Proposition 5.8. *Let I be a k -ideal of a semiring S . If IS_T is a k -strongly irreducible ideal of S_T , then $IS_T \cap S$ is a k -strongly irreducible ideal of S .*

Proof. Let IS_T be a k -strongly irreducible ideal of S_T . Suppose $A \cap B \subseteq IS_T \cap S$, where A, B are two k -ideals of S . Then

$$(A \cap B)S_T \subseteq (IS_T \cap S)S_T = IS_T,$$

by [2, Lemma 4]. This implies $AS_T \cap BS_T \subseteq IS_T$. So, $AS_T \subseteq IS_T$ or $BS_T \subseteq IS_T$, as IS_T is k -strongly irreducible. Therefore, $AS_T \cap S \subseteq IS_T \cap S$ or $BS_T \cap S \subseteq IS_T \cap S$, which further implies, $A \subseteq AS_T \cap S \subseteq IS_T \cap S$ or $B \subseteq BS_T \cap S \subseteq IS_T \cap S$, i.e., $A \subseteq IS_T \cap S$ or $B \subseteq IS_T \cap S$. \square

Theorem 5.9. *Let I be a k -strongly irreducible primary ideal of a semiring S such that $\mathcal{R}_k(I) \cap T = \emptyset$. Then IS_T is k -strongly irreducible.*

Proof. Let I be a k -strongly irreducible primary ideal of S . Suppose A, B be two k -ideals of S_T such that $A \cap B \subseteq IS_T$. Then

$$(A \cap S) \cap (B \cap S) \subseteq IS_T \cap S = I,$$

by Proposition 5.7. Let $x, x+y \in A \cap S$. Then $\frac{x}{1}, \frac{x+y}{1} \in A$. This implies $\frac{x}{1} + \frac{y}{1} \in A$. Therefore $\frac{y}{1} \in A$ as A is a k -ideal of S_T , which implies $y \in A \cap S$. So, $A \cap S$ and $B \cap S$ are k -ideals of S . Since I is k -strongly irreducible $A \cap S \subseteq I$ or $B \cap S \subseteq I$. Thus $A = (A \cap S)S_T \subseteq IS_T$ or $B = (B \cap S)S_T \subseteq IS_T$. \square

Theorem 5.10. *Assume that I is a k -primary ideal of a semiring S with $\mathcal{R}_k(I) \cap T = \emptyset$ and let IS_T be k -strongly irreducible ideal of S_T . Then I is k -strongly irreducible.*

Proof. Let IS_T be a k -strongly irreducible ideal of S_T . Since I is a k -primary ideal such that $\mathcal{R}_k(I) \cap T = \emptyset$, by Proposition 5.7 we have $IS_T \cap S = I$. Now, by Theorem 5.8, it follows that $I = IS_T \cap S$ is a k -strongly irreducible ideal of S . \square

6. IN SPECIAL SEMIRINGS

This section is bifurcated in scope. First, we investigate the behaviour of k -ideals across several distinguished classes of semirings, including Laskerian semirings, von Neumann regular semirings, unique factorisation semidomains, principal ideal semidomains, and weakly Noetherian semirings. The subsequent subsection is devoted to a focused analysis of k -ideals in the specialized setting of additively idempotent semirings.

We begin by recalling two important classes of semirings, obtained as natural generalizations of the corresponding ring-theoretic notions.

Definition 6.1. Let S be a semiring. Then

- (1) S is said to be *Laskerian* if every k -ideal has a k -primary decomposition.
- (2) S is said to be *von Neumann regular* if for all $a \in S$ there exists $x \in S$ such that $a = a^2x$.

In the results that follow, we analyse the interplay between k -strongly irreducible ideals and other distinguished subclasses of k -ideals, notably k -prime and k -primary ideals, across a range of semiring classes.

Proposition 6.2. Suppose S is a semiring.

- (1) If I is a k -strongly irreducible ideal of S , then I is a k -prime ideal of S if and only if $I = \mathcal{R}_k(I)$.
- (2) If S is a Laskerian semiring, then every k -strongly irreducible ideal is a k -primary ideal.
- (3) If S is a von Neumann regular semiring, then an ideal is k -strongly irreducible if and only if it is a k -primary ideal.

Proof. (1) If I is k -prime, then clearly $I = \mathcal{R}_k(I)$. Conversely, suppose $I = \mathcal{R}_k(I)$. Let $AB \subseteq I$, where A, B are two k -ideals of S . Then

$$A \cap B \subseteq \mathcal{R}_k(A \cap B) = \mathcal{R}_k(A) \cap \mathcal{R}_k(B) = \mathcal{R}_k(AB) \subseteq \mathcal{R}_k(I) = I.$$

Therefore, $A \subseteq I$ or $B \subseteq I$, as I is k -strongly irreducible. Hence I is k -prime.

- (2) Let I be a k -strongly irreducible ideal of S and suppose $\bigcap_{i=1}^n A_i$ be a k -primary decomposition of I .

Then $\bigcap_{i=1}^n A_i \subseteq I$. As I is k -strongly irreducible, we have $A_j \subseteq I$, for some $j \in \{1, 2, \dots, n\}$. This implies $A_j \subseteq I \subseteq A_j$, which further implies $I = A_j$. Thus I is k -primary.

(3) In a von Neumann regular semiring, k -prime and k -primary ideals are the same, by Theorem 3.9. Let I be a k -strongly irreducible ideal of S . To show that I is a k -primary ideal, it is enough to show that I is k -prime. Let $ab \in I$. Since S is a von Neumann regular semiring, there exist $r_1, r_2 \in S$ such that $a = a^2 r_1$, $b = b^2 r_2$. Suppose, $c \in \mathcal{C}_k(\langle a \rangle) \cap \mathcal{C}_k(\langle b \rangle) = \mathcal{C}_k(\langle a \cap b \rangle)$. Then $c + x_1 = x_2$, for some $x_1, x_2 \in \langle a \rangle \cap \langle b \rangle$. Now, $x_1 = am_1 = bm_2$, for $m_1, m_2 \in S$ and $x_2 = an_1 = bn_2$, for $n_1, n_2 \in S$. Then

$$x_1 = am_1 = a^2 r_1 m_1 = am_1 ar_1 = bm_2 ar_1 = abm_2 r_1.$$

Similarly, $x_2 = an_1 = a^2 r_1 n_1 = an_1 ar_1 = bn_2 ar_1 = abn_2 r_1$. Therefore, $c + x_1 = x_2$ implies $c + abm_2 r_1 = abn_2 r_1$. So, $c \in \mathcal{C}_k(\langle ab \rangle) \subseteq \mathcal{C}_k(I) = I$. Since I is k -strongly irreducible, we must have $a \in I$ or $b \in I$. \square

Remark 6.3. In a bounded distributive lattice, each k -ideal is a k -radical ideal. Therefore, by Proposition 6.2, every k -strongly irreducible ideal is k -prime. This confirms the result proved in [22, Theorem 5].

For rings, Azizi examined various properties of strongly irreducible ideals in unique factorisation domains [1, Theorem 2.2]. We now present subtractive analogous of these properties, extending the discussion to the present framework. The proof of the following lemma is identical to the corresponding result (dropping the ‘ k -part’) for rings.

Lemma 6.4. Let S be a unique factorisation semidomain and I be a k -proper ideal of S . Then

- (1) I is k -strongly irreducible if and only if $p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k} \in I$, where p_i are distinct prime elements of S and n_i are natural numbers, implies that $p_j^{n_j} \in I$, for some j , with $1 \leq j \leq k$.
- (2) If I is a non-zero principal k -ideal, then I is k -strongly irreducible if and only if the generator of I is a power of a prime element of S .

Theorem 6.5. Let S be a unique factorisation semidomain and let I be a k -proper ideal of S .

- (1) I is k -strongly irreducible if and only if for each $x, y \in S$, $\text{l.c.m.}\{x, y\} \in I$ implies that $x \in I$ or $y \in I$.
- (2) Every k -strongly irreducible ideal is a k -primary ideal.

Proof. (1) Let I be a k -strongly irreducible ideal of S and let $z := \text{l.c.m.}\{x, y\} \in I$, for some $x, y \in S$. Now $z \in \langle x \rangle \cap \langle y \rangle$. In particular, as z is the l.c.m., we have $\langle x \rangle \cap \langle y \rangle = \langle z \rangle \subseteq I$. Therefore, $\mathcal{C}_k(\langle x \rangle) \cap \mathcal{C}_k(\langle y \rangle) = \mathcal{C}_k(\langle z \rangle) \subseteq \mathcal{C}_k(I) = I$. This implies $x \in I$ or $y \in I$, by Theorem 3.11. Conversely, suppose that for each $x, y \in S$, $\text{l.c.m.}\{x, y\}$ implies $x \in I$ or $y \in I$. Let $\mathcal{C}_k(\langle x \rangle) \cap \mathcal{C}_k(\langle y \rangle) \subseteq I$, which implies

$$\langle x \rangle \cap \langle y \rangle \subseteq \mathcal{C}_k(\langle x \rangle \cap \langle y \rangle) = \mathcal{C}_k(\langle x \rangle) \cap \mathcal{C}_k(\langle y \rangle) \subseteq I.$$

Therefore $\text{l.c.m.}\{x, y\} \in \langle x \rangle \cap \langle y \rangle \subseteq I$. So, we get $x \in I$ or $y \in I$, by hypothesis. Hence, I is k -strongly irreducible.

(2) Suppose I is a k -strongly irreducible ideal of S and $x + I$ be a non-zero element of S/I such that $(x + I)(y + I) = 0 + I$, where $y + I \neq I$. This implies $xy + I = 0 + I$. Therefore, $xy \in I$. Let

$$x := p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k} q_1^{l_1} q_2^{l_2} \cdots q_r^{l_r}$$

$$y := p_1^{s_1} p_2^{s_2} \cdots p_k^{s_k} t_1^{m_1} t_2^{m_2} \cdots t_u^{m_u}$$

be prime decomposition of x and y respectively. Since $xy \in I$, by Lemma 6.4 we have one of the following:

- (i) for some $1 \leq j \leq k$, $p_j^{n_j+s_j} \in I$,
- (ii) for some $1 \leq j \leq r$, $q_j^{l_j} \in I$,
- (iii) for some $1 \leq j \leq u$, $t_j^{m_j} \in I$.

Let (i) hold and n be a natural number such that $n \geq l_j/n_j$. This implies $(n+1)n_j \geq n_j + s_j$. Therefore, $p_j^{n_j+s_j} | p_j^{(n+1)n_j}$, and clearly, $p_j^{(n+1)n_j} | x^{(n+1)}$. Then $p_j^{n_j+l_j} | x^{(n+1)}$. As $p_j^{n_j+l_j} \in I$, we have $x^{n+1} \in I$, which implies $x^{n+1} + I = I$. So $(x + I)^{n+1} = x^{n+1} + I = I$. Thus $x + I$ is nilpotent.

If (ii) holds, $q_j^{l_j} \in I$ implies $x \in I$. Then $x + I = I$, not possible. Similarly, (iii) cannot hold. Hence, I is a primary k -ideal of S . \square

As a consequence of the above results, we have the following corollaries.

Corollary 6.6. *Let S be a unique factorisation semidomain.*

- (1) *Every k -principal ideal of S is a k -strongly irreducible ideal if and only if it is a k -primary ideal.*
- (2) *Every k -strongly irreducible ideal of S can be generated by a set of prime powers.*

Corollary 6.7. *Let S be a semiring and let T be a multiplicatively closed subset of S .*

- (1) *If I is a k -strongly irreducible and a k -primary ideal of S which does not meet T , then $T^{-1}I$ is a k -strongly irreducible (and a k -primary) ideal of $T^{-1}S$.*
- (2) *If S is a unique factorisation semidomain or a von Neumann regular semiring, then there is a one-to-one correspondence between the k -strongly irreducible ideals of $T^{-1}S$ and the k -strongly irreducible ideals of S which do not meet T .*

Proof. (1) We first show that if I is a k -primary ideal of S and $I \cap T = \emptyset$, then $(T^{-1}I)^c = I$. Clearly $I \subseteq (T^{-1}I)^c$. Let $x \in (T^{-1}I)^c$. Then $\frac{x}{1} \in T^{-1}I$. This implies $\frac{x}{1} = \frac{i}{s}$, for some $i \in I$ and $s \in T$, which further implies $xs = i \in I$. Let $x \notin I$. Then $s^n \in I$, for some natural number n . This implies $s^n \in I \cap T$, a contradiction. Therefore $x \in I$. Thus $(T^{-1}I)^c = I$, which implies I is contained in C . Then it follows by Theorem 5.2 that $T^{-1}I$ is a k -strongly irreducible ideal of $T^{-1}S$. It remains to show that $T^{-1}I$ is a k -primary ideal. Suppose $xy \in T^{-1}I$. There must exist $u_1, u_2 \in S$ and $s_1, s_2 \in T$ such that $x = \frac{u_1}{s_1}$ and $y = \frac{u_2}{s_2}$. Then $u_1 u_2 \in I$, and hence, $u_1 \in I$ or $u_2^n \in I$. Thus $x \in T^{-1}I$ or $y^n \in I$.

(2) It follows from Theorem 6.5 and Proposition 6.2 that in a unique factorisation semidomain and in a von Neumann regular semiring, every k -strongly irreducible ideal is a k -primary ideal. Then, by (1), $T^{-1}I$ is a k -strongly irreducible ideal of $T^{-1}S$. Also, we have, by Theorem 5.2, for each k -strongly irreducible ideal I of $T^{-1}S$, I^c is a k -strongly irreducible ideal of S that does not meet T . \square

The converse of Lemma 3.6 fails to hold in general semirings. The following theorem shows that this converse is valid in the more restrictive setting of a *principal ideal semidomain*.

Theorem 6.8. *Each non-zero k -prime ideal of a principal ideal semidomain is k -maximal.*

Proof. Let S be a principal ideal semidomain and P be a k -prime ideal of S . Then $P = \langle p \rangle$, for some element p of S . Suppose Q is a k -ideal of S containing P . Then $Q = \langle q \rangle$, for some $q \in S$. Since $p \in Q$, we have $p = rq$, for some non-zero $r \in S$. As P is k -prime, $rq \in P$ implies $r \in P$ or $q \in P$. If $q \in P = \langle p \rangle$, we have $Q = P$. If $r \in P = \langle p \rangle$, then $r = pr' = rqr'$, for some $r' \in S$. As S is a semidomain and $r \neq 0$, $r = rqr'$ implies $qr' = 1$, which further implies $Q = S$. \square

Recall that a partially ordered ring $(R, +, \cdot, \leq)$ is ring with a partial order such that for all $a, b, c \in R$ and $a \leq b$ implies $a + c \leq b + c$, and $0 \leq a$, $0 \leq b$ implies $0 \leq ab$. The positive cone of $(R, +, \cdot, \leq)$ is the set $R^+ = \{x \in R \mid 0 \leq x\}$. The positive cone of a partially ordered semiring is always a *cancellative* and *conic* (or, *zero-sum free*) semiring, that is, $a + b = a + c$ implies $b = c$ and $a + b = 0$ implies $a = 0$ and $b = 0$.

Lemma 6.9. *Let S be a positive cone of a partially ordered ring R such that $D(S) = R$, where $D(S) := \{a - b \mid a, b \in S\}$. Then any principal ideal of S is a k -ideal.*

Proof. For an element $a \in S$, let $x + y \in \langle a \rangle$ and $x \in \langle a \rangle$. Therefore $x + y = at$ and $x = as$, for some $s, t \in S$. Since S is the positive cone of R , we have $0 \leq x, y$ and $y \leq x + y$, which further implies that $y = a(t - s) \in D(S)$ and $0 \leq t - s$. Therefore $t - s \in S$ and hence $y \in \langle a \rangle$. \square

The next theorem states an elementary property of a k -prime ideal in a *unique factorisation semidomain*.

Theorem 6.10. *In a UFSD every non-zero k -prime ideal of S contains a prime element.*

Proof. Let S be a UFSD and I be a non-zero k -prime ideal of S . Let $x \in I$ be a non-zero and non-unit element. Then $x = p_1 p_2 \cdots p_n$, where p_i 's are distinct prime elements of S for each i . Since I is a prime ideal of S , there exists an i such that $p_i \in I$. \square

The converse of the above theorem is true if S is a cancellative, partially ordered, conic semiring.

Theorem 6.11. *If S is a cancellative, partially ordered, conic semidomain and if every non-zero k -prime ideal of S contains a prime element, then S is a UFSD.*

Proof. Suppose S is a cancellative, partially ordered, conic semidomain and every non-zero k -prime ideal of S contains a prime element. We assume that S is not a UFSD. Let

$$T = \{u \in S \mid u \text{ is a unit}\} \bigcup \{p_1 p_2 \cdots p_n \in S\}.$$

Then T is a multiplicatively closed saturated subset of S . Now, S is a UFSD if and only if $T = S \setminus \{0\}$. As S is not a UFSD, there exists a non-zero $x \in S$ such that $x \notin T$. By Lemma 6.9, it is clear that $\langle x \rangle$ is a k -ideal. We claim that $\langle x \rangle \cap T = \emptyset$. As if $rx \in T$, for some $r \in S$, then $r \in T$. Therefore, $\langle x \rangle$ can be expanded to a k -prime ideal I of S which is disjoint from T . Then, by hypothesis, I contains a prime element, a contradiction. This completes the proof. \square

Let us recall that in a *weakly Noetherian semiring* every set of k -ideals satisfies the ascending chain condition. The following proposition reveals an interesting structural property of k -irreducible ideals in *weakly Noetherian semirings*.

Proposition 6.12. *Let S be a weakly Noetherian semiring. Then every k -irreducible ideal is k -primary.*

Proof. Let I be a k -irreducible ideal of S and $ab \in I$ with $b \notin I$. For each $n \geq 1$, let $I_n := \{r \in S \mid a^n r \in I\}$. Then each I_n is a k -ideal of S such that $I_n \subseteq I_{n+1}$, for all n . Therefore, there exists $m \in \mathbb{N}$ such that $I_n = I_{n+1}$, for all $n \geq m$ as S is weakly Noetherian. Let $A = \langle a^m \rangle + I$, $B = \langle b \rangle + I$. We show that $A \cap B = I$. Let $x \in A \cap B$. Then $x = a^m r + i_1$, for some $r \in S$ and $i_1 \in I$. Also $x = bs + i_2$, for some $s \in S$ and $i_2 \in I$. So, $ax = abs + ai_2 \in \langle ab \rangle + I$. Since $ab \in I$ $ax \in I$. Now $ax = a^{m+1}r + ai_1 \in I$. Then $a^{m+1}r \in I$, as I is a k -ideal.

This implies $r \in I_{m+1} = I_m$, i.e., $a^m r \in I$, which further implies $x \in I$. Since I is k -irreducible and $b \notin I$, we get $A = I$. Therefore, it follows that $a^m \in i$. Hence I is k -primary. \square

6.1. Additively idempotent semirings. In this final segment, we investigate variants of k -ideals and k -congruences in the milieu of additively idempotent semirings. It is worth emphasizing that additively idempotent semirings may deviate substantially from the class of subtractive semirings (cf. [12, Example 6.39]), thereby rendering the study of k -ideals particularly compelling. This perspective enables us to strengthen several results of [29, 30], where additively idempotent semirings are studied through the lens of B_1 -algebras. Here B_1 denotes the two-element Boolean algebra $\{0, 1\}$, commonly referred to as the *Boolean semifield*. We recall that, in additively idempotent semirings, the product of two k -ideals is again a k -ideal; see [24, Lemma 3.27]. This property fails in the broader class of general semirings; see [12, Example 6.43].

The following definition is a subtractive version of the definition of cancellation ideals, first introduced by LaGrassa in her thesis [27].

Definition 6.13. A non-zero k -ideal I of a semiring S is called a *k -cancellation ideal*, if $IJ = IK$ implies $J = K$, for all k -ideals J and K of S .

The following proposition characterizes k -cancellation ideals in terms of colon ideals and ideal inclusions in an additively idempotent setup.

Proposition 6.14. Let S be an additively idempotent semiring and I be a non-zero k -ideal of S . Then the following statements are equivalent:

- (1) I is a k -cancellation ideal of S .
- (2) $(IJ : I) = J$, for any k -ideal J of S .
- (3) $IJ \subseteq IK$ implies $J \subseteq K$, for all k -ideals J, K of S .

Proof. Let I, J, K be k -ideals of S .

(1) \Rightarrow (2): Obviously, $IJ \subseteq (IJ : I)$. Let $i_1 x_1 + i_2 x_2 + \dots + i_n x_n \in I(IJ : I)$, $i_k \in I$, and $x_k \in I$, for all k . $x_k I \subseteq IJ$, for all k . Therefore, $\sum_{k=1}^n i_k x_k \in IJ$. So, $I(IJ : I) \subseteq IJ$. Now, for all $j \in J$, we have $jI \subseteq IJ$. Therefore $IJ \subseteq I(IJ : I)$. Thus $IJ = I(IJ : I)$. Since S is an additively idempotent semiring, then by [24, Lemma 3.27], IJ is a k -ideal, furthermore by [15, Proposition 2.4], $(IJ : I)$ is a k -ideal. Since I is a k -cancellation ideal, we get $J = (IJ : I)$.

(2) \Rightarrow (3): Suppose, $IJ \subseteq IK$. Let $r \in (IJ : I)$. Then $rI \subseteq IJ \subseteq IK$. This implies $r \in (IK : I)$. So, $(IJ : I) \subseteq (IK : I)$, which further implies $J \subseteq K$.

(3) \Rightarrow (1): Let $IJ = IK$. Then $IJ \subseteq IK, IK \subseteq IJ$, which implies $J \subseteq K$ and $K \subseteq J$. Therefore $J = K$. \square

The next theorem provides a new characterization of k -prime ideals with respect to the natural partial order in any additively idempotent semiring.

Theorem 6.15. Let S be an additively idempotent semiring. Then a k -ideal is k -prime if and only if for $x, y \in S$ and $z \in I$, $xy \leq z$ implies that $x \in I$ or $y \in I$.

Proof. Suppose, for all $x, y \in S$ and $z \in I$, $xy \leq z$ implies $x \in I$ or $y \in I$. We show that I is k -prime. For that, let $y \in I$, $x + y \in I$. Then $x \leq x + y$, which implies $x \cdot 1_S \leq x + y$. Therefore, by assumption, $x \in I$ or $1_S \in I$. This implies $x \in I$ as I is an ideal. Now, let $xy \in I$. Then $xy \leq xy$ implies $x \in I$ or $y \in I$. Conversely, suppose that I is k -prime and for $x, y \in S$ and $z \in I$, $xy \leq z$. This implies $xy + z = z$, which further implies that $xy \in I$ as I is a k -ideal. Therefore, $x \in I$ or $y \in I$. \square

In similar spirit, the following theorem characterizes k -semiprime ideals with respect to the natural partial order of an additively idempotent semiring.

Theorem 6.16. *Let S be an additively idempotent semiring. A k -ideal I is k -semiprime if and only if for $z \in I$ and $x^2 \leq z$ implies that $x \in I$.*

Proof. Suppose I is a k -semiprime ideal of S . Let $x \in S$ and $z \in I$ such that $x^2 \leq z$. Then $x^2 + z = z$. We get $x^2 \in I$ as I is a k -ideal. Since I is k -semiprime, we have $I = \mathcal{R}_k(I)$. Let P be a k -prime ideal containing I . Then $x^2 \in P$ implies $x \in P$, which further implies $x \in \mathcal{R}_k(I) = I$. Conversely, suppose that for $x \in S$ and $z \in I$, $x^2 \leq z$ implies $x \in I$. Let A be a k -ideal of S such that $A^2 \subseteq I$. Let $x \in A$. Then $x^2 \in I$. Therefore, $x^2 \leq x^2$ implies $x \in I$, by our assumption. So, $x \in I$. We get $A \subseteq I$. Thus I is k -semiprime. \square

The following theorem characterizes k -strongly irreducible ideals in an additively idempotent UFSD with respect to the natural partial order and the l.c.m-operation.

Theorem 6.17. *Let S be an additively idempotent unique factorisation semidomain. Then I is a k -strongly irreducible ideal in S if and only if for $x, y \in S$ and $z \in I$, $\text{l.c.m}\{x, y\} \leq z$ implies $x \in I$ or $y \in I$.*

Proof. Suppose, for all $x, y \in S$ and $z \in I$, $\text{l.c.m}\{x, y\} \leq z$ implies $x \in I$ or $y \in I$. Let $\mathcal{C}_k(\langle x \rangle) \cap \mathcal{C}_k(\langle y \rangle) \subseteq I$. Then $\text{l.c.m}\{x, y\} \in \langle x \rangle \cap \langle y \rangle \subseteq I$. This implies $\text{l.c.m}\{x, y\} \in I$. Therefore $\text{l.c.m}\{x, y\} \leq \text{l.c.m}\{x, y\}$. So, we get $x \in I$ or $y \in I$, by our assumption. Conversely, let I be a k -strongly irreducible ideal of S . Suppose $x, y \in S$ and $z \in I$ such that $\text{l.c.m}\{x, y\} \leq z$. Then $\text{l.c.m}\{x, y\} + z = z$. We get $\text{l.c.m}\{x, y\} \in I$, as I is a k -ideal. Hence $x \in I$ or $y \in I$, S being a UFSD. \square

The rest of paper is devoted to study various congruences and associated ideals in the additively idempotent setting. Let us now recall the definition of prime congruences from [29, Definition 3.4].

Definition 6.18. Suppose S is an additively idempotent semiring. A congruence ρ on S is said to be *prime* if $(ab)\rho 0$ implies $a\rho 0$ or $b\rho 0$.

The next definition is the counterpart of the Bourne congruence associated with any ideal in an additively idempotent semiring.

Definition 6.19 (Section 3, [29]). Let S be a semiring and J be an ideal of S . Then ρ_J is a congruence on S defined as follows: $x\rho_J y$ if and only if there exists $z \in J$ such that $x + z = y + z$. this ρ_J is called *excellent congruence* on S . Moreover, the ideal J is called *saturated* if $J = \bar{J} := \{x \in S \mid x + z = z, \text{ for some } z \in J\}$

Remark 6.20. By a k -prime congruence, we mean a congruence that is a k -congruence as well as prime.

In [29, Theorem 3.7], Lescot showed that the map $J \rightarrow \bar{J}$ is a closure operator. We shall prove that \bar{J} coincides with the $\mathcal{C}_k(J)$. For that, we need the following lemma.

Lemma 6.21. *Let S be an additively idempotent semiring and I be an ideal of S . Then \bar{I} is a k -ideal of S .*

Proof. Let $x, x+y \in \bar{I}$. Then $x+z_1 = z_1$ and $x+y+z_2 = z_2$, for some $z_1, z_2 \in I$. This implies $x+z_1+x+y+z_2 = x+z_1+z_2$, which further implies $y+x+z_1+z_2 = x+z_1+z_2$. Therefore $y+z_1+z_2 = z_1+z_2$. So $y \in \bar{I}$. \square

Theorem 6.22. *Let S be an additively idempotent semiring. An ideal I is saturated if and only if it is a k -ideal.*

Proof. If I is saturated, then $I = \bar{I}$. By Lemma 6.21, I is a k -ideal. Conversely, suppose I is a k -ideal. As for $x \in I$, $x+x = x$, we have $x \in \bar{I}$. Therefore $I \subseteq \bar{I}$. Let $x \in \bar{I}$. Then $x+z = z$, for some $z \in I$. This implies $x \in I$. Thus $I = \bar{I}$. \square

The correspondence between k -ideals and k -congruences extends to some of their distinguished subclasses. In particular, we show that in an additively idempotent semiring there is a one-to-one correspondence between k -maximal congruences and k -maximal ideals. We begin by recalling the definition of a k -maximal congruence.

Definition 6.23. Let S be a semiring. A k -congruence θ is said to be k -maximal if $\theta \subseteq \theta'$ implies either $\theta = \theta'$ or $\theta' = S \times S$, for any k -congruence θ' on S .

The following theorem ensures the inclusion of the class of k -maximal congruences into the class of k -prime congruences, for any semiring S .

Theorem 6.24. Let S be a semiring. Then every k -maximal congruence is a k -prime congruence.

Proof. Let $\rho = K_A$ be a k -maximal congruence for some ideal A of S . Let $(uv, 0) \in K_A$ but $(u, 0) \notin K_A$. Then $u \notin A$, as $u + 0 = 0 + u$ will imply uK_A0 . Consider the ideal $B = \langle u \rangle + A$. If xK_Ay , then we have $x + z_1 = x + z_2$, for some $z_1, z_2 \in A$, which implies xK_By . Therefore $K_A \subseteq K_B$. By maximality of K_A , $K_B = S \times S$. In particular, $0K_B1$. Then $0 + us_1 + a = 1 + us_2 + b$, for some $s_1, s_2 \in S$ and $a, b \in A$. So, $us_1K_A(1 + us_2)$. Now

$$(uv)s_1 = v(us_1)K_Av(1 + us_2) = v + uv s_2.$$

Since uvK_A0 we have

$$0 = 0s_1K_A(uv)s_1K_Av + (uv)s_2K_Av + 0 \cdot s_2 = v.$$

Therefore, vK_A0 . \square

Han [19, Theorem 5.4] established a correspondence between k -prime ideals and k -prime congruences, and used this correspondence to show that the resulting spectra are homeomorphic. In the following theorem, we present a purely algebraic proof of the correspondence between k -maximal ideals and k -maximal congruences.

Theorem 6.25. Let S be an additively idempotent semiring and I be a k -ideal of S . Then K_I is a k -maximal congruence if and only if I is a k -maximal ideal.

Proof. Suppose I is a k -maximal ideal. Let $K_I \subseteq \rho$, for some k -congruence ρ on S . Let $x \in I$. Then $x + x = x$. This implies xK_I0 , which further implies $x\rho 0$. So $x \in I_\rho := \{y \in S \mid y\rho 0\}$. As I is k -maximal either $I = I_\rho$ or $I_\rho = S$. Let $I = I_\rho$. Now as ρ is a k -congruence $\rho = K_J$, for some ideal J of S . Then $x\rho y$ implies $x + z_1 = y + z_2$, for some $z_1, z_2 \in J$. Since $z_1, z_2 \in J$, we have $z_1\rho 0$ and $z_2\rho 0$. So, $z_1, z_2 \in I_\rho = I$. Therefore xK_Iy , i.e., $\rho = K_I$. If $I_\rho = S$, we have for all $x \in S$, $x\rho 0$. This implies $x\rho y$, for all $x, y \in S$. Thus $\rho = S \times S$. Conversely, suppose K_I is a k -maximal congruence and $I \subseteq J$, where J is a k -ideal of S . Then $K_I \subseteq K_J$. This implies $K_I = K_J$ or $K_J = S \times S$. Let $K_I = K_J$. For $x \in J$, $x + x = x$ implies xK_J0 , which further implies xK_I0 . Then $x + z_1 = z_2$, for some $z_1, z_2 \in I$. So $x \in I$, that is, $I = J$. If $K_J = S \times S$ for all $x \in S$, $(x, 0) \in K_J$. This implies $x + a = b$, for some $a, b \in J$. Therefore $x \in J$. Hence $J = S$. \square

The next theorem concludes that the class of k -prime congruences is contained in the class of k -irreducible congruences in an additively idempotent semiring S .

Theorem 6.26. Let S be an additively idempotent semiring. Every k -prime congruence is k -irreducible.

Proof. Let $\rho = K_J$ be a k -prime congruence for some ideal J of S . Let $K_A \cap K_B = K_J$, for some k -congruences K_A and K_B , where A and B are ideals of S . We claim that $\bar{A} \cap \bar{B} = \bar{J}$. Let $x \in \bar{A} \cap \bar{B}$. Since saturated closure and k -closure coincide, $\bar{A} \cap \bar{B} = \overline{A \cap B}$ (cf. Theorem 6.22). This implies $x + z = z$, for some $z \in A \cap B$. So $xK_A \cap K_B 0$. Then $xK_J 0$, that is, $x \in \bar{J}$. Again, if $y \in \bar{J}$, we have $yK_A \cap K_B 0$, which implies $y \in \bar{A} \cap \bar{B}$. Therefore $ab \in \bar{J}$ implies $abK_J 0$, where $a \in \bar{A}$, $b \in \bar{B}$. K_J being k -prime, $aK_J 0$ or $bK_J 0$. Thus $\bar{A} = \bar{J}$ or $\bar{B} = \bar{J}$. Let $\bar{A} = \bar{J}$. Now, since for $x \in A$, $x + x = x$ holds, we have $xK_A 0$. Therefore, $x \in \bar{A}$. So $A \subseteq \bar{A} = \bar{J}$. Let $xK_A y$. Then $x + z_1 = y + z_2$, where $z_1, z_2 \in A \subseteq \bar{J}$. This implies $z_1 + a = a$, $z_2 + b = b$, for some $a, b \in J$. Whence

$$x + z_1 + a + b = y + z_2 + a + b,$$

that is, $x + a + b = y + a + b$. This gives $xK_J y$. Therefore $K_A = K_J$. \square

The accompanying result establishes that, in an additively idempotent semiring, the congruence associated with any k -irreducible ideal is itself k -irreducible. The validity of the converse implication, however, remains an open problem.

Theorem 6.27. *Let S be an additively idempotent semiring. If I is a k -irreducible ideal, K_I is a k -irreducible congruence.*

Proof. I is a k -irreducible ideal. Let A_1, A_2 be two k -congruences such that $K_{A_1} \cap K_{A_2} = K_I$. We claim that $\mathcal{C}_K(A_1) \cap \mathcal{C}_K(A_2) = I$. Let $x \in \mathcal{C}_K(A_1) \cap \mathcal{C}_K(A_2)$. Then, there exist $z_1, z_2 \in A_1$ and $y_1, y_2 \in A_2$ such that $x + z_1 = z_2$ and $x + y_1 = x + y_2$. So $x \in K_{A_1} \cap K_{A_2} = K_I$. Therefore, $x \in I$ as I is a k -ideal. Again, for $a \in I$, $(a, 0) \in K_I = K_{A_1} \cap K_{A_2}$. Then $a + a_1 \in A_1$ and $a + a_2 \in A_2$, for some $a_1 \in A_1$ and $a_2 \in A_2$. It follows that $a \in \mathcal{C}_K(A_1) \cap \mathcal{C}_K(A_2)$. Thus $\mathcal{C}_K(A_1) \cap \mathcal{C}_K(A_2) = I$. So $\mathcal{C}_K(A_1) = I$ or $\mathcal{C}_K(A_2) = I$. Suppose $\mathcal{C}_K(A_1) = I$ and let $xK_{A_1}y$, which implies $x + c_1 = x + c_2$, for some $c_1, c_2 \in A_1$. Since $A_1 \subseteq I$, $c_1, c_2 \in I$, we have $(x, y) \in K_I$. Hence $K_{A_1} = K_I$. \square

The following theorem relaxes Theorem 6.27 by replacing k -irreducibility with the weaker notion of excellent irreducibility, arising from ideals whose saturated closures are k -irreducible.

Theorem 6.28. *Let S be an additively idempotent semiring and let \bar{J} be a k -irreducible ideal of S . then $\rho_{\bar{J}}$ is an excellent-irreducible congruence on S .*

Proof. Let $\rho_{J_1} \cap \rho_{J_2} = \rho_J$, where ρ_{J_1}, ρ_{J_2} are two excellent congruences on S . Define $I_{J_i} := \{x \in S \mid x\rho_{J_i}0\}$, for $i = 1, 2$. Both I_{J_1}, I_{J_2} are k -ideals of S . Let $x \in I_{J_1} \cap I_{J_2}$. Then $x\rho_{J_1}0 \cap x\rho_{J_2}0$. This implies $x\rho_J0$. Therefore $x \in \bar{J}$. For the reverse inclusion, let $y \in \bar{J}$. Then there exists $z \in J$ such that $y + z = z$, which implies $y\rho_J0$, that is, $y\rho_{J_1}0 \cap y\rho_{J_2}0$. Thus $y \in I_{J_1} \cap I_{J_2}$. Now, as \bar{J} is k -irreducible, $I_{J_1} = \bar{J}$ or $I_{J_2} = \bar{J}$. Suppose $I_{J_1} = \bar{J}$. Let $x \in J_1$. Then $x + x = x$ implies $x\rho_{J_1}0$. Therefore $x \in I_{J_1} = \bar{J}$. If $x\rho_{J_1}y$, then there exists $a \in J_1$ such that $x + a = y + a$. Now $a \in \bar{J}$ implies $a + z = z$, for some $z \in J$. So, $x + a + z = y + a + z$, i.e., $x + z = y + z$, whence $x\rho_Jy$. Hence $\rho_{J_1} = \rho_J$. \square

Next, we recall from [29, Definition 3.10], a proper subclass of prime ideals and define its subtractive counterpart.

Definition 6.29. An ideal I of a semiring S is said to be *absolutely prime* if $I \neq S$ and $(ab)\rho_I(ac)$ implies $a \in \bar{I}$ or $b\rho_Ic$. An ideal I is said to be *absolutely k -prime* if it is a k -ideal and absolutely prime.

Theorem 6.30. *Let S be an additively idempotent semiring. Then every absolutely k -prime ideal is k -prime.*

Proof. Let I be an absolutely k -prime ideal and $ab \in I$. Then $ab + ab = ab$ implies $(ab)\rho_I0$, that is, $(ab)\rho_I(a \cdot 0)$, which further implies $b\rho_I0$ or $a \in \bar{I} = I$. If $b\rho_I0$ we have $b + z = z$, for some $z \in I$, whence $b \in I$. Hence I is k -prime. \square

Theorem 6.31. *Let S be an additively idempotent semiring. Then every k -maximal ideal is absolutely k -prime.*

Proof. Let I be a k -maximal ideal. Then I is a k -ideal. We need to show that I is absolutely prime. Let $(ab)\rho_I(ac)$. Then $(ab)K_I(ac)$. Since I is a k -maximal ideal, K_I is a k -maximal congruence. This implies K_I is a k -prime congruence, that is, a prime congruence. Hence, by [23, Theorem 2.12], K_I is cancellative, which further implies bK_Ic or aK_I0 . If bK_Ic , then $b + z_1 = c + z_2$, for some $z_1, z_2 \in I$ implies $b + z_1 + z_1 + z_2 = c + z_2 + z_1 + z_2$, which gives $b + z_1 + z_2 = c + z_1 + z_2$. Therefore, $b\rho_Ic$. If aK_I0 , then $a \in I = \bar{I}$. \square

ACKNOWLEDGEMENT

The first author expresses sincere thanks to the Council of Scientific and Industrial Research (CSIR), Government of India, for the financial support (File No. 09/0096(17468)/2024-EMR-I). The third author is grateful to the University Grants Commission (India) for providing a Senior Research Fellowship (ID: 211610013222/ Joint CSIR-UGC NET JUNE 2021).

REFERENCES

- [1] A. Azizi, *Strongly irreducible ideals*, J. Aust. Math. Soc., **84** (2008), 145–154.
- [2] R. E. Atani and S. E. Atani, *Ideal theory in commutative semirings*, Bul. Acad. Stiin te Repub. Mold. Mat., (2008), 14–23.
- [3] S. E. Atani and A. G. Garfami, *Ideals in quotient semirings*, Chiang Mai J. Sci. **40** (2013), no. 1, 77–82.
- [4] J. Y. Abuhlail, S.N. Il'in, Y. Katsov, T.G. Nam, *Toward homological characterization of semirings by e-injective semimodules*, J. Algebra Appl. **17** (2018), no. 4, 1850059, 24 pp.
- [5] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley, 1969.
- [6] J. Y. Abuhlail and R. G. Noegraha, *Injective semimodules—revisited*, J. Algebra Appl. **21** (2022), no. 1, Paper No. 2250242, 28 pp.
- [7] P. Biswas, S. Bag and S. K. Sardar, *Some new class of ideals in semirings and their applications*, Filomat **38** (2024), no. 29, 10223–10237.
- [8] ———, *On the positive cone of rings of measurable functions*, Filomat **39** (2025), no. 22, 7877–7891.
- [9] P. Biswas, A. Goswami and S. K. Sardar, *Some functor calculus on semirings*, preprint, arXiv:2506.13378, (2025).
- [10] V. V. Chermnykh, J. Math. Sci. (N.Y.) **187** (2012), no. 2, 187–267; translated from Fundam. Prikl. Mat. **17** (2011/12), no. 3, 111–227.
- [11] K. Głązak, *A guide to the literature on semirings and their applications in mathematics and information sciences, With complete bibliography*, Kluwer Academic Publishers, 2002.
- [12] J. S. Golan, *Semirings and their applications*, Springer, 1999.
- [13] A. Goswami, On semisubtractive ideals of semirings, Adv. Stud. Euro-Tbil. Math. J. **18** (2025), no. 3, 207–223.
- [14] ———, On z -ideals and z -closure operations of semirings, I, Quaest. Math. **48** (2025), no. 3, 395–415.
- [15] ——— and T. A. Dube, *Some aspects of k -ideals of semirings*, Rend. Circ. Mat. Palermo (2) **73** (2024), no. 8, 3105–3117.
- [16] A. Goswami and A. Maatallah, Further remarks on z -ideals of semirings, Quaest. Math. **48** (2025), no. 10, 1533–1536.
- [17] H. P. Gumm and A. Ursini, *Ideals in universal algebras*, Algebra Universalis **19** (1984), no. 1, 45–54.
- [18] S. C. Han, Maximal k -ideals and r -ideals in semirings, J. Algebra Appl. **14** (2015), no. 10, 1250195, 13 pp.
- [19] ———, k -congruences and the Zariski topology in semirings, Hacet. J. Math. Stat. **50** (2021), no. 3, 699–709.
- [20] M. Henriksen, *Ideals in semirings with commutative addition*, Amer. Math. Soc. Notices, **6**(3) 31 (1958), 321.
- [21] S. N. Il'in, E. B. Katsov and T. G. Nam, *Toward homological structure theory of semimodules: on semirings all of whose cyclic semimodules are projective*, J. Algebra **476** (2017), 238–266.
- [22] K. Iséki, *Ideal theory of semiring*, Proc. Japan. Acad., **32** (1956), 554–559.
- [23] D. Joó and K. Mincheva, Prime congruences of additively idempotent semirings and a Nullstellensatz for tropical polynomials, Selecta Math. (N.S.) **24** (2018), no. 3, 2207–2233.
- [24] J. Jun, S. Ray, and J. Toller, *Lattices, spectral spaces, and closure operations on idempotent semirings*, J. Algebra, **594** (2022), 313–363.
- [25] E. B. Katsov, T. G. Nam and N. X. Tuyen, *On subtractive semisimple semirings*, Algebra Colloq. **16** (2009), no. 3, 415–426.
- [26] ———, *More on subtractive semirings: simpleness, perfectness, and related problems*, Comm. Algebra **39** (2011), no. 11, 4342–4356.
- [27] S. LaGrassa, *Semirings: ideals and polynomials*, PhD thesis, University of Iowa (1995).
- [28] D. R. LaTorre, *On h -ideals and k -ideals in hemirings*, Publ. Math. Debrecen, **12** (1965), 219–226.
- [29] P. Lescot, *Absolute algebra II—ideals and spectra*, J. Pure Appl. Algebra **215** (2011), no. 7, 1782–1790.
- [30] ———, *Prime and primary ideals in semirings*, Osaka J. Math., **52**(3) (2015), 721–737.
- [31] T. G. Nam and J. Zumbrägel, *On Steinberg algebras of Hausdorff ample groupoids over commutative semirings*, J. Pure Appl. Algebra **225** (2021), no. 4, Paper No. 106548, 22 pp.
- [32] P. Nasehpour, *Some remarks on semirings and their ideals*, Asian-Eur. J. Math. **12** (2019), no. 7, 2050002, 14 pp.
- [33] M. K. Sen and M. R. Adhikari, *On k -ideals of semirings*, Internat. J. Math. & Math. Sci., **15**(2) (1992), 347–350.
- [34] ———, *On maximal k -ideals of semirings*, Proc. Amer. Math. Soc. **118** (1993), 699–703.
- [35] N. Schwartz, *Strongly irreducible ideals and truncated valuations*, Comm. Algebra, **44**(3) (2016), 1055–1087.

- [36] H. S. Vandiver, *Note on a simple type of algebra in which the cancellation law of addition does not hold*, Bull. Am. Math. Soc., **40**(12) (1934), 914–920.
- [37] H. S. Vandiver, *On some simple types of semi-rings*, Am. Math. Monthly, **46**(1) (1939), 22–26.
- [38] H. J. Weinert, M. K. Sen, and M. R. Adhikari, *One-sided k -ideals and h -ideals in semirings*, Math. Pannon., **7** (1996), 147–162.

DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, 188, RAJA SUBODH CHANDRA MALLICK RD, JADAVPUR, KOLKATA, WEST BENGAL 700032, INDIA.

Email address: `pubalis.math.rs@jadavpuruniversity.in`

[1] DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF JOHANNESBURG, P.O. BOX 524, AUCKLAND PARK 2006, SOUTH AFRICA. [2] NATIONAL INSTITUTE FOR THEORETICAL AND COMPUTATIONAL SCIENCES (NITHECS), SOUTH AFRICA.

Email address: `agoswami@uj.ac.za`

DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, 188, RAJA SUBODH CHANDRA MALLICK RD, JADAVPUR, KOLKATA, WEST BENGAL 700032, INDIA

Email address: `pronayb.math.rs@jadavpuruniversity.in, pronaybiswas1729@gmail.com`

DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, 188, RAJA SUBODH CHANDRA MALLICK RD, JADAVPUR, KOLKATA, WEST BENGAL 700032, INDIA.

Email address: `sujitk.sardar@jadavpuruniversity.in`