

REPRESENTATIONS OF CODE LOOPS BY BINARY CODES

ROSEMARY MIGUEL PIRES, ALEXANDRE GRISHKOV, AND MARINA RASSKAZOVA

ABSTRACT. Code loops are Moufang loops constructed from doubly even binary codes. Then, given a code loop L , we ask which doubly even binary code V produces L . In this sense, V is called a representation of L . In this article we define and show how to determine all minimal and reduced representations of nonassociative code loops of rank 3 and 4.

1. INTRODUCTION

Code loops were introduced by Robert Griess [2] from doubly even binary codes as follows.

Let $K = \mathbb{F}_2 = \{0, 1\}$ be a field with two elements and K^n be an n -dimensional vector space over K . Let $u = (u_1, \dots, u_n)$ and $v = (v_1, \dots, v_n)$ in K^n and define $|v| = |\{i \mid v_i = 1\}|$ (the Hamming weight) and $|u \cap v| = |\{i \mid u_i = v_i = 1\}|$. A *doubly even binary code* is a subspace $V \subseteq K^n$ such that $|v| \equiv 0 \pmod{4}$.

Let V be a doubly even binary code and the function $\phi : V \times V \rightarrow \{1, -1\}$, called *factor set*, defined by:

$$\begin{aligned}\phi(v, v) &= (-1)^{\frac{|v|}{4}}, \\ \phi(v, w) &= (-1)^{\frac{|v \cap w|}{2}} \phi(w, v), \\ \phi(0, v) &= \phi(v, 0) = 1, \\ \phi(v + w, u) &= \phi(v, w + u) \phi(v, w) \phi(w, u) (-1)^{|v \cap w \cap u|}.\end{aligned}$$

In order to define a code loop, let V be a doubly even binary code and $\phi : V \times V \rightarrow \{1, -1\}$ be a factor set. Consider the set $L(V) = \{1, -1\} \times V$ and define a product ‘.’ in $L(V)$ in the following way

$$\begin{aligned}v.w &= \phi(v, w)(v + w), \\ v.(-w) &= (-v).w = -(v.w).\end{aligned}$$

With this product, we obtain a Moufang loop called *code loop*. Equivalently, a Moufang loop L such that L/A is an elementary abelian 2-group for some central subgroup A of order 2 is called a code loop. We say that $L(V)$ has *rank* m if the dimension of the K -vector space V is equal to m . Note that $v \in L(V)$ and $-v \in L(V)$ mean the elements $(1, v)$ and $(-1, v)$, respectively.

It’s important to recall that code loops have some useful properties related to the commutator, associator, and square associated with their elements that we will use to get the main results presented in this article. Chein and Goodaire [1] proved that code

2010 *Mathematics Subject Classification.* Primary: 20N05.

Key words and phrases. code loops, characteristic vectors, representations.

loops have a unique nonidentity square, a unique nonidentity commutator, and a unique nonidentity associator. In other words, for any $u, v, w \in V$:

$$(1.1) \quad v^2 = (-1)^{\frac{|v|}{4}} 0,$$

$$(1.2) \quad [u, v] = u^{-1}v^{-1}uv = (-1)^{\frac{|u \cap v|}{2}} 0,$$

$$(1.3) \quad (u, v, w) = ((uv)w)((u(vw))^{-1}) = (-1)^{|u \cap v \cap w|} 0.$$

Let V be a doubly even binary code and $v \in V$. Note that, if $v^2 = -1$, then $|v| \equiv 4 \pmod{8}$. Otherwise, we have $|v| \equiv 0 \pmod{8}$.

Now let $u, v \in V$. As a consequence of the definition of doubly even binary code, we obtain $|u \cap v| \equiv 0 \pmod{2}$. Then in the case $[u, v] = -1$, we obtain $|u \cap v| \equiv 2 \pmod{4}$ and in the other case, $|u \cap v| \equiv 0 \pmod{4}$.

So far, we've shown how to construct code loops from doubly even binary codes, but we would like to investigate the other direction. In other words, given a certain code loop L , we would like to determine the doubly even codes V such that $L \simeq L(V)$.

A *representation* of a given code loop L is a doubly even binary code $V \subseteq K^m$ such that $L \simeq L(V)$ and the number m is called the *degree of the representation*. So the degree of the representation is the length of the code. We notice that there are many different representations for the same code loop. When necessary, we denote the degree of V by $\deg(V)$.

In this article we determine all minimal representations of nonassociative code loops of rank 3 and 4 (Sections 3 and 4). We will also show how to determine reduced representations. In order to do this, first, it is necessary to know all these code loops, up to isomorphism. In [3], these code loops were classified using the concept of characteristic vector associated with them. We recall the main theorems of classification of nonassociative code loops of rank 3 and 4 in Section 2.

2. NONASSOCIATIVE CODE LOOPS OF RANK 3 AND 4

Let L be a code loop with generator set $X = \{x_1, \dots, x_n\}$ and center $\{1, -1\}$. Then we define the characteristic vector of L , denoted by $\lambda_X(L)$ or $\lambda(L)$, by

$$\lambda(L) = (\lambda_1, \dots, \lambda_n; \lambda_{12}, \dots, \lambda_{1n}, \dots, \lambda_{(n-1)n}; \lambda_{123}, \dots, \lambda_{12n}, \dots, \lambda_{(n-2)(n-1)n}),$$

where $\lambda_i, \lambda_{ij}, \lambda_{ijk} \in \mathbb{F}_2$, $(-1)^{\lambda_i} = x_i^2$, $(-1)^{\lambda_{ij}} = [x_i, x_j]$ and $(-1)^{\lambda_{ijk}} = (x_i, x_j, x_k)$.

Here, $[x, y]$ denotes the commutator of x and y , and (x, y, z) denotes the associator of x, y and z . The center of a loop L , denoted by $\mathcal{Z}(L)$, is the set

$$\mathcal{Z}(L) = \{x \in L \mid [x, y] = (x, y, z) = 1, \forall y, z \in L\}.$$

Once a generating set X is fixed, we obtain a unique characteristic vector $\lambda_X(L)$, which depends on the choice of the elements in X and on the order of its elements. Different minimal generating sets may produce different characteristic vectors for the same code loop, as illustrated by the following example:

Let $L = \mathbb{Z}_2 \times \mathbb{Z}_4 = \{0, 1\} \times \{0, 1, 2, 3\}$ and let $Z = \{(0, 0), (0, 2)\}$. Then $Z \leq \mathcal{Z}(L)$ (since L is an abelian group), and L/Z is the Klein group, so L is a code loop of rank 2. We can consider various (ordered) minimal generating subsets of L . With $X = (a, b) = ((0, 1), (1, 0))$ we obtain $a + a = (0, 2)$ and $b + b = (0, 0)$, while with $X' = (a', b') = ((0, 1), (1, 1))$ we obtain $a' + a' = (0, 2)$ and $b' + b' = (0, 2)$. Note that b has order 2, while b' has order 4, so the characteristic vectors associated with each generating set are different.

The fact that the characteristic vector uniquely determines a code loop was established in [3]. By definition of a characteristic vector, the number of possible such vectors is 2^m , where $m = n + \frac{n(n-1)}{2} + \frac{n(n-1)(n-2)}{6}$. In [3], the authors constructed a free loop \mathcal{F}_n in the variety generated by code loops and showed how to establish a correspondence between a vector with coordinates in \mathbb{F}_2 and a nonassociative code loop. However, different vectors may determine the same nonassociative code loop up to isomorphism. Moreover, using the natural action of $GL_n(2)$ on the set of characteristic vectors, it was proved that the orbits of this action determine the isomorphism classes of nonassociative code loops. Therefore, there is a bijective correspondence between the set of isomorphism classes of nonassociative code loops of rank n and the set of $GL_n(2)$ -orbits of characteristic vectors of such loops (see [3, Corollary 2.11, p. 172] and [3, Proposition 2.12, p. 172]).

We can determine a representation V of L just by knowing the characteristic vector $\lambda(L)$ associated with L . Indeed, the method consists of finding a basis of V with vectors v_1, \dots, v_n corresponding to x_1, \dots, x_n , respectively. The squares, commutators and associators of the generators of L will determine the respective squares, commutators and associators of the basis elements of V and, from Equations (1.1), (1.2) and (1.3), we get the weights of v_i , $v_i \cap v_j$ and $v_i \cap v_j \cap v_k$, that is, $|v_i|$, $|v_i \cap v_j|$ and $|v_i \cap v_j \cap v_k|$, for each $i, j, k \in \{1, \dots, n\}$. To determine the possible vectors of a basis of V , it is enough to analyze these weights, as we will see in the next section. We usually use the following notation: $t_i = |v_i|$, $t_{ij} = |v_i \cap v_j|$ and $t_{ijk} = |v_i \cap v_j \cap v_k|$.

Now, we assume that L is a nonassociative code loop of rank 3 with generators a, b, c . Then the associator (a, b, c) is equal to -1 , otherwise, by Moufang Theorem, L would be a group. The characteristic vector associated to L is given by $\lambda(L) = (\lambda_1, \dots, \lambda_6)$ (or simply $\lambda(L) = (\lambda_1 \dots \lambda_6)$) where $a^2 = (-1)^{\lambda_1}$, $b^2 = (-1)^{\lambda_2}$, $c^2 = (-1)^{\lambda_3}$, $[a, b] = (-1)^{\lambda_4}$, $[a, c] = (-1)^{\lambda_5}$ and $[b, c] = (-1)^{\lambda_6}$. The omitted seventh coordinate is equal to one.

As discussed above, we know that two code loops with the same characteristic vector are isomorphic. This follows directly from [3, Corollary 2.11, p.172] and [3, Proposition 2.12, p.172]. To illustrate and justify this in a concrete case, consider the nonassociative code loops of rank 3. By definition of the characteristic vector, and since we are considering nonassociative loops of this rank, there are exactly 64 such vectors corresponding to these loops. According to [3, Proposition 3.2, p.175], these vectors are divided into 5 equivalence classes (orbits) under the natural action of $GL_3(2)$. Each orbit corresponds to a unique nonassociative code loop given by Theorem 2.1 below, where each listed characteristic vector is a representative of the corresponding orbit.

Theorem 2.1 ([3]). *Consider C_1^3, \dots, C_5^3 the code loops with the following characteristic vectors:*

$$\lambda(C_1^3) = (111111), \quad \lambda(C_2^3) = (000000), \quad \lambda(C_3^3) = (000111),$$

$$\lambda(C_4^3) = (110000), \quad \lambda(C_5^3) = (100000).$$

Then any two loops from the list $\{C_1^3, \dots, C_5^3\}$ are not isomorphic and every nonassociative code loop of rank 3 is isomorphic to one of this list.

From the above classification, we can now describe the congruence classes of t_i , t_{ij} , and t_{ijk} for each loop, which encode the intersection properties of the basis vectors in a representation. The values listed in the next remark correspond to the case of code loops of rank 3.

Remark 2.2. According to Theorem 2.1 and Equations (1.1), (1.2) and (1.3), any representation of the loops C_1^3, \dots, C_5^3 has a basis $\{v_1, v_2, v_3\}$ that satisfies the following properties, respectively:

1. $t_1 \equiv t_2 \equiv t_3 \equiv 4 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 2 \pmod{4}$;
2. $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 0 \pmod{4}$;
3. $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 2 \pmod{4}$;
4. $t_1 \equiv t_2 \equiv 4 \pmod{8}$, $t_3 \equiv 0 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 0 \pmod{4}$;
5. $t_1 \equiv 4 \pmod{8}$, $t_2 \equiv t_3 \equiv 0 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 0 \pmod{4}$.

For any representation, since $(v_1, v_2, v_3) = -1$, we also have that $t_{123} \equiv 1 \pmod{2}$. Therefore it doesn't happen $v_j \subset v_k$, $j \neq k$ or $v_j \cap v_k = \emptyset$. If $v_1 \subset v_2$, for example, we would have $t_{123} \equiv t_{13} \equiv 0 \pmod{2}$, which would be absurd because t_{123} is odd. If $v_2 \cap v_3 = \emptyset$, for example, we would have $t_{123} \equiv t_1 \equiv 0 \pmod{4}$, which also can not happen.

Let L be a nonassociative code loop of rank 4. By Lemma 3.4 of [3], we can assume that L has a generating set $X = \{a, b, c, d\}$ such that the nucleus of L is generated by d , that is, $N(L) = \mathbb{F}_2 d$. Since L is nonassociative, we have $(a, b, c) = -1$. By definition, the characteristic vector associated to L has 14 coordinates. However, since $(a, b, d) = (a, c, d) = (b, c, d) = 1$ and $(a, b, c) = -1$, we can omit the coordinates corresponding to the values of these associators and thus consider the characteristic vector of L of the form $\lambda_X(L) = (\lambda_1, \dots, \lambda_{10})$ (or simply $\lambda(L) = (\lambda_1 \dots \lambda_{10})$), where

$$\begin{aligned} a^2 &= (-1)^{\lambda_1}, & b^2 &= (-1)^{\lambda_2}, & c^2 &= (-1)^{\lambda_3}, & d^2 &= (-1)^{\lambda_4}, \\ [a, b] &= (-1)^{\lambda_5}, & [a, c] &= (-1)^{\lambda_6}, & [a, d] &= (-1)^{\lambda_7}, \\ [b, c] &= (-1)^{\lambda_8}, & [b, d] &= (-1)^{\lambda_9}, & [c, d] &= (-1)^{\lambda_{10}}. \end{aligned}$$

Similarly to the rank 3 case, the set of characteristic vectors of nonassociative code loops of rank 4 can be partitioned into orbits under the natural action of $GL_4(2)$, each corresponding to a unique loop up to isomorphism [4]. Theorem 2.3 below lists one representative from each orbit and thus provides the classification of code loops of rank 4.

Theorem 2.3 ([3]). *Consider C_1^4, \dots, C_{16}^4 the code loops with the following characteristic vectors. Every nonassociative code loop of rank 4 is isomorphic to one from the list. Moreover, no two of those loops are isomorphic to each other.*

L	$\lambda(L)$	L	$\lambda(L)$
C_1^4	(1110110100)	C_9^4	(0100001000)
C_2^4	(0000000000)	C_{10}^4	(0001111000)
C_3^4	(0000110100)	C_{11}^4	(0001001000)
C_4^4	(0010100000)	C_{12}^4	(0000001100)
C_5^4	(0000010100)	C_{13}^4	(0110111100)
C_6^4	(1111110100)	C_{14}^4	(0001001100)
C_7^4	(0001000000)	C_{15}^4	(1001001100)
C_8^4	(0000001000)	C_{16}^4	(0001111100)

The next remark lists the congruence classes of t_i , t_{ij} , and t_{ijk} that will be used to calculate representations of code loops of rank 4.

Remark 2.4. As a consequence of the Theorem 2.3 we will obtain, for each code loop, representations with basis $\{v_1, v_2, v_3, v_4\}$ that satisfies the following properties:

- If $\lambda_i = 0$, then $t_i \equiv 0 \pmod{8}$. Otherwise, we have $t_i \equiv 4 \pmod{8}$.
- If $\lambda_{ij} = 0$, then $t_{ij} \equiv 0 \pmod{4}$. Otherwise, we have $t_{ij} \equiv 2 \pmod{4}$.
- $t_{123} \equiv 1 \pmod{2}$ and $t_{1j4} \equiv 0 \pmod{2}$.

Remark 2.5. From the characteristic vector we do not obtain the absolute values of the weights t_i and t_{ij} , but only their congruence classes: $t_i \pmod{8}$ from (1.1) and $t_{ij} \pmod{4}$ from (1.2). These congruences provide information about the vectors in a representation V and the properties of their intersections. The possible congruence classes for t_i , t_{ij} and t_{ijk} for the code loops of rank 3 and 4 are listed in Remarks 2.2 and 2.4. The exact values of these weights are analyzed in the next sections, where we consider all possibilities (up to a fixed bound) compatible with the above congruences, in order to determine the minimal representations.

3. REPRESENTATIONS OF CODE LOOPS

Let L be a code loop of rank n with generators $B = \{x_1, \dots, x_n\}$ and $V = \text{Span}_{\mathbb{F}_2} \{v_1, \dots, v_n\}$ a representation of L with degree m . As in [3] we identify the \mathbb{F}_2 -space \mathbb{F}_2^m with $P_m = \{\sigma \subseteq I_m\}$, then $v_i \subseteq I_m$. Let suppose that $\bigcup_{i=1}^n v_i = I_m$. Then we have an equivalence on I_m : $i \sim j \iff \{i \in v_p \iff j \in v_p\}$. Consider the corresponding partition of I_m in equivalence classes: $I_m = \bigcup_{k=1}^t I_m^k$.

The same partition we can obtain in the following way. Let $\sigma \subseteq I_n$ and

$$(3.1) \quad v^\sigma = \left(\bigcap_{i \in \sigma} v_i \right) \setminus \left(\bigcup_{j \notin \sigma} v_j \right).$$

Lemma 1. In notation above, we have:

- i) $v^\sigma \cap v^\tau = \emptyset$, $\sigma \neq \tau$;
- ii) $\bigcup_{\sigma \subseteq I_n} v^\sigma = I_m$;
- iii) $\{v^\sigma \neq \emptyset \mid \sigma \subseteq I_n\} = \{I_m^k \mid k = 1, \dots, t\}$.

Proof.

- i) Let $i \in v^\sigma \cap v^\tau$, and $j \in \sigma \setminus \tau$. Hence, by (3.1), $i \in v_k$, for all $k \in \sigma$. Since $j \neq \tau$, then $i \notin v_j$ by (3.1). As $j \in \sigma$ we get a contradiction.
- ii) Let $i \in I_m$, and define $\sigma = \{k \in I_n \mid i \in v_k\}$. Then for $q \notin \sigma$ we have $i \notin v_q$, and $i \in v^\sigma$.
- iii) Let $i \sim j$ and $i \in v^\sigma$, then $i \in v_k$, $k \in \sigma$, hence $j \in v_k$. Since $i \in v^\sigma$, $i \notin v_p$, $p \notin \sigma$. As $i \sim j$, $j \notin v_p$. Hence $j \in v^\sigma$. It means that $I_m^k \subseteq v^\sigma$, if $i, j \in I_m^k$. Let $k \notin I_m^k$, $k \in v^\sigma$, then there exists v_q such that $i \notin v_q$, and $j \notin v_q$, $k \in v_q$. By $k \in v^\sigma$, $k \in v_q$ and (3.1) we get that $q \in \sigma$. On the other hand, since $i \notin v_q$ and $i \in v^\sigma$ hence $q \notin \sigma$. Contradiction.

□

Lemma 2. The partition $I_m = \bigcup_{k=1}^t I_m^k$ does not depend on the choice of the system of generators v_1, \dots, v_n .

Proof. We have $L = V \cup -V$, where $V = \text{Span}_{\mathbb{F}_2} \{v_1, \dots, v_n\}$. Recall that in V the operation is $v + w = (v \setminus w) \cup (w \setminus v)$. Let $v \in V$ and $i \in v \cap I_m^k$. We need to prove that $I_m^k \subseteq v$. Suppose that $j \in I_m^k \setminus v$. Let $v = \sum_{k \in \sigma \subseteq I_k} v_k$. If $|\sigma| = 1$, then $v = v_k$, hence $i \in v$, $j \notin v$ and $i \not\sim j$. Contradiction with $i, j \in I_m^k$.

Let $|\sigma| = s$ and for all $|\tau| < s$ we have $i \in v_\tau = \sum_{k \in \tau} v_k$ if and only if $j \in v_\tau$.

We have $v_\sigma = v_\tau + v_l$, $\tau = \sigma \setminus l$, $|\tau| < s$. Let $i \in v_\sigma$ and $i \in v_\tau$, then $i \notin v_l$, since $v_\sigma = (v_\tau \setminus v_l) \cup (v_l \setminus v_\tau)$. By $i \sim j$ we get $j \notin v_l$ and $j \in v_\tau$, hence $j \in v_\sigma$. Let $i \in v_\sigma$, $i \notin v_\tau$, then $i \in v_l$. By induction $j \in v_l$ and $j \notin v_\tau$. Hence $j \in v_\sigma$. \square

We will consider only representations such that, for any equivalence class X , we have $|X| < 8$. The choice is arbitrary and it only serves the purpose of making the enumeration of all reduced representations tractable. We call these representations by **reduced representations**.

Definition 3.1. A reduced representation V is **minimal** if whenever $L(V)$ is isomorphic to $L(W)$ then $\deg(V) \leq \deg(W)$, where W is another representation.

By (3.1), we will get all the equivalence classes on I_m and we will show that it's possible to find all reduced and minimal representations of degree m for code loops of rank 3 and 4 using these classes. To do this, it is necessary to find the cardinality of these equivalence classes and show how these values can be obtained just by knowing the weights, t_i, t_{ij} and t_{ijk} , with $i, j, k \in \{1, 2, 3, 4\}$. For each choice of these weights, there is a different representation. This choice must be in accordance with the properties mentioned in the Remark 2.2 and 2.4.

Definition 3.2. Let V be a m -degree representation of a code loop L and X_1, \dots, X_r the equivalence classes on I_m . We define the **type** of V as the vector $(|X_1|, \dots, |X_r|)$ such that $|X_1| \leq |X_2| \leq \dots \leq |X_r|$. When necessary, we can write $(|X_1| \dots |X_r|)$.

Definition 3.3. Let V_1 and V_2 be doubly even binary codes of \mathbb{F}_2^m . We say that V_1 and V_2 are equivalent codes if and only if there is a bijection $\varphi \in S_m$ such that $V_1^\varphi = V_2$.


3.1. Code Loops of rank 3. Let C_i^3 , $i = 1, \dots, 5$, be the nonassociative code loops of rank 3 and $\lambda(C_i^3)$ the associated characteristic vector as shown in Section 2. We also denote the corresponding representations by V_i^3 and assume that $V_i^3 = \text{Span}_{\mathbb{F}_2} \{v_1, v_2, v_3\}$.

Let m be the degree of V_i^3 and $I_m = \{1, 2, \dots, m\}$, such that $v_1, v_2, v_3 \subset I_m$ and $v_1 \cup v_2 \cup v_3 = I_m$. According to Lemma 1, the sets below determine all equivalence classes on I_m (see Fig. 1):

$$\begin{aligned} X_{123} &= v_1 \cap v_2 \cap v_3, \quad X_{12} = (v_1 \cap v_2) \setminus v_3, \quad X_{13} = (v_1 \cap v_3) \setminus v_2, \quad X_{23} = (v_2 \cap v_3) \setminus v_1, \\ X_1 &= v_1 \setminus (v_2 \cup v_3), \quad X_2 = v_2 \setminus (v_1 \cup v_3), \quad X_3 = v_3 \setminus (v_1 \cup v_2). \end{aligned}$$

Now, note that the generators of V_i^3 depend on the equivalence classes since we can write them as follows:

$$v_1 = X_{123} \cup X_{12} \cup X_{13} \cup X_1, \quad v_2 = X_{123} \cup X_{12} \cup X_{23} \cup X_2, \quad v_3 = X_{123} \cup X_{13} \cup X_{23} \cup X_3.$$

FIGURE 1. Equivalence classes on I_m for code loops of rank 3

So if we find the equivalence classes on I_m , we will determine the generators of the representation. As the elements of the representation depend on the weights of v_i , $v_i \cap v_j$ ($i, j \in \{1, 2, 3\}$) and $v_1 \cap v_2 \cap v_3$, we can establish a relationship between the cardinalities of the classes and these weights.

Denote the cardinalities of the sets X_{123} , X_{ij} and X_i by x_{123} , x_{ij} and x_i , respectively; $i, j = 1, 2, 3$, and assume that $x_{123} = t_{123}$. These cardinalities and the weights are related through the following system in the variables x_{12} , x_{13} , x_{23} , x_1 , x_2 and x_3 :

$$(3.2) \quad \begin{cases} t_{123} + x_{12} = t_{12} \\ t_{123} + x_{13} = t_{13} \\ t_{123} + x_{23} = t_{23} \\ t_{123} + x_{12} + x_{13} + x_1 = t_1 \\ t_{123} + x_{12} + x_{23} + x_2 = t_2 \\ t_{123} + x_{13} + x_{23} + x_3 = t_3 \end{cases}$$

Then the solutions are given by:

$$(3.3) \quad \begin{cases} x_{12} = t_{12} - t_{123} \\ x_{13} = t_{13} - t_{123} \\ x_{23} = t_{23} - t_{123} \\ x_1 = t_1 - t_{12} - t_{13} + t_{123} \\ x_2 = t_2 - t_{12} - t_{23} + t_{123} \\ x_3 = t_3 - t_{13} - t_{23} + t_{123} \end{cases}$$

Therefore, to determine a representation of a code loop of rank 3, first choose values for t_{123} , t_{ij} and t_i , $i, j = 1, 2, 3$; after look for the solution for the Linear System (3.2). Since we want equivalence classes whose cardinality is strictly less than 8, we consider solutions $v = (x_{12}, x_{13}, x_{23}, x_1, x_2, x_3)$ such that $0 \leq x_i \leq 7$ and $0 \leq x_{ij} \leq 7$, with $i, j = 1, 2, 3$,

Remark 3.4. It follows from 7 equivalence classes of at most 7 elements that:

- $t_{123} = 1, 3, 5$ or 7 ;

- $t_{ij} = 2, 4, 6, 8, 10, 12$ or 14 ;
- $t_i = 4, 8, 12, 16, 20, 24$ or 28 ;
- $7 \leq m \leq 49$.

Note that the degree m and the type of V_i^3 are obtained from these soluções. In fact, $m = x_{123} + x_{12} + x_{13} + x_{23} + x_1 + x_2 + x_3$ and the type is a vector determined by putting the coordinates of $(x_{123}, x_{12}, x_{13}, x_{23}, x_1, x_2, x_3)$ in ascending order.

After finding the system's solutions, that is, the cardinalities of the equivalence classes over I_m , it will be necessary to label its elements. By choice, we label the elements as follows:

$$\begin{aligned} X_{123} &= \{1, \dots, t_{123}\}, \\ X_{12} &= \{t_{123} + 1, \dots, t_{123} + x_{12}\}, \\ X_{13} &= \{t_{123} + x_{12} + 1, \dots, t_{123} + x_{12} + x_{13}\}, \\ X_1 &= \{t_{123} + x_{12} + x_{13} + 1, \dots, t_{123} + x_{12} + x_{13} + x_1\}, \\ X_{23} &= \{t_{123} + x_{12} + x_{13} + x_1 + 1, \dots, t_{123} + x_{12} + x_{13} + x_1 + x_{23}\}, \\ X_2 &= \{t_{123} + x_{12} + x_{13} + x_{23} + x_1 + 1, \dots, t_{123} + x_{12} + x_{13} + x_{23} + x_1 + x_2\}, \\ X_3 &= \{t_{123} + x_{12} + x_{13} + x_{23} + x_1 + x_2 + 1, \dots, t_{123} + x_{12} + x_{13} + x_{23} + x_1 + x_2 + x_3\}. \end{aligned}$$

3.2. Code Loops of Rank 4. Let V a m -degree reduced representation with basis $\{v_1, v_2, v_3, v_4\}$ corresponding to a code loop of rank 4. Here we also use the notation t or t_{1234} to denote the weight of the 4 generators, that is, $t = t_{1234} = |v_1 \cap v_2 \cap v_3 \cap v_4|$.

As we know, the weights of the generators and their intersections are determined through the characteristic vector associated with the code loop. So, for our first step, take a characteristic vector associated with a code loop of rank 4. Since $t_{123} \equiv 1 \pmod{2}$, $t_{ij4} \equiv 0 \pmod{2}$, $t_{ij} \equiv 0 \pmod{2}$ and $t_i \equiv 0 \pmod{4}$ and V is reduced, then we will get $1 \leq t_{123} \leq 13$, $0 \leq t_{ij4} \leq 14$, $2 \leq t_{ij} \leq 28$ ($i, j = 1, 2, 3$; $i \neq j$), $0 \leq t_{i4} \leq 28$ and $4 \leq t_i \leq 56$.

Remark 3.5. Consider $I_m = \{1, 2, \dots, m\}$, such that $v_1, v_2, v_3, v_4 \subset I_m$ and $v_1 \cup v_2 \cup v_3 \cup v_4 = I_m$. According to Lemma 1, the equivalence classes on $I_m = \{1, 2, \dots, m\}$ are obtained by calculating the following sets (see Fig. 2):

$$\begin{aligned} X_{1234} &= v_1 \cap v_2 \cap v_3 \cap v_4, & X_{123} &= (v_1 \cap v_2 \cap v_3) \setminus v_4, & X_{124} &= (v_1 \cap v_2 \cap v_4) \setminus v_3, \\ X_{134} &= (v_1 \cap v_3 \cap v_4) \setminus v_2, & X_{234} &= (v_2 \cap v_3 \cap v_4) \setminus v_1, & X_{12} &= (v_1 \cap v_2) \setminus (v_3 \cup v_4), \\ X_{13} &= (v_1 \cap v_3) \setminus (v_2 \cup v_4), & X_{14} &= (v_1 \cap v_4) \setminus (v_2 \cup v_3), & X_{23} &= (v_2 \cap v_3) \setminus (v_1 \cup v_4), \\ X_{24} &= (v_2 \cap v_4) \setminus (v_1 \cup v_3), & X_{34} &= (v_3 \cap v_4) \setminus (v_1 \cup v_2), & X_1 &= v_1 \setminus (v_2 \cup v_3 \cup v_4), \\ X_2 &= v_2 \setminus (v_1 \cup v_3 \cup v_4), & X_3 &= v_3 \setminus (v_1 \cup v_2 \cup v_4), & X_4 &= v_4 \setminus (v_1 \cup v_2 \cup v_3). \end{aligned}$$

As done in previous section, denote by x_{1234} , x_{ijk} , x_{ij} or x_i ; $i, j, k = 1, 2, 3, 4$; the cardinality of this sets. Also assume $t_{1234} = x_{1234}$. The second step to finding reduced representations is to choose values for t_{1234} , t_{ijk} , t_{ij} and t_i , with $i, j, k = 1, 2, 3, 4$, that satisfy the properties presented in the Remark 2.4.

In this case, we can also relate these weights and the cardinalities of the sets listed in the Remark 3.5 through the following linear system of 14 equations in the variables x_{123} , x_{124} , x_{134} , x_{234} , x_{12} , x_{13} , x_{14} , x_{23} , x_{24} , x_{34} , x_1 , x_2 , x_3 and x_4 :

	X_2	X_{24}	X_4
v_1	X_1	X_{12}	X_{124}
$v_1 \cap v_3$	X_{13}	X_{123}	X_{1234}
v_3	X_3	X_{23}	X_{234}
			X_{34}

 FIGURE 2. Equivalence classes on I_m for code loops of rank 4

$$(3.4) \quad \left\{ \begin{array}{l} t + x_{123} = t_{123} \\ t + x_{124} = t_{124} \\ t + x_{134} = t_{134} \\ t + x_{234} = t_{234} \\ t + x_{123} + x_{124} + x_{12} = t_{12} \\ t + x_{123} + x_{134} + x_{13} = t_{13} \\ t + x_{124} + x_{134} + x_{14} = t_{14} \\ t + x_{123} + x_{234} + x_{23} = t_{23} \\ t + x_{124} + x_{234} + x_{24} = t_{24} \\ t + x_{134} + x_{234} + x_{34} = t_{34} \\ t + x_{123} + x_{12} + x_{124} + x_{134} + x_{13} + x_{14} + x_1 = t_1 \\ t + x_{123} + x_{12} + x_{124} + x_{234} + x_{23} + x_{24} + x_2 = t_2 \\ t + x_{123} + x_{13} + x_{134} + x_{234} + x_{23} + x_{34} + x_3 = t_3 \\ t + x_{124} + x_{134} + x_{234} + x_{14} + x_{24} + x_{34} + x_4 = t_4 \end{array} \right.$$

The third step is to solve this system. Solving it, we obtain:

$$\left\{ \begin{array}{l} x_{123} = t_{123} - t \\ x_{124} = t_{124} - t \\ x_{134} = t_{134} - t \\ x_{234} = t_{234} - t \\ x_{12} = t_{12} - t_{123} - t_{124} + t \\ x_{13} = t_{13} - t_{123} - t_{134} + t \\ x_{14} = t_{14} - t_{124} - t_{134} + t \\ x_{23} = t_{23} - t_{123} - t_{234} + t \\ x_{24} = t_{24} - t_{124} - t_{234} + t \\ x_{34} = t_{34} - t_{134} - t_{234} + t \\ x_1 = t_1 - t_{12} - t_{13} + t_{123} - t_{14} + t_{124} + t_{134} - t \\ x_2 = t_2 - t_{12} - t_{23} + t_{123} - t_{24} + t_{234} + t_{124} - t \\ x_3 = t_3 - t_{13} - t_{23} + t_{123} - t_{34} + t_{234} + t_{134} - t \\ x_4 = t_4 - t_{14} - t_{24} + t_{124} - t_{34} + t_{134} + t_{234} - t \end{array} \right.$$

Remember that the reduced representations will be get for all those solutions that satisfies the following conditions, for all $i, j, k \in \{1, 2, 3, 4\}$:

$$0 \leq x_i \leq 7, \quad 0 \leq x_{ij} \leq 7, \quad 0 \leq x_{ijk} \leq 7, \quad 0 \leq x_{1234} \leq 7.$$

With those solutions, the sets $X_{1234}, X_{123}, X_{124}, X_{134}, X_{234}, X_{12}, X_{13}, X_{14}, X_{23}, X_{24}, X_{34}, X_1, X_2, X_3$ and X_4 are determined following the labeling chosen in the Table 1. The order chosen to determine these sets is arbitrary. Note that only non-empty sets determine the equivalence classes on I_m , where m is the degree of V . At last, find the generators v_1, v_2, v_3 and v_4 of V writing:

$$\begin{aligned} v_1 &= X_{1234} \cup X_{123} \cup X_{124} \cup X_{134} \cup X_{12} \cup X_{13} \cup X_{14} \cup X_1, \\ v_2 &= X_{1234} \cup X_{123} \cup X_{124} \cup X_{234} \cup X_{12} \cup X_{23} \cup X_{24} \cup X_2, \\ v_3 &= X_{1234} \cup X_{123} \cup X_{134} \cup X_{234} \cup X_{13} \cup X_{23} \cup X_{34} \cup X_3, \\ v_4 &= X_{1234} \cup X_{124} \cup X_{134} \cup X_{234} \cup X_{14} \cup X_{24} \cup X_{34} \cup X_4. \end{aligned}$$

As an illustration, we will present a reduced representation, denoted by V_{16}^4 , for the code loop with characteristic vector (0001111100) . We will use the following notation:

$$\begin{aligned} u &= (t, t_{123}, t_{124}, t_{134}, t_{234}, t_{12}, t_{13}, t_{14}, t_{23}, t_{24}, t_{34}, t_1, t_2, t_3) \\ v &= (x_{123}, x_{124}, x_{134}, x_{234}, x_{12}, x_{13}, x_{14}, x_{23}, x_{24}, x_{34}, x_1, x_2, x_3) \end{aligned}$$

If $u = [0, 1, 0, 2, 0, 2, 6, 2, 6, 0, 4, 8, 8, 16, 4]$, then $v = (1, 0, 2, 0, 1, 3, 0, 5, 0, 2, 1, 1, 3, 0)$ is the solution of the System (3.4). Thus the generators of V_{16}^4 are given by:

$$\begin{aligned} v_1 &= \{1, 2, 3, 4, 5, 6, 7, 8\}, \\ v_2 &= \{1, 4, 9, 10, 11, 12, 13, 14\}, \\ v_3 &= \{1, 2, 3, 5, 6, 7, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19\}, \\ v_4 &= \{2, 3, 15, 16\}. \end{aligned}$$

TABLE 1. Labeling elements of equivalence classes on I_m .

X_{1234}	$\{\} \text{ if } x_{1234} = 0$ $\{1, \dots, x_{1234}\} \text{ if } x_{1234} \neq 0$	
X_{123}	$\{\} \text{ if } x_{123} = 0$ $\{x_{1234} + 1, \dots, n_{123}\}$	$n_{123} = x_{1234} + x_{123}$
X_{124}	$\{\} \text{ if } x_{124} = 0$ $\{n_{123} + 1, \dots, n_{124}\} \text{ if } x_{124} \neq 0$	$n_{124} = n_{123} + x_{124}$
X_{12}	$\{\} \text{ if } x_{12} = 0$ $\{n_{134} + 1, \dots, n_{12}\}$	$n_{12} = n_{124} + x_{12}$
X_{134}	$\{\} \text{ if } x_{134} = 0$ $\{n_{12} + 1, \dots, n_{134}\} \text{ if } x_{134} \neq 0$	$n_{134} = n_{12} + x_{134}$
X_{13}	$\{\} \text{ if } x_{13} = 0$ $\{n_{134} + 1, \dots, n_{13}\}$	$n_{13} = n_{134} + x_{13}$
X_{14}	$\{\} \text{ if } x_{14} = 0$ $\{n_{13} + 1, \dots, n_{14}\} \text{ if } x_{14} \neq 0$	$n_{14} = n_{13} + x_{14}$
X_1	$\{\} \text{ if } x_1 = 0$ $\{n_{14} + 1, \dots, n_1\}$	$n_1 = n_{14} + x_1$
X_{234}	$\{\} \text{ if } x_{234} = 0$ $\{n_1 + 1, \dots, n_{234}\} \text{ if } x_{234} \neq 0$	$n_{234} = n_1 + x_{234}$
X_{23}	$\{\} \text{ if } x_{23} = 0$ $\{n_{234} + 1, \dots, n_{23}\}$	$n_{23} = n_{234} + x_{23}$
X_{24}	$\{\} \text{ if } x_{24} = 0$ $\{n_{23} + 1, \dots, n_{24}\} \text{ if } x_{24} \neq 0$	$n_{24} = n_{23} + x_{24}$
X_2	$\{\} \text{ if } x_2 = 0$ $\{n_{24} + 1, \dots, n_2\}$	$n_2 = n_{24} + x_2$
X_{34}	$\{\} \text{ if } x_{34} = 0$ $\{n_2 + 1, \dots, n_{34}\} \text{ if } x_{34} \neq 0$	$n_{34} = n_2 + x_{34}$
X_3	$\{\} \text{ if } x_3 = 0$ $\{n_{34} + 1, \dots, n_3\}$	$n_3 = n_{34} + x_3$
X_4	$\{\} \text{ if } x_4 = 0$ $\{n_3 + 1, \dots, n_4\} \text{ if } x_4 \neq 0$	$n_4 = n_3 + x_4$

4. MINIMAL REPRESENTATIONS

Below, we present in the theorems 4.1 and 4.3 the representations of nonassociative code loops of rank 3 and 4 where the degree of each representation is minimal and we determine the degrees of these representations. Here we use the notation $\langle v_1, v_2, \dots, v_n \rangle$ to denote the vector subspace of the space \mathbb{F}_2^m generated by v_1, v_2, \dots, v_n . We also use the notations introduced in Section 3.2.

We define the length of the vector $v_i + v_j$ to be the cardinality of the set $v_i \cup v_j$, that is, the length of $v_i + v_j$ is the number $l_{ij} = t_i + t_j - t_{ij}$. Analogously we define the length of the vector $v_i + v_j + v_k$ to be the cardinality of the set $v_i \cup v_j \cup v_k$, that is, the length of $v_i + v_j + v_k$ is the number $l_{ijk} = t_i + t_j + t_k - (t_{ij} + t_{ik} + t_{jk}) + t_{ijk}$.

Theorem 4.1. *The code loops C_1^3, \dots, C_5^3 have the following minimal representations V_1, \dots, V_5 , which are given by*

$$\begin{aligned} V_1 &= \langle (1, 2, 3, 4), (1, 2, 5, 6), (1, 3, 5, 7) \rangle, \\ V_2 &= \langle (1, 2, 3, 4, 5, 6, 7, 8), (1, 2, 3, 4, 9, 10, 11, 12), (1, 5, 6, 7, 9, 10, 11, 13) \rangle, \\ V_3 &= \langle (1, 2, 3, 4, 5, 6, 7, 8), (1, 2, 3, 4, 5, 6, 9, 10), (1, 2, 3, 4, 5, 7, 9, 11) \rangle, \\ V_4 &= \langle (1, 2, 3, 4), (1, 2, 5 - 14), (1, 3, 5, 6, 7, 8, 9, 10, 11, 15, 16, 17) \rangle, \\ V_5 &= \langle (1 - 12), (1 - 8, 13, 14, 15, 16), (1, 2, 3, 4, 5, 9, 10, 11, 13, 14, 15, 17) \rangle. \end{aligned}$$

Proof. We consider the \mathbb{F}_2 -subspaces V_1, \dots, V_5 of $\mathbb{F}_2^7, \mathbb{F}_2^{13}, \mathbb{F}_2^{11}, \mathbb{F}_2^{17}, \mathbb{F}_2^{17}$, respectively, as above. First we will see that each space V_i is a representation of C_i^3 , that is, a doubly even code of \mathbb{F}_2^m , for some m and that $C_i^3 \simeq L(V_i)$, $i = 1, \dots, 5$.

Since the elements of V_5 are $v_0 = 0$, $v_1 = (1 - 12)$, $v_2 = (1 - 8, 13 - 16)$, $v_3 = (1 - 5, 9 - 11, 13 - 15, 17)$, $v_4 = v_1 + v_2 = (9 - 16)$, $v_5 = v_1 + v_3 = (6, 7, 8, 12, 13, 14, 15, 17)$, $v_6 = v_2 + v_3 = (6 - 11, 16, 17)$ and $v_7 = v_1 + v_2 + v_3 = (1 - 5, 12, 16, 17)$, we see, clearly, that all the vectors have weight with multiplicity 4 and the weight of the intersection of each pair of vectors is even. Thus, V_5 is a doubly even code. Analogously, we prove that V_i , $i = 1, \dots, 4$ is a doubly even code.

Now, the isomorphism $C_i^3 \simeq L(V_i)$ follows directly from Theorem 2.1 of Classification of Code Loops of rank 3. First, we just need to calculate the characteristic vector associated to $L(V_i)$ and note that it corresponds to the same code loop C_i^3 (see [[3], Proposition 3.2]).

As example, we calculate the characteristic vector associate to $L(V_5) = \{1, -1\} \times V_5$. We have $v_i^2 = (-1)^{\frac{|v_i|}{4}} = -1$ and $[v_i, v_j] = (-1)^{\frac{|v_i \cap v_j|}{2}} = 1$, for $i, j = 1, 2, 3$. Hence $\lambda(L(V_5)) = (111000)$. Since this vectors produces the code loop C_5^3 , by Theorem 2.1 we conclude that $C_5^3 \simeq L(V_5)$.

We are going to demonstrate now that each V_i , $i = 1, \dots, 5$, up to isomorphism, is the unique minimal representation of the code loop C_i^3 . We consider $X = \{a, b, c\}$ a set of generators for C_i^3 such that $\lambda = \lambda_X(C_i^3)$ is the corresponding characteristic vector, and we assume that V is a minimal representation of C_i^3 , where v, w, u are the elements of the basis of V which corresponds to a, b, c . We use the notation $t = |v \cap w \cap u|$. Remember that $t \equiv 1 \pmod{2}$, because $(a, b, c) = -1$.

Case $i = 1$: In this case suppose that $\deg(V) \leq 7$. The characteristic vector is $\lambda = (111111)$, then $|v| \equiv |w| \equiv |u| \equiv 4 \pmod{8}$. We can suppose that $v = (1, 2, 3, 4)$, but $[a, b] = -1$, then $|v \cap w| \equiv 2 \pmod{4}$, and hence, $|v \cap w| = 2$. Analogously, we obtain $|v \cap u| = 2$. Let $w = (1, 2, 5, 6)$, hence we also obtain $|w \cap u| = 2$ and then, $t = 1$. Hence, $u = (1, 3, 5, 7)$. Therefore, $V = V_1$.

Case $i = 2$: The characteristic vector is $\lambda = (000000)$, then $|v| \equiv |w| \equiv |u| \equiv 0 \pmod{8}$. Since $\deg(V) \leq 13$, then $|v| = 8$. Analogously, $|w| = |u| = 8$. Let $v = (1-8)$. As $|v \cap w| \equiv 0 \pmod{4}$, hence $|v \cap w| = 4$. Analogously, $|v \cap u| = |w \cap u| = 4$.

Let $w = (1-4, 9-12)$. We have two possibilities for t : $t = 1$ or $t = 3$. Case $t = 3$, we will have $\deg(V) = 15$, a contradiction. Hence, $t = 1$ and $u = (1, 5-7, 9-11, 13)$. Therefore, $V = V_2$.

Case $i = 3$: We consider $\lambda = (000111)$ and $\deg(V) \leq 11$. Analogously to the previous case, we have $|v| = |w| = |u| = 8$. In this case, we have $|v \cap w| = |v \cap u| = |w \cap u| = 6$. We suppose $v = (1-8)$ and $w = (1-6, 9, 10)$. We have three possibilities for t : 1, 3 or 5. Case $t \leq 3$ we will have $|v| > 8$, a contradiction. Hence, $t = 5$ and we can assume $u = (1-5, 7, 9, 11)$. Therefore, $V = V_3$.

Case $i = 4$: In this case, consider the characteristic vector $\lambda = (111110)$. Since $|v| \equiv |w| \equiv |u| \equiv 4 \pmod{8}$ and $|w \cap u| \equiv 0 \pmod{4}$ then $|w| = |u| = 12$ and $|w \cap u| = 8$. Let $v = (1, 2, 3, 4)$, so $|v \cap w| = |v \cap u| = 2$ and $t = 1$. Then we can assume $w = (1, 2, 5-14)$ and $u = (1, 3, 5-11, 15-17)$. Therefore, $V = V_4$.

Now, we suppose $v = (1-12)$. If $|v \cap w| \leq 6$ then $\deg(V) \geq 18$. In fact, we have $|v \cap w| = 2, 6$ or 10 . If $|v \cap w| = 2$ we will have $|v + w| = 20$, a contradiction. If $|v \cap w| = 6$ we will have $|v + w| = 12$ and hence, $\deg(V) \geq 18$, a contradiction. Therefore, $|v \cap w| = 10$. Analogously we obtain $|v \cap u| = 10$. Without loss of generality, we suppose $w = (1, 2, 5-14)$. Since $t \equiv 1 \pmod{2}$, then the possibilities for t are 1, 3 or 7. In any case we will have $|v| \geq 13$, which is a contradiction. Then, there is not this last possibility for v .

Case $i = 5$: Consider the characteristic vector given by $\lambda = (111000)$, so we have $|v| \equiv |w| \equiv |u| \equiv 4 \pmod{8}$. Assuming $v = (1, 2, 3, 4)$ and since $[a, b] = 1$, we will have $|v \cap w| \equiv 0 \pmod{4}$. Case $v \cap w \neq \emptyset$ we will have $|v \cap w| = 4$, and hence $v \subset w$, which is a contradiction, because this give us $t \equiv 0 \pmod{2}$, which does not occur. The case $v \cap w = \emptyset$ also does not occur, since we must have nonempty intersection between v, w and u .

Therefore $|v| \geq 12$. Analogously, we prove that $|w| \geq 12$ and $|u| \geq 12$.

The representation V is minimal and $\deg(V) \leq 17$, so we obtain $|v| = |w| = |u| = 12$. Without loss of generality, we suppose $v = (1-12)$ and $|w \cap v| = 4$ or 8 . If $|v \cap w| = 4$, then $|v + w| = 16$ and hence, $\deg(V) \geq |v \cap w| + |v + w| = 20$, which is a contradiction.

Therefore, $v \cap w = (1-8)$ and $w = (1-8, 13-16)$. Analogously, we have $|v \cap u| = |w \cap u| = 8$.

If $t \leq 3$, then $|v| \geq |v \cap w \cap u| + |(v \cap w) \setminus (v \cap w \cap u)| + |(v \cap u) \setminus (v \cap w \cap u)| \geq 3+5+5 = 13$, a contradiction. Therefore, $t \geq 5$.

If $t = 7$, then $\deg(V) \geq 19$. To prove this, we consider $u = (i_1, \dots, i_{12})$, where $i_1, \dots, i_7 \in v \cap w \cap u$, but since $v \cap w = (1-8)$ then $i_1, \dots, i_7 \in v \cap w$. Suppose $u = (1-7, i_8, \dots, i_{12})$. Since $|u \cap v| = 8$ and $|u \cap w| = 8$, so $i_8 \in \{9, 10, 11, 12\}$ whereas $i_9 \in \{13, 14, 15, 16\}$. We choose $i_8 = 10$ and $i_9 = 13$. Hence $i_j \notin \{9, 11, 12, 14, 15, 16\}$, $j = 10, 11, 12$. Hence, a possibility for u it will be $u = (1-7, 10, 13, 17-19)$ so that $\deg(V) \geq 19$, a contradiction with the fact that V is a minimal representation. Thus, $t = 5$ and $u = (1-5, 9-11, 13-15, 17)$. Therefore, $V = V_5$.

□

Corollary 4.2. *Each minimal representation of the code loops C_1^3, \dots, C_5^3 has the following types, respectively:*

$$(1111111), (1111333), (1111115), (1111337), (1113335).$$

In order to present the Representation Theorem for nonassociative code loops of rank 4, note that, according to Theorem 2.3, we have exactly 16 code loops of rank 4, namely, $C_1^4, C_2^4, \dots, C_{16}^4$. For each C_i^4 , $i = 1, \dots, 16$, we have to find $V_i \subseteq \mathbb{F}_2^m$ doubly even code of minimal degree m such that $C_i^4 \cong L(V_i)$.

Theorem 4.3. *The code loops C_1^4, \dots, C_{16}^4 have the following minimal representations V_1, \dots, V_{16} , which are given by:*

$$\begin{aligned}
 V_1 &= \langle (1, 2, 3, 4), (1, 2, 5, 6), (1, 3, 5, 7), (1 - 8) \rangle, \\
 V_2 &= \langle (1 - 8), (1 - 4, 9 - 12), (1, 5, 6, 7, 9, 10, 11, 13), (1, 2, 5, 8, 9, 12, 13, 14) \rangle, \\
 V_3 &= \langle (1 - 8), (1 - 6, 9, 10), (1, 2, 3, 4, 5, 7, 9, 11), (1, 6 - 12) \rangle, \\
 V_4 &= \langle (1 - 8), (1 - 6, 9, 10), (1, 2, 3, 7, 9, 11 - 17), (1, 4, 7, 8, 9, 10, 11, 18) \rangle, \\
 V_5 &= \langle (1 - 8), (1, 2, 3, 4, 9, 10, 11, 12), (1, 5, 9, 13 - 17), (1, 2, 5, 6, 9, 10, 13, 18) \rangle, \\
 V_6 &= \langle (1, 2, 3, 4), (1, 2, 5, 6), (1, 3, 5, 7), (8, 9, 10, 11) \rangle, \\
 V_7 &= \langle (1 - 8), (1, 2, 3, 4, 9, 10, 11, 12), (1, 2, 3, 5, 9, 13, 14, 15), (1 - 16) \rangle, \\
 V_8 &= \langle (1 - 8), (1, 2, 3, 4, 9 - 12), (1, 2, 3, 5, 9, 13, 14, 15), (1, 2, 9, 10, 13, 14, 16, 17) \rangle, \\
 V_9 &= \langle (1 - 8), (1, 2, 3, 4, 9 - 16), (1, 5, 6, 7, 9, 10, 11, 17), (5, 6, 9, 10, 12, 13, 18, 19) \rangle, \\
 V_{10} &= \langle (1 - 8), (1, 2, 9 - 14), (1, 3, 9, 10, 11, 15, 16, 17), (1 - 6, 9 - 18) \rangle, \\
 V_{11} &= \langle (1 - 8), (1, 2, 3, 4, 9 - 12), (1, 2, 3, 5, 9, 13, 14, 15), (6, 7, 16, 17) \rangle, \\
 V_{12} &= \langle (1 - 8), (1, 2, 3, 4, 9 - 12), (1, 2, 3, 5, 9, 10, 11, 13), (1, 2, 9, 10, 14, 15, 16, 17) \rangle, \\
 V_{13} &= \langle (1 - 8), (1, 2, 9, 10), (1, 3, 9, 11), (4, 5, 12, 13, 14, 15, 16, 17) \rangle, \\
 V_{14} &= \langle (1 - 8), (1, 2, 3, 4, 9 - 12), (1, 2, 3, 5, 9, 10, 11, 13), (1, 2, 9, 10) \rangle, \\
 V_{15} &= \langle (1 - 12), (1, 2, 3, 4, 13 - 16), (1, 2, 3, 5, 13, 14, 15, 17), (1, 2, 13, 14) \rangle, \\
 V_{16} &= \langle (1 - 8), (1, 2, 9, 10, 11, 12, 13, 14), (1, 3, 9, 10, 11, 12, 13, 15), (4, 5, 16, 17) \rangle.
 \end{aligned}$$

Proof. Each V_i , $i = 1, \dots, 16$, is clearly a doubly even binary code. Now, to prove that $C_i^4 \cong L(V_i)$ we just need to find the characteristic vector associated to $L(V_i)$ and apply the Theorem 2.3 (Classification of code loop of rank 4). Therefore, V_i is a representation of C_i^4 , $i = 1, \dots, 16$.

Now, we are going to prove, up to isomorphism, that V_i is the unique minimal representation of C_i^4 , $i = 1, \dots, 16$. We suppose that $V = \langle v_1, v_2, v_3, v_4 \rangle$ is a minimal representation of C_i^4 . To determine V we calculate t_i , t_{ij} , t_{ijk} and $t = t_{1234}$ using the following properties:

$$\begin{aligned}
 (4.1) \quad & t \leq t_{ij4} \leq t_{ij} - t_{123} + t \quad (i, j = 1, 2, 3; i \neq j) \\
 & t_{124} + t_{134} - t \leq t_{14} \leq t_1 - t_{12} - t_{13} + t_{123} + t_{124} + t_{134} - t \\
 & t_{124} + t_{234} - t \leq t_{24} \leq t_2 - t_{12} - t_{23} + t_{123} + t_{124} + t_{234} - t \\
 & t_{134} + t_{234} - t \leq t_{34} \leq t_3 - t_{13} - t_{23} + t_{123} + t_{134} + t_{234} - t \\
 & t_4 \geq t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t
 \end{aligned}$$

Let's prove the unicity of each V_i , for all $i = 1, \dots, 16$, up to isomorphism. For a better visualization of these representations, see the diagrams presented in the A.

Case $i = 1$: In this case, the characteristic vector is $\lambda_1 = (1110110100)$. Since $\deg(V) \leq 8$, $t_1 \equiv t_2 \equiv t_3 \equiv 4 \pmod{8}$ and $t_{12} \equiv t_{13} \equiv t_{23} \equiv 2 \pmod{4}$, then $t_1 = t_2 = t_3 = 4$, $t_{12} = t_{13} = t_{23} = 2$ and $t_{123} = 1$. So we can assume $v_1 = (1, 2, 3, 4)$, $v_2 = (1, 2, 5, 6)$ and $v_3 = (1, 3, 5, 7)$.

Next, we find v_4 using the fact that $t_4 \equiv 0 \pmod{8}$ and $t_{14} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$. It is necessary to analyse two possible values for t : 0 and 1.

Case $t = 0$, we have $t_{ij4} = 0$ and thus, $t_{ij} = 0$, for $i, j = 1, 2, 3$ and $i \neq j$, according to Properties (4.1). It means that $\{1, \dots, 7\} \cap v_4 = \emptyset$. Therefore, $\deg(V) > 7 + t_4 \geq 15$ and then, we don't have minimal reduced representation in this case. Case $t = 1$, we

have $t_{ij4} = 2$ and thus, $t_{i4} = 4$, that is, $v_i \subset v_4$, $i = 1, 2, 3$. Then, since V is minimal, we obtain $t_4 = 8$ and we can assume $v_4 = (1-8)$. Therefore $V = V_1$.

Case $i = 2$: In this case, the characteristic vector is $\lambda_2 = (0000000000)$, so $t_i \equiv 0 \pmod{8}$ and $t_{ij} \equiv 0 \pmod{4}$ ($i \neq j$). Since $\deg(V) \leq 14$, then $t_i = 8$ and $t_{ij} = 4$; $i, j = 1, 2, 3$. Indeed, it's clear that $t_1 = 8$ and $t_{12} = t_{13} = 4$. If we assume that $t_2 = 16$ or $t_3 = 16$, we will have $l_{12} = |v_1 \cup v_2| = 20 > \deg(V)$ and $l_{13} = |v_1 \cup v_3| = 20 > \deg(V)$, which contradicts the minimality of the degree of V . Thus $t_2 = t_3 = 8$ and $t_{23} = 4$. We also obtain that $t_{123} = 1$, since if $t_{123} = 3$ gives us $l_{123} = |v_1 \cup v_2 \cup v_3| = 15 > \deg(V)$.

Therefore: $v_1 = (1-8)$, $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$ and $v_3 = (1, 5, 6, 7, 9, 10, 11, 13)$.

To find v_4 , we notice that $l_{123} = 13$ and $\{1, \dots, 13\} \not\subseteq v_4$, because otherwise, $t_4 \geq 16$ and $\deg(V) > 14$. Thus $t_4 = 8$ and v_4 has 7 elements in the set $\{1, \dots, 13\}$ and a new element, that is, $v_4 = (i_1, \dots, i_7, i_8 = 14)$, $i_j \in \{1, \dots, 13\}$, $j \in \{1, \dots, 7\}$. Furthermore, we need to find the weights t, t_{i4} and t_{ij4} that satisfy the following equation:

$$(4.2) \quad t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t + 1 = 8.$$

We have two possibilities for t : 0 or 1. Accordingly with Properties (4.1) and the minimality of V , we conclude that there is a unique solution to the Equation (4.2):

$$t = 1, \quad t_{ij4} = 2 \text{ and } t_{i4} = 4.$$

This way we get $v_4 = (1, 2, 5, 8, 9, 12, 13, 14)$ and $V = V_2$.

Case $i = 3$: $\lambda_3 = (0000110100)$; $t_1 \equiv t_2 \equiv t_3 \equiv t_4 \equiv 0 \pmod{8}$, $t_{12} \equiv t_{13} \equiv t_{23} \equiv 2 \pmod{4}$ and $t_{14} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

Since $\deg(V) \leq 12$, we have that $t_1 = 8$, $t_{12} = 2$ or 6 and $t_{13} = 2$ or 6 . We assume that $v_1 = (1-8)$. Case $t_{12} = 2$, we could consider, for instance, $v_2 = (1, 2, 9, 10, 11, 12, 13, 14)$, but this would give us $\deg(V) \geq 14$. Then, $t_{12} = 6$. As we want to find V with the smallest possible degree, we have that $t_2 = 8$ and we can assume $v_2 = (1, 2, 3, 4, 5, 6, 9, 10)$. In a similar way, we get $t_{13} = 6$, $t_3 = 8$ and $t_{23} = 6$.

Using the fact that $t_1 > t_{12} + t_{13} - t_{123}$, we obtain that $t_{123} = 5$. We consider $v_3 = (1, 2, 3, 4, 5, 7, 9, 11)$. So $v_1 \cup v_2 \cup v_3 = \{1, \dots, 11\}$.

To find v_4 , first we note that $\{1, \dots, 11\} \not\subseteq v_4$, $\{1, \dots, 11\} \cap v_4 \neq \emptyset$ and $t_4 = 8$, because otherwise we would have $\deg(V) > 12$. This way, we only have two possibilities for v_4 :

- (1) $v_4 = (i_1, \dots, i_7, i_8 = 12)$, with $i_1, \dots, i_7 \in \{1, \dots, 11\}$;
- (2) $v_4 = (i_1, \dots, i_8)$, with $i_1, \dots, i_8 \in \{1, \dots, 11\}$.

The next step is to determine the weights t, t_{ij4} and t_{i4} ($i, j = 1, 2, 3$) that satisfy the Properties (4.1), the minimality of V and one of the following equations:

$$(4.3) \quad t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t + 1 = 8$$

$$(4.4) \quad t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 8$$

We have six possibilities for t : 0, 1, 2, 3, 4 or 5. According to Table 2, we do not have a representation in cases $t = 0, 2, 3$ and 5. For $t = 0$, the Equations (4.3) and (4.4) has no solutions. For $t = 2, 3$ and $t = 5$, we have a contradiction related to the weights t_{i4} . For $t = 4$, there is a non-minimal representation and for $t = 1$, we obtain a solution to the Equation (4.3), so we can consider $v_4 = (1, 6, 7, 8, 9, 10, 11, 12)$ and, thus, $V = V_3$.

TABLE 2. V_3 .

$t = 0: 0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0$	$t = 2: 2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2$	$t = 4: 4 \leq t_{ij4} \leq 5 \Rightarrow t_{ij4} = 4$
$0 \leq t_{i4} \leq 1 \Rightarrow t_{i4} = 0$	$2 \leq t_{i4} \leq 3$	$4 \leq t_{i4} \leq 5 \Rightarrow t_{i4} = 4$
$t = 1: 1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2$	$t = 3: 3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4$	$t = 5: 5 \leq t_{ij4} \leq 6 \Rightarrow t_{ij4} = 6$
$3 \leq t_{i4} \leq 5 \Rightarrow t_{i4} = 4$	$5 \leq t_{i4} \leq 6$	$7 \leq t_{i4} \leq 8 \Rightarrow t_{i4} = 8$

Case $i = 4$: $\lambda_4 = (0010100000)$; $t_1 \equiv t_2 \equiv t_4 \equiv 0 \pmod{8}$; $t_3 \equiv 4 \pmod{8}$; $t_{12} \equiv 2 \pmod{4}$; $t_{13} \equiv t_{14} \equiv t_{23} \equiv t_{24} \equiv 0 \pmod{4}$.

First, we note that $t_{12} \leq 6$. For instance, if $t_{12} = 10$, we will have $t_1 \geq 16$ and $t_2 \geq 16$, and so $l_{12} = |v_1 \cup v_2| \geq 22$ and thus will be greater than the degree of V . Since $t_{12} \leq 6$ and $l_{12} \leq \deg(V)$, so we have $t_1 = t_2 = 8$ and so $t_{13} = t_{23} = 4$ and $t_{123} = 1$ or 3 . We can consider $v_1 = (1, 2, 3, 4, 5, 6, 7, 8)$.

Case $t_{12} = 2$, we will have $t_{123} = 1$. Since $t_3 > t_{13} + t_{23} - t_{123}$, we will obtain $t_3 \geq 12$. This gives us that $\deg(V) > 18$, because $l_{123} = t_3 + 7 \geq 19$. Therefore $t_{12} = 6$ and, we can take $v_2 = (1, 2, 3, 4, 5, 6, 9, 10)$. In this case we obtain $t_{123} = 3$, because $t_{12} + t_{13} - t_{123} < 8$. As we have $t_3 > t_{13} + t_{23} - t_{123}$, that is, $t_3 > 5$, and $l_{123} < \deg(V)$, then $t_3 = 12$. We consider $v_3 = (1, 2, 3, 7, 9, 11, 12, 13, 14, 15, 16, 17)$.

Now, to determine the generator v_4 , note que $v_1 \cup v_2 \cup v_3 = \{1, \dots, 17\} \not\subseteq v_4$. So we have two possibilities for v_4 : $v_4 = (i_1, \dots, i_{15}, 18)$ or $v_4 = (i_1, \dots, i_7, 18)$, with $i_j \in \{1, \dots, 17\}$, $j = 1, \dots, 15$. We need to find the values t_{i4}, t_{ij4} and t such that:

$$(4.5) \quad t_4 = t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t + 1.$$

We have four possibilities for t : $0, 1, 2$ or 3 . Accordingly with Properties (4.1) and the minimality of V , we conclude that there is a unique solution to the Equation (4.5):

$$t = 1, \quad t_{ij4} = 2, \quad t_{i4} = 4 \text{ and } t_4 = 8.$$

This way we get $v_4 = (1, 4, 7, 8, 9, 10, 11, 18)$ and $V = V_4$.

Case $i = 5$: $\lambda_5 = (0000010100)$; $t_1 \equiv t_2 \equiv t_3 \equiv t_4 \equiv 0 \pmod{8}$; $t_{12} \equiv t_{14} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$; $t_{13} \equiv t_{23} \equiv 2 \pmod{4}$.

Since $l_{13} < 18$ and $l_{23} < 18$, we obtain $t_1 = t_2 = t_3 = 8$, $t_{13} \leq 6$ and $t_{23} \leq 6$. If $t_{13} = 10$, for instance, we will have $t_1 \geq 16$, $t_3 \geq 16$ and $l_{13} \geq 22 \geq \deg(V)$, which contradicts the minimality of the degree of V . We also have $t_{12} = 4$ and so, $t_{123} = 1$ or 3 . As $t_1 > t_{12} + t_{13} - t_{123}$, $t_2 > t_{12} + t_{23} - t_{123}$ and $t_3 > t_{13} + t_{23} - t_{123}$, that is, $t_{13} - t_{123} < 4$, $t_{23} - t_{123} < 4$ and $t_{23} + t_{13} < t_3 + t_{123}$, we obtain $t_{123} = 1$ and $t_{13} = t_{23} = 2$. Thus, it can be considered $v_1 = (1, 8)$, $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$ and $v_3 = (1, 5, 9, 13, 14, 15, 16, 17)$.

Note that $v_1 \cup v_2 \cup v_3 = \{1, \dots, 17\} \not\subseteq v_4$, then $t_4 \leq 16$. Thus $v_4 = (i_1, \dots, i_{15}, 18)$ or $v_4 = (i_1, \dots, i_7, 18)$ where $i_j \in \{1, \dots, 17\}$. By (4.1), this means that the weights t_{i4}, t_{ij4} and t need to satisfy one of the following equations:

$$(4.6) \quad t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 15$$

$$(4.7) \quad t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 7$$

According to Table (3), there is a solution only in the case where $t_{124} = t_{134} = t_{234} = 2$ and $t_{14} = t_{24} = t_{34} = 4$. So consider $v_4 = (1, 2, 5, 6, 9, 10, 13, 18)$ to get $V = V_5$.

TABLE 3. V_5 .

$t = 0$	$t = 1$
$0 \leq t_{124} \leq 3 \Rightarrow t_{124} = 0 \text{ or } 2$	$2 \leq t_{124} \leq 4 \Rightarrow t_{124} = 2 \text{ or } 4$
$0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$	$1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$
$0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$	$1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$
$t_{124} = t_{134} = t_{234} = 0$	$t_{124} = t_{134} = t_{234} = 2$
$0 \leq t_{14} \leq 3 \Rightarrow t_{14} = 0$	$3 \leq t_{14} \leq 6 \Rightarrow t_{14} = 4$
$0 \leq t_{24} \leq 3 \Rightarrow t_{24} = 0$	$3 \leq t_{24} \leq 6 \Rightarrow t_{24} = 4$
$0 \leq t_{34} \leq 5 \Rightarrow t_{34} = 0 \text{ or } 4$	$3 \leq t_{34} \leq 8 \Rightarrow t_{34} = 4 \text{ or } 8$
$t_{124} = 2; t_{134} = t_{234} = 0$	$t_{124} = 4; t_{134} = t_{234} = 2$
$2 \leq t_{14} \leq 5 \Rightarrow t_{14} = 4$	$5 \leq t_{14} \leq 8 \Rightarrow t_{14} = 8$
$2 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4$	$5 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8$
$0 \leq t_{34} \leq 5 \Rightarrow t_{34} = 0 \text{ or } 4$	$3 \leq t_{34} \leq 8 \Rightarrow t_{34} = 4 \text{ or } 8$

Case $i = 6$: $\lambda_6 = (1111110100)$; $t_1 \equiv t_2 \equiv t_3 \equiv t_4 \equiv 4 \pmod{8}$; $t_{12} \equiv t_{13} \equiv t_{23} \equiv 2 \pmod{4}$; $t_{14} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

As $\deg(V) \leq 11$, then $t_1 = 4$, which implies $t_{12} = t_{13} = 2$ and $t_{123} = 1$. Furthermore, the fact that $l_{12} < 11$ and $l_{13} < 11$ gives us $t_2 = t_3 = 4$ and $t_{23} = 2$. Thus, consider $v_1 = (1, 2, 3, 4)$, $v_2 = (1, 2, 5, 6)$ and $v_3 = (1, 3, 5, 7)$. To find v_4 , note that the minimality of the degree of V implies that $\{1, \dots, 7\} \not\subseteq v_4$, and so, $t_4 = 4$. Thus, $t_{i4} = 0$, and this way, $t = t_{ij4} = 0$. Therefore, $\{1, \dots, 7\} \cap v_4 = \emptyset$, which implies $v_4 = (8, 9, 10, 11)$ and $V = V_6$.

Case $i = 7$: $\lambda_7 = (0001000000)$; $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$; $t_4 \equiv 4 \pmod{8}$; $t_{12} \equiv t_{13} \equiv 0 \pmod{4}$; $t_{23} \equiv t_{14} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

As $l_{ij} < 16$, we have $t_i = 8$ and $t_{ij} = 4$, for $i, j = 1, 2, 3$. If, for instance, $t_1 = t_2 = 16$ and $t_{12} = 12$, we will have $l_{12} = 20$. Suppose $v_1 = (1-8)$ and $v_2 = (2, 3, 4, 9, 10, 11, 12)$.

Note that there are two possibilities for t_{123} : 1 or 3. Both cases were analyzed and the results are shown in the Table 4.

We assume $v_3 = (1, 5, 6, 7, 9, 10, 11, 13)$ in the first case and $v_3 = (1, 2, 3, 5, 9, 13, 14, 15)$ in the second case. By item (1.1.1.), we have $\{1, \dots, 13\} \cap v_4 = \emptyset$. If $t_4 = 4$ and $v_4 = (14, 15, 16, 17)$, for instance, we would obtain a representation with a degree greater than 16. In items (1.1.2.), (1.2.1.), (1.2.2.), (2.1.), (2.2.) and (2.3.) we also get non-minimal representations. In items (1.1.3.), (1.2.3.) and cases similar to these, we have a contradiction with the hypothesis $t_{14} \equiv 0 \pmod{4}$. Finally, the minimal representation is determined in the last case because $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 15$. Then, if $t_4 = 16$, we can consider $v_4 = (1, \dots, 16)$ and therefore, $V = V_7$.

TABLE 4. V_7 .

Case 1: $t_{123} = 1$	
1.1. $t = 0: 0 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2 \text{ or } 4$	1.2. $t = 1: 1 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 2 \text{ or } 4$
1.1.1. $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{i4} \leq 1 \Rightarrow t_{i4} = 0$	1.2.1. $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 3 \leq t_{i4} \leq 4 \Rightarrow t_{i4} = 4$
1.1.2. $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 4 \leq t_{i4} \leq 5 \Rightarrow t_{i4} = 4$	1.2.2. $t_{124} = t_{134} = t_{234} = 4$ $\Rightarrow 7 \leq t_{i4} \leq 8 \Rightarrow t_{i4} = 8$
1.1.3. $t_{124} = 2; t_{134} = t_{234} = 0$ $\Rightarrow 2 \leq t_{i4} \leq 3$	1.2.3. $t_{124} = 4; t_{134} = t_{234} = 2$ $\Rightarrow 5 \leq t_{i4} \leq 6$
Case 2: $t_{123} = 3$	
2.1. $t = 0: 0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0$ $0 \leq t_{i4} \leq 3 \Rightarrow t_{i4} = 0$	2.3. $t = 2: 2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2$ $2 \leq t_{i4} \leq 5 \Rightarrow t_{i4} = 4$
2.2. $t = 1: 1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2$ $3 \leq t_{i4} \leq 6 \Rightarrow t_{i4} = 4$	2.4. $t = 3: 3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4$ $5 \leq t_{i4} \leq 8 \Rightarrow t_{i4} = 8$

Case $i = 8$: $\lambda_8 = (0000001000)$; $t_1 \equiv t_2 \equiv t_3 \equiv t_4 \equiv 0 \pmod{8}$; $t_{12} \equiv t_{13} \equiv t_{23} \equiv 0 \pmod{4}$; $t_{24} \equiv t_{34} \equiv 0 \pmod{4}$; $t_{14} \equiv 2 \pmod{4}$.

As $l_{ij} < 17$, we need to have $t_i = 8$ and $t_{ij} = 4$, for $i, j = 1, 2, 3$. Assume $v_1 = (1 - 8)$ and $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$. There two cases to analyse (see Table 5).

In case 1, we assume that $v_3 = (1, 5, 6, 7, 9, 10, 11, 13)$. In (1.1.1.), if $0 \leq t_{i4} \leq 1$ then $t_{i4} = 0$ and thus, $\{1, \dots, 13\} \cap v_4 = \emptyset$. Soon $t_4 \geq 8$ and $\deg(V) \geq 21$, that is, we will obtain a non-minimal representation. In other cases, we find contradictions related to the weights t_{i4} . The other cases are analyzed in a similar way. On the other hand, in case 2, we assume $v_3 = (1, 2, 3, 5, 9, 13, 14, 15)$. If $t = 0$, we will have $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 2$. Then, if $t_4 = 8$ and if we consider $v_4 = (i_1, i_2, 16, \dots, 21)$, with $i_1, i_2 \in \{1, \dots, 15\}$ and $i_1, i_2 \in v_1 \cap v_4$, we will have a non-minimal representation. If $t = 1$ and $t = 3$, we will also find non-minimal representations with degree equal to 21 and 18, respectively. If $t = 2$, then $t_4 \geq t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 6$. If we assume $t_4 = 8$ and $v_4 = (1, 2, 9, 10, 13, 14, 16, 17)$, we obtain $V = V_8$.

TABLE 5. V_8 .

Case 1: $t_{123} = 1$	
1.1. $t = 0$: $0 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 0$ or 2	1.2. $t = 1$: $1 \leq t_{124} \leq 4 \Rightarrow t_{124} = 2$ or 4
1.1.1. $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{i4} \leq 1 \Rightarrow t_{i4} = 0$	1.2.1. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$
1.1.2. $t_{124} = 2$; $t_{134} = t_{234} = 0 \Rightarrow 2 \leq t_{24} \leq 3$	1.2.2. $t_{124} = 2$; $t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$
1.1.3. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 4 \leq t_{14} \leq 5$	1.2.3. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 7 \leq t_{14} \leq 8$
Case 2: $t_{123} = 3$	
2.1. $t = 0$: $0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0$	2.3. $t = 2$: $2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2$
$0 \leq t_{i4} \leq 3 \Rightarrow t_{14} = 2$; $t_{24} = t_{34} = 0$	$2 \leq t_{i4} \leq 5 \Rightarrow t_{14} = 2$; $t_{24} = t_{34} = 4$
2.2. $t = 1$: $1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2$	2.4. $t = 3$: $3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4$
$3 \leq t_{i4} \leq 6 \Rightarrow t_{14} = 6$; $t_{24} = t_{34} = 4$	$5 \leq t_{i4} \leq 8 \Rightarrow t_{14} = 6$; $t_{24} = t_{34} = 8$

Case $i = 9$: $\lambda_9 = (0100001000)$; $t_1 \equiv t_3 \equiv t_4 \equiv 0 \pmod{8}$; $t_2 \equiv 4 \pmod{8}$; $t_{14} \equiv 2 \pmod{4}$; $t_{12} \equiv t_{13} \equiv t_{23} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

We affirm that $t_1 = 8$, $t_2 = 12$ and $t_{12} = 4$. Indeed, first note that $l_{12} < 19$ implies $t_1 \leq 16$. If $t_1 = 8$, it is clear that $t_{12} = 4$ and $t_2 > 4$. Since $l_{12} < 19$, so $t_2 < 15$, which implies that $t_2 = 12$. If $t_1 = 16$, we will have $t_{12} = 4, 8$ or 12. But, in order to have $l_{12} < 19$, we should have $t_2 - t_{12} < 3$, which does not happen for any value of t_2 and t_{12} . Let $v_1 = (1 - 8)$ and $v_2 = (1, 2, 3, 4, 9, 10, 11, 12, 13, 14, 15, 16)$. Note that $t_{13} = 4$ and $t_3 = 8$, because $t_1 = 8$ and $l_{13} < 19$. So, $t_{23} = 4$.

Now, let's find v_3 and v_4 . The possible cases for the remaining weights are shown in Table 6. In case 1, we assume $v_3 = (1, 5, 6, 7, 9, 10, 11, 17)$. For items (1.1.1.), (1.1.2.) - (1.1.7.), (1.2.1.) - (1.2.4.), (1.2.7.) and (1.2.8), we have contradictions related to the weights t_{i4} . In items (1.1.3.), (1.2.5.) and (1.2.6.), we obtain representations of degree equal to 21, 20 and 22, respectively. In item (1.1.8.), since we have $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 6$, we can consider $v_4 = (i_1, \dots, i_6, 18, 19)$, with $\{i_1, \dots, i_6\} \subset \{1, \dots, 17\}$, and, therefore, $V = V_9$. In case 2, we can consider $v_3 = (1, 2, 3, 5, 9, 17, 18, 19)$. In this case, if there is another representation of degree 19, we must have $v_4 \subset \{1, \dots, 19\}$. Soon $t_4 = 8$ or $t_4 = 16$, and $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 8$ or $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 16$. After analyzing all the possibilities for t , we see that these equations have no solution.

TABLE 6. V_9 .

Case 1: $t_{123} = 1$	
1.1. $t = 0: 0 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 0 \text{ or } 2$	1.2. $t = 1: 1 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 2 \text{ or } 4$
1.1.1. $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$	1.2.1. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$
1.1.2. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 4 \leq t_{14} \leq 5$	1.2.2. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 7 \leq t_{14} \leq 8$
1.1.3. $t_{124} = 2; t_{134} = t_{234} = 0$ $\Rightarrow 2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 2 \leq t_{24} \leq 7 \Rightarrow t_{24} = 4;$ $0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$	1.2.3. $t_{124} = t_{134} = 2; t_{234} = 4 \Rightarrow 3 \leq t_{14} \leq 4$
1.1.4. $t_{124} = t_{134} = 2; t_{234} = 0 \Rightarrow 4 \leq t_{14} \leq 5$	1.2.4. $t_{124} = t_{134} = 4; t_{234} = 2 \Rightarrow 7 \leq t_{14} \leq 8$
1.1.5. $t_{124} = t_{234} = 2; t_{134} = 0 \Rightarrow 2 \leq t_{34} \leq 3$	1.2.5. $t_{124} = 2; t_{134} = t_{234} = 4$ $\Rightarrow 5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 5 \leq t_{24} \leq 10 \Rightarrow t_{24} = 8;$ $7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$
1.1.6. $t_{124} = t_{134} = 0; t_{234} = 2 \Rightarrow 0 \leq t_{14} \leq 1$	1.2.6. $t_{124} = 4; t_{134} = t_{234} = 2$ $\Rightarrow 5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 5 \leq t_{24} \leq 10 \Rightarrow t_{24} = 8;$ $3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$
1.1.7. $t_{124} = t_{234} = 0; t_{134} = 2 \Rightarrow 2 \leq t_{34} \leq 3$	1.2.7. $t_{124} = t_{234} = 4; t_{134} = 2 \Rightarrow 7 \leq t_{14} \leq 8$
1.1.8. $t_{124} = 0; t_{134} = t_{234} = 2$ $\Rightarrow 2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 2 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$	1.2.8. $t_{124} = t_{234} = 2; t_{134} = 4 \Rightarrow 5 \leq t_{34} \leq 6$
Case 2: $t_{123} = 3$	
2.1. $t = 0: 0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0$ $0 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 0 \leq t_{34} \leq 3 \Rightarrow t_{34} = 0;$ $0 \leq t_{24} \leq 7 \Rightarrow t_{24} = 0 \text{ or } 4$	2.3. $t = 2: 2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2$ $2 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 2 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4;$ $2 \leq t_{24} \leq 9 \Rightarrow t_{24} = 4 \text{ or } 8$
2.2. $t = 1: 1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2$ $3 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 3 \leq t_{34} \leq 6 \Rightarrow t_{34} = 4;$ $3 \leq t_{24} \leq 10 \Rightarrow t_{24} = 4 \text{ or } 8$	2.4. $t = 3: 3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4$ $5 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 5 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8;$ $5 \leq t_{24} \leq 12 \Rightarrow t_{24} = 8 \text{ or } 12$

Case $i = 10$: $\lambda_{10} = (0001111000)$; $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$; $t_4 \equiv 4 \pmod{8}$;
 $t_{12} \equiv t_{13} \equiv 2 \pmod{4}$; $t_{14} \equiv 2 \pmod{4}$; $t_{23} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

Let's prove that $t_1 = t_2 = 8$ and $t_{12} = 2$. If $t_1 = 8$, then $t_{12} = 2$ or 6. Case $t_{12} = 6$, $t_{12} < 19$ implies $t_2 = 8$ or 16. There isn't representations in this case. Indeed, if $t_2 = 8$ and $t_{23} < 19$, then $t_{23} = 4$ and $t_3 = 8$. As $t_{12} + t_{23} - t_{123} < t_2$, so $t_{123} = 3$, which implies $t_{13} > 2$. This is an absurd, because we need to have $t_{12} + t_{13} - t_{123} < t_1 = 8$. The analysis for $t_2 = 16$ in this case is analogous. Therefore, $t_{12} = 2$. Furthermore, $t_{12} < 19$ implies $t_2 = 8$. Assume $v_1 = (1 - 8)$ and $v_2 = (1, 2, 9, 10, 11, 12, 13, 14)$. If $t_1 = 16$, then $t_2 - t_{12} < 3$, which only has a solution if $t_2 = 8$ and $t_{12} = 6$. There is also no representation in this case.

By $t_{12} = 2$, $t_2 = 8$ and $t_{23} < 19$, note that $t_{123} = 1$, $t_{23} = 4$ and $t_3 = 8$. We use the fact that $t_{13} + t_{23} - t_{123} < t_3$ to obtain $t_{13} = 2$. Then, we can assume $v_3 = (1, 3, 9, 10, 11, 15, 16, 17)$.

To find v_4 , we analyze the possibilities for the weights t, t_{i4} and t_{ij4} , presented in Table 7. In items (1.2.), (2.1.) and (2.2.), we have representations of degree 21, 20 and 22, respectively. As last, in item (1.1.), we have $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 2$, so $t_4 = 4$ and we can consider $v_4 = (4, 5, 18, 19)$. Therefore, $V = V_{10}$.

TABLE 7. V_{10} .

1. $t = 0$	2. $t = 1$
$0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$	$1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$
$0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$	$1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$
$0 \leq t_{234} \leq 3 \Rightarrow t_{234} = 0 \text{ or } 2$	$1 \leq t_{234} \leq 4 \Rightarrow t_{234} = 2 \text{ or } 4$
1.1. $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 3 \Rightarrow t_{24} = 0;$ $0 \leq t_{34} \leq 3 \Rightarrow t_{34} = 0$	2.1. $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 3 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 3 \leq t_{24} \leq 6 \Rightarrow t_{24} = 4;$ $3 \leq t_{34} \leq 6 \Rightarrow t_{34} = 4$
1.2. $t_{124} = t_{134} = 0; t_{234} = 2$ $\Rightarrow 0 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 2 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $2 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$	2.2. $t_{124} = t_{134} = 2; t_{234} = 4$ $\Rightarrow 3 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 5 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8;$ $5 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$

Case $i = 11$: $\lambda_{11} = (0001001000)$; $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$; $t_4 \equiv 4 \pmod{8}$; $t_{14} \equiv 2 \pmod{4}$; $t_{12} \equiv t_{13} \equiv t_{23} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

Firt, note that $t_1 < 16$. If $t_1 = 16$, for instance, we will have $t_{12} = 4, 8$ or 12 . Case $t_{12} = 4$: since $l_{12} < 17$ so $t_2 < 5$, which is an absurd. Case $t_{12} = 8$ or 12 : $t_2 \geq 16$, which give us $l_{12} > 17$, which can't happen either. Therefore, $t_1 = 8$ and, this way $t_{12} = t_{13} = 4$. By $l_{12} < 17$, we get $t_2 = 8$. Analogously, $t_3 = 8$ and $t_{23} = 4$. Assume $v_1 = (1-8)$ and $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$.

Let's find v_3 and v_4 . According to Table 8, there are two cases to analyze: $t_{123} = 1$ and $t_{123} = 3$. In case 1, we don't have representations. In case 2, we can consider $v_3 = (1, 2, 3, 5, 9, 13, 14, 15)$. In item (2.2.) we also don't have a representation. In last two cases (2.3. and 2.4.) we have non-minimal representations. But if $t = 0$, that is, in item (2.1.), we have $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 2$. Then if $t_4 = 4$ and $v_4 = (6, 7, 16, 17)$ we obtain $V = V_{11}$. If $t_4 = 12$, we will obtain a representation with degree greater than 17.

TABLE 8. V_{11} .

Case 1: $t_{123} = 1$	
1.1. $t = 0: 0 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 0 \text{ or } 2$	1.2. $t = 1: 1 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 2 \text{ or } 4$
$t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$	$t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$
$t_{124} = t_{134} = t_{234} = 2 \Rightarrow 4 \leq t_{14} \leq 5$	$t_{124} = t_{134} = t_{234} = 4 \Rightarrow 7 \leq t_{14} \leq 8$
$t_{124} = 2; t_{134} = t_{234} = 0 \Rightarrow 2 \leq t_{24} \leq 3$	$t_{124} = t_{134} = 2; t_{234} = 4 \Rightarrow 3 \leq t_{14} \leq 4$
$t_{124} = t_{134} = 2; t_{234} = 0 \Rightarrow 4 \leq t_{14} \leq 5$	$t_{124} = t_{134} = 4; t_{234} = 2 \Rightarrow 7 \leq t_{14} \leq 8$
$t_{124} = t_{234} = 2; t_{134} = 0 \Rightarrow 2 \leq t_{34} \leq 3$	$t_{124} = 2; t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$
$t_{124} = t_{134} = 0; t_{234} = 2 \Rightarrow 0 \leq t_{14} \leq 1$	$t_{124} = 4; t_{134} = t_{234} = 2 \Rightarrow 5 \leq t_{24} \leq 6$
$t_{124} = t_{234} = 0; t_{134} = 2 \Rightarrow 2 \leq t_{34} \leq 3$	$t_{124} = t_{234} = 4; t_{134} = 2 \Rightarrow 5 \leq t_{34} \leq 6$
$t_{124} = 0; t_{134} = t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$	$t_{124} = t_{234} = 2; t_{134} = 4 \Rightarrow 5 \leq t_{34} \leq 6$
Case 2: $t_{123} = 3$	
2.1. $t = 0: 0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0$	2.3. $t = 2: 2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2$
$0 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 3 \Rightarrow t_{24} = 0;$ $0 \leq t_{34} \leq 3 \Rightarrow t_{34} = 0$	$2 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 2 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $2 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$
2.2. $t = 1: 1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2$	2.4. $t = 3: 3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4$
$3 \leq t_{24} \leq 6$	$5 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 5 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8;$ $5 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$

Case $i = 12$: $\lambda_{12} = (0000001100)$; $t_1 \equiv t_2 \equiv t_3 \equiv t_4 \equiv 0 \pmod{8}$; $t_{12} \equiv t_{13} \equiv t_{24} \equiv 0 \pmod{4}$; $t_{34} \equiv 0 \pmod{4}$; $t_{14} \equiv t_{23} \equiv 2 \pmod{4}$.

Similarly to what was done in the previous case, we obtain $t_1 = t_2 = t_3 = 8$ and $t_{12} = t_{13} = 4$. Let $v_1 = (1-8)$ and $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$. Note that $t_{23} = 2$ or 6 . Furthermore, since $t_{12} + t_{23} - t_{123} < 8$, then $t_{23} - t_{123} < 4$. Therefore there is two cases to analyze (see Table 9).

Note that it is only possible to find representations in the following cases: (1.1.2.), (1.1.3.), (1.2.2.), (1.2.3.), (2.1.1.), (2.2.1.), (2.4.2.) and (2.3.2.). In the last item we find the minimal representation we want. Indeed: In case 1, consider $v_3 = (1, 5, 6, 7, 9, 13, 14, 15)$. For (1.1.2.): $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 4$. If, for instance, we consider $v_4 = (i_1, \dots, i_4, 16, 17, 18, 19)$, with $\{i_1, \dots, i_4\} \subset \{1, \dots, 15\}$, we will have a non-minimal representation with degree equal to 19. For (1.2.2.): $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 11$. We also obtain a non-minimal representation. For items (1.1.3.) and (1.2.3.): the analysis is analogous to the previous item.

In case 2, let $v_3 = (1, 2, 3, 5, 9, 10, 11, 13)$. For (2.1.1.), (2.2.1.) and (2.4.2.): we obtain a non-minimal representation of degree equal to 19, 20 and 18, respectively. For (2.3.2.): $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 4$ and $t_4 = 8$. If we consider $v_4 = (1, 2, 9, 10, 14, 15, 16, 17)$ we get $V = V_{12}$.

TABLE 9. V_{12} .

Case 1: $t_{123} = 1$ and $t_{23} = 2$	
1.1. $t = 0$ $0 \leq t_{124} \leq 3 \Rightarrow t_{124} = 0$ or 2 $0 \leq t_{134} \leq 3 \Rightarrow t_{134} = 0$ or 2 $0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$	1.2. $t = 1$ $1 \leq t_{124} \leq 4 \Rightarrow t_{124} = 2$ or 4 $1 \leq t_{134} \leq 4 \Rightarrow t_{134} = 2$ or 4 $1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$
1.1.1. $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$	1.2.1. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$
1.1.2. $t_{124} = t_{234} = 0; t_{134} = 2$ $\Rightarrow 2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 3 \Rightarrow t_{24} = 0;$ $2 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$	1.2.2. $t_{124} = t_{234} = 2; t_{134} = 4$ $\Rightarrow 5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 3 \leq t_{24} \leq 6 \Rightarrow t_{24} = 4;$ $5 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$
1.1.3. $t_{124} = 2; t_{134} = t_{234} = 0$ $\Rightarrow 2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 2 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $0 \leq t_{34} \leq 3 \Rightarrow t_{34} = 0$	1.2.3. $t_{124} = 4; t_{134} = t_{234} = 2$ $\Rightarrow 5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 5 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8;$ $3 \leq t_{34} \leq 6 \Rightarrow t_{34} = 4$
1.1.4. $t_{124} = t_{134} = 2; t_{234} = 0 \Rightarrow 4 \leq t_{14} \leq 5$	1.2.4. $t_{124} = t_{134} = 4; t_{234} = 2 \Rightarrow 7 \leq t_{14} \leq 8$
Case 2: $t_{123} = 3$ and $t_{23} = 6$	
2.1. $t = 0$ $0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$ $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$ $0 \leq t_{234} \leq 3 \Rightarrow t_{234} = 0$ or 2	2.3. $t = 2$ $2 \leq t_{124} \leq 3 \Rightarrow t_{124} = 2$ $2 \leq t_{134} \leq 3 \Rightarrow t_{134} = 2$ $2 \leq t_{234} \leq 5 \Rightarrow t_{234} = 2$ or 4
2.1.1. $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0;$ $0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$	2.3.1. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$
2.1.2. $t_{124} = t_{134} = 0; t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$	2.3.2. $t_{124} = t_{134} = 2; t_{234} = 4$ $\Rightarrow 2 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 4 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$
2.2. $t = 1$ $1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$ $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$ $1 \leq t_{234} \leq 4 \Rightarrow t_{234} = 2$ or 4	2.4. $t = 3$ $3 \leq t_{124} \leq 4 \Rightarrow t_{124} = 4$ $3 \leq t_{134} \leq 4 \Rightarrow t_{134} = 4$ $3 \leq t_{234} \leq 6 \Rightarrow t_{234} = 4$ or 6
2.2.1. $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 3 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4;$ $3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$	2.4.1. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$
2.2.2. $t_{124} = t_{134} = 2; t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$	2.4.2. $t_{124} = t_{134} = 4; t_{234} = 6$ $\Rightarrow 5 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 7 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8;$ $7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$

Case $i = 13$: $\lambda_{13} = (0110111100)$; $t_1 \equiv t_4 \equiv 0 \pmod{8}$; $t_2 \equiv t_3 \equiv 4 \pmod{8}$; $t_{24} \equiv t_{34} \equiv 0 \pmod{4}$; $t_{12} \equiv t_{13} \equiv t_{14} \equiv t_{23} \equiv 2 \pmod{4}$.

As $\deg(V) = 17$, then $t_1 = 8$ and so, $t_{12} = 2$ or 6 and $t_{13} = 2$ or 6 . If $t_1 = 16$, we will have $t_2 - t_{12} < 1$, because $l_{12} < 17$, but this can't happen. Since $t_1 = 8$ and $l_{12} < 17$,

we must have $t_2 - t_{12} < 9$. Therefore, there are two cases to analyze: (1) $t_2 = 12$ and $t_{12} = 6$; (2) $t_2 = 4$ and $t_{12} = 2$ (see Table 10). In case 1, there are two subcases to analyze, because we want $l_{23} = t_2 + t_3 - t_{23}$ to be less than 17, that is, $t_3 - t_{23} < 5$. By item (1.1.), we find $t_{13} = 6$ and $t_{123} = 5$, because $l_{13} = t_1 + t_3 - t_{13} < 17$ and $t_{12} + t_{13} - t_{123} < 8$. So let $v_1 = (1 - 8)$, $v_2 = (1 - 6, 9 - 14)$ and $v_3 = (1 - 5, 7, 9 - 13, 15)$. It is only possible to find (non-minimal) representations in items (1.1.3.c) and (1.1.4.c). In any other item, we find some contradiction in the hypotheses related to the weights t_{i4} . In item (1.2.) we have $t_{13} = 2$, so $t_{123} = 1$, because $l_{13} = t_1 + t_3 - t_{13} < 17$ and t_{12} . Let $v_1 = (1 - 8)$, $v_2 = (1 - 6, 9 - 14)$ and $v_3 = (1, 7, 9, 15)$. There are non-minimal representations for items (1.2.1.c) and (1.2.2.c).

In case 2, we have $t_{23} = 2$ and $t_{123} = 1$. We want $l_{23} = t_2 + t_3 - t_{23}$ and $l_{13} = t_1 + t_3 - t_{13}$ to be smaller than 17, then $t_3 = 4$ or 12 and $t_3 - t_{13} < 9$. There are two possible solutions: (2.1.) $t_3 = 4$ and $t_{13} = 2$ and (2.2.) $t_3 = 12$ and $t_{13} = 6$. For item (2.1.), suppose $v_1 = (1 - 8)$, $v_2 = (1, 2, 9, 10)$ and $v_3 = (1, 3, 9, 11)$. By items (2.1.1.), (2.2.1.c) and (2.2.2.c), there are non-minimal representations. In item (2.1.2.), since we have $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 2$, so $t_4 \geq 8$. If $t_4 = 8$ and $v_4 = (4, 5, 12 - 17)$, with $v_1 \cap v_4 = \{4, 5\}$, we obtain $V = V_{13}$.

TABLE 10. V_{13} .

Case 1: $t_2 = 12$ and $t_{12} = 6$	
1.1.	$t_3 = 12$ and $t_{23} = 10$
1.1.1.	$t = 0$: $0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$; $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$; $0 \leq t_{234} \leq 5 \Rightarrow t_{234} = 0, 2$ or 4 ; $0 \leq t_{14} \leq 1$
1.1.2.	$t = 1$: $1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$; $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$; $1 \leq t_{234} \leq 6 \Rightarrow t_{234} = 2, 4$ or 6 ; $3 \leq t_{14} \leq 4$
1.1.3.	$t = 2$: $2 \leq t_{124} \leq 3 \Rightarrow t_{124} = 2$; $2 \leq t_{134} \leq 3 \Rightarrow t_{134} = 2$; $2 \leq t_{234} \leq 7 \Rightarrow t_{234} = 2, 4$ or 6 (a) $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$ (b) $t_{124} = t_{134} = 2$; $t_{234} = 6 \Rightarrow 6 \leq t_{24} \leq 7$ (c) $t_{124} = t_{134} = 2$; $t_{234} = 4$; $2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2$; $4 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4$; $4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$
1.1.4.	$t = 3$: $3 \leq t_{124} \leq 4 \Rightarrow t_{124} = 4$; $3 \leq t_{134} \leq 4 \Rightarrow t_{134} = 4$; $3 \leq t_{234} \leq 8 \Rightarrow t_{234} = 4, 6$ or 8 (a) $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$ (b) $t_{124} = t_{134} = 4$; $t_{234} = 8 \Rightarrow 9 \leq t_{24} \leq 10$ (c) $t_{124} = t_{134} = 4$; $t_{234} = 6$; $5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6$; $7 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8$; $7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$
1.1.5.	$t = 4$: $4 \leq t_{124} \leq 5 \Rightarrow t_{124} = 2$ or 4 ; $4 \leq t_{134} \leq 5 \Rightarrow t_{134} = 2$; $4 \leq t_{234} \leq 9 \Rightarrow t_{234} = 2$; $4 \leq t_{14} \leq 5$
1.1.6.	$t = 5$: $5 \leq t_{124} \leq 6 \Rightarrow t_{124} = 2$ or 4 ; $5 \leq t_{134} \leq 6 \Rightarrow t_{134} = 2$; $5 \leq t_{234} \leq 10 \Rightarrow t_{234} = 2$; $7 \leq t_{14} \leq 8$
1.2.	$t_3 = 4$ and $t_{23} = 2$
1.2.1.	$t = 0$: $0 \leq t_{124} \leq 5 \Rightarrow t_{124} = 0, 2$ or 4 ; $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$; $0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$ (a) $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$ (b) $t_{124} = 4$; $t_{134} = t_{234} = 0 \Rightarrow 4 \leq t_{14} \leq 5$ (c) $t_{124} = 2$; $t_{134} = t_{234} = 0$; $2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2$; $2 \leq t_{24} \leq 7 \Rightarrow t_{24} = 4$; $0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$

Continues on the next page

Table 10 (continued)	
Case 1: $t_2 = 12$ and $t_{12} = 6$	
1.2.2. $t = 1$:	
$1 \leq t_{124} \leq 6 \Rightarrow t_{124} = 2, 4 \text{ or } 6$; $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$; $1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$ (a) $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$ (b) $t_{124} = 6$; $t_{134} = t_{234} = 2 \Rightarrow 7 \leq t_{14} \leq 8$ (c) $t_{124} = 4$; $t_{134} = t_{234} = 2$; $5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6$; $5 \leq t_{24} \leq 10 \Rightarrow t_{24} = 8$; $3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$	
Case 2: $t_2 = 4$ and $t_{13} = 2$	
2.1. $t_3 = 4$ and $t_{13} = 2$	
2.1.1. $t = 0$:	
$0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0 \Rightarrow 0 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2$; $0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0$; $0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$	
2.1.2. $t = 1$:	
$1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2 \Rightarrow 3 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6$; $3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4$; $3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$	
2.2. $t_3 = 12$ and $t_{13} = 6$	
2.2.1. $t = 0$:	
$0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$; $0 \leq t_{134} \leq 5 \Rightarrow t_{134} = 0, 2 \text{ or } 4$; $0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$ (a) $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$ (b) $t_{124} = t_{234} = 0$; $t_{134} = 4 \Rightarrow 4 \leq t_{14} \leq 5$ (c) $t_{124} = t_{234} = 0$; $t_{134} = 2$; $2 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2$; $0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0$; $2 \leq t_{34} \leq 7 \Rightarrow t_{34} = 4$	
2.2.2. $t = 1$:	
$1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$; $1 \leq t_{134} \leq 6 \Rightarrow t_{134} = 2, 4 \text{ or } 6$; $1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$ (a) $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$ (b) $t_{124} = t_{234} = 2$; $t_{134} = 6 \Rightarrow 8 \leq t_{14} \leq 9$ (c) $t_{124} = t_{234} = 2$; $t_{134} = 4$; $5 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6$; $3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4$; $5 \leq t_{34} \leq 10 \Rightarrow t_{34} = 8$	

Case $i = 14$: $\lambda_{14} = (0001001100)$; $t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}$; $t_4 \equiv 4 \pmod{8}$; $t_{14} \equiv t_{23} \equiv 2 \pmod{4}$; $t_{12} \equiv t_{13} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

Since $\deg(V) \leq 13$, then $t_1 = t_2 = t_3 = 8$, $t_{12} = t_{13} = 4$ and $t_{23} = 6$ ($l_{23} < 13$). Using the facts $t_{12} + t_{23} - t_{123} < 8$ and $t_{123} < t_{12} = t_{13}$, we get $t_{123} = 3$. So let $v_1 = (1 - 8)$, $v_2 = (1, 2, 3, 4, 9, 10, 11, 12)$ and $v_3 = (1, 2, 3, 5, 9, 10, 11, 13)$.

There are four possibilities for the weight t (see Table 11). By items (1.1.), (2.1.) and (4.1.), there are non-minimal representations of degree equal to 15, 16 and 14, respectively. By item (3.1.), as $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 4$, we can consider $t_4 = 4$ and $v_4 = (1, 2, 9, 10)$. Thus, $V = V_{14}$. In other items there are no representations.

Case $i = 15$: $\lambda_{15} = (1001001100)$:] $t_1 \equiv t_4 \equiv 4 \pmod{8}$; $t_2 \equiv t_3 \equiv 0 \pmod{8}$; $t_{14} \equiv t_{23} \equiv 2 \pmod{4}$; $t_{12} \equiv t_{13} \equiv t_{24} \equiv t_{34} \equiv 0 \pmod{4}$.

It is clear that $t_1 = 12$. Besides, $t_2 = 8$ and $t_{12} = 4$, because $l_{12} < 17$, that is, $t_2 - t_{12} < 5$. Assume $v_1 = (1 - 12)$ and $v_2 = (1, 2, 3, 4, 13, 14, 15, 16)$. Analogously, we get $t_3 = 8$ and $t_{13} = 4$. We also have $t_{23} = 6$, and so $t_{123} = 3$. If $t_{23} = 2$, then $t_{123} = 1$, which implies $l_{123} = 19 > \deg(V)$. Assume $v_3 = (1, 2, 3, 5, 13, 14, 15, 17)$.

As usual, our last step is to find v_4 . According to Table 12, there are non-minimal representations by items (1.1.), (2.1.) and (4.1.). By item (3.1.), if $t_{14} = 6$, we will have a non-minimal representation. But if $t_{14} = 2$, then $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 4$. If $t_4 = 4$ and $v_4 = (1, 2, 13, 14)$, we obtain $V = V_{15}$.

TABLE 11. V_{14} .

1. $t = 0$	3. $t = 2$
$0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$ $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$ $0 \leq t_{234} \leq 3 \Rightarrow t_{234} = 0 \text{ or } 2$	$2 \leq t_{124} \leq 3 \Rightarrow t_{124} = 2$ $2 \leq t_{134} \leq 3 \Rightarrow t_{134} = 2$ $2 \leq t_{234} \leq 5 \Rightarrow t_{234} = 2 \text{ or } 4$
1.1. $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{14} \leq 3 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0;$ $0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$	3.1. $t_{124} = t_{134} = 2; t_{234} = 4$ $\Rightarrow 2 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 4 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4;$ $4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$
1.2. $t_{124} = 2; t_{134} = t_{234} = 0 \Rightarrow 2 \leq t_{24} \leq 3$	3.2. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$
2. $t = 1$	4. $t = 3$
1 $\leq t_{124} \leq 2 \Rightarrow t_{124} = 2$ $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$ $1 \leq t_{234} \leq 4 \Rightarrow t_{234} = 2 \text{ or } 4$	3 $\leq t_{124} \leq 4 \Rightarrow t_{124} = 2;$ $3 \leq t_{134} \leq 4 \Rightarrow t_{134} = 2;$ $3 \leq t_{234} \leq 6 \Rightarrow t_{234} = 4 \text{ or } 6$
2.1. $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 3 \leq t_{14} \leq 6 \Rightarrow t_{14} = 6; 3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4;$ $3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$	4.1. $t_{124} = t_{134} = 4; t_{234} = 6$ $\Rightarrow 5 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 7 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8;$ $7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$
2.2. $t_{124} = t_{134} = 2; t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$	4.2. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$

TABLE 12. V_{15} .

1. $t = 0$	3. $t = 2$
$0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$ $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$ $0 \leq t_{234} \leq 3 \Rightarrow t_{234} = 0 \text{ or } 2$	$2 \leq t_{124} \leq 3 \Rightarrow t_{124} = 2$ $2 \leq t_{134} \leq 3 \Rightarrow t_{134} = 2$ $2 \leq t_{234} \leq 5 \Rightarrow t_{234} = 2 \text{ or } 4$
1.1. $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0; 0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0;$ $0 \leq t_{14} \leq 7 \Rightarrow t_{14} = 2 \text{ or } 6;$	3.1. $t_{124} = t_{134} = 2; t_{234} = 4$ $\Rightarrow 4 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4; 4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4;$ $2 \leq t_{14} \leq 9 \Rightarrow t_{14} = 2 \text{ or } 6$
1.2. $t_{124} = t_{134} = 0; t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$	3.2. $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$
2. $t = 1$	4. $t = 3$
1 $\leq t_{124} \leq 2 \Rightarrow t_{124} = 2$ $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$ $1 \leq t_{234} \leq 4 \Rightarrow t_{234} = 2 \text{ or } 4$	3 $\leq t_{124} \leq 4 \Rightarrow t_{124} = 4;$ $3 \leq t_{134} \leq 4 \Rightarrow t_{134} = 4;$ $3 \leq t_{234} \leq 6 \Rightarrow t_{234} = 4 \text{ or } 6$
2.1. $t_{124} = t_{134} = t_{234} = 2$ $3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4; 3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4;$ $3 \leq t_{14} \leq 10 \Rightarrow t_{14} = 6 \text{ or } 10$	4.1. $t_{124} = t_{134} = 4; t_{234} = 6$ $\Rightarrow 7 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8; 7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8;$ $5 \leq t_{14} \leq 12 \Rightarrow t_{14} = 6 \text{ or } 10$
2.2. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$	4.2. $t_{124} = t_{134} = t_{234} = 4 \Rightarrow 3 \leq t_{24} \leq 6$

Case $i = 16$:

$$\lambda_{16} = (000111100) :: t_1 \equiv t_2 \equiv t_3 \equiv 0 \pmod{8}; t_4 \equiv 4 \pmod{8}; t_{24} \equiv t_{34} \equiv 0 \pmod{4};$$

$$t_{12} \equiv t_{13} \equiv t_{23} \equiv t_{24} \equiv 2 \pmod{4}.$$

We have $t_{12} \leq 6$, otherwise $t_1 \geq 16$, $t_2 \geq 16$ and $\deg(V) \geq 22$. Similarly, $t_{13} \leq 6$ and $t_{23} \leq 6$. Since $t_1 + t_2 - t_{12} \leq 17$ and $t_1 + t_3 - t_{13} \leq 17$, soon $t_1 = t_2 = t_3 = 8$. Let $v_1 = (1 - 8)$.

Let's do the analysis in two cases: $t_{12} = 2$ and $t_{12} = 6$.

In case 1, consider $v_2 = (1, 2, 9 - 14)$. In this case, $t_{123} = 1$. If $t_{13} = t_{23} = 2$, we will have $\deg(V) \geq 19$, because $t_{123} = 19$. If $t_{13} = t_{23} = 6$, we will have $t_3 > 11$, because $t_{13} + t_{23} - t_{123} = 11$. Thus we have only two subcases to analyze: (1.1.) $t_{13} = 2$ and $t_{23} = 6$; and (1.2.) $t_{13} = 6$ and $t_{23} = 2$. See Table 13.

For item (1.1.), suppose $v_3 = (1, 3, 9, 10, 11, 12, 13, 15)$. By item (1.1.1.a), where $t = 0$; $t_{124} = t_{134} = t_{234} = 0$ and $t_{14} = 2$, $t_{24} = t_{34} = 0$, we obtain $t_{14} + t_{24} + t_{34} - t_{124} - t_{134} - t_{234} + t = 2$. If $t_4 = 4$, we can consider $v_4 = (4, 5, 16, 17)$. Therefore, $V = V_{16}$. Note that

in any other item, if there is a representation, it has a degree greater than 17, that is, it is not minimal. We found non-minimal representations through the analysis of the following items: (1.1.1.c), (1.1.2.a) and (1.1.2.c).

In case 2, we have $t_{123} = 1$ or 5, since we need to have $t_1 < t_{12} + t_{13} - t_{123}$. With an analysis to the previous case, we observe that we do not have representations in this case. For item (2.1.), use that $t_1 = t_2 = 8$ to get $t_{13} = t_{23} = 2$ and for item (2.2.), note that $t_{13} = t_{23} = 6$.

TABLE 13. V_{16} .

Case 1: $t_{12} = 2$	
1.1. $t_{13} = 2$ and $t_{23} = 6$	
1.1.1. $t = 0$ $0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$ $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$ $0 \leq t_{234} \leq 5 \Rightarrow t_{234} = 0, 2$ or 4 (a) $t_{124} = t_{134} = t_{234} = 0$ $\Rightarrow 0 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 0 \leq t_{24} \leq 1 \Rightarrow t_{24} = 0; 0 \leq t_{34} \leq 1 \Rightarrow t_{34} = 0$ (b) $t_{124} = t_{134} = 0; t_{234} = 2 \Rightarrow 2 \leq t_{24} \leq 3$ (c) $t_{124} = t_{134} = 0; t_{234} = 4$ $\Rightarrow 0 \leq t_{14} \leq 5 \Rightarrow t_{14} = 2; 4 \leq t_{24} \leq 5 \Rightarrow t_{24} = 4; 4 \leq t_{34} \leq 5 \Rightarrow t_{34} = 4$	1.1.2. $t = 1$ $1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$ $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$ $1 \leq t_{234} \leq 6 \Rightarrow t_{234} = 2, 4$ or 6 (a) $t_{124} = t_{134} = t_{234} = 2$ $\Rightarrow 3 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 3 \leq t_{24} \leq 4 \Rightarrow t_{24} = 4; 3 \leq t_{34} \leq 4 \Rightarrow t_{34} = 4$ (b) $t_{124} = t_{134} = 2; t_{234} = 4 \Rightarrow 5 \leq t_{24} \leq 6$ (c) $t_{124} = t_{134} = 2; t_{234} = 6$ $\Rightarrow 3 \leq t_{14} \leq 8 \Rightarrow t_{14} = 6; 7 \leq t_{24} \leq 8 \Rightarrow t_{24} = 8; 7 \leq t_{34} \leq 8 \Rightarrow t_{34} = 8$
1.2. $t_{13} = 6$ and $t_{23} = 2$	
1.2.1. $t = 0$ $0 \leq t_{124} \leq 1 \Rightarrow t_{124} = 0$ $0 \leq t_{134} \leq 5 \Rightarrow t_{134} = 0, 2$ or 4 $0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$ (a) $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$ (b) $t_{124} = t_{234} = 0; t_{134} = 2 \Rightarrow 2 \leq t_{34} \leq 3$ (c) $t_{124} = t_{234} = 0; t_{134} = 4 \Rightarrow 4 \leq t_{14} \leq 5$	1.2.2. $t = 1$ $1 \leq t_{124} \leq 2 \Rightarrow t_{124} = 2$ $1 \leq t_{134} \leq 6 \Rightarrow t_{134} = 2, 4$ or 6 $1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$ (a) $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$ (b) $t_{124} = t_{234} = 2; t_{134} = 4 \Rightarrow 5 \leq t_{34} \leq 6$ (c) $t_{124} = t_{234} = 2; t_{134} = 6 \Rightarrow 7 \leq t_{14} \leq 8$
Case 2: $t_{12} = 6$	
2.1. $t_{123} = 1$	
2.1.1. $t = 0$ $0 \leq t_{124} \leq 5 \Rightarrow t_{124} = 0, 2$ or 4 $0 \leq t_{134} \leq 1 \Rightarrow t_{134} = 0$ $0 \leq t_{234} \leq 1 \Rightarrow t_{234} = 0$ (a) $t_{124} = t_{134} = t_{234} = 0 \Rightarrow 0 \leq t_{14} \leq 1$ (b) $t_{124} = 2; t_{134} = t_{234} = 0 \Rightarrow 2 \leq t_{24} \leq 3$ (c) $t_{124} = 4; t_{134} = t_{234} = 0 \Rightarrow 4 \leq t_{14} \leq 5$	2.1.2. $t = 1$ $1 \leq t_{124} \leq 6 \Rightarrow t_{124} = 2, 4$ or 6 $1 \leq t_{134} \leq 2 \Rightarrow t_{134} = 2$ $1 \leq t_{234} \leq 2 \Rightarrow t_{234} = 2$ (a) $t_{124} = t_{134} = t_{234} = 2 \Rightarrow 3 \leq t_{14} \leq 4$ (b) $t_{124} = 4; t_{134} = t_{234} = 2 \Rightarrow 5 \leq t_{24} \leq 6$ (c) $t_{124} = 6; t_{134} = t_{234} = 2 \Rightarrow 7 \leq t_{14} \leq 8$
2.2. $t_{123} = 5$	
(a) $t = 0: 0 \leq t_{ij4} \leq 1 \Rightarrow t_{ij4} = 0 \Rightarrow 0 \leq t_{14} \leq 1$ (b) $t = 1: 1 \leq t_{ij4} \leq 2 \Rightarrow t_{ij4} = 2 \Rightarrow 3 \leq t_{14} \leq 4$ (c) $t = 2: 2 \leq t_{ij4} \leq 3 \Rightarrow t_{ij4} = 2 \Rightarrow 2 \leq t_{24} \leq 3$	(d) $t = 3: 3 \leq t_{ij4} \leq 4 \Rightarrow t_{ij4} = 4 \Rightarrow 5 \leq t_{24} \leq 6$ (e) $t = 4: 4 \leq t_{ij4} \leq 5 \Rightarrow t_{ij4} = 4 \Rightarrow 4 \leq t_{14} \leq 5$ (f) $t = 5: 5 \leq t_{ij4} \leq 6 \Rightarrow t_{ij4} = 6 \Rightarrow 7 \leq t_{14} \leq 8$

□


As a direct consequence of the Theorem 4.3, we have the following corollary.

Corollary 4.4. *Each minimal representation of the code loops C_1^4, \dots, C_{16}^4 has the following degree and type, respectively:*

i	$\deg(V_i)$	type of V_i	i	$\deg(V_i)$	type of V_i
1	8	(11111111)	2	14	(11111111222)
3	12	(111111114)	4	18	(11111111226)
5	18	(111111112224)	6	11	(11111114)
7	17	(11113334)	8	17	(11111122223)
9	19	(11111222233)	10	19	(111223333)
11	17	(111122333)	12	17	(1111112234)
13	17	(111111236)	14	13	(111111223)
15	17	(111111227)	16	17	(111112235)

Note that in the case of code loops of rank 3 and 4 the type of code loop define this loop up to isomorphism. May be it is true in general case.






Let's see a illustration. According to Corollary 4.4, the minimal representation V_3 has 9 equivalence classes. These classes are presented in the following diagram:





ACKNOWLEDGMENTS


For financial support, the first author thanks National Council for Scientific and Technological Development - CNPq (grant 406932/2023-9); the second author thanks FAPESP (grant 2018/23690-6), CNPq (grants 307593/2023-1 and 406932/2023-9) and in accordance with the state task of the IM SB RAS, project FWNF-2022-003; and the third author thanks FAPESP (grant 2021/12820-9) and CNPq (grant 406932/2023-9). The authors are grateful to the anonymous referees for their valuable suggestions.

APPENDIX A. DIAGRAMS: MINIMAL REPRESENTATIONS OF CODE LOOPS OF RANK 4









References

- [1] O. Chein and E.G. Goodaire, *Moufang Loops with a Unique Nonidentity Commutator (Associator, Square)*, J. Algebra 130 (1990), 369-384.
- [2] R. L. Griess Jr., *Code loops*, J. Algebra 100 (1986), 224-234.
- [3] A. Grishkov and R. Miguel Pires, *Variety of loops generated by code loops*, International Journal of Algebra and Computation, Volume 28, No. 1 (2018), 163-177.
- [4] R. Miguel Pires, *Loops de código: automorfismos e representações*. Tese de Doutorado, Instituto de Matemática e Estatística, Universidade de São Paulo, São Paulo (2011).

(Miguel Pires) INSTITUTO DE CIÊNCIAS EXATAS, UNIVERSIDADE FEDERAL FLUMINENSE, RUA DESEMBARGADOR ELLIS HERMYDIO FIGUEIRA, ATERRADO, VOLTA REDONDA, RJ, BRAZIL, CEP 27213-145

Email address, Pires: `rosemarypires@id.uff.br`

(Grishkov) INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO, 1010 BUTANTÃ, SÃO PAULO, SP, BRAZIL, CEP 05508-090 AND SOBOLEV INSTITUTE OF MATHEMATICS, OMSK, RUSSIA

Email address, Grishkov: `shuragri@gmail.com`

(Rasskazova) CENTRO DE MATEMÁTICA, COMPUTAÇÃO E COLEÇÃO, UNIVERSIDADE FEDERAL DO ABC, AVENIDA DOS ESTADOS, 5001, BAIRRO BANGU, SANTO ANDRÉ, SP, BRASIL, CEP 09280-560; AND SIBERIAN STATE AUTOMOBILE AND HIGHWAY UNIVERSITY, PROSPEKT MIRA, 5, OMSK, OMSK OBLAST, RÚSSIA, 644080

Email address, Rasskazova: `marina.rasskazova@ufabc.edu.br`