

ON FIVE QUESTIONS OF BOURIN AND LEE: SYMMETRIC MODULI AND AN EULER OPERATOR IDENTITY

TENG ZHANG

ABSTRACT. We answer five questions posed by Bourin and Lee on symmetric moduli and related orbit inequalities in [12], and thereby obtain a sequence of sharp results for matrices and compact operators. We first show that the isometry-orbit identity behind the matrix weighted parallelogram law cannot be extended beyond the parameter range $0 \leq x \leq 1$, and that a counterexample already exists in dimension one. Next, we prove that the exponent 2 in the Bourin–Lee unitary-orbit estimate for the quadratic symmetric modulus is optimal in every dimension $n \geq 2$ by constructing an explicit 2×2 counterexample for all $p > 2$. We then construct a compact operator Z for which the associated singular-value inequality fails for every $p > 2$, in fact for a fixed choice of indices. We also settle a Thompson-type triangle problem for symmetric moduli: the inequality fails for the arithmetic symmetric modulus but holds for the quadratic symmetric modulus. Finally, we develop isometry-orbit refinements of an Euler operator identity and derive sharp Clarkson–McCarthy type inequalities for Schatten p -norms, together with further consequences for unitarily invariant norms and singular values.

1. INTRODUCTION

Throughout this paper, \mathcal{H} denotes a complex Hilbert space, and \mathbb{K} the ideal of compact operators on \mathcal{H} . For $Z \in \mathbb{K}$, we write $\{\mu_m^\downarrow(Z)\}_{m \geq 1}$ for the singular values of Z , namely the eigenvalues of $|Z| = (Z^*Z)^{1/2}$ (counted with multiplicity) arranged in nonincreasing order. Let $\mathbb{M}_{m,n}$ be the space of complex $m \times n$ matrices, and write $\mathbb{M}_n := \mathbb{M}_{n,n}$. Let $X \in \mathbb{M}_n$. We adopt the convention that $\mu_m^\downarrow(X) = 0$ for all $m > n$, so that $\{\mu_m^\downarrow(X)\}_{m \geq 1}$ can be regarded as an infinite nonincreasing sequence, consistent with the compact-operator notation. We denote by X^\top its transpose, and by X^* its conjugate transpose (adjoint). For scalars $d_1, \dots, d_r \in \mathbb{C}$, we denote by $\text{diag}(d_1, \dots, d_r)$ the $r \times r$ diagonal matrix whose diagonal entries are d_1, \dots, d_r (and all off-diagonal entries are 0). More generally, if $X_1 \in \mathbb{M}_{n_1}, \dots, X_k \in \mathbb{M}_{n_k}$, then $\text{diag}(X_1, \dots, X_k) := X_1 \oplus \dots \oplus X_k$ denotes the block-diagonal matrix with diagonal blocks X_1, \dots, X_k .

1.1. A problem on matrix weighted parallelogram law. For $A, B \in \mathbb{M}_n$ and $x \in \mathbb{R}$, we write $A\nabla_x B := (1-x)A + xB$ for the weighted arithmetic mean of A and B . We begin by recalling a unitary-orbit refinement of the weighted parallelogram identity due to Bourin and Lee [12].

Theorem 1.1 ([12, Corollary 2.2]). *Let $A, B \in \mathbb{M}_n$ and $0 \leq x \leq 1$. Then there exist isometries $U, V, S, T \in \mathbb{M}_{2n,n}$ such that*

$$|A \oplus B|^2 = U |A\nabla_x B|^2 U^* + V |B\nabla_x A|^2 V^* + x(1-x) \left\{ S |A - B|^2 S^* + T |A - B|^2 T^* \right\}.$$

Taking traces in Theorem 1.1 yields

$$\|A \oplus B\|_2^2 = \|A\nabla_x B\|_2^2 + \|B\nabla_x A\|_2^2 + 2x(1-x)\|A - B\|_2^2$$

2020 *Mathematics Subject Classification.* 15A60, 47A30, 47A57, 15A42, 47B10.

Key words and phrases. Symmetric modulus; unitary orbits; Euler identity; Thompson inequality; Clarkson–McCarthy type inequalities.

for $0 \leq x \leq 1$. Since the right-hand side of that trace identity is a polynomial in x , it extends to all $x \in \mathbb{R}$. This motivates the following question.

Question 1.2 ([12, Question 2.3]). *Does Theorem 1.1 hold for every $x \in \mathbb{R}$?*

We first answer Question 1.2 in the negative: the restriction $0 \leq x \leq 1$ in Theorem 1.1 cannot be removed and therefore is sharp (see Theorem 1.3 below).

Theorem 1.3. *Fix $x \in \mathbb{R} \setminus [0, 1]$. Then Theorem 1.1 fails for this x and some $A, B \in \mathbb{M}_n$. In fact, a counterexample exists already for $n = 1$.*

1.2. Symmetric moduli and three related questions. For two Hermitian matrices $A, B \in \mathbb{M}_n$, we write $A \leq B$ if $B - A$ is positive semidefinite. While for $A, B \in \mathbb{M}_n$ one has the triangle inequality

$$\|A + B\|_1 \leq \|A\|_1 + \|B\|_1,$$

simple examples show that the inequality

$$|A + B| \leq |A| + |B|$$

need not hold in the positive semidefinite order. However, Thompson [20] observed that there exist unitaries $U, V \in \mathbb{M}_n$ such that

$$|A + B| \leq U|A|U^* + V|B|V^*, \quad (1.1)$$

which, upon taking traces, yields the triangle inequality as a consequence. Such ‘‘unitary-orbit’’ inequalities have since been generalized and extended to a variety of settings; see the surveys by Bourin, Harada and Lee [7, 9].

In this paper, we generalize (1.1) to the setting of rectangular matrices (see Lemma 5.2) in order to address Question 1.11.

Recall the Cartesian decomposition of a matrix $Z \in \mathbb{M}_n$:

$$Z = \operatorname{Re} Z + i \operatorname{Im} Z, \quad \operatorname{Re} Z := \frac{Z + Z^*}{2}, \quad \operatorname{Im} Z := \frac{Z - Z^*}{2i}.$$

Note that $|\operatorname{Re} Z|^2 + |\operatorname{Im} Z|^2 = (|Z|^2 + |Z^*|^2)/2$. Bourin and Lee [12, Corollary 4.3] obtained the following unitary-orbit estimate for the quadratic symmetric modulus, which can be regarded as a matrix analogue of the scalar inequality $|z| \leq |a| + |b|$ for a complex number $z = a + ib$.

Theorem 1.4 (Bourin–Lee). *Let $Z \in \mathbb{M}_n$. Then there exist unitary matrices $U, V \in \mathbb{M}_n$ such that*

$$\sqrt{\frac{|Z|^2 + |Z^*|^2}{2}} \leq U|\operatorname{Re} Z|U^* + V|\operatorname{Im} Z|V^*.$$

They also posed the following question concerning the optimality of the exponent 2.

Question 1.5 ([12, Question 4.6]). *Let $p > 0$. Suppose that, for every $Z \in \mathbb{M}_n$, one can find unitary matrices $U, V \in \mathbb{M}_n$ such that*

$$\left(\frac{|Z|^p + |Z^*|^p}{2} \right)^{1/p} \leq U|\operatorname{Re} Z|U^* + V|\operatorname{Im} Z|V^*. \quad (1.2)$$

Must we necessarily have $p \leq 2$?

Our second result answers Question 1.5 affirmatively in every noncommutative dimension $n \geq 2$ by constructing a 2×2 counterexample for $p > 2$.

Theorem 1.6. *Let $n \geq 2$ and $p > 0$. If for every $Z \in \mathbb{M}_n$ there exist unitary matrices $U, V \in \mathbb{M}_n$ such that (1.2) holds, then necessarily $p \leq 2$.*

Remark 1.7. For $n = 1$, the statement of Question 1.5 is different: for a complex number z , the left-hand side of (1.2) equals $|z|$ for every $p > 0$, while the right-hand side equals $|\Re z| + |\Im z| \geq |z|$. Hence (1.2) holds for all $p > 0$ in dimension 1. The necessity $p \leq 2$ is therefore a genuinely noncommutative phenomenon requiring $n \geq 2$.

Bourin and Lee [12] further raised the following question on singular values.

Remark 1.8. In [12, Corollary 4.7], the indices on the right-hand side appear with a typographical slip. Starting from the operator inequality in [12, Corollary 4.3] and applying the Weyl inequality for singular values,

$$\mu_{1+j+k}^\downarrow(A + B) \leq \mu_{1+j}^\downarrow(A) + \mu_{1+k}^\downarrow(B) \quad (A, B \geq 0),$$

one obtains the corrected bound

$$\mu_{1+j+k}^\downarrow\left(\sqrt{\frac{|Z|^2 + |Z^*|^2}{2}}\right) \leq \mu_{1+j}^\downarrow(\operatorname{Re} Z) + \mu_{1+k}^\downarrow(\operatorname{Im} Z). \quad (1.3)$$

Accordingly, the natural corrected version of [12, Question 4.8] asks whether there exists $Z \in \mathbb{K}$ such that for every $p > 2$ one has

$$\mu_{1+j+k}^\downarrow\left(\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}\right) > \mu_{1+j}^\downarrow(\operatorname{Re} Z) + \mu_{1+k}^\downarrow(\operatorname{Im} Z)$$

for some integers $j, k \geq 0$.

Question 1.9 ([12, Question 4.8]). *Does there exist $Z \in \mathbb{K}$ such that for every $p > 2$ one has*

$$\mu_{1+j+k}^\downarrow\left(\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}\right) > \mu_{1+j}^\downarrow(\operatorname{Re} Z) + \mu_{1+k}^\downarrow(\operatorname{Im} Z) \quad (1.4)$$

for some pair of integers $j, k \geq 0$?

We answer Question 1.9 in the affirmative.

Theorem 1.10. *There exists $Z \in \mathbb{K}$ and fixed integers $j, k \geq 0$ such that for every $p > 2$, (1.4) holds. In our construction one may take $(j, k) = (2, 0)$.*

Let

$$|Z|_{\text{sym}} := \frac{|Z| + |Z^*|}{2}, \quad |Z|_{\text{qsym}} := \left(\frac{|Z|^2 + |Z^*|^2}{2}\right)^{1/2}$$

be the arithmetic symmetric modulus or the quadratic symmetric modulus of $Z \in \mathbb{M}_n$, respectively.

Now, we turn to a related problem raised by Bourin and Lee concerning Thompson-type triangle inequalities for symmetric moduli, compared with (1.1).

Question 1.11 ([12, Question 4.9]). *Denote by $|Z|_{(q)\text{sym}}$ either the arithmetic symmetric modulus or the quadratic symmetric modulus of $Z \in \mathbb{M}_n$. Let $X, Y \in \mathbb{M}_n$. Does the Thompson triangle type inequality*

$$|X + Y|_{(q)\text{sym}} \leq U|X|_{(q)\text{sym}}U^* + V|Y|_{(q)\text{sym}}V^*. \quad (1.5)$$

hold for some unitary $U, V \in \mathbb{M}_n$?

Our next theorem provides a complete answer to Question 1.11: the inequality (1.5) fails for the arithmetic symmetric modulus but holds for the quadratic symmetric modulus.

Theorem 1.12. *Let $X, Y \in \mathbb{M}_n$.*

(i) In general, there do not exist unitaries U, V such that

$$|X + Y|_{\text{sym}} \leq U|X|_{\text{sym}}U^* + V|Y|_{\text{sym}}V^*.$$

(ii) There exist unitaries U, V such that

$$|X + Y|_{\text{qsym}} \leq U|X|_{\text{qsym}}U^* + V|Y|_{\text{qsym}}V^*.$$

1.3. Matrix analogues of Euler's quadrilateral identity. Finally, Bourin and Lee [12] asked for matrix analogues of Euler's quadrilateral identity.

Question 1.13 ([12, Question 4.10]). *What are the matrix versions, if any, of Euler's quadrilateral identity*

$$\|x + y + z\|^2 + \|x\|^2 + \|y\|^2 + \|z\|^2 = \|x + y\|^2 + \|y + z\|^2 + \|z + x\|^2$$

for three points $x, y, z \in \mathbb{C}^n$?

A direct computation yields the following Euler-type operator identity.

Proposition 1.14 (Euler operator identity). *Let $A, B, C \in \mathbb{M}_n$. Then*

$$|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2 = |A + B|^2 + |B + C|^2 + |C + A|^2. \quad (1.6)$$

We recall that an *isometry* is a rectangular matrix $W \in \mathbb{M}_{m,n}$ such that $W^*W = I_n$. Bourin and Lee [12] suggested exploring isometry/unitary-orbit refinements of Euler-type identities. Our following three results give isometry-orbit refinements of (1.6).

Theorem 1.15. *Let $A, B, C \in \mathbb{M}_n$. Then there exist isometries $U_{ij} \in \mathbb{M}_{4n,n}$, $1 \leq i, j \leq 4$, such that*

$$\begin{aligned} |A + B|^2 \oplus |B + C|^2 \oplus |C + A|^2 \oplus 0 &= \frac{1}{4} \sum_{j=1}^4 U_{1j} |A + B + C|^2 U_{1j}^* + \frac{1}{4} \sum_{j=1}^4 U_{2j} |A|^2 U_{2j}^* \\ &\quad + \frac{1}{4} \sum_{j=1}^4 U_{3j} |B|^2 U_{3j}^* + \frac{1}{4} \sum_{j=1}^4 U_{4j} |C|^2 U_{4j}^*. \end{aligned} \quad (1.7)$$

Theorem 1.16. *Let $A, B, C \in \mathbb{M}_n$. Then there exist three isometries $U_1, U_2, U_3 \in \mathbb{M}_{3n,n}$ such that*

$$|A + B|^2 \oplus |B + C|^2 \oplus |A + C|^2 = \frac{1}{3} \sum_{k=1}^3 U_k (|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2) U_k^*.$$

Theorem 1.17. *Let $A, B, C \in \mathbb{M}_n$. Then there exist four isometries $U_1, U_2, U_3, U_4 \in \mathbb{M}_{4n,n}$ such that*

$$|A + B + C|^2 \oplus |A|^2 \oplus |B|^2 \oplus |C|^2 = \frac{1}{4} \sum_{k=1}^4 U_k (|A + B|^2 + |B + C|^2 + |A + C|^2) U_k^*.$$

Naturally, we ask the following question.

Question 1.18. *Does there exist isometries $U_1, U_2, U_3, U_4 \in \mathbb{M}_{3n,n}$ such that*

$$\begin{aligned} |A + B|^2 \oplus |B + C|^2 \oplus |A + C|^2 &= U_1 |A + B + C|^2 U_1^* + U_2 |A|^2 U_2^* \\ &\quad + U_3 |B|^2 U_3^* + U_4 |C|^2 U_4^* ? \end{aligned}$$

Unfortunately, the answer to Question 1.18 is negative. In fact, take $A = B = C = I_n$. Then

$$|A + B|^2 = |2I_n|^2 = 4I_n, \quad |A + B + C|^2 = |3I_n|^2 = 9I_n, \quad |A|^2 = |B|^2 = |C|^2 = I_n.$$

Hence the desired identity would become

$$4I_{3n} = 9U_1U_1^* + U_2U_2^* + U_3U_3^* + U_4U_4^*.$$

Since all terms on the right-hand side are positive semidefinite, we have

$$4I_{3n} \geq 9U_1U_1^*.$$

But $U_1U_1^*$ is an orthogonal projection (of rank n), so $\|9U_1U_1^*\|_\infty = 9$ while $\|4I_{3n}\|_\infty = 4$, a contradiction. Therefore such isometries cannot exist in general.

1.4. Sharp Clarkson–McCarthy type inequalities on Euler’s identity. For a Hilbert space \mathcal{H} , let S_p denote the Schatten p -class. For $p > 0$, we say that $X \in S_p$ if X is a compact operator on \mathcal{H} whose singular values $(\mu_k^\downarrow(X))_{k \geq 1}$ satisfy $\sum_{k \geq 1} \mu_k^\downarrow(X)^p < \infty$. In this case, the Schatten p -(quasi)norm is defined by

$$\|X\|_p := \left(\sum_{k \geq 1} \mu_k^\downarrow(X)^p \right)^{1/p}.$$

(When $0 < p < 1$, $\|\cdot\|_p$ is a quasi-norm.)

Building on the Euler operator identity (1.6), we obtain the following sharp Clarkson–McCarthy type inequalities.

Theorem 1.19. *Let $A, B, C \in S_p$. Then for $p \geq 2$,*

$$\|A + B\|_p^p + \|B + C\|_p^p + \|C + A\|_p^p \leq 2^{p-2} (\|A + B + C\|_p^p + \|A\|_p^p + \|B\|_p^p + \|C\|_p^p).$$

For $0 < p \leq 2$, the inequality is reversed. Moreover, the constant 2^{p-2} is optimal: equality holds for all $p > 0$ by taking $A = B = -C \neq 0$.

As a complement to Theorem 1.19, numerical evidence suggests the following sharp inequality. For a weaker bound with coefficient $3^{\frac{p}{2}-1}$, see Theorem 7.10.

Conjecture 1.20. *Let $A, B, C \in S_p$. Then for $p \geq 2$,*

$$\|A + B + C\|_p^p + \|A\|_p^p + \|B\|_p^p + \|C\|_p^p \leq \frac{3^{p-1} + 1}{2^p} (\|A + B\|_p^p + \|B + C\|_p^p + \|C + A\|_p^p).$$

For $0 < p \leq 2$, the inequality is reversed. Moreover, the constant $\frac{3^{p-1} + 1}{2^p}$ is optimal: equality holds for all $p > 0$ by taking $A = B = C \neq 0$.

Classical *Hlawka-type* inequalities compare norms (or their powers) of vector sums with those of their pairwise sums. A standard form of Hlawka’s inequality [17] asserts that if V is an inner product space and $x, y, z \in V$, then

$$\|x + y\| + \|y + z\| + \|z + x\| \leq \|x\| + \|y\| + \|z\| + \|x + y + z\|,$$

where $\|\cdot\|$ denotes the norm induced by the inner product; equivalently, the inequality holds for any norm satisfying the parallelogram identity. Numerous refinements and variants replace $\|\cdot\|$ by powers $\|\cdot\|^p$ and/or adjust the constants.

Theorem 1.19 and Conjecture 1.20 can be viewed as a Schatten– p (i.e. noncommutative L_p) analogue: it features a sharp p -dependent constant and exhibits a change in behavior at $p = 2$, as is typical for inequalities governed by the uniform convexity/concavity of L_p spaces.

As a further refinement of Theorem 1.19, we establish the following sharp mixed ℓ_q – ℓ_p Clarkson–McCarthy type inequalities.

Theorem 1.21. *Let $A, B, C \in S_p$. Then for $1 < p \leq 2$,*

$$\|A + B + C\|_p^q + \|A\|_p^q + \|B\|_p^q + \|C\|_p^q \leq 2^{1-\frac{q}{p}} \left(\|A + B\|_p^p + \|B + C\|_p^p + \|C + A\|_p^p \right)^{q/p},$$

$$\|A + B\|_p^q + \|B + C\|_p^q + \|C + A\|_p^q \leq 2^{1-\frac{q}{p}} \left(\|A + B + C\|_p^p + \|A\|_p^p + \|B\|_p^p + \|C\|_p^p \right)^{q/p},$$

where q is the conjugate exponent of p . For $p \geq 2$, these two inequalities are reversed. Moreover, the constant $2^{1-\frac{q}{p}}$ is optimal: equality holds for all $p > 1$ by taking $A = B = -C \neq 0$.

Remark 1.22. Setting $C = -B$ in Theorems 1.19 and 1.21 recovers the classical Clarkson–McCarthy inequality of [16]. For a complete characterization of the validity of Clarkson–McCarthy type inequalities, see the author’s paper [21, Corollary 1.14] and the following Theorem 1.23 (also see [21, Theorem 1.16]). For further results and developments on Clarkson–McCarthy type inequalities, we refer the reader to [5, 6, 9, 10, 15, 21, 22].

We now record the following mixed-norm inequality (Theorem 1.23), which strengthens the result in the author’s earlier work [21, Theorem 1.16]. Unlike the approach in [21], which applies the three-lines theorem to a suitable trace functional and then invokes a duality argument, we present a new proof based on a vector-valued Riesz–Thorin type complex interpolation scheme.

Theorem 1.23 (Mixed $\ell_q(S_p)$ – $\ell_p(S_p)$ estimates). *Let $U = [u_{ij}] \in \mathbb{M}_{s,t}$ and set*

$$\mu := \max_{1 \leq i \leq s, 1 \leq j \leq t} |u_{ij}|.$$

For $1 < p < \infty$, let q be the conjugate exponent of p , i.e. $\frac{1}{p} + \frac{1}{q} = 1$. Define the linear map on t -tuples

$$T_U : \ell_p^t(S_p) \longrightarrow \ell_q^s(S_p), \quad T_U(Z_1, \dots, Z_t) := \left(\sum_{j=1}^t u_{1j} Z_j, \dots, \sum_{j=1}^t u_{sj} Z_j \right).$$

We will regard T_U as an operator between ℓ_p - and ℓ_q -sums when equipped with the corresponding norms.

(i) Assume that U is a contraction, i.e. $\|U\|_\infty \leq 1$ (in particular, this holds if $U^*U = I_t$ or $UU^* = I_s$). Then for $1 < p \leq 2$,

$$\left(\sum_{i=1}^s \left\| \sum_{j=1}^t u_{ij} Z_j \right\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\sum_{j=1}^t \|Z_j\|_p^p \right)^{1/p}, \quad (1.8)$$

and the inequality is reversed for $p \geq 2$ if $U^*U = I_t$.

(ii) Assume in addition that $U^*U = I_t$ (i.e. U is an isometry). Then for $1 < p \leq 2$,

$$\left(\sum_{j=1}^t \|Z_j\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\sum_{i=1}^s \left\| \sum_{j=1}^t u_{ij} Z_j \right\|_p^p \right)^{1/p}, \quad (1.9)$$

and the inequality is reversed for $p \geq 2$.

Theorem 1.23 has several interesting consequences. For example, Audenaert–Kittaneh’s conjecture [1, Section 8.1, Conjecture 7] (proved by the author in [22]) states that if $A_1, \dots, A_n \in S_p$, then for $1 < p \leq 2$,

$$\left\| \sum_{i=1}^n A_i \right\|_p^q + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^q \leq n \left(\sum_{i=1}^n \|A_i\|_p^p \right)^{q/p}, \quad (1.10)$$

where q is the conjugate exponent of p ; for $p \geq 2$ the inequality is reversed. Moreover, the constant n is optimal: equality holds for all $1 < p < \infty$ by taking $A_1 = \dots = A_n \neq 0$. This inequality can be recovered from Theorem 1.23 by choosing the isometry

$$U = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & -1 & \cdots & 0 \\ 1 & 0 & \ddots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -1 \end{pmatrix} \in \mathbb{M}_{\frac{n(n-1)+2}{2}, n},$$

whose rows encode the combinations $\sum_{i=1}^n A_i$ and $A_i - A_j$. Moreover, Theorem 1.23 yields the following sharp reverse-type companion to (1.10). For more details, see Subsection 7.4.

Corollary 1.24. *Let $A_1, \dots, A_n \in S_p$. Then for $1 < p \leq 2$,*

$$\sum_{i=1}^n \|A_i\|_p^q \leq n^{-q/p} \left(\left\| \sum_{i=1}^n A_i \right\|_p^p + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^p \right)^{\frac{q}{p}}, \quad (1.11)$$

where q stands for the conjugate exponent of p . For $p \geq 2$, (1.11) is reversed. Moreover, the constant $n^{-q/p}$ is optimal: equality holds for all $1 < p < \infty$ by taking $A_1 = \dots = A_n \neq 0$.

As further consequences of Euler's identity (1.6), we next derive sharp Clarkson–McCarthy type inequalities for general unitarily invariant norms, together with a unitary-orbit refinement.

Theorem 1.25. *Let $A, B, C \in \mathbb{M}_n$. Then for $p \geq 2$,*

$$\| |A + B|^p + |B + C|^p + |C + A|^p \| \leq 2^{p-2} \| |A + B + C|^p + |A|^p + |B|^p + |C|^p \|.$$

For $0 < p \leq 2$, the inequality is reversed. Moreover, the constant 2^{p-2} is optimal: equality holds for all $p > 0$ by taking $A = B = -C \neq 0$.

Theorem 1.26. *Let $A, B, C \in \mathbb{M}_n$. Then for $p \geq 2$, there exist unitaries $U_1, U_2, U_3 \in \mathbb{M}_n$ such that*

$$U_1 |A + B|^p U_1^* + U_2 |B + C|^p U_2^* + U_3 |C + A|^p U_3^* \leq 2^{p-2} (|A + B + C|^p + |A|^p + |B|^p + |C|^p).$$

For $0 < p \leq 2$, the inequality is reversed. Moreover, the constant 2^{p-2} is optimal: equality holds for all $p > 0$ by taking $A = B = -C \neq 0$.

Organization of this paper. Section 2 gives a negative answer to the extension problem for the matrix-weighted parallelogram law: for each $x \notin [0, 1]$ we construct a counterexample, proving Theorem 1.3. In Section 3, we prove Theorem 1.6 and show that the range $p \leq 2$ in the Bourin–Lee unitary-orbit inequality (Theorem 1.4) is optimal by exhibiting an explicit 2×2 obstruction for every $p > 2$. Section 4 establishes Theorem 1.10 via a compact-operator construction that forces the failure of the corresponding singular-value inequality for all $p > 2$, and hence confirms the optimality of (1.3). In Section 5, we resolve the Thompson-type triangle problem for symmetric moduli (Theorem 1.12). Finally, Sections 6–8 develop isometry-orbit refinements of an Euler operator identity (Theorems 1.15, 1.16, 1.17), derive Clarkson–McCarthy type inequalities, and draw further consequences for unitarily invariant norms and singular values.

2. PROOF OF THEOREM 1.3

Proof of Theorem 1.3. Let $n = 1$ and take scalar matrices

$$A = 1, \quad B = \frac{x-1}{x} \quad (\text{note that } x \neq 0 \text{ since } x \notin [0, 1]).$$

Then

$$A\nabla_x B = (1-x)A + xB = (1-x) + x \cdot \frac{x-1}{x} = 0,$$

while

$$B\nabla_x A = (1-x)B + xA = (1-x)\frac{x-1}{x} + x = \frac{2x-1}{x} \neq 0,$$

because $x \notin [0, 1]$ implies $x \neq \frac{1}{2}$. Moreover,

$$A - B = 1 - \frac{x-1}{x} = \frac{1}{x} \neq 0.$$

Assume, for contradiction, that Theorem 1.1 holds for this x . Since $x \notin [0, 1]$, we have $x(1-x) < 0$. With $A\nabla_x B = 0$, the identity in Theorem 1.1 would reduce to

$$|A \oplus B|^2 = V |B\nabla_x A|^2 V^* + x(1-x) \left\{ S |A - B|^2 S^* + T |A - B|^2 T^* \right\},$$

for some isometries $V, S, T \in \mathbb{M}_{2,1}$. Moving the last term to the left and using $-x(1-x) > 0$, we obtain

$$|A \oplus B|^2 + (-x(1-x)) \left\{ S |A - B|^2 S^* + T |A - B|^2 T^* \right\} = V |B\nabla_x A|^2 V^*.$$

Here $|A \oplus B|^2 = \text{diag}(|A|^2, |B|^2) = \text{diag}(1, |(x-1)/x|^2)$ is positive definite, and the bracketed term is positive semidefinite; hence the whole left-hand side is positive definite, thus has rank 2.

On the other hand, since $V \in M_{2,1}$ is an isometry, VV^* is a rank-one projection; moreover $|B\nabla_x A|^2 > 0$. Therefore the right-hand side $V |B\nabla_x A|^2 V^*$ has rank 1.

This is impossible. Hence Theorem 1.1 cannot hold for this x . Since $x \in \mathbb{R} \setminus [0, 1]$ was arbitrary, the claim follows. \square

3. PROOF OF THEOREM 1.6

3.1. Trace necessary condition. Assume (1.2) holds for a given $Z \in \mathbb{M}_n$ and some unitaries U, V . Then, taking traces yields

$$\text{Tr} \left(\frac{|Z|^p + |Z^*|^p}{2} \right)^{1/p} \leq \text{Tr} |\text{Re } Z| + \text{Tr} |\text{Im } Z|. \quad (3.1)$$

Indeed, all terms in (1.2) are positive semidefinite, and for $0 \leq A \leq B$ we have $\text{Tr } A \leq \text{Tr } B$. Moreover, the right-hand side of (1.2) has trace independent of the choice of U, V by unitary invariance. Consequently, if (3.1) fails for some Z , then (1.2) fails for that Z for all choices of U, V .

Thus, to disprove (1.2) for $p > 2$, it suffices to find, for a given p , a matrix Z violating (3.1).

3.2. A 2×2 counterexample for $p > 2$. Fix $\theta \in (0, \pi/2)$ and set $c = \cos \theta$, $s = \sin \theta$. Define

$$Z_\theta = \begin{pmatrix} c & 0 \\ -s & 0 \end{pmatrix} \in \mathbb{M}_2. \quad (3.2)$$

Note that Z_θ is real, hence $Z_\theta^* = Z_\theta^T$.

Claim 3.1. *For every $p > 0$,*

$$\mathrm{Tr} \left(\frac{|Z_\theta|^p + |Z_\theta^*|^p}{2} \right)^{1/p} = \left(\frac{1+c}{2} \right)^{1/p} + \left(\frac{1-c}{2} \right)^{1/p}. \quad (3.3)$$

Proof. Compute

$$Z_\theta^* Z_\theta = \begin{pmatrix} c \\ -s \end{pmatrix} (c, -s) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} =: P.$$

Hence $|Z_\theta| = (Z_\theta^* Z_\theta)^{1/2} = P$. Similarly,

$$Z_\theta Z_\theta^* = \begin{pmatrix} c^2 & -cs \\ -cs & s^2 \end{pmatrix} =: Q.$$

A direct computation shows $Q^2 = Q$. Moreover $Q = Q^*$ (indeed Q is real symmetric), hence Q is an orthogonal projection. Therefore $|Z_\theta^*| = (Z_\theta Z_\theta^*)^{1/2} = Q$.

Since P and Q are projections, $P^p = P$ and $Q^p = Q$ for every $p > 0$. Thus

$$\left(\frac{|Z_\theta|^p + |Z_\theta^*|^p}{2} \right)^{1/p} = \left(\frac{P + Q}{2} \right)^{1/p}.$$

We compute the eigenvalues of $(P + Q)/2$. We have

$$P + Q = \begin{pmatrix} 1+c^2 & -cs \\ -cs & s^2 \end{pmatrix}, \quad \mathrm{Tr}(P + Q) = 2, \quad \det(P + Q) = s^2.$$

Hence the characteristic polynomial is $\lambda^2 - 2\lambda + s^2 = 0$, so the eigenvalues are $\lambda_\pm = 1 \pm \sqrt{1-s^2} = 1 \pm c$. Therefore the eigenvalues of $(P + Q)/2$ are $(1 \pm c)/2$, and those of $((P + Q)/2)^{1/p}$ are $((1 \pm c)/2)^{1/p}$. Taking the trace yields (3.3). \square

Claim 3.2. *For Z_θ in (3.2),*

$$\mathrm{Tr} |\mathrm{Re} Z_\theta| + \mathrm{Tr} |\mathrm{Im} Z_\theta| = 1 + s. \quad (3.4)$$

Proof. We compute

$$\mathrm{Re} Z_\theta = \frac{Z_\theta + Z_\theta^*}{2} = \begin{pmatrix} c & -s/2 \\ -s/2 & 0 \end{pmatrix}, \quad \mathrm{Im} Z_\theta = \frac{Z_\theta - Z_\theta^*}{2i} = \begin{pmatrix} 0 & -is/2 \\ is/2 & 0 \end{pmatrix}.$$

For $\mathrm{Re} Z_\theta$, the characteristic polynomial is

$$\lambda^2 - c\lambda - \frac{s^2}{4},$$

whose roots are

$$\lambda_\pm = \frac{c \pm \sqrt{c^2 + s^2}}{2} = \frac{c \pm 1}{2}.$$

Since $c \in (0, 1)$, we have $\lambda_+ > 0$ and $\lambda_- < 0$, hence

$$\mathrm{Tr} |\mathrm{Re} Z_\theta| = |\lambda_+| + |\lambda_-| = \frac{1+c}{2} + \frac{1-c}{2} = 1.$$

For $\mathrm{Im} Z_\theta$, the eigenvalues are $\pm s/2$, hence $\mathrm{Tr} |\mathrm{Im} Z_\theta| = s$. Adding yields (3.4). \square

Combining (3.1), Claims 3.1 and 3.2, we see that if (1.2) were to hold for Z_θ , then necessarily

$$\left(\frac{1+c}{2} \right)^{1/p} + \left(\frac{1-c}{2} \right)^{1/p} \leq 1 + s. \quad (3.5)$$

Claim 3.3. *Let $p > 2$. Then there exists $\theta_0 \in (0, \pi/2)$ such that for all $\theta \in (0, \theta_0)$ the inequality (3.5) fails.*

Proof. Define

$$\Phi(\theta) := \left(\frac{1 + \cos \theta}{2} \right)^{1/p} + \left(\frac{1 - \cos \theta}{2} \right)^{1/p} - (1 + \sin \theta).$$

As $\theta \rightarrow 0^+$ we have the standard expansions

$$\sin \theta = \theta + O(\theta^3), \quad \cos \theta = 1 - \frac{\theta^2}{2} + O(\theta^4).$$

Hence

$$\frac{1 + \cos \theta}{2} = 1 - \frac{\theta^2}{4} + O(\theta^4) \implies \left(\frac{1 + \cos \theta}{2} \right)^{1/p} = 1 + O(\theta^2),$$

and

$$\begin{aligned} \frac{1 - \cos \theta}{2} &= \frac{\theta^2}{4} + O(\theta^4) = \frac{\theta^2}{4}(1 + O(\theta^2)) \\ &\implies \left(\frac{1 - \cos \theta}{2} \right)^{1/p} = 4^{-1/p} \theta^{2/p} (1 + O(\theta^2)) = 4^{-1/p} \theta^{2/p} + O(\theta^{2/p+2}). \end{aligned}$$

Combining these estimates gives

$$\Phi(\theta) = 4^{-1/p} \theta^{2/p} - \theta + O(\theta^2) + O(\theta^{2/p+2}) \quad (\theta \rightarrow 0^+).$$

Dividing by θ yields

$$\frac{\Phi(\theta)}{\theta} = 4^{-1/p} \theta^{2/p-1} - 1 + O(\theta) + O(\theta^{2/p+1}) \xrightarrow[\theta \rightarrow 0^+]{} +\infty,$$

since $p > 2$ implies $2/p - 1 < 0$. Therefore $\Phi(\theta) > 0$ for all sufficiently small θ , i.e. (3.5) fails for all $\theta \in (0, \theta_0)$ for some $\theta_0 > 0$. \square

Proof of Theorem 1.6. Assume $p > 2$. Consider the 2×2 matrix Z_θ defined in (3.2). If (1.2) held for all $Z \in \mathbb{M}_2$, then it would hold for Z_θ and imply the trace inequality (3.1). Claims 3.1 and 3.2 reduce (3.1) to (3.5), which fails for all sufficiently small θ by Claim 3.3. Hence (1.2) cannot hold for all $Z \in \mathbb{M}_2$ when $p > 2$.

If $n > 2$, embed Z_θ as $Z_\theta \oplus 0_{n-2} \in \mathbb{M}_n$; the same trace obstruction applies, so (1.2) fails in \mathbb{M}_n as well. This contradicts the hypothesis, so necessarily $p \leq 2$. \square

4. PROOF OF THEOREM 1.10

Proof of Theorem 1.10. Let $H = \ell^2(\mathbb{N})$ with its standard orthonormal basis $(e_n)_{n \geq 1}$, and define a finite-rank operator $Z \in \mathbb{K}$ by

$$Ze_1 = e_2, \quad Ze_2 = e_3, \quad Ze_n = 0 \quad (n \geq 3).$$

Equivalently, Z is the unilateral shift truncated to the first three coordinates, that is,

$$Z = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \oplus 0 \oplus 0 \oplus \dots.$$

We first compute $|Z|$ and $|Z^*|$. Since

$$Z^*Z = \text{diag}(1, 1, 0, 0, 0, \dots), \quad ZZ^* = \text{diag}(0, 1, 1, 0, 0, \dots),$$

we get

$$|Z| = (Z^*Z)^{1/2} = \text{diag}(1, 1, 0, 0, 0, \dots), \quad |Z^*| = (ZZ^*)^{1/2} = \text{diag}(0, 1, 1, 0, 0, \dots).$$

Hence, for every $p > 0$,

$$\frac{|Z|^p + |Z^*|^p}{2} = \text{diag}\left(\frac{1}{2}, 1, \frac{1}{2}, 0, 0, \dots\right),$$

and therefore

$$\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p} = \text{diag}\left(2^{-1/p}, 1, 2^{-1/p}, 0, 0, \dots\right).$$

In particular, the singular values of this operator satisfy

$$\mu_1^\downarrow = 1, \quad \mu_2^\downarrow = 2^{-1/p}, \quad \mu_3^\downarrow = 2^{-1/p}, \quad \mu_m^\downarrow = 0 \ (m \geq 4).$$

Next, compute the real and imaginary parts:

$$\text{Re } Z = \frac{Z + Z^*}{2} = \begin{pmatrix} 0 & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \oplus 0 \oplus 0 \oplus \dots, \quad \text{Im } Z = \frac{Z - Z^*}{2i}.$$

A direct computation shows that $\text{Re } Z$ has eigenvalues 0 and $\pm 2^{-1/2}$, hence

$$\mu^\downarrow(\text{Re } Z) = \left(2^{-1/2}, 2^{-1/2}, 0, 0, 0, \dots\right).$$

Moreover, $\text{Im } Z$ is unitarily similar to $\text{Re } Z$. Indeed, for the 3×3 leading block set $D = \text{diag}(1, -i, -1)$ and extend it by $\oplus I$ on the remaining coordinates. Then one checks that

$$\text{Im } Z = D(\text{Re } Z)D^*.$$

Hence $\text{Re } Z$ and $\text{Im } Z$ have the same singular values.

$$\mu^\downarrow(\text{Im } Z) = \left(2^{-1/2}, 2^{-1/2}, 0, 0, 0, \dots\right).$$

Therefore,

$$\mu_1^\downarrow(\text{Im } Z) = 2^{-1/2}, \quad \mu_3^\downarrow(\text{Re } Z) = 0.$$

Choose $j = 2$ and $k = 0$, so that $1 + j + k = 3$, $1 + j = 3$, and $1 + k = 1$. Then for every $p > 2$,

$$\mu_{1+j+k}^\downarrow\left(\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}\right) = \mu_3^\downarrow\left(\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}\right) = 2^{-1/p},$$

while

$$\mu_{1+j}^\downarrow(\text{Re } Z) + \mu_{1+k}^\downarrow(\text{Im } Z) = \mu_3^\downarrow(\text{Re } Z) + \mu_1^\downarrow(\text{Im } Z) = 0 + 2^{-1/2} = 2^{-1/2}.$$

Since $p > 2$ implies $2^{-1/p} > 2^{-1/2}$, we conclude that

$$\mu_{1+j+k}^\downarrow\left(\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}\right) > \mu_{1+j}^\downarrow(\text{Re } Z) + \mu_{1+k}^\downarrow(\text{Im } Z).$$

□

Remark 4.1. The construction is intentionally minimal and answers the *existence* question in Question 1.9. Here Z has rank 2, and the operator $\left(\frac{|Z|^p + |Z^*|^p}{2}\right)^{1/p}$ is diagonal and easy to read off explicitly. The strict inequality is obtained by selecting indices $(j, k) = (2, 0)$ for which the corresponding singular value of $\text{Re } Z$ vanishes at level $1 + j = 3$, while $\text{Im } Z$ contributes only its top singular value $2^{-1/2}$, yet the third singular value on the left-hand side equals $2^{-1/p} > 2^{-1/2}$ whenever $p > 2$.

5. RECTANGULAR THOMPSON INEQUALITY AND PROOF OF THEOREM 1.12

In this section, we establish a rectangular version of Thompson's inequality, which will be the main tool in the proof of Theorem 1.12 (ii). The key idea is to reduce the rectangular setting to the square one by means of the polar decomposition: for a rectangular matrix $X \in \mathbb{M}_{m,n}$, one can write $X = W|X|$ with a partial isometry W whose initial projection coincides with the support projection of $|X|$. After recording this elementary fact (Lemma 5.1), we apply the classical (square) Thompson inequality to the compressed matrices W^*A and W^*B and then compare $|W^*A|, |W^*B|$ with $|A|, |B|$, respectively, to obtain the desired unitary-orbit domination. Finally, we use the rectangular Thompson inequality to settle Question 1.11, thereby proving Theorem 1.12.

Lemma 5.1. *Let $X \in \mathbb{M}_{m,n}$. Then there exists a partial isometry $W \in \mathbb{M}_{m,n}$ such that*

$$X = W|X|, \quad |X| = (X^*X)^{1/2},$$

and

$$W^*W = \text{supp}(|X|), \quad W^*W \leq I_n, WW^* \leq I_m, \quad (W^*W)|X| = |X|,$$

where $\text{supp}(\cdot)$ means the support projection.

Proof. Take a singular value decomposition $X = U\Sigma V^*$, where $U \in \mathbb{M}_m$ and $V \in \mathbb{M}_n$ are unitary and

$$\Sigma = \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} \in \mathbb{M}_{m,n}, \quad D = \text{diag}(s_1, \dots, s_r), \quad s_1 \geq \dots \geq s_r > 0, \quad r = \text{rank}(X).$$

Then

$$X^*X = V(\Sigma^*\Sigma)V^*, \quad \Sigma^*\Sigma = \begin{pmatrix} D^2 & 0 \\ 0 & 0 \end{pmatrix},$$

hence

$$|X| = (X^*X)^{1/2} = V \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} V^*.$$

Define $W := U \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} V^* \in \mathbb{M}_{m,n}$. Clearly, $W^*W \leq I_n, WW^* \leq I_m$ and

$$W|X| = U \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} V^* \cdot V \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} V^* = U \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} V^* = U\Sigma V^* = X.$$

Moreover,

$$(W^*W)|X| = P|X| = \left(V \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} V^* \right) \left(V \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} V^* \right) = V \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix} V^* = |X|.$$

That is, W^*W is exactly the support projection of $|X|$. □

Lemma 5.1 allows us to choose the partial isometry in the polar decomposition so that it acts as the identity on the range of $|X|$. This normalization is exactly what we need to turn $|A+B|$ into the absolute value of a *square* matrix, to which the classical Thompson inequality applies.

Lemma 5.2 (Rectangular Thompson inequality). *Let $A, B \in \mathbb{M}_{m,n}$. Then there exist unitary matrices $U, V \in \mathbb{M}_n$ such that*

$$|A+B| \leq U|A|U^* + V|B|V^*. \tag{5.1}$$

Proof. By using Lemma 5.1, there exists a partial isometry W such that $A + B = W|A + B|$, W^*W is the support projection of $|A + B|$ and $W^*W|A + B| = |A + B|$. Therefore

$$W^*(A + B) = W^*(W|A + B|) = |A + B|.$$

In particular,

$$|A + B| = |W^*(A + B)| = |W^*A + W^*B|. \quad (5.2)$$

Now W^*A and W^*B are $n \times n$ (square) matrices. Applying Thompson's inequality (1.1) (the square case) to W^*A and W^*B , there exist unitaries $U, V \in \mathbb{M}_n$ such that

$$|W^*A + W^*B| \leq U|W^*A|U^* + V|W^*B|V^*. \quad (5.3)$$

Since the left-hand side equals $|A + B|$, it remains to compare $|W^*A|$ with $|A|$.

Since $WW^* \leq I_m$, we have

$$|W^*A|^2 = (W^*A)^*(W^*A) = A^*WW^*A \leq A^*A = |A|^2.$$

By the operator monotonicity of $t \mapsto t^{1/2}$ on $[0, \infty)$, this implies $|W^*A| \leq |A|$. Similarly, $|W^*B| \leq |B|$. Conjugating preserves the order, hence

$$U|W^*A|U^* \leq U|A|U^*, \quad V|W^*B|V^* \leq V|B|V^*.$$

Substituting into (5.3) and combining with (5.2) yields (5.1). \square

With Lemma 5.2 in hand, we now prove Theorem 1.12. Part (i) is verified by a concrete counterexample for the symmetric modulus, while part (ii) follows from an application of Lemma 5.2 to a suitable linear embedding that converts the quadratic symmetrization into a rectangular modulus.

Proof of Theorem 1.12. (i) Take

$$X = \begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \in M_2(\mathbb{R}).$$

A direct computation gives

$$|X|_{\text{sym}} = \frac{1}{2\sqrt{5}} \begin{pmatrix} 5 & 2 \\ 2 & 5 \end{pmatrix}, \quad |Y|_{\text{sym}} = \frac{1}{2} I_2, \quad |X + Y|_{\text{sym}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}.$$

Hence

$$\| |X + Y|_{\text{sym}} \|_{\infty} = \frac{3}{\sqrt{2}}, \quad \| |X|_{\text{sym}} \|_{\infty} = \frac{7}{2\sqrt{5}}, \quad \| |Y|_{\text{sym}} \|_{\infty} = \frac{1}{2},$$

so

$$\| |X + Y|_{\text{sym}} \|_{\infty} > \| |X|_{\text{sym}} \|_{\infty} + \| |Y|_{\text{sym}} \|_{\infty} \quad \left(\text{i.e. } \frac{3}{\sqrt{2}} > \frac{7}{2\sqrt{5}} + \frac{1}{2} \right).$$

If the inequality in Question 1.11 held for some unitaries U, V , then taking operator norms would yield

$$\| |X + Y|_{\text{sym}} \|_{\infty} \leq \| U|X|_{\text{sym}}U^* + V|Y|_{\text{sym}}V^* \|_{\infty} \leq \| |X|_{\text{sym}} \|_{\infty} + \| |Y|_{\text{sym}} \|_{\infty},$$

a contradiction. This proves (i).

(ii) Define the linear map $\Gamma : \mathbb{M}_n \rightarrow \mathbb{M}_{2n,n}$ by

$$\Gamma(Z) := \begin{pmatrix} Z \\ Z^* \end{pmatrix}.$$

Then

$$\Gamma(Z)^* \Gamma(Z) = Z^*Z + ZZ^*, \quad \text{hence} \quad |\Gamma(Z)| = (Z^*Z + ZZ^*)^{1/2} = \sqrt{2}|Z|_{\text{qsym}}.$$

Moreover, $\Gamma(X+Y) = \Gamma(X) + \Gamma(Y)$. Applying rectangular Thompson inequality (Lemma 5.2) yields unitaries $U, V \in \mathbb{M}_n$ such that

$$|\Gamma(X+Y)| \leq U|\Gamma(X)|U^* + V|\Gamma(Y)|V^*.$$

Dividing by $\sqrt{2}$ and using $|\Gamma(Z)| = \sqrt{2}|Z|_{\text{qsym}}$ gives

$$|X+Y|_{\text{qsym}} \leq U|X|_{\text{qsym}}U^* + V|Y|_{\text{qsym}}V^*,$$

as desired. \square

6. EULER-TYPE IDENTITIES AND ISOMETRY ORBITS

In this section, we develop an operator-valued Euler identity and its consequences in the language of isometry orbits. Our goal is twofold. First, we give a proof of Theorem 1.15 by encoding the Euler-type relation into a unitary conjugation of suitable block matrices and then decomposing the resulting positive operator into isometric compressions of its blocks. Second, we collect a convenient isometry decomposition principle for positive block matrices, which will be used later to derive orbit-dominance statements from positivity and symmetry.

We start with a Pythagoras-type theorem for partitioned matrices due to Bourin and Lee [11], which allows us to express the square modulus of a block matrix as a sum of isometric conjugations of the square moduli of its blocks.

Lemma 6.1 ([11, Theorem 2.1]). *Let $m, n \geq 1$ and let $T = [T_{ij}]_{i,j=1}^m$ be an $m \times m$ block matrix with $T_{ij} \in \mathbb{M}_n$. Then there exist isometries $W_{ij} \in \mathbb{M}_{mn,n}$ such that*

$$|T|^2 = \sum_{i=1}^m \sum_{j=1}^m W_{ij} |T_{ij}|^2 W_{ij}^*.$$

Remark 6.2. The standard $m \times m$ grid partitioning is row/column compatible in the sense of [11], so the above lemma applies directly to block matrices $T = [T_{ij}]$ written with respect to such a partition.

We now apply Lemma 6.1 to a block matrix obtained from a Hadamard conjugation. This provides an isometric-orbit refinement of the Euler operator identity and yields Theorem 1.15.

Proof of Theorem 1.15. Let

$$X := \begin{pmatrix} A+B+C \\ A \\ B \\ C \end{pmatrix}, \quad Y := \begin{pmatrix} A+B \\ B+C \\ C+A \\ 0 \end{pmatrix} \in \mathbb{M}_{4n,n}.$$

Consider the 4×4 Hadamard matrix

$$H = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ -1 & 1 & 1 & -1 \end{pmatrix}, \quad H^*H = I_4,$$

and set $\mathcal{U} := H \otimes I_n \in \mathbb{M}_{4n}$. A direct block computation gives $X = \mathcal{U}Y$. Hence $X^*X = Y^*Y$, i.e. (1.6).

Let

$$\Delta_Y := \begin{pmatrix} A+B & 0 & 0 & 0 \\ 0 & B+C & 0 & 0 \\ 0 & 0 & C+A & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \in \mathbb{M}_{4n}.$$

Set

$$T := \mathcal{U} \Delta_Y \mathcal{U}^* = \frac{1}{2} \begin{pmatrix} A+B+C & A & B & C \\ A & A+B+C & -C & -B \\ B & -C & A+B+C & -A \\ C & -B & -A & A+B+C \end{pmatrix}.$$

Since \mathcal{U} is unitary,

$$|T|^2 = |\mathcal{U} \Delta_Y \mathcal{U}^*|^2 = \mathcal{U} |\Delta_Y|^2 \mathcal{U}^* = \mathcal{U} \operatorname{diag}(|A+B|^2, |B+C|^2, |C+A|^2, 0) \mathcal{U}^*.$$

Apply Lemma 6.1 with $m = 4$ to the 4×4 block matrix $T = [T_{ij}]$. We obtain isometries $W_{ij} \in \mathbb{M}_{4n,n}$ such that

$$|T|^2 = \sum_{i=1}^4 \sum_{j=1}^4 W_{ij} |T_{ij}|^2 W_{ij}^*.$$

Every block T_{ij} equals $(A+B+C)/2, \pm A/2, \pm B/2$ or $\pm C/2$. Hence

$$|T_{ij}|^2 \in \left\{ \frac{1}{4}|A+B+C|^2, \frac{1}{4}|A|^2, \frac{1}{4}|B|^2, \frac{1}{4}|C|^2 \right\}.$$

Moreover, each of $(A+B+C)/2, A/2, B/2, C/2$ appears exactly four times among the sixteen blocks. Grouping the corresponding four terms and renaming the isometries, we obtain isometries $V_{ij} \in \mathbb{M}_{4n,n}$ such that

$$|T|^2 = \frac{1}{4} \sum_{j=1}^4 V_{1j} |A+B+C|^2 V_{1j}^* + \frac{1}{4} \sum_{j=1}^4 V_{2j} |A|^2 V_{2j}^* + \frac{1}{4} \sum_{j=1}^4 V_{3j} |B|^2 V_{3j}^* + \frac{1}{4} \sum_{j=1}^4 V_{4j} |C|^2 V_{4j}^*.$$

Conjugating by \mathcal{U}^* yields

$$|\Delta_Y|^2 = \frac{1}{4} \sum_{j=1}^4 (\mathcal{U}^* V_{1j}) |A+B+C|^2 (\mathcal{U}^* V_{1j})^* + \cdots + \frac{1}{4} \sum_{j=1}^4 (\mathcal{U}^* V_{4j}) |C|^2 (\mathcal{U}^* V_{4j})^*.$$

Finally, each $\mathcal{U}^* V_{ij}$ is still an isometry in $\mathbb{M}_{4n,n}$ since

$$(\mathcal{U}^* V_{ij})^* (\mathcal{U}^* V_{ij}) = V_{ij}^* \mathcal{U} \mathcal{U}^* V_{ij} = V_{ij}^* V_{ij} = I_n.$$

Renaming $U_{ij} := \mathcal{U}^* V_{ij}$ gives (1.7). \square

Next, we record a useful lemma of Bourin and Lee [9], which decomposes a 2×2 positive block matrix into a sum of two unitary orbits of its diagonal compressions. This lemma will serve as a building block for a higher-dimensional isometry decomposition.

Lemma 6.3 ([9, Lemma 3.4]). *Let $\begin{pmatrix} X & Y \\ Y^* & Z \end{pmatrix} \in \mathbb{M}_{2n}$ be positive semidefinite, written in $n \times n$ blocks. Then there exist unitary matrices $U, V_0 \in \mathbb{M}_{2n}$ such that*

$$\begin{pmatrix} X & Y \\ Y^* & Z \end{pmatrix} = U \begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix} U^* + V_0 \begin{pmatrix} 0 & 0 \\ 0 & Z \end{pmatrix} V_0^*.$$

As an immediate consequence of Lemma 6.3, one obtains an isometry decomposition for positive block matrices with an arbitrary number of blocks. For the reader's convenience, we include a self-contained proof.

Lemma 6.4. *Let $m \geq 2$ and let $H = [H_{ij}]_{i,j=1}^m \in \mathbb{M}_{mn}$ be positive semidefinite, written in $n \times n$ blocks. Then there exist isometries $V_1, \dots, V_m \in \mathbb{M}_{mn,n}$ (i.e., $V_k^* V_k = I_n$) such that*

$$H = \sum_{k=1}^m V_k H_{kk} V_k^*. \tag{6.1}$$

Proof. Since $H \geq 0$, let $R := H^{1/2}$, so $R = R^*$ and $H = R^2 = RR^* = R^*R$. Partition R into block columns $R = [R_1 \cdots R_m]$. Then

$$H = RR^* = \sum_{k=1}^m R_k R_k^*. \quad (6.2)$$

Step 1: Identify the diagonal blocks. By block multiplication, the (k, k) diagonal block of $H = R^*R$ is

$$H_{kk} = R_k^* R_k \in \mathbb{M}_n^+.$$

In particular, $|R_k| := (R_k^* R_k)^{1/2} = H_{kk}^{1/2}$.

Step 2: Polar decomposition and a partial isometry. Take the polar decomposition of each R_k :

$$R_k = U_k |R_k| = U_k H_{kk}^{1/2},$$

where $U_k \in \mathbb{M}_{mn,n}$ is a partial isometry satisfying

$$U_k^* U_k = P_k, \quad P_k := \text{supp}(|R_k|) = \text{supp}(H_{kk})$$

(the orthogonal projection onto $\overline{\text{Ran}}(|R_k|)$). Note that $H_{kk}^{1/2}(I_n - P_k) = 0$.

Step 3: Extend U_k to a genuine isometry V_k . If $P_k = I_n$, then $U_k^* U_k = I_n$ and U_k is already an isometry; set $V_k := U_k$.

Assume now $P_k \neq I_n$. Set $d_k := \text{rank}(P_k) \leq n$. Then $\dim \text{Ran}(I_n - P_k) = n - d_k$. Also, since U_k is a partial isometry with initial projection P_k , its range projection $Q_k := U_k U_k^*$ has rank d_k , hence

$$\dim \text{Ran}(I_{mn} - Q_k) = mn - d_k \geq n - d_k,$$

using $m \geq 2 \Rightarrow mn \geq n$.

Therefore, there exists an isometry

$$W_k : \text{Ran}(I_n - P_k) \longrightarrow \text{Ran}(I_{mn} - Q_k),$$

i.e. a matrix $W_k \in \mathbb{M}_{mn,n}$ such that

$$W_k^* W_k = I_n - P_k, \quad W_k W_k^* \leq I_{mn} - Q_k.$$

(For example, choose orthonormal bases of the two subspaces and map one to the other.)

Now define

$$V_k := U_k + W_k \in \mathbb{M}_{mn,n}.$$

Then we claim the cross terms vanish. First, since $Q_k := U_k U_k^*$ is the range projection of U_k , we have $Q_k U_k = U_k$, hence $U_k^*(I_{mn} - Q_k) = 0$. On the other hand, $W_k W_k^* \leq I_{mn} - Q_k$ implies $\text{Ran}(W_k) \subseteq \text{Ran}(I_{mn} - Q_k)$, equivalently $(I_{mn} - Q_k)W_k = W_k$. Therefore

$$U_k^* W_k = U_k^*(I_{mn} - Q_k)W_k = 0.$$

Similarly, U_k has initial projection $P_k = U_k^* U_k$, so $U_k P_k = U_k$ and hence $(I_n - P_k)U_k^* = 0$. Since $W_k^* W_k = I_n - P_k$, we have $\text{Ran}(W_k^*) \subseteq \text{Ran}(I_n - P_k)$, equivalently $W_k^* = W_k^*(I_n - P_k)$. Thus

$$W_k^* U_k = W_k^*(I_n - P_k)U_k = 0.$$

Consequently,

$$V_k^* V_k = (U_k + W_k)^*(U_k + W_k) = U_k^* U_k + W_k^* W_k = P_k + (I_n - P_k) = I_n,$$

so V_k is an isometry in $\mathbb{M}_{mn,n}$.

Moreover, since $H_{kk}^{1/2}(I_n - P_k) = 0$, we still have

$$V_k H_{kk}^{1/2} = (U_k + W_k) H_{kk}^{1/2} = U_k H_{kk}^{1/2} = R_k.$$

Step 4: Conclude the decomposition. From $R_k = V_k H_{kk}^{1/2}$ we deduce

$$R_k R_k^* = V_k H_{kk}^{1/2} H_{kk}^{1/2} V_k^* = V_k H_{kk} V_k^*.$$

Summing over k and using (6.2) gives

$$H = \sum_{k=1}^m R_k R_k^* = \sum_{k=1}^m V_k H_{kk} V_k^*,$$

which is exactly (6.1). \square

We are now ready to prove Theorem 1.16 and Theorem 1.17. In both cases, we conjugate an appropriate direct sum by a Fourier matrix $F_m \otimes I_n$ to obtain a positive block matrix with constant diagonal blocks, and then apply Lemma 6.4 to conclude the desired isometry-orbit domination.

Proof of Theorem 1.16. First, we compute Fourier conjugations of two direct sums. Let $\omega = e^{2\pi i/3}$ and set

$$W_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} I & I & I \\ I & \omega I & \omega^2 I \\ I & \omega^2 I & \omega I \end{pmatrix} = F_3 \otimes I_n \in \mathbb{M}_{3n}.$$

For

$$D := |A + B|^2 \oplus |B + C|^2 \oplus |A + C|^2 = \text{diag}(D_1, D_2, D_3), \quad \begin{cases} D_1 = |A + B|^2, \\ D_2 = |B + C|^2, \\ D_3 = |A + C|^2, \end{cases}$$

one has

$$W_3 D W_3^* = \frac{1}{3} \begin{pmatrix} D_1 + D_2 + D_3 & D_1 + \omega^2 D_2 + \omega D_3 & D_1 + \omega D_2 + \omega^2 D_3 \\ D_1 + \omega D_2 + \omega^2 D_3 & D_1 + D_2 + D_3 & D_1 + \omega^2 D_2 + \omega D_3 \\ D_1 + \omega^2 D_2 + \omega D_3 & D_1 + \omega D_2 + \omega^2 D_3 & D_1 + D_2 + D_3 \end{pmatrix}. \quad (6.3)$$

In particular, every diagonal block of $W_3 D W_3^*$ equals $(D_1 + D_2 + D_3)/3$.

Clearly, $W_3 D W_3^* \geq 0$ and, by (6.3), each diagonal block equals $(D_1 + D_2 + D_3)/3$. A direct expansion shows

$$D_1 + D_2 + D_3 = |A + B|^2 + |B + C|^2 + |A + C|^2 = |A + B + C|^2 + |A|^2 + |B|^2 + |C|^2.$$

Denote this common sum by T . Hence $W_3 D W_3^*$ is a 3×3 block positive matrix with constant diagonal block $T/3$. Applying (6.1) to $H = W_3 D W_3^*$ yields isometries $V_1, V_2, V_3 \in \mathbb{M}_{3n,n}$ such that

$$W_3 D W_3^* = \sum_{k=1}^3 V_k (T/3) V_k^*.$$

Conjugating by W_3^* and setting $U_k = W_3^* V_k$ (still isometries) gives the claim. \square

Proof of Theorem 1.17. First, let $\eta = e^{2\pi i/4} = i$ and set

$$W_4 = \frac{1}{2} \begin{pmatrix} I & I & I & I \\ I & \eta I & \eta^2 I & \eta^3 I \\ I & \eta^2 I & \eta^4 I & \eta^6 I \\ I & \eta^3 I & \eta^6 I & \eta^9 I \end{pmatrix} = F_4 \otimes I_n \in \mathbb{M}_{4n},$$

so that $\eta^2 = -1$, $\eta^3 = -i$, $\eta^4 = 1$. For

$$E := |A + B + C|^2 \oplus |A|^2 \oplus |B|^2 \oplus |C|^2 = \text{diag}(E_1, E_2, E_3, E_4),$$

with $E_1 = |A + B + C|^2$, $E_2 = |A|^2$, $E_3 = |B|^2$, $E_4 = |C|^2$, we obtain

$$W_4 E W_4^* = \frac{1}{4} [M_{rs}]_{r,s=1}^4, \quad M_{rs} = E_1 + \eta^{-(r-s)} E_2 + \eta^{-2(r-s)} E_3 + \eta^{-3(r-s)} E_4. \quad (6.4)$$

In particular, every diagonal block of $W_4 E W_4^*$ equals $(E_1 + E_2 + E_3 + E_4)/4$.

Clearly, $W_4 E W_4^* \geq 0$ and, by (6.4), each diagonal block equals $(E_1 + E_2 + E_3 + E_4)/4$. As in the previous proof,

$$E_1 + E_2 + E_3 + E_4 = |A + B + C|^2 + |A|^2 + |B|^2 + |C|^2 = |A + B|^2 + |B + C|^2 + |A + C|^2.$$

Denote this sum by S . Thus $W_4 E W_4^*$ is a 4×4 block positive matrix with constant diagonal block $S/4$. Applying (6.1) to $H = W_4 E W_4^*$ yields isometries $V_1, \dots, V_4 \in \mathbb{M}_{4n,n}$ such that

$$W_4 E W_4^* = \sum_{k=1}^4 V_k (S/4) V_k^*.$$

Conjugating by W_4^* and setting $U_k = W_4^* V_k$ completes the proof. \square

7. SHARP CLARKSON–MCCARTHY TYPE INEQUALITIES ON EULER’S IDENTITY

In this section, we exploit Euler’s operator identity (1.6) and the associated isometry-unitary-orbit refinements obtained in Section 6 to derive sharp Clarkson–McCarthy type inequalities. We treat two parallel frameworks: first, inequalities for general unitarily invariant norms; second, stronger unitary-orbit dominations, from which Schatten p -norm inequalities follow by taking traces. Finally, we introduce a vector-valued complex interpolation scheme and deduce a mixed $\ell_q(S_p)$ – $\ell_p(S_p)$ estimate, which will be used to prove the mixed-exponent Euler-type inequalities later in the paper.

7.1. Unitarily invariant norm case. We begin with a Jensen-type principle for unitarily invariant norms, which allows us to convert the square-sum identity (1.6) into p -power estimates via the function $t \mapsto t^{p/2}$.

We will use the following Jensen-type inequalities for unitarily invariant norms.

Lemma 7.1 ([14, Lemma 2.1]). *Let $m \geq 1$. Let $A_1, \dots, A_m \in \mathbb{M}_n$ be positive semidefinite and let $\alpha_1, \dots, \alpha_m$ be positive real numbers such that $\sum_{j=1}^m \alpha_j = 1$. Then*

(1) *for every convex function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$,*

$$\begin{aligned} \left\| f\left(\sum_{j=1}^m \alpha_j A_j\right) \right\| &\leq \left\| \sum_{j=1}^m \alpha_j f(A_j) \right\|, \\ \left\| \sum_{j=1}^m f(A_j) \right\| &\leq \left\| f\left(\sum_{j=1}^m A_j\right) \right\|; \end{aligned}$$

(2) *for every concave function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$,*

$$\begin{aligned} \left\| \sum_{j=1}^m \alpha_j f(A_j) \right\| &\leq \left\| f\left(\sum_{j=1}^m \alpha_j A_j\right) \right\|, \\ \left\| f\left(\sum_{j=1}^m A_j\right) \right\| &\leq \left\| \sum_{j=1}^m f(A_j) \right\|. \end{aligned}$$

Proof of Theorem 1.25. Set $r := p/2$ and $f(t) := t^r$ on $[0, \infty)$. Note that $f(0) = 0$.

Define the positive semidefinite matrices

$$X_1 := |A + B|^2, \quad X_2 := |B + C|^2, \quad X_3 := |C + A|^2,$$

and

$$Y_1 := |A + B + C|^2, \quad Y_2 := |A|^2, \quad Y_3 := |B|^2, \quad Y_4 := |C|^2.$$

By Euler's identity (1.6),

$$X_1 + X_2 + X_3 = Y_1 + Y_2 + Y_3 + Y_4 =: S.$$

Moreover, $|A + B|^p = X_1^r = f(X_1)$, etc., and $|A|^p = Y_2^r = f(Y_2)$, etc.

Step 1: case 1: $p \geq 2$. That is, $r \geq 1$. Then f is convex.

Applying Lemma 7.1(1) *the second inequality* to X_1, X_2, X_3 yields

$$\|f(X_1) + f(X_2) + f(X_3)\| \leq \|f(X_1 + X_2 + X_3)\| = \|f(S)\|.$$

Next, applying Lemma 7.1(1) *the first inequality* with $n = 4$ and $\alpha_j = 1/4$ to Y_1, Y_2, Y_3, Y_4 gives

$$\left\| f\left(\frac{1}{4}S\right) \right\| \leq \left\| \frac{1}{4}(f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)) \right\|.$$

Using $f(tX) = t^r f(X)$ for $t > 0$ and the absolute homogeneity of $\|\cdot\|$, we obtain

$$\frac{1}{4^r} \|f(S)\| \leq \frac{1}{4} \|f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)\|.$$

Hence

$$\|f(S)\| \leq 4^{r-1} \|f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)\| = 2^{p-2} \||A + B + C|^p + |A|^p + |B|^p + |C|^p\|.$$

Combining the above inequalities yields the desired estimate for $p \geq 2$.

Step 2: $0 < p \leq 2$. That is, $0 < r \leq 1$. Then f is concave.

Applying Lemma 7.1(2) *the second inequality* to X_1, X_2, X_3 yields

$$\|f(S)\| = \|f(X_1 + X_2 + X_3)\| \leq \|f(X_1) + f(X_2) + f(X_3)\|.$$

Next, applying Lemma 7.1(2) *the first inequality* with $n = 4$ and $\alpha_j = 1/4$ to Y_1, Y_2, Y_3, Y_4 gives

$$\left\| \frac{1}{4}(f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)) \right\| \leq \left\| f\left(\frac{1}{4}S\right) \right\|.$$

Using homogeneity again,

$$\frac{1}{4} \|f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)\| \leq \frac{1}{4^r} \|f(S)\|,$$

and therefore

$$2^{p-2} \||A + B + C|^p + |A|^p + |B|^p + |C|^p\| = 4^{r-1} \|f(Y_1) + f(Y_2) + f(Y_3) + f(Y_4)\| \leq \|f(S)\|.$$

Combining with $\|f(S)\| \leq \|f(X_1) + f(X_2) + f(X_3)\|$ yields the reversed inequality for $0 < p \leq 2$.

Step 3: optimality. Taking $A = B = -C \neq 0$, we have $A + B = 2A$, $B + C = 0$, $C + A = 0$, and $A + B + C = A$. Thus

$$\||A + B|^p + |B + C|^p + |C + A|^p\| = \||2A|^p\| = 2^p \||A|^p\|,$$

while

$$2^{p-2} \||A + B + C|^p + |A|^p + |B|^p + |C|^p\| = 2^{p-2} \||4A|^p\| = 2^p \||A|^p\|.$$

Hence equality holds for all $p > 0$, and the constant 2^{p-2} is best possible. \square

7.2. Unitary orbit case. We next turn to a unitary-orbit version of Jensen's inequality for monotone convex/concave functions. These orbit inequalities yield matrix-order dominations, and hence deduce Schatten p -norm consequences.

The following unitary-orbit inequalities on monotone concave/convex functions will be used in the sequel. Lemma 7.2 (1) *the first inequality* is obtained by a straightforward induction on n from the corresponding two-variable inequality [9, Corollary 3.2]. Lemma 7.2 (2) *the first inequality* follows similarly from [9, Theorem 3.1] (equivalently, [2, Theorem 2.1]). Lemma 7.2 (1) *the second inequality* follows from the isometric-column Jensen inequality [9, Corollary 2.4] by choosing $Z_i = \sqrt{\alpha_i} I$, so that $\sum_{i=1}^n Z_i^* Z_i = I$ and $\sum_{i=1}^n Z_i^* A_i Z_i = \sum_{i=1}^n \alpha_i A_i$. Similarly, Lemma 7.2 (2) *the second inequality* follows from the same result, using the fact that the inequality in [9, Corollary 2.4] reverses for concave functions.

Lemma 7.2. *Let $A_1, \dots, A_n \in \mathbb{M}_n$ be positive semidefinite and let $\alpha_1, \dots, \alpha_n$ be positive real numbers such that $\sum_{j=1}^n \alpha_j = 1$. Then*

(1) *for every monotone convex function $f : [0, \infty) \rightarrow \mathbb{R}$ with $f(0) \leq 0$, there exist unitaries $U, U_1, \dots, U_n \in \mathbb{M}_n$ such that*

$$\begin{aligned} f\left(\sum_{j=1}^n A_j\right) &\geq \sum_{j=1}^n U_j f(A_j) U_j^*, \\ f\left(\sum_{j=1}^n \alpha_j A_j\right) &\leq U \left(\sum_{j=1}^n \alpha_j f(A_j)\right) U^*; \end{aligned}$$

(2) *for every monotone concave function $f : [0, \infty) \rightarrow \mathbb{R}$ with $f(0) \geq 0$, there exist unitaries $U, U_1, \dots, U_n \in \mathbb{M}_n$ such that*

$$\begin{aligned} f\left(\sum_{j=1}^n A_j\right) &\leq \sum_{j=1}^n U_j f(A_j) U_j^*, \\ f\left(\sum_{j=1}^n \alpha_j A_j\right) &\geq U \left(\sum_{j=1}^n \alpha_j f(A_j)\right) U^*. \end{aligned}$$

We remark that Theorems 1.19, 1.26, and 7.10 can in fact be derived directly from the results of the author in [21, Theorem 1.12 and Corollary 1.14]. For the reader's convenience, and to keep the present paper self-contained, we nevertheless include complete proofs.

Proof of Theorem 1.26. Let $r := p/2$ and define $f(t) := t^r$ on $[0, \infty)$. Note that f is monotone. If $p \geq 2$, then $r \geq 1$ and f is convex with $f(0) = 0 \leq 0$. If $0 < p \leq 2$, then $0 < r \leq 1$ and f is concave with $f(0) = 0 \geq 0$.

Set

$$X_1 := |A + B|^2, \quad X_2 := |B + C|^2, \quad X_3 := |C + A|^2,$$

and

$$Y_1 := |A + B + C|^2, \quad Y_2 := |A|^2, \quad Y_3 := |B|^2, \quad Y_4 := |C|^2.$$

By Euler's identity (1.6),

$$S := X_1 + X_2 + X_3 = Y_1 + Y_2 + Y_3 + Y_4.$$

Moreover, $f(X_1) = |A + B|^p$, $f(X_2) = |B + C|^p$, $f(X_3) = |C + A|^p$, and $f(Y_1) = |A + B + C|^p$, $f(Y_2) = |A|^p$, $f(Y_3) = |B|^p$, $f(Y_4) = |C|^p$.

Step 1: case $p \geq 2$. Apply Lemma 7.2(1) *the first inequality* to X_1, X_2, X_3 . Then there exist unitaries $W_1, W_2, W_3 \in \mathbb{M}_n$ such that

$$\sum_{j=1}^3 W_j f(X_j) W_j^* \leq f(S). \quad (7.1)$$

Next apply Lemma 7.2(1) *the second inequality* to Y_1, Y_2, Y_3, Y_4 with weights $\alpha_j = 1/4$. Then there exists a unitary $U \in \mathbb{M}_n$ such that

$$f\left(\frac{1}{4}S\right) \leq U \left(\frac{1}{4} \sum_{k=1}^4 f(Y_k)\right) U^*.$$

Using $f(tX) = t^r f(X)$ for $t > 0$, we obtain

$$f(S) = 4^r f\left(\frac{1}{4}S\right) \leq 4^{r-1} U \left(\sum_{k=1}^4 f(Y_k)\right) U^*.$$

Conjugating by U^* yields

$$U^* f(S) U \leq 4^{r-1} \sum_{k=1}^4 f(Y_k). \quad (7.2)$$

Now conjugate (7.1) by U^* and set $U_j := U^* W_j$ to get

$$\sum_{j=1}^3 U_j f(X_j) U_j^* \leq U^* f(S) U.$$

Combining with (7.2) gives

$$\sum_{j=1}^3 U_j f(X_j) U_j^* \leq 4^{r-1} \sum_{k=1}^4 f(Y_k) = 2^{p-2}(|A+B+C|^p + |A|^p + |B|^p + |C|^p),$$

i.e.

$$U_1|A+B|^p U_1^* + U_2|B+C|^p U_2^* + U_3|C+A|^p U_3^* \leq 2^{p-2}(|A+B+C|^p + |A|^p + |B|^p + |C|^p).$$

Step 2: case $0 < p \leq 2$. Now apply Lemma 7.2(2) *the first inequality* to X_1, X_2, X_3 to get

$$f(S) \leq \sum_{j=1}^3 W_j f(X_j) W_j^*$$

for some unitaries W_j . Also apply Lemma 7.2(2) *the second inequality* to Y_1, \dots, Y_4 with $\alpha_k = 1/4$ to get a unitary U such that

$$f(S) = 4^r f\left(\frac{1}{4}S\right) \geq 4^{r-1} U \left(\sum_{k=1}^4 f(Y_k)\right) U^*.$$

Conjugating and combining exactly as above (setting $U_j := U^* W_j$) yields the reversed inequality

$$\sum_{j=1}^3 U_j f(X_j) U_j^* \geq 4^{r-1} \sum_{k=1}^4 f(Y_k) = 2^{p-2}(|A+B+C|^p + |A|^p + |B|^p + |C|^p).$$

Step 3: optimality. Take $A = B = -C \neq 0$. Then $A+B = 2A$, $B+C = 0$, $C+A = 0$, and $A+B+C = A$. Hence the inequality becomes an equality for all $p > 0$, and the constant $2^{p-2} = 4^{r-1}$ is best possible. \square

Taking traces in the unitary-orbit inequality (or, equivalently, taking the trace norm in the unitarily invariant norm statement) immediately yields the Schatten p -norm formulation.

Proof of Theorem 1.19. Taking trace norms in Theorem 1.25, or simply taking traces in Theorem 1.26, and then passing to general Schatten class operators by approximating with finite-rank operators, yields Theorem 1.19. \square

7.3. Complex interpolation theory. We now introduce the tools from complex interpolation needed in this paper; for the basic theory of interpolation, we refer the interested reader to [3, Chapters 1–5].

Definition 7.3. Let A_0 and A_1 be two topological vector spaces. We say that A_0 and A_1 are *compatible* if there exists a Hausdorff topological vector space \mathcal{Z} such that A_0 and A_1 are subspaces of \mathcal{Z} . In this case, we can form their sum $A_0 + A_1$ and their intersection $A_0 \cap A_1$.

Definition 7.4. Let X and Y be normed spaces, and let $T : X \rightarrow Y$ be a bounded linear operator. The *operator norm* of T is defined by

$$\|T\|_{X \rightarrow Y} := \sup_{\substack{x \in X \\ x \neq 0}} \frac{\|Tx\|_Y}{\|x\|_X}.$$

Definition 7.5. Let (X_0, X_1) be a compatible Banach couple and $0 < \theta < 1$. The space $(X_0, X_1)_{[\theta]}$ is defined as

$$(X_0, X_1)_{[\theta]} := \{f(\theta) : f \in \mathcal{F}(X_0, X_1)\},$$

where $\mathcal{F}(X_0, X_1)$ denotes the space of all bounded continuous functions $f : \{z \in \mathbb{C} : 0 \leq \Re z \leq 1\} \rightarrow X_0 + X_1$ which are analytic on $\{z \in \mathbb{C} : 0 < \Re z < 1\}$, satisfy $f(it) \in X_0$ for all $t \in \mathbb{R}$ and $f(1+it) \in X_1$ for all $t \in \mathbb{R}$, and are bounded on the boundary lines in the respective norms of X_0 and X_1 . The norm on $(X_0, X_1)_{[\theta]}$ is given by

$$\|x\|_{(X_0, X_1)_{[\theta]}} := \inf \{\|f\|_{\mathcal{F}(X_0, X_1)} : f(\theta) = x\}.$$

We recall the abstract complex interpolation theorem for Banach couples; see [3, Theorem 4.1.2].

Theorem 7.6 ([3]). *Let (X_0, X_1) and (Y_0, Y_1) be two compatible couples of Banach spaces, and let $0 < \theta < 1$. Suppose that*

$$T : X_0 + X_1 \longrightarrow Y_0 + Y_1$$

is a linear operator such that $T : X_j \rightarrow Y_j$ is bounded for $j = 0, 1$. Then T is bounded from $(X_0, X_1)_{[\theta]}$ to $(Y_0, Y_1)_{[\theta]}$, and

$$\|T\|_{(X_0, X_1)_{[\theta]} \rightarrow (Y_0, Y_1)_{[\theta]}} \leq \|T\|_{X_0 \rightarrow Y_0}^{1-\theta} \|T\|_{X_1 \rightarrow Y_1}^{\theta}.$$

We will also use the classical Riesz–Thorin interpolation theorem. For L^p spaces, see [3, Theorem 5.1.1]. For certain noncommutative L^p spaces, such as the Schatten p -classes S_p , a Riesz–Thorin interpolation theorem is also known to hold; see [18, Eq. (2.1)], since matrix algebras are semifinite von Neumann algebras. See also [13, Theorem 13.1], [19, Theorems 2.9–2.10], and [19, Remark 1, p. 23].

Theorem 7.7 ([18]). *Assume that $1 \leq p_0, p_1 \leq \infty$ and $0 < \theta < 1$. Then*

$$(S_{p_0}, S_{p_1})_{[\theta]} = S_p \quad (\text{with equal norms}),$$

if

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

We shall also use the complex interpolation formula for ℓ_p -sums; see [3, Theorem 5.6.3, p. 123].¹

Theorem 7.8 ([3]). *Let (X_0, X_1) be a compatible couple of Banach spaces. Assume that $1 \leq p_0, p_1 \leq \infty$ and $0 < \theta < 1$. For each $m \in \mathbb{N}$ and $k = 0, 1$, define $\ell_{p_k}^m(X_k)$ as the space of m -tuples $a = (a_1, \dots, a_m)$ with $a_j \in X_k$, equipped with the norm*

$$\|a\|_{\ell_{p_k}^m(X_k)} = \begin{cases} \left(\sum_{j=1}^m \|a_j\|_{X_k}^{p_k} \right)^{1/p_k}, & p_k < \infty, \\ \max_{1 \leq j \leq m} \|a_j\|_{X_k}, & p_k = \infty. \end{cases}$$

Then

$$(\ell_{p_0}^m(X_0), \ell_{p_1}^m(X_1))_{[\theta]} = \ell_p^m((X_0, X_1)_{[\theta]}) \quad (\text{with equal norms}),$$

if

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

Combining Theorem 7.7 and Theorem 7.8 gives the following.

Theorem 7.9. *Assume that $1 \leq p_0, p_1, q_0, q_1 \leq \infty$ and $0 < \theta < 1$. Then for each $m \in \mathbb{N}$,*

$$(\ell_{p_0}^m(S_{q_0}), \ell_{p_1}^m(S_{q_1}))_{[\theta]} = \ell_p^m(S_q) \quad (\text{with equal norms}),$$

if

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1} \quad \text{and} \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}.$$

With the interpolation tools in hand, we proceed to prove Theorem 1.23. We first obtain (1.8) for $1 < p \leq 2$ via complex interpolation between the endpoints $(p, q) = (1, \infty)$ and $(2, 2)$, and then deduce (1.9) from (1.8) applied to U^* .

Proof of Theorem 1.23. We first prove (1.8) for $1 < p \leq 2$ by complex interpolation between the endpoints $(p, q) = (1, \infty)$ and $(p, q) = (2, 2)$, and then deduce (1.9) from (1.8) applied to U^* .

Step 1: endpoint 1: $\ell_1^t(S_1) \rightarrow \ell_\infty^s(S_1)$. For each i ,

$$\left\| \sum_{j=1}^t u_{ij} Z_j \right\|_1 \leq \sum_{j=1}^t |u_{ij}| \|Z_j\|_1 \leq \mu \sum_{j=1}^t \|Z_j\|_1.$$

Taking the maximum over i yields

$$\|T_U(Z)\|_{\ell_\infty^s(S_1)} \leq \mu \|Z\|_{\ell_1^t(S_1)},$$

hence

$$\|T_U\|_{\ell_1^t(S_1) \rightarrow \ell_\infty^s(S_1)} \leq \mu. \quad (7.3)$$

¹Following [3, p. 121, Section 5.6], let A be a Banach space, let $s \in \mathbb{R}$ and $q > 0$. Denote by $\dot{\ell}_q^s(A)$ the space of all bi-infinite sequences $a = (a_\nu)_{\nu \in \mathbb{Z}}$ with $a_\nu \in A$ such that

$$\|a\|_{\dot{\ell}_q^s(A)} := \left(\sum_{\nu \in \mathbb{Z}} (2^{\nu s} \|a_\nu\|_A)^q \right)^{1/q} < \infty.$$

In particular, taking $s = 0$ and fixing $m \in \mathbb{N}$, we may identify an m -tuple (a_1, \dots, a_m) with an element of this sequence space by extending it to a bi-infinite sequence $a = (a_\nu)_{\nu \in \mathbb{Z}}$ via $a_\nu = 0$ for all $\nu \notin \{1, \dots, m\}$; this agrees with our definition in Theorem 7.8.

Step 2: endpoint 2: $\ell_2^t(S_2) \rightarrow \ell_2^s(S_2)$. Equip S_2 with the Hilbert–Schmidt inner product $\langle X, Y \rangle = \text{Tr}(Y^* X)$, so that $\|X\|_2^2 = \langle X, X \rangle$. A direct computation gives

$$\|T_U(Z)\|_{\ell_2^s(S_2)}^2 = \sum_{j,k=1}^t (U^* U)_{jk} \langle Z_k, Z_j \rangle = \text{Tr}(U^* U G), \quad G = [\langle Z_k, Z_j \rangle]_{j,k=1}^t \geq 0.$$

If $\|U\|_\infty \leq 1$, then $U^* U \leq I_t$, and therefore

$$\|T_U(Z)\|_{\ell_2^s(S_2)}^2 \leq \sum_{j=1}^t \|Z_j\|_2^2 = \|Z\|_{\ell_2^t(S_2)}^2,$$

so that

$$\|T_U\|_{\ell_2^t(S_2) \rightarrow \ell_2^s(S_2)} \leq 1. \quad (7.4)$$

Step 3: interpolation. Fix $0 < \theta < 1$. Applying Theorem 7.6 to the couples

$$X_0 = \ell_1^t(S_1), \quad X_1 = \ell_2^t(S_2), \quad Y_0 = \ell_\infty^s(S_1), \quad Y_1 = \ell_2^s(S_2),$$

and using (7.3)–(7.4), we obtain

$$\|T_U\|_{(X_0, X_1)_{[\theta]} \rightarrow (Y_0, Y_1)_{[\theta]}} \leq \mu^{1-\theta}. \quad (7.5)$$

By Theorem 7.9,

$$(X_0, X_1)_{[\theta]} = \ell_p^t(S_p), \quad (Y_0, Y_1)_{[\theta]} = \ell_q^s(S_p),$$

where

$$\frac{1}{p} = 1 - \frac{\theta}{2}, \quad \frac{1}{q} = \frac{\theta}{2},$$

so that q is the conjugate exponent of p and $1 < p \leq 2$. Thus (7.5) yields

$$\|T_U(Z)\|_{\ell_q^s(S_p)} \leq \mu^{1-\theta} \|Z\|_{\ell_p^t(S_p)} = \mu^{\frac{2}{p}-1} \|Z\|_{\ell_p^t(S_p)},$$

which is exactly (1.8).

Step 4: the case $p \geq 2$ (reverse inequality in (1.8)). Assume in addition that $U^* U = I_t$ (as required in the statement for the reverse direction). Let q be the conjugate exponent of p , so that $1 < q \leq 2$.

Let $Z = (Z_1, \dots, Z_t) \in \ell_p^t(S_p)$. By duality,

$$\|Z\|_{\ell_p^t(S_p)} = \sup \left\{ |\langle Z, W \rangle| : W \in \ell_q^t(S_q), \|W\|_{\ell_q^t(S_q)} \leq 1 \right\}, \quad \langle Z, W \rangle := \sum_{j=1}^t \text{Tr}(W_j^* Z_j).$$

For such a W , since $U^* U = I_t$ we have $T_{U^*} \circ T_U = \text{Id}$, hence $Z = T_{U^*}(T_U(Z))$ and therefore

$$\langle Z, W \rangle = \langle T_{U^*}(T_U(Z)), W \rangle = \langle T_U(Z), T_U(W) \rangle.$$

By Hölder's inequality for Schatten classes and for ℓ_p -sums,

$$|\langle T_U(Z), T_U(W) \rangle| \leq \|T_U(Z)\|_{\ell_q^s(S_p)} \|T_U(W)\|_{\ell_p^s(S_q)}.$$

Since $1 < q \leq 2$ and $\|U\|_\infty \leq 1$, we may apply the already proved estimate (1.8) (with q in place of p and p in place of q) to obtain

$$\|T_U(W)\|_{\ell_p^s(S_q)} \leq \mu^{\frac{2}{q}-1} \|W\|_{\ell_q^t(S_q)} = \mu^{1-\frac{2}{p}} \|W\|_{\ell_q^t(S_q)}.$$

Hence, for $\|W\|_{\ell_q^t(S_q)} \leq 1$,

$$|\langle Z, W \rangle| \leq \mu^{1-\frac{2}{p}} \|T_U(Z)\|_{\ell_q^s(S_p)}.$$

Taking the supremum over all such W yields

$$\|Z\|_{\ell_p^t(S_p)} \leq \mu^{1-\frac{2}{p}} \|T_U(Z)\|_{\ell_q^s(S_p)},$$

equivalently,

$$\|T_U(Z)\|_{\ell_q^s(S_p)} \geq \mu^{\frac{2}{p}-1} \|Z\|_{\ell_p^t(S_p)}.$$

This is exactly the reversed inequality in (1.8) for $p \geq 2$.

Step 5: the reverse-direction estimate. Assume now that $U^*U = I_t$. Consider the adjoint matrix $U^* \in \mathbb{M}_{t,s}$ and the associated map

$$T_{U^*} : \ell_p^s(S_p) \longrightarrow \ell_p^t(S_p), \quad T_{U^*}(W_1, \dots, W_s) := \left(\sum_{i=1}^s \overline{w_{i1}} W_i, \dots, \sum_{i=1}^s \overline{w_{it}} W_i \right).$$

Then $T_{U^*} \circ T_U = \text{Id}$ on $\ell_p^t(S_p)$. Applying (1.8) to U^* (note that $\max_{i,j} |(U^*)_{ij}| = \mu$) gives, for $1 < p \leq 2$,

$$\|T_{U^*}(W)\|_{\ell_q^s(S_p)} \leq \mu^{\frac{2}{p}-1} \|W\|_{\ell_p^s(S_p)} \quad (W \in \ell_p^s(S_p)).$$

Taking $W = T_U(Z)$ and using $T_{U^*}(T_U(Z)) = Z$ yields (1.9). The case $p \geq 2$ again follows by duality. \square

Proof of Theorem 1.21. Fix $1 < p \leq 2$ and let q be the conjugate exponent of p , i.e. $1/p + 1/q = 1$. Set

$$X_1 := A+B, \quad X_2 := B+C, \quad X_3 := C+A, \quad Y_1 := A+B+C, \quad Y_2 := A, \quad Y_3 := B, \quad Y_4 := C.$$

Step 1: A suborthogonal coefficient matrix with U^ isometric.* Consider the real 3×4 matrix

$$U := \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \end{pmatrix}.$$

Then

$$\begin{pmatrix} X_1 \\ X_2 \\ X_3 \end{pmatrix} = U \begin{pmatrix} Y_1 \\ Y_2 \\ Y_3 \\ Y_4 \end{pmatrix}, \quad \text{i.e.} \quad X_i = \sum_{j=1}^4 u_{ij} Y_j \quad (i = 1, 2, 3).$$

Moreover, a direct computation gives $UU^* = I_3$; hence U^* is an isometry. Also $\max_{i,j} |u_{ij}| = 1/2 =: \mu$.

*Step 2: The relations $X = UY$ and $Y = U^*X$.* By construction,

$$\begin{pmatrix} X_1 \\ X_2 \\ X_3 \end{pmatrix} = U \begin{pmatrix} Y_1 \\ Y_2 \\ Y_3 \\ Y_4 \end{pmatrix}.$$

Moreover, a direct computation gives $Y = U^*X$ and explicitly

$$Y_1 = \frac{X_1 + X_2 + X_3}{2}, \quad Y_2 = \frac{X_1 - X_2 + X_3}{2}, \quad Y_3 = \frac{X_1 + X_2 - X_3}{2}, \quad Y_4 = \frac{-X_1 + X_2 + X_3}{2},$$

which are easily verified from $X_1 = A + B$, $X_2 = B + C$, $X_3 = C + A$.

Step 3: Apply Theorem 1.23. Since $\|U\|_\infty \leq 1$, Theorem 1.23 (i) gives for $1 < p \leq 2$,

$$\left(\|X_1\|_p^q + \|X_2\|_p^q + \|X_3\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\|Y_1\|_p^p + \|Y_2\|_p^p + \|Y_3\|_p^p + \|Y_4\|_p^p \right)^{1/p}.$$

Raising both sides to the power q and using $\mu = 1/2$ yields

$$\|X_1\|_p^q + \|X_2\|_p^q + \|X_3\|_p^q \leq 2^{1-\frac{q}{p}} \left(\|Y_1\|_p^p + \|Y_2\|_p^p + \|Y_3\|_p^p + \|Y_4\|_p^p \right)^{q/p},$$

which is the second inequality.

Applying the same estimate to U^* and using $Y = U^*X$ gives

$$\left(\|Y_1\|_p^q + \|Y_2\|_p^q + \|Y_3\|_p^q + \|Y_4\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\|X_1\|_p^p + \|X_2\|_p^p + \|X_3\|_p^p \right)^{1/p},$$

and raising to q again yields the first inequality with the same constant $2^{1-\frac{q}{p}}$.

Step 4: optimality. Taking $A = B = -C \neq 0$ gives $A + B = 2A$ and $B + C = C + A = 0$, while $A + B + C = A$. Thus equality holds for all $p > 1$, and the constant $2^{1-\frac{q}{p}}$ is optimal. \square

7.4. Derivation of Audenaert–Kittaneh’s inequality and its sharp complement. In this subsection, we show that the Audenaert–Kittaneh inequality (1.10) and the complementary estimate (1.11) both follow from Theorem 1.23 by a suitable choice of the coefficient matrix U .

Let $n \geq 2$ and set

$$s := 1 + \binom{n}{2}.$$

Fix an enumeration of the pairs (i, j) with $1 \leq i < j \leq n$:

$$(i_1, j_1), (i_2, j_2), \dots, (i_{\binom{n}{2}}, j_{\binom{n}{2}}).$$

Define

$$U = \frac{1}{\sqrt{n}} \begin{pmatrix} \mathbf{1}^\top \\ (e_{i_1}^\top - e_{j_1}^\top) \\ (e_{i_2}^\top - e_{j_2}^\top) \\ \vdots \\ (e_{i_{\binom{n}{2}}}^\top - e_{j_{\binom{n}{2}}}^\top) \end{pmatrix} \in \mathbb{M}_{s,n},$$

where $\mathbf{1} = (1, \dots, 1)^\top \in \mathbb{C}^n$ and e_k denotes the k th standard basis vector in \mathbb{C}^n . Then

$$\mu := \max_{\alpha, j} |u_{\alpha j}| = \frac{1}{\sqrt{n}}.$$

We claim U is an isometry, i.e. $U^*U = I_n$ (hence $\|U\|_\infty = 1$ and U is a contraction). Indeed,

$$U^*U = \frac{1}{n} \left(\mathbf{1}\mathbf{1}^* + \sum_{1 \leq i < j \leq n} (e_i - e_j)(e_i - e_j)^* \right),$$

and one checks the identity

$$\sum_{1 \leq i < j \leq n} (e_i - e_j)(e_i - e_j)^* = nI_n - \mathbf{1}\mathbf{1}^*.$$

Therefore $U^*U = \frac{1}{n}(\mathbf{1}\mathbf{1}^* + nI_n - \mathbf{1}\mathbf{1}^*) = I_n$.

Given $A_1, \dots, A_n \in S_p$, consider the n -tuple $Z := (A_1, \dots, A_n) \in \ell_p^n(S_p)$. By construction,

$$T_U(Z) = \left(\frac{1}{\sqrt{n}} \sum_{k=1}^n A_k, \frac{1}{\sqrt{n}}(A_{i_1} - A_{j_1}), \dots, \frac{1}{\sqrt{n}}(A_{i_{\binom{n}{2}}} - A_{j_{\binom{n}{2}}}) \right) \in \ell_q^s(S_p).$$

Consequently, for $r \in \{p, q\}$ we have

$$\sum_{\alpha=1}^s \left\| \sum_{j=1}^n u_{\alpha j} A_j \right\|_p^r = \frac{1}{n^{r/2}} \left(\left\| \sum_{k=1}^n A_k \right\|_p^r + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^r \right). \quad (7.6)$$

In particular, (7.6) with $r = q$ gives the left side of (1.8), and with $r = p$ gives the right side of (1.9).

Step 1: Audenaert–Kittaneh inequality from Theorem 1.23(i). Let $1 < p \leq 2$ and let q be the conjugate exponent of p . Since U is a contraction, we may apply (1.8) to $Z = (A_1, \dots, A_n)$:

$$\left(\sum_{\alpha=1}^s \left\| \sum_{j=1}^n u_{\alpha j} A_j \right\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\sum_{j=1}^n \|A_j\|_p^p \right)^{1/p}.$$

Using (7.6) with $r = q$ and $\mu = 1/\sqrt{n}$, we obtain

$$\frac{1}{\sqrt{n}} \left(\left\| \sum_{k=1}^n A_k \right\|_p^q + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^q \right)^{1/q} \leq \left(\frac{1}{\sqrt{n}} \right)^{\frac{2}{p}-1} \left(\sum_{j=1}^n \|A_j\|_p^p \right)^{1/p}.$$

Multiplying by \sqrt{n} and noting that

$$\sqrt{n} \left(\frac{1}{\sqrt{n}} \right)^{\frac{2}{p}-1} = n^{1-\frac{1}{p}}, \quad \text{and} \quad q \left(1 - \frac{1}{p} \right) = 1,$$

we get

$$\left\| \sum_{k=1}^n A_k \right\|_p^q + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^q \leq n \left(\sum_{j=1}^n \|A_j\|_p^p \right)^{q/p},$$

which is exactly (1.10). For $p \geq 2$, the inequality is reversed by Theorem 1.23(i) since $U^*U = I_n$.

Step 2: the complementary estimate from Theorem 1.23(ii). Since $U^*U = I_n$, we may apply (1.9) to the same U and Z . For $1 < p \leq 2$,

$$\left(\sum_{j=1}^n \|A_j\|_p^q \right)^{1/q} \leq \mu^{\frac{2}{p}-1} \left(\sum_{\alpha=1}^s \left\| \sum_{j=1}^n u_{\alpha j} A_j \right\|_p^p \right)^{1/p}.$$

Using (7.6) with $r = p$ and $\mu = 1/\sqrt{n}$ yields

$$\left(\sum_{j=1}^n \|A_j\|_p^q \right)^{1/q} \leq \left(\frac{1}{\sqrt{n}} \right)^{\frac{2}{p}-1} \cdot \left(\frac{1}{\sqrt{n}} \right) \left(\left\| \sum_{k=1}^n A_k \right\|_p^p + \sum_{1 \leq i < j \leq n} \|A_i - A_j\|_p^p \right)^{1/p}.$$

Since $\left(\frac{1}{\sqrt{n}} \right)^{\frac{2}{p}-1} \cdot \frac{1}{\sqrt{n}} = n^{-1/p}$, raising both sides to the power q gives (1.11). For $p \geq 2$, the inequality is reversed by Theorem 1.23(ii).

Step 3: optimality of the constant in (1.11). Let $A_1 = \dots = A_n = A \neq 0$. Then $\sum_{k=1}^n A_k = nA$ and $A_i - A_j = 0$, hence

$$\text{LHS of (1.11)} = n\|A\|_p^q, \quad \text{RHS of (1.11)} = n^{-q/p} \|nA\|_p^q = n^{-q/p} \cdot n^q \|A\|_p^q = n\|A\|_p^q.$$

Thus equality holds for all $1 < p < \infty$, and the constant $n^{-q/p}$ is best possible.

7.5. A weaker bound on Conjecture 1.20. We record a non-sharp estimate with the explicit constant $3^{\frac{p}{2}-1}$, which follows directly from the Euler-type identities; however, this coefficient is not optimal.

Theorem 7.10. *Let $A, B, C \in S_p$. Then for $p \geq 2$,*

$$\|A + B + C\|_p^p + \|A\|_p^p + \|B\|_p^p + \|C\|_p^p \leq 3^{\frac{p}{2}-1} (\|A + B\|_p^p + \|B + C\|_p^p + \|C + A\|_p^p).$$

For $0 < p \leq 2$, the inequality is reversed.

Proof. Set $r := p/2$ and $f(t) := t^r$ on $[0, \infty)$. Let

$$X_1 := |A + B|^2, \quad X_2 := |B + C|^2, \quad X_3 := |C + A|^2,$$

and

$$Y_1 := |A + B + C|^2, \quad Y_2 := |A|^2, \quad Y_3 := |B|^2, \quad Y_4 := |C|^2.$$

By Euler's identity (1.6),

$$S := X_1 + X_2 + X_3 = Y_1 + Y_2 + Y_3 + Y_4.$$

Assume first that $p \geq 2$, i.e. $r \geq 1$, so that f is convex and $f(0) = 0$. Using the trace form of the Jensen/Rotfel'd type inequality for convex f (equivalently, Lemma 7.1 (1) with the trace norm), we have

$$\mathrm{Tr}\left(\sum_{k=1}^4 f(Y_k)\right) \leq \mathrm{Tr}(f(S)). \quad (7.7)$$

Next apply Jensen's inequality with weights $\alpha_j = 1/3$ to X_1, X_2, X_3 :

$$\mathrm{Tr}\left(f\left(\frac{1}{3}S\right)\right) \leq \mathrm{Tr}\left(\frac{1}{3} \sum_{j=1}^3 f(X_j)\right).$$

Since $f(tX) = t^r f(X)$ for $t > 0$, this becomes

$$3^{-r} \mathrm{Tr}(f(S)) \leq \frac{1}{3} \mathrm{Tr}\left(\sum_{j=1}^3 f(X_j)\right),$$

or equivalently

$$\mathrm{Tr}(f(S)) \leq 3^{r-1} \mathrm{Tr}\left(\sum_{j=1}^3 f(X_j)\right). \quad (7.8)$$

Combining (7.7)–(7.8) gives

$$\mathrm{Tr}\left(\sum_{k=1}^4 |Y_k|^r\right) = \mathrm{Tr}\left(\sum_{k=1}^4 f(Y_k)\right) \leq 3^{r-1} \mathrm{Tr}\left(\sum_{j=1}^3 f(X_j)\right) = 3^{r-1} \mathrm{Tr}\left(\sum_{j=1}^3 |X_j|^r\right).$$

Finally, passing to general Schatten class operators by approximating with finite-rank operators, we obtain exactly the desired inequality with constant $3^{r-1} = 3^{\frac{p}{2}-1}$.

For $0 < p \leq 2$ (i.e. $0 < r \leq 1$), the function f is concave and all inequalities above reverse, yielding the reversed statement. \square

Remark 7.11. The constant $3^{\frac{p}{2}-1}$ in Theorem 7.10 is generally not optimal; numerical evidence suggests that the sharp constant should be the one proposed in Conjecture 1.20.

8. UNITARY-ORBIT REFINEMENTS OF EULER'S IDENTITY AND SPECTRAL CONSEQUENCES

8.1. A unitary-orbit bound for the Euler modulus. To convert the square-sum identity (1.6) into a linear (modulus) estimate in unitary orbits, we use the following unitary subadditivity result for concave functions due to Aujla and Bourin [2, Theorem 2.1], which is also included in Lemma 7.2 (2) *the first inequality*.

Lemma 8.1 (Aujla–Bourin). *Let $H, K \in \mathbb{M}_n$ be positive semidefinite and let $f : [0, \infty) \rightarrow [0, \infty)$ be monotone concave with $f(0) \geq 0$. Then there exist unitaries $U, V \in \mathbb{M}_n$ such that*

$$f(H + K) \leq Uf(H)U^* + Vf(K)V^*.$$

We now apply Lemma 8.1 with the concave function $f(t) = t^{1/2}$ to the Euler square-sum relation (1.6). This yields a three-term unitary-orbit domination of the Euler modulus by the pairwise moduli.

Theorem 8.2. *Let $A, B, C \in \mathbb{M}_n$. Then there exist unitaries $U, V, W \in \mathbb{M}_n$ such that*

$$\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} \leq U|A + B|U^* + V|B + C|V^* + W|C + A|W^*. \quad (8.1)$$

Proof. By (1.6),

$$\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} = \sqrt{|A + B|^2 + |B + C|^2 + |C + A|^2}.$$

Apply Lemma 8.1 with $f(t) = \sqrt{t}$ to $H = |A + B|^2$ and $K = |B + C|^2 + |C + A|^2$ to get unitaries U_1, V_1 with

$$\sqrt{H + K} \leq U_1|A + B|U_1^* + V_1\sqrt{K}V_1^*.$$

Apply Lemma 8.1 again to $|B + C|^2$ and $|C + A|^2$ to obtain unitaries U_2, V_2 such that

$$\sqrt{K} \leq U_2|B + C|U_2^* + V_2|C + A|V_2^*.$$

Substitute and absorb conjugation by V_1 into U_2, V_2 to obtain (8.1). \square

As a first immediate consequence, taking any unitarily invariant norm in (8.1) and using the triangle inequality yields a norm inequality.

Corollary 8.3. *Let $A, B, C \in \mathbb{M}_n$. Then*

$$\left\| \sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} \right\| \leq \|A + B\| + \|B + C\| + \|C + A\|.$$

We next record a singular-value refinement, obtained by combining Theorem 8.2 with the Weyl inequality for sums of positive semidefinite matrices.

Corollary 8.4. *Let $A, B, C \in \mathbb{M}_n$. Then for all integers $j, k, \ell \in \{0, 1, \dots, n - 1\}$,*

$$\mu_{1+j+k+\ell}^{\downarrow} \left(\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} \right) \leq \mu_{1+j}^{\downarrow}(A + B) + \mu_{1+k}^{\downarrow}(B + C) + \mu_{1+\ell}^{\downarrow}(C + A).$$

Proof. By Theorem 8.2, there exist unitaries $U, V, W \in \mathbb{M}_n$ such that

$$\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} \leq U|A + B|U^* + V|B + C|V^* + W|C + A|W^*.$$

Set $X := U|A + B|U^*$, $Y := V|B + C|V^*$ and $Z := W|C + A|W^*$. Then $X, Y, Z \geq 0$ and $\mu_j^{\downarrow}(X) = \mu_j^{\downarrow}(|A + B|) = \mu_j^{\downarrow}(A + B)$, and similarly for Y, Z .

We use the following singular-value inequality (see, e.g., [4]): for $P, Q \geq 0$ and $r, s \geq 0$,

$$\mu_{1+r+s}^{\downarrow}(P + Q) \leq \mu_{1+r}^{\downarrow}(P) + \mu_{1+s}^{\downarrow}(Q). \quad (8.2)$$

Applying (8.2) to $P = X$ and $Q = Y + Z$ gives

$$\mu_{1+j+k+\ell}^{\downarrow}(X + Y + Z) \leq \mu_{1+j}^{\downarrow}(X) + \mu_{1+k+\ell}^{\downarrow}(Y + Z).$$

Next apply (8.2) to $P = Y$ and $Q = Z$ to obtain

$$\mu_{1+k+\ell}^\downarrow(Y + Z) \leq \mu_{1+k}^\downarrow(Y) + \mu_{1+\ell}^\downarrow(Z).$$

Combining the last two inequalities yields

$$\mu_{1+j+k+\ell}^\downarrow(X + Y + Z) \leq \mu_{1+j}^\downarrow(X) + \mu_{1+k}^\downarrow(Y) + \mu_{1+\ell}^\downarrow(Z).$$

Finally, since

$$\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2} \leq X + Y + Z$$

and both sides are positive semidefinite, the monotonicity of singular values gives

$$\mu_{1+j+k+\ell}^\downarrow\left(\sqrt{|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2}\right) \leq \mu_{1+j+k+\ell}^\downarrow(X + Y + Z),$$

which implies the desired estimate. \square

8.2. Symmetric norms, antinorms, and Ky Fan consequences. Finally, since Euler's identity (1.6) is an exact equality of positive operators, it also yields further consequences for symmetric norms and antinorms (in the sense of [8]) on the positive semidefinite cone of \mathbb{M}_n , and in particular for Ky Fan sums of singular values.

Corollary 8.5. *Let $A, B, C \in \mathbb{M}_n$.*

(a) *For every symmetric (unitarily invariant) norm $\|\cdot\|$,*

$$\|\|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2\| \leq \||A + B|^2\| + \||B + C|^2\| + \||C + A|^2\|.$$

(b) *For every symmetric antinorm $\|\cdot\|_!$,*

$$\left\| |A + B + C|^2 + |A|^2 + |B|^2 + |C|^2 \right\|_! \geq \||A + B|^2\|_! + \||B + C|^2\|_! + \||C + A|^2\|_!.$$

Applying Corollary 8.5 to Ky Fan m -norms and Ky Fan m -antinorms gives the following eigenvalue-sum inequalities.

Corollary 8.6. *Let $A, B, C \in \mathbb{M}_n$. Then for $m = 1, \dots, n$,*

$$\begin{aligned} \sum_{j=1}^m \mu_j^\downarrow(|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2) &\leq \sum_{j=1}^m \mu_j^\downarrow(|A + B|^2) + \sum_{j=1}^m \mu_j^\downarrow(|B + C|^2) + \sum_{j=1}^m \mu_j^\downarrow(|C + A|^2), \\ \sum_{j=1}^m \mu_j^\uparrow(|A + B + C|^2 + |A|^2 + |B|^2 + |C|^2) &\geq \sum_{j=1}^m \mu_j^\uparrow(|A + B|^2) + \sum_{j=1}^m \mu_j^\uparrow(|B + C|^2) + \sum_{j=1}^m \mu_j^\uparrow(|C + A|^2). \end{aligned}$$

ACKNOWLEDGMENTS

The author is very grateful to Professor J.-C. Bourin for reading this manuscript and for his valuable comments. This work is supported by the China Scholarship Council, the Young Elite Scientists Sponsorship Program for PhD Students (China Association for Science and Technology), and the Fundamental Research Funds for the Central Universities at Xi'an Jiaotong University (Grant No. xzy022024045).

REFERENCES

- [1] K. M. R. Audenaert and F. Kittaneh, Problems and conjectures in matrix and operator inequalities, in *Études opératorielles*, Banach Center Publ. **112** (Polish Acad. Sci. Inst. Math., Warsaw, 2017), 15–31. doi:10.4064/bc112-0-2
- [2] J. S. Aujla and J.-C. Bourin, Eigenvalue inequalities for convex and log-convex functions, *Linear Algebra Appl.* **424** (2007), 25–35. doi:10.1016/j.laa.2006.02.027
- [3] J. Bergh and J. Löfström, *Interpolation Spaces: An Introduction*, Grundlehren der mathematischen Wissenschaften, vol. 223, Springer, Berlin–Heidelberg, 1976. doi:10.1007/978-3-642-66451-9
- [4] R. Bhatia, *Matrix Analysis*, Graduate Texts in Mathematics, vol. 169, Springer, New York, 1997. doi:10.1007/978-1-4612-0653-8
- [5] R. Bhatia and J. A. R. Holbrook, On the Clarkson–McCarthy inequalities, *Math. Ann.* **281** (1988), 7–12. doi:10.1007/BF01449211
- [6] R. Bhatia and F. Kittaneh, Clarkson inequalities with several operators, *Bull. Lond. Math. Soc.* **36** (2004), 820–832. doi:10.1112/S0024609304003467
- [7] J.-C. Bourin, T. Harada, and E.-Y. Lee, Subadditivity inequalities for compact operators, *Canad. Math. Bull.* **57** (2014), no. 1, 25–36. doi:10.4153/CMB-2012-009-9
- [8] J.-C. Bourin and F. Hiai, Norm and anti-norm inequalities for positive semi-definite matrices, *Internat. J. Math.* **22** (2011), 1121–1138. doi:10.1142/S0129167X1100715X
- [9] J.-C. Bourin and E.-Y. Lee, Unitary orbits of Hermitian operators with convex or concave functions, *Bull. Lond. Math. Soc.* **44** (2012), 1085–1102. doi:10.1112/blms/bds080
- [10] J.-C. Bourin and E.-Y. Lee, Clarkson–McCarthy inequalities with unitary and isometry orbits, *Linear Algebra Appl.* **601** (2020), 170–179. doi:10.1016/j.laa.2020.04.019
- [11] J.-C. Bourin and E.-Y. Lee, A Pythagorean theorem for partitioned matrices, *Proc. Amer. Math. Soc.* **152** (2024), no. 10, 4075–4086. doi:10.1090/proc/15677
- [12] J.-C. Bourin and E.-Y. Lee, Matrix parallelogram laws and symmetric moduli, *Internat. J. Math.* **31** (2026), no. 6, 2650018. doi:10.1142/S0129167X26500187
- [13] I. C. Gohberg and M. G. Kreĭn, *Introduction to the Theory of Linear Nonselfadjoint Operators in Hilbert Space*, Translations of Mathematical Monographs, vol. 18, American Mathematical Society, Providence, RI, 1969. doi:10.1090/mmono/018
- [14] O. Hirzallah and F. Kittaneh, Non-commutative Clarkson inequalities for n -tuples of operators, *Integral Equations Operator Theory* **60** (2008), 369–379. doi:10.1007/s00020-008-1565-x
- [15] D. J. Kečkić, Continuous generalization of Clarkson–McCarthy inequalities, *Banach J. Math. Anal.* **13** (2019), no. 1, 26–46. doi:10.1215/17358787-2018-0014
- [16] J. McCarthy, C_p , *Israel J. Math.* **5** (1967), 249–271. doi:10.1007/BF02771613
- [17] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993. doi:10.1007/978-94-017-1043-5
- [18] G. Pisier and Q. Xu, Non-commutative L^p -spaces, in *Handbook of the Geometry of Banach Spaces*, Vol. 2, W. B. Johnson and J. Lindenstrauss (eds.), Elsevier, 2003, pp. 1459–1517. doi:10.1016/S1874-5849(03)80041-4
- [19] B. Simon, *Trace Ideals and Their Applications*, 2nd ed., Mathematical Surveys and Monographs, vol. 120, American Mathematical Society, Providence, RI, 2005. doi:10.1090/surv/120
- [20] R. C. Thompson, Convex and concave functions of singular values of matrix sums, *Pacific J. Math.* **66** (1976), no. 1, 285–290. doi:10.2140/pjm.1976.66.285
- [21] T. Zhang, Full characterization of existence of Clarkson–McCarthy type inequalities, *arXiv* preprint (2024), arXiv:2410.21961v3.
- [22] T. Zhang, Proof of Audenaert–Kittaneh’s conjecture, *Israel J. Math.* (2025). doi:10.1007/s11856-025-2890-3