

Totally Δ -Modular Tree Decompositions of Graphic Matrices for Integer Programming

Caleb McFarland*† 1

¹School of Mathematics, Georgia Institute of Technology

June 2025

Abstract

We introduce the tree-decomposition-based parameter *totally Δ -modular treewidth* (TDM-treewidth) for matrices with two nonzero entries per row. We show how to solve integer programs whose matrices have bounded TDM-treewidth when variables are bounded. This extends previous graph-based decomposition parameters for matrices with at most two nonzero entries per row to include matrices with entries outside of $\{-1, 0, 1\}$. We also give an analogue of the Grid Theorem of Robertson and Seymour for matrices of bounded TDM-treewidth in the language of rooted signed graphs.

1 Introduction

We are interested in solving integer programs of the form

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

where $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Z}^m$, $w \in \mathbb{Z}^n$, and $\ell, u \in (\mathbb{Z} \cup \{\infty\})^n$. We may assume that A has at least two nonzero entries per row by possibly changing ℓ, u . Integer programs generalize many combinatorial optimization problems, see [Sch03]. It is well known that solving integer programs is NP-hard in general, so there has been much research on what conditions on A allow for a polynomial time algorithm. Examples include when A has a constant number of columns (see [LJ83] and improvements in [Kan87, Dad12]), a constant number of rows (see [Pap81] and improvements in [EW19]), or different block structures [CEH⁺21, CEP⁺20, EHK⁺19, BKP⁺24, CKL⁺25]).

The most classical example is when A is totally unimodular, see [Sch03]. To generalize this, a matrix is called *totally Δ -modular* if every square submatrix of A has determinant bounded by Δ in absolute value. Hence $\Delta = 1$ is exactly when A is totally unimodular. The following conjecture seeks to generalize the unimodular case to all Δ .

Conjecture 1.1 ([She96]). For any constant Δ , the integer program $\max\{w^T x : Ax \leq b, x \in \mathbb{Z}^n\}$ can be solved in polynomial time when A is totally Δ -modular.

Artmann, Weismantel, and Zenklusen [AWZ17] answered the conjecture in the affirmative for $\Delta = 2$, but $\Delta \geq 3$ remains open. Fiorini, Joret, Weltge, and Yuditsky [FJWY25] gave an algorithm for solving MAXIMUM WEIGHTED INDEPENDENT SET on graphs which do not contain many vertex-disjoint odd cycles. They then used this algorithm to solve totally Δ -modular integer programs with two nonzero entries per row. Kober [Kob25] extends this to totally Δ -modular integer programs with two nonzero entries per row after

*Supported in part by the National Science Foundation under Grant No. DMS- 2452111.

†Supported in part by the Georgia Tech ARC-ACO Fellowship.

removing a constant number of rows and columns. Choi, Gorsky, Kim, McFarland, and Wiederrect [CGK⁺25] extend the work of Fiorini et. al. to graphs which “decompose” into parts which do not contain many vertex-disjoint odd cycles. They again use this algorithm for MAXIMUM WEIGHTED INDEPENDENT SET to solve the analogous class of integer programs when the coefficients are in $\{-1, 0, 1\}$. Choi et. al. also give a grid theorem for when matrices with two nonzero entries per row and entries in $\{-1, 0, 1\}$ decompose into parts which are totally Δ -modular using signed graphs. We extend their work to matrices with entries outside of $\{-1, 0, 1\}$. First, we give a grid theorem which describes the obstructions to decomposing a matrix into totally Δ -modular parts; this theorem is stated in terms of rooted signed graphs. Then we show how to use such a decomposition to solve integer programs.

A *rooted signed graph* is a triple (G, γ, K) where G is a graph, $\gamma : E(G) \rightarrow \mathbb{Z}_2$ represents the parity of edges, and $K \subseteq V(G)$ is the set of roots. The odd cycle packing number (OCP) of a (rooted) signed graph, denoted $\text{OCP}(G, \gamma)$, is the size of a maximum collection of pairwise vertex-disjoint cycles so that the edges of each cycle sum to 1 under the weighting γ . Given a matrix A with exactly two nonzero entries per row, we can associate to A a rooted signed graph $G_\bullet^+(A) = (G, \gamma, K)$ where G has edge-vertex incidence matrix with the same support as A , γ keeps track of which rows have entries with the same or different signs, and K is the set of columns that contain an entry outside of $\{-1, 0, 1\}$. By $G(A)$ we denote the graph G , and by $G^+(A)$ we denote the (unrooted) signed graph (G, γ) . See Section 2 for precise definitions.

Informally, a matrix with two nonzero entries per row has bounded *totally Δ -modular treewidth* if it has a tree decomposition such that each bag is totally Δ -modular. In a rooted signed graph, this corresponds to each bag having few elements in K and bounded odd cycle packing number. However, the precise definition has technical restrictions on the adhesions. For each bag we have an apex set, the “protector”, and adhesions outside this apex set all have size 0 or 1 depending on whether the protector is “strong”. This requirement is to allow for dynamic programming: If a bag has an unbounded number of children and each contribute new vertices in their adhesion, this can pose a problem for dynamic programming. The reason that some protectors need not be strong is because we may eventually assume all variables of the integer program are in $\{0, 1\}$, see Section 7 for more details. We now formally define the totally Δ -modular treewidth (TDM-treewidth) of a rooted signed graph (and thus a matrix with two nonzero entries per row).

Definition 1.2 (TDM-treewidth). Let (G, γ, K) be a rooted signed graph. Given a tree decomposition (T, β) of (G, γ, K) , a *protector* for a bag $\beta(t)$ is a set $\alpha(t) \subseteq \beta(t)$ that contains all the roots in the bag and such that all adhesions to t have size at most one after deleting $\alpha(t)$. That is, $K \cap \beta(t) \subseteq \alpha(t)$ and $|(\beta(t) \cap \beta(t')) \setminus \alpha(t)| \leq 1$ for all $t, t' \in E(T)$. A protector is *strong* if all the adhesions to t are left empty after deleting $\alpha(t)$, i.e. if $\beta(t) \cap \beta(t') \subseteq \alpha(t)$ for all $t, t' \in E(T)$.

A *TDM-tree-decomposition* of a rooted signed graph (G, γ, K) is a tuple (T, β, α, J) where T is a tree, J is a (possibly empty) subtree of T , and $\beta, \alpha : V(T) \rightarrow 2^{V(G)}$ all satisfy

1. (T, β) is a tree-decomposition of G ,
2. $\alpha(t)$ is a protector for $\beta(t)$ for all $t \in V(T)$ and a strong protector when $t \in V(J)$,
3. $K \subseteq \bigcup_{j \in V(J)} \beta(j)$,

The *width* of a TDM-tree-decomposition is

$$\max_{t \in V(T)} |\alpha(t)| + \text{OCP}(G[\beta(t) \setminus \alpha(t)], \gamma).$$

The *TDM-treewidth* of (G, γ, K) , denoted $\text{TDM-tw}(G, \gamma, K)$, is the minimum width of a TDM-tree-decomposition.

The definition for TDM-treewidth arises naturally out of other width parameters for rooted graphs and signed graphs, see Section 5. See Section 7 for possible modifications of the definition and further discussion.

We are now ready to state our main algorithmic result, where we assume that w, b, ℓ , and u can be represented in $\text{poly}(n)$ many bits.

Theorem 1.3. Let d, k be nonnegative integers, and let A be a matrix with two nonzero entries per row whose associated rooted signed graph has TDM-treewidth at most k . Then for any $w \in \mathbb{Z}^n, b \in \mathbb{Z}^m$, and $\ell, u \in \mathbb{Z}^m$ with $\|u - \ell\|_\infty \leq d$, we can solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

in $d^{\mathcal{O}(k)} n^{f(k)}$ time for some computable f .

We prove Theorem 1.3 in Section 6 via dynamic programming and a reduction to MAXIMUM WEIGHTED INDEPENDENT SET. We note that in the above theorem we do not require a bound on $\|A\|_\infty$, the largest entry of A in absolute value. We do however require a bound on $\|u - \ell\|_\infty$, which is natural for many combinatorial problems. This need arises out of the lack of a proximity result like the one of Cook et. al. [CGST86, Theorem 1] (see also [PWW20, CKPW22]) for matrices which are only “locally” totally Δ -modular. In particular, the algorithm is still poly-time for $d \in \text{poly}(n)$, which is in line with the bound of $d = n\Delta$ we would get for totally Δ -modular matrices by the previously mentioned proximity result of Cook et. al.

We also give a grid theorem for having bounded TDM-treewidth. We first define the relevant grids. The *cylindrical* $(n \times m)$ -grid is $P_n \square C_m$, the Cartesian (or box) product of the path on n vertices P_n and the cycle C_m . Let γ_0, γ_1 be the functions which are identically equal to 0 or 1 respectively.

Definition 1.4 (Parity Handle \mathcal{H}_k and parity vortex \mathcal{V}_k). Let W be the cylindrical $(k \times 4k)$ -grid which we embed naturally in the plane such that the cycles C_{4k} are concentric. Let $X = \{x_1, \dots, x_{2k}\}$ consist of every other vertex along the outer face in clockwise order.

The *unsigned, unrooted parity handle* \mathbf{H}_k of order k is the graph obtained from W by adding the edges $x_i x_{2k-i+1}$ for all $i \in [k]$. The *parity handle* \mathcal{H}_k of order k is the rooted signed graph $(\mathbf{H}_k, \gamma_1, \emptyset)$.

The *unsigned, unrooted parity vortex* \mathbf{V}_k of order k is the graph obtained from W by adding the edges $x_{2i-1} x_{2i}$ for all $i \in [k]$. The *parity vortex* \mathcal{V}_k of order k is the rooted signed graph $(\mathbf{V}_k, \gamma_1, \emptyset)$.

See Figure 1a for examples

Definition 1.5 (Rooted Grid \mathcal{W}_k). Let \mathbf{W}_k be the $(k \times k)$ -grid and let $K_{\mathbf{W}_k}$ be the vertex set of the first row of \mathbf{W}_k . The rooted grid \mathcal{W}_k of order k is the rooted signed graph $(\mathbf{W}_k, \gamma_0, K_{\mathbf{W}_k})$. See Figure 1b for an example.

We can now state our main structural result.

Theorem 1.6. There exists a polynomial $f : \mathbb{N} \rightarrow \mathbb{N}$ such that for every integer $k \geq 1$ and every rooted signed graph (G, γ, K) ,

1. if (G, γ, K) contains one of $\mathcal{H}_k, \mathcal{V}_k$, or \mathcal{W}_k as a minor, then $\text{TDM-tw}(G, \gamma, K) \geq \Omega(k)$, and
2. if $\text{TDM-tw}(G, \gamma, K) \geq f(k)$, then G contains one of $\mathcal{H}_k, \mathcal{V}_k$, or \mathcal{W}_k as a minor.

Theorem 1.6 and Theorem 1.3 give the following immediate corollary.


Corollary 1.7. Let d, k be nonnegative integers, and let A be a matrix with two nonzero entries per row whose associated rooted signed graph forbids $\mathcal{H}_k, \mathcal{V}_k$, and \mathcal{W}_k as minors. Then for all $w \in \mathbb{Z}^n, b \in \mathbb{Z}^m$, and $\ell, u \in \mathbb{Z}^m$ with $\|u - \ell\|_\infty \leq d$, we can solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$


in $d^{\text{poly}(k)} n^{f(k)}$ time for some computable f .

Our work is towards the following problem posed in [CGK⁺25].

Question 1.8 (Question 10.10 [CGK⁺25]). What is the most general class of matrices with two nonzero entries per row such that we can solve the integer program $\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$ in polynomial time for all matrices A in the class?



(a) The parity handle of order 5 (left) and the parity vortex of order 5 (right). Both graphs are shifting equivalent to the graph with odd edges exactly equal to the red edges.



(b) The rooted grid of order 8. The vertices in red are the rooted vertices $K_{\mathbb{W}_k}$. All cycles of the rooted grid are even.

This question seeks to generalize the below question which is central to algorithmic graph theory.

Question 1.9. What is the most general class of graphs for which we can solve MAXIMUM WEIGHTED INDEPENDENT SET (MIS) in polynomial time for all graphs in the class?

In order to make the above questions tractable, it is natural to only consider classes which are closed under certain operations. For example, if we ask Question 1.9 for hereditary graph classes, then there has been a long line of research on solving MIS over graph classes defined by forbidden induced subgraphs [GLS84, Min80, Ale04, LM08, CLB81, FHT93].

More relevant to this work is when we force the classes to be closed under the minor relation. Then Question 1.9 is answered by the famous Grid Theorem of Robertson and Seymour [RS86], which implies that MIS can be solved on a minor-closed class of graphs if and only if the class excludes a planar graph. If we allow our matrices in Question 1.8 to be in $\{0, 1\}^{m \times n}$ and use the typical minor relation on $G(A)$, then Question 1.8 is equivalent to Question 1.9.

If we instead allow our matrices in Question 1.8 to be over $\{-1, 0, 1\}$, it is natural to associate such matrices with signed graphs and to force classes to be closed under the signed graph minor relation (see Section 2.1 for relevant definitions). Historically, signed graphs have been studied in relation to matroids and geometry [Zas82, Zas91, Zas13], but they have recently found relations to integer programming [CGK⁺25, Kob25]. Question 1.8 for signed graph minor closed classes of matrices with entries in $\{-1, 0, 1\}$ then again reduces to Question 1.9 for graph classes closed under odd-minors, see [CGK⁺25, Theorem 10.3] based on [FJWY25, Section 3]. We discuss the relation between odd-minors and signed graphs in Section 2.2. Choi et. al. [CGK⁺25] showed that MIS (and hence the analogous integer programming problem) is solvable over classes which exclude both a planar graph with at most two odd faces and a planar graph with all odd cycles touching a common face (that is, \mathbf{H}_k and \mathbf{V}_k for some k) as an odd-minor. This generalized an array of previous work on the MIS problem in odd-minor-closed graph classes [EGHK21, JDKW21, JdK21, GW23, JMST25, FJWY25]. Other notable work on the MIS in odd-minor-closed graph classes include [Ger89, Taz12]. The problem is only known to be hard when the class contains all planar graphs, and so an exact hardness threshold is not known for this case.

We instead pose Question 1.8 for matrices which may have entries outside of $\{-1, 0, 1\}$. We again force our classes to be minor closed, but it is not obvious which minor operation to choose. A matrix with two nonzero entries per row is totally Δ -modular if and only if it has bounded entries, a bounded number of columns with entries outside $\{-1, 0, 1\}$, and the associated signed graph has bounded odd cycle packing (see Lemma 2.9). Motivated by this equivalence, we propose representing such matrices as rooted signed graphs as defined above. We can then naturally encode the minor relation as *rooted signed graph minors* (see Section 2.3 for relevant definitions), which allows us to apply results on rooted minors and rooted tree decompositions [MSW17, JS24, HLMR24].

The best known results for matrices with two nonzero entries per row and entries possibly outside $\{-1, 0, 1\}$ are given by the follow results which more generally apply to matrices with an arbitrary number of nonzero entries per row. The *primal graph* of a matrix A is a graph with vertex set equal to the columns of A where two columns are adjacent if and only if there exists a row where both columns are nonzero. If we assume the domain size $\|u - \ell\|_\infty$ is bounded, then we can solve the integer program when the primal graph has bounded treewidth [JK15, Fre90]. If we instead bound $\|A\|_\infty$, then we can solve the integer program when the corresponding graph has bounded treedepth¹ [EHK⁺19, GOR17, GO18]. Integer programming remains NP-hard on matrices with $\|A\|_\infty$ bounded and where the primal graph has bounded treewidth [GOR17], so treedepth can not be relaxed to treewidth. Similarly, integer programming remains NP-hard on matrices where the primal graph has bounded treedepth and $\|A\|_\infty$ is unbounded [DEG⁺21]. If we consider the matroid represented by A , integer programming remains NP-hard even when the matroid has bounded branch-width and the entries are in $\{-1, 0, 1\}$ [CG07]. However, all of these reductions (as well as others [KPW20, EGK⁺19, BLP⁺25]) rely on matrices with at least 3 nonzero entries per row, and the only known hardness result for matrices with at most two nonzero entries per row is the reduction to MAXIMUM WEIGHTED INDEPENDENT SET.

¹A graph has bounded *treedepth* if it has a tree decomposition (T, β) of bounded width such that T has bounded height.

The above results imply that if $\|u - \ell\|_\infty$ is bounded and if A is a matrix with two nonzero entries per row such that $G(A)$ has bounded treewidth, then we can solve the integer program in polynomial time. [Theorem 1.3](#) generalizes both this result and the result of Choi et. al. [CGK⁺25] (see [Theorem 4.8](#)) when $\|u - \ell\|_\infty$ is bounded.

The paper is organized as follows. In [Section 2](#), we give the relevant background on signed graphs, rooted graphs, and their connection to totally Δ -modular matrices. In [Section 3](#), we discuss K -free treewidth, as introduced in [JS24], and a grid theorem for rooted minors. In [Section 4](#), we discuss OCP-treewidth, as introduced in [CGK⁺25], and its applications to integer programming. In [Section 5](#), we discuss the connection between K -free treewidth, OCP-treewidth, and TDM-treewidth. We use this connection to give an FPT-approximation algorithm for the TDM-treewidth of a graph, and we prove [Theorem 1.6](#). In [Section 6](#), we prove [Theorem 1.3](#). Finally, in [Section 7](#), we discuss potential future work and open questions.

2 Preliminaries

We use $[k]$ to denote the set $\{1, 2, \dots, k\}$. By $\|A\|_\infty$ we denote the largest absolute value among all entries of the matrix A . All graphs may have parallel edges but not loops. We denote by $G[X]$ the graph induced on X for $X \subseteq V(G)$. For V_1, V_2 a partition of $V(G)$, we denote by $[V_1, V_2]$ the set of all edges with one end in V_1 and one end in V_2 . We call such a $[V_1, V_2]$ an *edge cut*.

Definition 2.1 (tree decomposition). A *tree decomposition* of a graph G is a pair (T, β) where T is a tree and $\beta : V(T) \rightarrow 2^{V(G)}$ such that

1. $\bigcup_{t \in V(T)} \beta(t) = V(G)$,
2. for every edge $e \in E(G)$, there exists $t \in V(T)$ such that both ends of e are contained in $\beta(t)$, and
3. for every $v \in V(G)$, the subgraph of T induced by $\{t \in V(T) : v \in \beta(t)\}$ is connected.

The *width* of a tree decomposition (T, β) is $\max_{t \in V(T)} |\beta(t)| - 1$. The *treewidth* of a graph G is the minimum width of a tree decomposition of G . We call $\beta(t)$ the *bag* of t . Each edge $tt' \in E(T)$ corresponds to an *adhesion* $\beta(t) \cap \beta(t')$.

Definition 2.2 (grid). The $k \times k$ grid is the graph with vertex set $[k]^2$ such that (i, j) is adjacent to (i', j') if and only if $|i - i'| + |j - j'| = 1$.

2.1 Signed graphs

A signed graph is a pair (G, γ) where G is a graph and $\gamma : E(G) \rightarrow \mathbb{Z}_2$. For $e \in E(G)$, if $\gamma(e) = 0$ the edge is said to be *even*, and if $\gamma(e) = 1$ the edge is said to be *odd*. The *parity* of a path or cycle C in G is $\sum_{e \in E(C)} \gamma(e)$, and a cycle is said to be *even* or *odd* if the parity is 0 or 1 respectively. The *odd cycle packing number* (OCP) of a signed graph (G, γ) , denoted $\text{OCP}(G, \gamma)$ is the size of a maximum collection of pairwise vertex-disjoint odd cycles with respect to γ .

We commonly use γ_1 to denote the function which maps every edge to $1 \in \mathbb{Z}_2$ where the domain of the function is clear from context. Similarly we use γ_0 to denote the function which maps everything to $0 \in \mathbb{Z}_2$.

Given a signed graph (G, γ) , *shifting at a vertex* $v \in V(G)$ replaces $\gamma(e)$ with $\gamma(e) + 1$ for each non-loop edge e incident to v . We say (G, γ') is obtained from (G, γ) by *shifting* if it is obtained by a sequence of shiftings at vertices. Note that if (G, γ') is a shifting of (G, γ) , then every cycle in (G, γ') has the same parity as in (G, γ) . The converse is also true: if (G, γ') and (G, γ) have the same parity for every cycle, then (G, γ') can be obtained from (G, γ) by shifting. Hence it is often natural to think of signed graphs under the equivalence relation given by shifting.

A signed graph (H, γ_H) is a *(signed graph) minor* of (G, γ_G) if it can be obtained by vertex deletion, edge deletion, shifting, and contracting even edges. Because we may only contract even edges, every cycle in (H, γ_H) naturally corresponds to a cycle in (G, γ_G) of the same parity. Equivalently, (H, γ_H) is a minor of (G, γ_G) if there exists a *minor model* of (H, γ_H) in (G, γ_G) . That is, if there exist

1. a mapping φ_V from $V(H)$ to pairwise vertex disjoint trees in G ,
2. an injection $\varphi_E : E(H) \rightarrow E(G)$ such that $\varphi_E(uv)$ has one end in $\varphi_V(u)$ and one end in $\varphi_V(v)$ for each $uv \in E(H)$, and
3. a shifting (G, γ'_G) of (G, γ_G) such that
 - (a) for all $v \in V(H)$, the edges of the tree $\varphi_V(v)$ all have label 0, and
 - (b) for every $e \in E(H)$, $\gamma'_G(\varphi_E(e)) = \gamma_H(e)$.

Note that if (H, γ_H) is a signed graph minor of (G, γ_G) then H is a minor of G in the usual sense.

A signed graph (G, γ_G) contains a signed graph (H, γ_H) as a *subdivision* if there exists $\gamma'_H, \varphi_V, \varphi_E$ such that

1. (H, γ'_H) is a shifting of (H, γ_H) ,
2. $\varphi_V : V(H) \rightarrow V(G)$ is an injection, and
3. φ_E is a mapping from edges of H to internally vertex-disjoint paths in G such that
 - (a) $\varphi_E(uv)$ is a path with ends $\varphi_V(u), \varphi_V(v)$ and
 - (b) the parity of the path $\varphi_E(uv)$ in (G, γ) is equal to $\gamma'_H(uv)$.

Note that if (G, γ_G) contains (H, γ_H) as a subdivision, then (G, γ_G) contains (H, γ_H) as a minor.

We call the cycles of length 4 in a grid the *cells*. A signed $n \times m$ grid is said to be *even* if all of its cells are even. Note that if (W, γ) is an even grid, then (W, γ_0) is a shifting. We now show that in any signed grid, we can always find a large even grid.




Figure 2: The construction in the proof of Lemma 2.3.

Lemma 2.3. Let $k \in \mathbb{N}$. Let G be the $k^2 \times k^2$ grid and let W be the $k \times k$ grid. Then (G, γ) contains (W, γ_0) as a subdivision for every γ .

Proof. Enumerate the vertices of G as $V(G) = \{(i, j) : i, j \in [k^2]\}$ as in [Definition 2.2](#). Similarly let the vertices of W be enumerated (a, b) for $a, b \in [k]$. Let $f(x) = (k+1)(x-1) + 1$. We will map the vertex (a, b) in W to the vertex $(f(a), f(b))$ in G . For every $b \in [k-1]$, the edge between (a, b) and $(a, b+1)$ will be mapped to the path in G with vertex set $\{f(a), f(b)+\ell : 0 \leq \ell \leq k+1\}$. For every $a \in [k-1]$, the edge between $(a, 1)$ and $(a+1, 1)$ will be mapped to the path with vertex set $\{f(a)+\ell, 1 : 0 \leq \ell \leq k+1\}$. It remains to map the edges between (a, b) and $(a+1, b)$ for $b \geq 2$.

For $a \in \{2, 3, \dots, k\}$ and $b \in [k-1]$, let $H_{a,b}$ be the $k \times k$ subgrid of G with opposite corners at $(f(a)+1, f(b-1)+1)$ and $(f(a+1)-1, f(b)-1)$. We may assume that each $H_{a,b}$ contains an odd cell, otherwise we take $W = H_{a,b}$. We claim that between any two corners of the grid $H_{a,b}$, there exists a path in $H_{a,b}$ of both parities. Indeed, we make take two vertex-disjoint paths from the two corners to the vertices of an odd cell. Then by deciding which way to traverse through the odd cell we may decide the parity of the resulting path.

We fix $a \in [k-1]$ and define the mapping from the edge between (a, b) and $(a, b+1)$ in order of increasing b . The first two vertices of the path are $(f(a)+1, f(b))$ and $(f(a)+1, f(b)-1)$. The last two vertices of the path are $(f(a+1)-1, f(b)-1)$ and $(f(a+1)-1, f(b))$. Note that $(f(a)+1, f(b)-1)$ and $(f(a+1)-1, f(b)-1)$ are two corners of the grid $H_{a,b}$. Thus we may complete the path through $H_{a,b}$ with a parity of our choosing. We choose the path that forces the cell with vertices $(a, b-1), (a, b), (a+1, b), (a+1, b-1)$ to get mapped to an even cycle. Because the mapping from all other edges of the cell are fixed when choosing the edge between (a, b) and $(a, b+1)$, such a choice is always possible. By choosing the paths in this way, every cell is mapped to an even cycle. \square

The above lemma implies that (G, γ) forbids an even grid as a signed graph minor if and only if G forbids a grid as a minor. More precisely, we get the following corollary.

Corollary 2.4. For any γ , the following holds. If G forbids the $k \times k$ grid as a minor, then (G, γ) forbids the $k \times k$ even grid as a signed graph minor. Conversely, if (G, γ) forbids the $k \times k$ even grid as a signed graph minor, then G forbids the $k^2 \times k^2$ grid as a minor.

2.2 Odd-minors and signed graphs

A graph G contains a graph H as an *odd-minor* if H can be obtained from G by deleting vertices, deleting edges, and contracting edge cuts. Equivalently, H is an odd-minor of G if there exists

1. a mapping φ_V from $V(H)$ to pairwise vertex disjoint trees in G ,
2. an injection $\varphi_E : E(H) \rightarrow E(G)$ such that $\varphi_E(uv)$ has one end in $\varphi_V(u)$ and one end in $\varphi_V(v)$ for each $uv \in E(H)$, and
3. a 2-coloring $c : V(G) \rightarrow [2]$ such that
 - (a) for all $v \in V(H)$, c is a proper 2-coloring when restricted to the tree $\varphi_V(v)$, and
 - (b) for every $e \in E(H)$, $\varphi_E(e)$ is monochromatic.

Odd-minors have been studied extensively both as it relates to coloring [[GGR⁺09](#), [Ste22](#), [KSSW25](#)] and algorithms [[Ger89](#), [Taz12](#), [GW23](#), [JMST25](#), [CGK⁺25](#)]. Odd-minors and signed graph minors are essentially equivalent due to the following observations. Recall that γ_1 is the function which maps every edge to $1 \in \mathbb{Z}_2$.

Observation 2.5. G contains H as an odd-minor if and only if (G, γ_1) contains (H, γ_1) as a signed graph minor.

Proof. Suppose G contains H as an odd-minor. Consider the sequence of deletions and contractions. Clearly whenever we delete a vertex or edge in G we may do the same in (G, γ_1) . Whenever we contract an edge cut $[V_1, V_2] \subseteq E(G)$, we can shift in (G, γ_1) at every vertex in V_1 such that the even edges are exactly the edges $[V_1, V_2]$. We may then contract the edges in the signed graph.

Suppose (G, γ_1) contains (H, γ_1) as a signed graph minor. Consider the minor model φ_V, φ_E of (H, γ_1) in (G, γ_1) . Let $c : V(G) \rightarrow [2]$ denote whether a vertex was shifted at in G in order to make the edges of $\varphi_V(v)$ have label 0 for every $v \in V(H)$. Then on every tree $\varphi_V(v)$, c is a proper 2-coloring. Because (G, γ_1) and (H, γ_1) both have all edges odd, every edge in the image of φ_E must be monochromatic under c . Thus we obtain a minor model for H in G as an odd-minor. \square

Observation 2.6. Let H be a graph of minimum degree 3. Let (G, γ) be a signed graph, and let G' be the graph formed from G by subdividing every even edge exactly once. Then (G, γ) contains (H, γ_1) as a signed graph minor if and only if G' contains H as an odd-minor.

Proof. Suppose (G, γ) contains (H, γ_1) as a minor. Clearly (G', γ_1) contains (G, γ) as a minor by shifting at every subdividing vertex and contracting one of the incident edges. Thus (G', γ_1) contains (H, γ_1) as a minor, and so G' contains H as an odd-minor by Observation 2.5.

Suppose G' contains H as an odd-minor. Then (G', γ_1) contains (H, γ_1) as a minor. Because H has minimum degree 3, every subdividing vertex must be contracted or deleted when forming H . We may reorder the operations such that all subdividing vertices are removed first. After performing the operations on the subdividing vertices, the resulting graph is a minor of (G, γ) . Therefore (H, γ_1) is a minor of (G, γ) . \square

2.3 Rooted (signed) graphs

A *rooted graph* is a pair (G, K) where G is a graph and $K \subseteq V(G)$ denotes the set of *roots*. When we contract an edge in a rooted graph, the resulting vertex is a root if and only if one of the ends of the contracted edge was a root. A rooted graph (H, K_H) is a rooted minor of (G, K_G) if it can be obtained by vertex deletion, edge deletion, contraction, and removing vertices from the set of roots.

A *rooted signed graph* is a triple (G, γ, K) where (G, γ) is a signed graph and $K \subseteq V(G)$. A rooted signed graph (H, γ_H, K_H) is a rooted minor of (G, γ_G, K_G) if it can be obtained via vertex deletion, edge deletion, shifting, contracting even edges, and removing vertices from the set of roots.

We note that Lemma 2.3 and Corollary 2.4 apply more generally for the rooted grid.

Lemma 2.7. Let $k \in \mathbb{N}$. Let $(\mathbf{W}_{k^2}, K_{\mathbf{W}_{k^2}})$ be the $k^2 \times k^2$ grid rooted at the first row of \mathbf{W}_{k^2} . Then $(\mathbf{W}_{k^2}, \gamma, K_{\mathbf{W}_{k^2}})$ contains \mathcal{W}_k as a rooted grid minor for every γ .

Proof. The proof is exactly as in Lemma 2.3, noting that for any grid minor found by the proof, there exists k vertex disjoint paths from the first row of the grid minor to the first row of \mathbf{W}_{k^2} . \square

Corollary 2.8. For any γ , the following holds. If (G, K) forbids $(\mathbf{W}_k, K_{\mathbf{W}_k})$ as a rooted minor, then (G, γ, K) forbids \mathcal{W}_k as a rooted signed graph minor. Conversely, if (G, γ, K) forbids \mathcal{W}_k as a rooted signed graph minor, then (G, K) forbids $(\mathbf{W}_{k^2}, K_{\mathbf{W}_{k^2}})$ as a rooted minor.

2.4 Matrices

Let A be a matrix with exactly two nonzero entries per row. By $G(A)$ we denote the graph whose edge-vertex incidence matrix is the support of A . That is, $G(A)$ has vertex set equal to the columns of A and edge set equal to the rows of A , and vertex v is incident to edge e if and only if $A_{e,v}$ is nonzero. By $G^+(A)$ we denote the signed graph $(G(A), \gamma)$ where $\gamma(e) = 0$ if the entries of row e have different signs, and $\gamma(e) = 1$ if the entries of row e have the same sign. By $G_\bullet^+(A)$ we denote the rooted signed graph $(G(A), \gamma, K)$ where $(G(A), \gamma) = G^+(A)$ and K is the set of columns that contain an entry outside of $\{-1, 0, 1\}$.

A matrix $A \in \mathbb{Z}^{m \times n}$ is *totally Δ -modular* if every subdeterminant of A is contained in $\{-\Delta, -\Delta + 1, \dots, \Delta\}$. If A has two nonzero entries per row, this is equivalent to bounding $\|A\|_\infty$, the number of

columns with entries outside $\{-1, 0, 1\}$, and the odd cycle packing of the corresponding graph, as given by the below lemma.

Lemma 2.9 (Lemma 10.2 [CGK⁺25]). Let $A \in \mathbb{Z}^{m \times n}$ have two nonzero entries per row and let $(G, \gamma, K) = G_\bullet^+(A)$. Then if A is totally Δ -modular,

1. $\|A\|_\infty \leq \Delta$,
2. $|K| \leq 2 \log_2 \Delta$, and
3. $\text{OCP}(G, \gamma) \leq \log_2 \Delta$.

Conversely, if Δ is the largest absolute value of a subdeterminant of A , then $\Delta \leq 2^{\text{OCP}(G, \gamma)} \|A\|_\infty^{|K|}$.

3 K -free treewidth

Jansen and Swennenhuus [JS24] introduced K -free treewidth as a notion of treewidth specialized for rooted graphs. This notion will be central in decomposing our rooted signed graph with respect to the roots. We note this differs from other notions of treewidth with respect to rooted graphs, such as annotated treewidth or bidimensionality [SST25, PTW25].

Definition 3.1 (Tree K -free-decomposition). Let G be a graph and $K \subseteq V(G)$. A *tree K -free-decomposition* of G is a triple (T, β, L) where T is a tree, $L \subseteq V(G) \setminus K$, and $\beta : V(T) \rightarrow 2^{V(G)}$ such that

1. (T, β) is a tree decomposition of G , and
2. for each $v \in L$, there exists a unique $t \in V(T)$ such that $v \in \beta(t)$, and t is a leaf of T .

The *width* of a tree K -free-decomposition (T, β, L) is $\max\{0, \max_{t \in V(T)} |\beta(t) \setminus L| - 1\}$. The *K -free treewidth* of a graph G , denoted $\text{tw}_K(G, K)$ is the minimum width of a tree K -free-decomposition of G .

Intuitively, a graph has low K -free treewidth if their exists a tree decomposition where internal bags must be small, but leaf bags can contain many vertices outside of K .

Jansen and Swennenhuus [JS24] also gave an FPT 5-approximation for K -free treewidth.

Theorem 3.2 (Theorem 3.4 [JS24]). There exists an algorithm that, given an n -vertex graph G , $K \subseteq V(G)$, and a non-negative integer k , either computes a tree K -free-decomposition (T, β) of G of width at most $5k + 5$ with $|V(T)| \in \mathcal{O}(n)$ nodes, or correctly returns that $\text{tw}_K(G, K) > k$. The algorithm runs in time $2^{\mathcal{O}(k)} \text{poly}(n)$.

Marx, Seymour, and Wollan [MSW17] proved a grid theorem for rooted minors in the language of tangles, but we present the statement in terms of K -free treewidth as given in [HLMR24].

Theorem 3.3 (Theorem 9 [HLMR24]). There exists a function $f_{3.3} : \mathbb{N} \rightarrow \mathbb{N}$ such that for every positive integer k , every graph G , and every set $K \subseteq V(G)$, if (G, K) does not contain $(\mathbf{W}_k, K_{\mathbf{W}_k})$ as a rooted-minor, then the K -free treewidth of G is at most $f_{3.3}(k)$. Furthermore $f_{3.3} \in \mathcal{O}(k^{36+o(1)})$.

We also show that the approximate converse is true. First, we note the following observation.

Observation 3.4. If (H, K_H) is a rooted minor of (G, K_G) , then $\text{tw}_K(H, K_H) \leq \text{tw}_K(G, K_G)$.

Proof. Clearly removing vertices from K_G and deleting vertices or edges from G can only decrease the K -free treewidth. If (T, β, L) is a tree K_G -free-decomposition of (G, K_G) and we contract an edge $uv \in E(G)$ to obtain the rooted graph (H, K_H) , then by identifying the two vertices in each bag of the tree we obtain a tree K_H -free-decomposition of (H, K_H) of only smaller width. The new vertex is in L if and only if both ends of the contracted edge were in L . \square

We now show the approximate converse of [Theorem 3.3](#).

Lemma 3.5. If (G, K) contains $(\mathbf{W}_{3k+1}, K_{\mathbf{W}_{3k+1}})$ as a rooted minor, then $\text{tw}_K(G, K) \geq k$.

Proof. By [Observation 3.4](#), it suffices to show that the K -free treewidth of $(\mathbf{W}_{3k+1}, K_{\mathbf{W}_{3k+1}})$ is at least k . Let $(\mathbf{W}_{3k+1}, K_{\mathbf{W}_{3k+1}}) = (W, K)$. Note that K is highly connected set. That is, for any $A, B \subseteq K$, there are $\min\{|A|, |B|\}$ vertex disjoint (A, B) -paths in W . Suppose for contradiction there exists a tree K -free-decomposition (T, β, L) of (W, K) of width less than k . We may assume that T has maximum degree at most 3 as follows. For any vertex $t \in V(T)$ with degree larger than 3, create a new vertex t' adjacent to t and make two of the neighbors of t be adjacent to t' instead of t . By setting $\beta(t') = \beta(t)$, we obtain a tree K -free-decomposition of width less than k . By repeating this process we may assume that T has maximum degree at most 3.

Each edge $t_1t_2 \in E(T)$ corresponds to a separation (G_1, G_2) in G of order at most k . That is, if T_1, T_2 are the components of $T - t_1t_2$ containing t_1, t_2 respectively, then $V(G_1) = \bigcup_{t \in V(T_1)} \beta(t)$, $V(G_2) = \bigcup_{t \in V(T_2)} \beta(t)$, and there are no edges in G with one end in $V(G_1) \setminus V(G_2)$ and on end in $V(G_2) \setminus V(G_1)$. Because K is highly connected, there exists a unique side of the $i \in [2]$ such that G_i contains at most k vertices from K . We can orient the edges of T to point away from this side, and because T is acyclic, there exists a vertex $t \in V(T)$ which is a sink. Let T_1, \dots, T_ℓ be the subtrees of T whose union is T and such that each T_i, T_j pairwise intersect exactly at t . Note that $\ell = d(t) \leq 3$. For $i \in [\ell]$, let $A_i = \bigcup_{t' \in V(T_i)} \beta(t')$. Then $\bigcup_{i \in [\ell]} A_i = V(G)$ and each A_i contains at most k of the vertices of K , contradicting that $|K| = 3k + 1$. \square

4 (tame) OCP-treewidth

Odd-Cycle-Packing-treewidth (or OCP-treewidth) was introduced by Choi, Gorsky, Kim, McFarland, and Wiederrect [\[CGK⁺25\]](#) as a variant of treewidth concerned with the odd cycle packing number of each bag. For notational convenience we will only be concerned with tame OCP-tree-decompositions, and we do not define OCP-tree-decompositions in general. The minimum width of a tame OCP-tree-decomposition is functionally equivalent to the minimum width of an OCP-tree-decomposition, see [\[CGK⁺25, Theorem 4.2\]](#), but the definition of a tame OCP-tree-decomposition matches more closely with the definition of a TDM-tree-decomposition.

Definition 4.1 (tame OCP-tree-decomposition). A *tame OCP-tree-decomposition* of a graph G is a triple (T, β, α) where T is tree and $\beta, \alpha: V(T) \rightarrow 2^{V(G)}$ such that

1. (T, β) is a tree decomposition of G ,
2. $\alpha(t) \subseteq \beta(t)$ for each $t \in V(T)$, and
3. for each $tt' \in E(T)$, $|(\beta(t) \cap \beta(t')) \setminus \alpha(t)| \leq 1$.

The *tame width* (*t-width*) of a tame OCP-tree-decomposition (T, β, α) is:

$$\max_{t \in V(T)} |\alpha(t)| + \text{OCP}(G[\beta(t) \setminus \alpha(t)]).$$

We note that this definition of width differs by at most 1 from the usual definition of the width for a tame OCP-tree-decomposition, but is easier to work with for tame decompositions. The *tOCP-treewidth* of a graph G , denoted $\text{tOCP-tw}(G)$, is the minimum tame width of an tame OCP-tree-decomposition of G .

We can extend this definition to signed graphs (G, γ) by letting the odd cycle packing number in the above definition refer to odd cycle packing number of the signed graph. Note that $\text{tOCP-tw}(G, \gamma_1) = \text{tOCP-tw}(G)$. Similarly we can extend the definition to rooted signed graphs by ignoring the set of roots. Note that for a rooted signed graph (G, γ, \emptyset) , the sets $\alpha(t)$ in the above definition are protectors for the bags $\beta(t)$. In particular, if (T, β, α, J) is a TDM-tree-decomposition of (G, γ, K) , then (T, β, α) is a tame OCP-tree-decomposition of (G, γ) .

Choi et. al. noted that (tame) OCP-treewidth is odd-minor-monotone, but we show more generally that the tOCP-treewidth of signed graphs is minor monotone with respect to signed graph minors.

Lemma 4.2. Let (H, γ_H) be a minor of (G, γ_G) . Then $\text{tOCP-tw}(H, \gamma_H) \leq \text{tOCP-tw}(G, \gamma_G)$.

Proof. Clearly deleting an edge or vertex can only decrease the tOCP-treewidth. Shifting also does not change the tOCP-treewidth because it does not change the odd cycle packing of any subgraph. Thus we may assume that (H, γ_H) is obtained from (G, γ_G) by contracting a single even edge. Let (T, β, α) be a tame OCP-tree-decomposition of (G, γ_G) . We then replace both ends of the edge with the new vertex in every $\beta(t), \alpha(t)$ for $t \in V(T)$ to obtain a tame OCP-tree-decomposition of (H, γ_H) of only smaller t-width. We note that contracting an even edge can only decrease the odd cycle packing number of every subgraph. \square

The following lemma will be helpful for translating between decompositions for unsigned graphs and signed graphs.

Lemma 4.3. Let (G', γ') be obtained from (G, γ) by replacing an edge with a path of the same parity. Then $\text{tOCP-tw}(G', \gamma') = \text{tOCP-tw}(G, \gamma)$. Furthermore there exists an algorithm which takes as input (G', γ') , (G, γ) , and a tame OCP-tree-decomposition of (G', γ') of t-width k and returns a tame OCP-tree-decomposition of (G, γ) of t-width k with the same number of bags in time $g(k)\text{poly}(|V(G')|)$.

Proof. Suppose (G', γ') is obtained from (G, γ) by replacing the edge $xy \in E(G)$ by the path $P = xv_1v_2 \dots v_\ell y$ where $\gamma'(e) = \gamma(e)$ for all $e \in E(G) \cap E(G')$ and $\sum_{e \in E(P)} \gamma'(e) = \gamma(xy)$. Let (T, β, α) be a tame OCP-tree-decomposition for (G, γ) . There is some bag $\beta(t)$ for which $x, y \in \beta(t)$. Let $\beta'(t) = \beta(t) \cup V(P)$. For all $t' \in V(T) \setminus \{t\}$, let $\beta'(t') = \beta(t')$. Then (T, β', α) is a tame OCP-tree-decomposition of (G', γ') of the same t-width. Note that $\text{OCP}(G'[\beta'(t) \setminus \alpha(t)], \gamma') = \text{OCP}(G[\beta(t) \setminus \alpha(t)], \gamma)$. Hence $\text{tOCP-tw}(G', \gamma') \leq \text{tOCP-tw}(G, \gamma)$. We note that (G, γ) is a minor of (G', γ') , and so $\text{tOCP-tw}(G, \gamma) \leq \text{tOCP-tw}(G', \gamma')$.

Because (G, γ) is obtained from (G', γ') by only shifting and contracting edges in P , one can simply identify the internal vertices of P with one of its endpoints in each bag to algorithmically obtain a tame OCP-tree-decomposition for (G, γ) as in the proof of Lemma 4.2. \square

Choi et. al. [CGK⁺25] also gave an FPT approximation for tOCP-treewidth, and they gave an excluded odd-minor characterization for having bounded tOCP-treewidth.

Theorem 4.4 (Theorem 4.2 [CGK⁺25]). There exists an algorithm that, given an n -vertex graph G and a non-negative integer k , either computes a tame OCP-tree-decomposition (T, β, α) for G of t-width at most $f_{4.4}(k)$ and with $|V(T)| \in \mathcal{O}(n)$, or correctly returns that $\text{tOCP-tw}(G) > k$. The algorithm runs in time $\mathcal{O}_t(n^6)$. Furthermore, $f_{4.4}(k) \in \text{poly}(k)$.

Theorem 4.5 (Theorem 1.6 and Theorem 3.4 [CGK⁺25]). There exists a polynomial $f_{4.5} : \mathbb{N} \rightarrow \mathbb{N}$ such that for every positive integer k and every graph G ,

1. if G contains \mathbf{H}_k or \mathbf{V}_k as an odd-minor, then $\text{tOCP-tw}(G) \geq \frac{k}{2} - 1$, and
2. if $\text{tOCP-tw}(G) \geq f_{4.5}(k)$, then G contains \mathbf{H}_k or \mathbf{V}_k as an odd-minor.

The above imply the same results for signed graphs by utilizing Observation 2.6 and Lemma 4.3.

Corollary 4.6. There exists an algorithm that, given an n -vertex, m -edge signed graph (G, γ) and a non-negative integer k , either computes a tame OCP-tree-decomposition (T, β, α) for (G, γ) of t-width at most $f_{4.4}(k)$ and with $|V(T)| \in \mathcal{O}(n+m)$, or correctly return that $\text{tOCP-tw}(G, \gamma) = \text{tOCP-tw}(G', \gamma_1) = \text{tOCP-tw}(G') > k$ by Lemma 4.3. Note that (T, β, α) is a tame OCP-tree-decomposition of (G', γ_1) of t-width $f_{4.4}(k)$. Then by Lemma 4.3, we can compute a tame OCP-tree-decomposition of (G, γ) of the same t-width.

We can augment the decomposition such that $|V(T)| \in \mathcal{O}(n)$ as follows. For any edge $t_1t_2 \in E(T)$ such that $\beta(t_1) \subseteq \beta(t_2)$, we contract the edge t_1t_2 in T to create the new vertex t . We then set $\beta(t) = \beta(t_2)$ and

$\alpha(t) = \alpha(t_2)$. Note that for any vertex t' which was originally adjacent to t_1 in T , $\beta(t_1) \cap \beta(t') = \beta(t_2) \cap \beta(t')$, and so the resulting decomposition still satisfies $|(\beta(t) \cap \beta(t')) \setminus \alpha(t)| \leq 1$ for all t' adjacent to t . The resulting decomposition has $|V(T)| \in \mathcal{O}(n)$. To see this, root T at an arbitrary vertex r . It follows that for any vertex $t \in V(T)$ which is not the root, the bag of t contains a new vertex v not contained in the bag of t 's parent such that $\{x \in V(T) : v \in \beta(x)\}$ is contained in the subtree rooted at t . This defines an injection from $V(T) \setminus \{r\}$ to $V(G)$, and so $|V(T)| \in \mathcal{O}(n)$. \square

Corollary 4.7. There exists a polynomial $f_{4.5} : \mathbb{N} \rightarrow \mathbb{N}$ such that for every positive integer k and every signed graph (G, γ) ,

1. if (G, γ) contains (\mathbf{H}_k, γ_1) or (\mathbf{V}_k, γ_1) as a signed graph minor, then $\text{tOCP-tw}(G, \gamma) \geq k/2 - 1$, and
2. if $\text{tOCP-tw}(G, \gamma) \geq f_{4.5}(k)$, then (G, γ) contains (\mathbf{H}_k, γ_1) or (\mathbf{V}_k, γ_1) as a signed graph minor.

Proof. Let G' be the graph obtained from (G, γ) by subdividing every even edge exactly once. Then by Lemma 4.3, $\text{tOCP-tw}(G') = \text{tOCP-tw}(G', \gamma_1) = \text{tOCP-tw}(G, \gamma)$. The result then follows immediately from Theorem 4.5 and Observation 2.6. \square

Fiorini et. al. [FJWY25] gave a connection between solving integer programs over matrices in $\{-1, 0, 1\}^{m \times n}$ with two nonzero entries per row and solving MAXIMUM WEIGHTED INDEPENDENT SET. Choi et. al. [CGK⁺25] used this to solve integer programs with coefficient matrix A such that the associated signed graph $G^+(A)$ has bounded (tame) OCP-treewidth.

Theorem 4.8 (Theorem 1.3 [CGK⁺25]). Let $A \in \{-1, 0, 1\}^{m \times n}$ have two nonzero entries per row such that the associated signed graph has tOCP-treewidth at most k . Then there exists a computable function $f_{4.8}(k)$ such that we can solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

in $n^{f_{4.8}(k)}$ time, where $w \in \mathbb{Z}^n$, $b \in \mathbb{Z}^m$, and $\ell, u \in (\mathbb{Z} \cup \{-\infty, \infty\})^n$.

Our main algorithmic result Theorem 1.3 is an extension of Theorem 4.8 to when A has entries outside of $\{-1, 0, 1\}$.

5 Properties of TDM-treewidth

In this section we show how TDM-treewidth is a combination of K -free treewidth and tOCP-treewidth. We use these connections to prove Theorem 1.6 and give an FPT-approximation for the TDM-treewidth of a graph. We later use these results to prove Theorem 1.3 in Section 6.

First, we show that TDM-treewidth is minor monotone with respect to rooted signed graphs.

Lemma 5.1. If (H, γ_H, K_H) is a minor of (G, γ_G, K_G) , then $\text{TDM-tw}(H, \gamma_H, K_H) \leq \text{TDM-tw}(G, \gamma_G, K_G)$.

Proof. We may assume that (H, γ_H, K_H) is formed from (G, γ_G, K_G) via a single vertex deletion, a single edge deletion, shifting at a single vertex, contracting one even edge, or removing a vertex from the set of roots. Let (T, β, α, J) be a TDM-tree-decomposition for (G, γ_G, K_G) . If (H, γ_H, K_H) is formed by an edge deletion, shifting, or removing a vertex from the set of roots, then (T, β, α, J) is a TDM-tree-decomposition for (H, γ_H, K_H) of the same width. Note that shifting does not change the odd cycle packing number of any subgraph. If (H, γ_H, K_H) is formed by vertex deletion, then by removing that vertex from the image of $\beta(t), \alpha(t)$ for every $t \in V(T)$, we obtain a TDM-tree-decomposition for (H, γ_H, K_H) of only smaller width. Suppose that (H, γ_H, K_H) is formed by contracting the edge uv to create the new vertex v' . We then set replace each instance of u or v in any of $\beta(t), \alpha(t)$ with v' in order to obtain a TDM-tree-decomposition of (H, γ_H, K_H) of only smaller width. Note that contracting an even edge can only decrease the odd cycle packing number. \square

We now more explicitly state the relation between the width parameters discussed thus far.

Lemma 5.2. Let (G, γ, K) be a rooted signed graph. Then we have the following relationship between parameters.

$$\max\{\text{tw}_K(G, K) + 1, \text{tOCP-tw}(G, \gamma)\} \leq \text{TDM-tw}(G, \gamma, K) \leq \text{tw}_K(G, K) + 1 + \text{tOCP-tw}(G, \gamma)$$

Proof. Let (J, β_J, L) be a tree K -free-decomposition of (G, K) of width at most $k - 1$ and suppose that $\text{tOCP-tw}(G, \gamma) = r$. For every leaf j of J , $\text{tOCP-tw}(G[\beta(j) \cap L], \gamma) \leq \text{tOCP-tw}(G, \gamma) = r$. Let $\ell(J)$ denote the set of leaves of J . For each $j \in \ell(J)$, let (T_j, β_j, α_j) be a tame OCP-tree-decomposition of $(G[\beta(j) \cap L], \gamma)$ of t-width at most r . We now construct the TDM-tree-decomposition of (G, γ, K) as follows. Let T be a tree formed from the disjoint union of J and T_j for $j \in \ell(J)$ by adding an edge from each leaf $j \in \ell(J)$ to an arbitrary vertex in T_j . For each $j \in V(J)$, let $\beta(j) = \alpha(j) = \beta_J(j) \setminus L$. For each $t \in V(T_j)$, let $\beta(t) = \beta_j(t) \cup (\beta_J(j) \setminus L)$ and $\alpha(t) = \alpha_j(t) \cup (\beta_J(j) \setminus L)$. Then (T, β, α, J) is a TDM-tree-decomposition of (G, γ, K) of width at most $k + r$.

Suppose now that (T, β, α, J) is a TDM-tree-decomposition of (G, γ, K) of width at most k . Then (T, β, α) is a tame OCP-tree-decomposition of (G, γ) of t-width at most k , so it suffices to show $\text{tw}_K(G, K) \leq k - 1$. We augment (T, β, α, J) as follows. For every $j \in V(J) \setminus \ell(J)$ with $\beta(j) \neq \alpha(j)$, we add a new leaf t to J (and thus T) adjacent only to j such that $\beta(t) = \beta(j)$ and $\alpha(t) = \alpha(j)$. We then set $\beta(j)$ to be equal to $\alpha(j)$. After performing this augmentation we still have a TDM-tree-decomposition of width at most k but now with the extra property that $\beta(j) = \alpha(j)$ for every $j \in V(J) \setminus \ell(J)$. We then take $L = V(G) \setminus \bigcup_{j \in V(J)} \alpha(j)$. Root T at an arbitrary vertex in $V(J)$. For $j \in \ell(J)$, let T_j be the subtree of T consisting of all vertices whose unique path to the root includes j and let $\beta'(j) = \bigcup_{t \in V(T_j)} \beta(t)$. For $j \in V(J) \setminus \ell(J)$, let $\beta'(j) = \beta(j)$. Then (J, β', L) is a tree K -free-decomposition of (G, K) of width at most $k - 1$. \square

The above proof naturally gives us an FPT-approximation for $\text{TDM-tw}(G, \gamma, K)$.

Theorem 5.3. There exists an algorithm that, given an n -vertex, m -edge rooted signed graph (G, γ, K) and a non-negative integer k , either computes a TDM-tree-decomposition (T, β, α, J) of (G, γ, K) of width at most $f_{5.3}(k)$ with $|V(T)| \in \mathcal{O}(n)$, or correctly returns that $\text{TDM-tw}(G, \gamma, K) > k$. The algorithm runs in time $g(k)\text{poly}(m, n)$. Furthermore $f_{5.3}(k) \in \text{poly}(k)$.

Proof. We first apply Theorem 3.2 to either obtain a tree K -free-decomposition (J, β_J, L) of (G, K) of width at most $5k$ with $|J| \in \mathcal{O}(n)$, or we correctly determine that $\text{tw}_K(G, K) > k - 1$. By Lemma 5.2 we have $\text{TDM-tw}(G, \gamma, K) \geq \text{tw}_K(G, K) + 1$, and so in the second case we may correctly return that $\text{TDM-tw}(G, \gamma, K) > k$.

Hence we may assume that we obtain the tree K -free-decomposition (J, β_J, L) . Then for each leaf $j \in V(J)$, we apply Corollary 4.6 to either obtain a tame OCP-tree-decomposition of $(G[\beta(j) \cap L], \gamma)$ of width at most $f_{4.4}(k)$, or we correctly determine that $\text{tOCP-tw}(G[\beta(j) \cap L], \gamma) > k$. Again in the second case we correctly determine that $\text{TDM-tw}(G, \gamma, K) \geq \text{tOCP-tw}(G, \gamma) \geq \text{tOCP-tw}(G[\beta(j) \cap L], \gamma) > k$ by Lemma 5.2. Thus we may assume that we obtain a tame OCP-tree-decomposition (T_j, β_j, α_j) of $(G[\beta(j) \cap L], \gamma)$ of width at most k and with $|V(T_j)| \in \mathcal{O}(|\beta(j) \cap L|)$. The number of vertices over all such trees T_j is in $\mathcal{O}(|L|)$. We then construct a TDM-tree-decomposition (T, β, α, J) of (G, γ, K) as in the proof of Lemma 5.2. The resulting decomposition has width at most $5k + 1 + f_{4.4}(k)$ and $|V(T)| \in \mathcal{O}(n)$. \square

Lemma 5.2 and Corollary 2.8 immediately imply Theorem 1.6, which we restate below for convenience.

Theorem 1.6. There exists a polynomial $f : \mathbb{N} \rightarrow \mathbb{N}$ such that for every integer $k \geq 1$ and every rooted signed graph (G, γ, K) ,

1. if (G, γ, K) contains one of $\mathcal{H}_k, \mathcal{V}_k$, or \mathcal{W}_k as a minor, then $\text{TDM-tw}(G, \gamma, K) \geq \Omega(k)$, and
2. if $\text{TDM-tw}(G, \gamma, K) \geq f(k)$, then G contains one of $\mathcal{H}_k, \mathcal{V}_k$, or \mathcal{W}_k as a minor.

Proof. Suppose that (G, γ, K) forbids $\mathcal{H}_k, \mathcal{V}_k$, and \mathcal{W}_k as a minor. By Corollary 2.8, (G, K) forbids $(\mathbf{W}_{k^2}, K_{\mathbf{W}_{k^2}})$ as a rooted minor. Then by Theorem 3.3, $\text{tw}_K(G, K) \leq f_{3.3}(k^2)$. Similarly by Corollary 4.7, $\text{tOCP-tw}(G, \gamma) \leq f_{4.5}(k)$. Hence the result follows from Lemma 5.2.

Suppose instead that (G, γ, K) contains one of $\mathcal{H}_k, \mathcal{V}_k$, or \mathcal{W}_k as a minor. If (G, γ, K) contains \mathcal{W}_k as a minor, then (G, K) contains $(\mathbf{W}_k, K_{\mathbf{W}_k})$ as a minor. Then by Lemma 3.5 and Lemma 5.2, $\text{TDM-tw}(G, \gamma, K) \geq \text{tw}_K(G) + 1 \geq \Omega(k)$. If (G, γ, K) contains one of $\mathcal{H}_k, \mathcal{V}_k$ as a minor, then (G, γ) contains one of $(\mathbf{H}_k, \gamma_1), (\mathbf{V}_k, \gamma_1)$ as a signed graph minor. Thus by Corollary 4.7 and Lemma 5.2 we have $\text{TDM-tw}(G, \gamma, K) \geq \text{tOCP-tw}(G, \gamma) \geq \Omega(k)$. \square

6 Solving integer programs with large entries

In this section we prove Theorem 1.3, which seeks to extend Theorem 4.8 to matrices with entries outside $\{-1, 0, 1\}$ under condition that variables have bounded domain size, and there is no “highly linked” set of columns with entries outside $\{-1, 0, 1\}$. We restate Theorem 1.3 below for convenience.

Theorem 1.3. Let d, k be nonnegative integers, and let A be a matrix with two nonzero entries per row whose associated rooted signed graph has TDM-treewidth at most k . Then for any $w \in \mathbb{Z}^n, b \in \mathbb{Z}^m$, and $\ell, u \in \mathbb{Z}^m$ with $\|u - \ell\|_\infty \leq d$, we can solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

in $d^{\mathcal{O}(k)} n^{f(k)}$ time for some computable f .

Proof. Let $G_+^+(A) = (G, \gamma, K)$. For each $v \in V(G)$, let x_v, ℓ_v, u_v be the associated entries in x, ℓ, u respectively. By Lemma 5.2, $\text{tw}_K(G, K) \leq k - 1$. Then by Theorem 3.2, we obtain a tree K -free-decomposition (T, β, L) of (G, K) of width at most $5k$ with $|V(T)| \in \mathcal{O}(n)$ in time $2^{\mathcal{O}(k)} \text{poly}(n)$. We may assume that T contains a non-leaf vertex, otherwise we subdivide the single edge in T to create a new vertex t and set $\beta(t) = \beta(t') \setminus L$ for one of the other vertices t' of T . We root T at an arbitrary non-leaf vertex r . For each $t \in V(T)$, we denote by T_t the subtree of T consisting of all vertices whose unique path to the root includes t . We denote by IP_t the above integer program restricted only to variables x_v for $v \in \bigcup_{t' \in V(T_t)} \beta(t')$. We keep the constraints which include only the variables we restricted to.

We construct a dynamic programming table $p[t, \eta_t]$ where for each $t \in V(T)$ and each assignment $\eta_t : \{x_v : v \in \beta(t) \setminus L\} \rightarrow \mathbb{Z}$ of x_v to a value in $[\ell_v, u_v]$, $p[t, \eta_t]$ is the optimal objective value of IP_t under the partial assignment η_t , or we store $-\infty$ if the partial assignment can not be extended to a feasible solution. We note that $p[t, \eta_t]$ is finite if the partial assignment can be extended to a feasible solution. We compute $p[t, \eta_t]$ from the leaves of T towards the root. The final solution is given by

$$\max_{\eta_r} p[r, \eta_r].$$

As there are d^{5k} choices for η_r , this can be computed in time $\mathcal{O}(d^{5k})$.

Fix a vertex $t \in V(T)$ and fix η_t . If t is a leaf, then because

$$\text{tOCP-tw}(G[\beta(t) \cap L], \gamma) \leq \text{tOCP-tw}(G, \gamma) \leq \text{TDM-tw}(G, \gamma, K) \leq k$$

by Lemma 5.2, we can compute $p[t, \eta_t]$ in $n^{f_{4.8}(k)}$ time by Theorem 4.8.

We may then assume that t is not a leaf and we have computed $p[t', \eta_{t'}]$ for every t' a child of t and every $\eta_{t'}$. For each child t' of t , we compute

$$s[t', \eta_t] = \max \left\{ p[t', \eta_{t'}] - \sum_{v \in \beta(t') \cap \beta(t)} w_v \eta_{t'}(x_v) : \eta_{t'}(x_v) = \eta_t(x_v) \text{ for all } v \in \beta(t) \cap \beta(t') \right\}.$$

We note that $s[t', \eta_t]$ is uniquely determined by η_t restricted to $\beta(t) \cap \beta(t')$. Thus we can compute $s[t', \eta_t]$ for all η_t when we compute $p[t', \eta_{t'}]$ for all $\eta_{t'}$ in time $\mathcal{O}_k(d^{5k})$. This can be done by iterating over all choices

of η_t restricted to $\beta(t) \cap \beta(t')$, and then for each choice iterating over all ways to extend it to $\eta_{t'}$ by choosing the assignment of x_v for $v \in \beta(t') \setminus \beta(t)$.

We then set

$$p[t, \eta] = \sum_{v \in \beta(t)} w_v \eta(x_v) + \sum_{t' \text{ child of } t} s[t', \eta].$$

if the assignment η does not violate any constraint induced by the bag of t , otherwise we set $p[t, \eta] = -\infty$. The correctness of the above algorithm follows immediately by induction. \square

7 Further discussion

In this section we discuss some of the assumptions made in various theorems and whether they can be relaxed.

7.1 On $\{0, 1\}$ -variables and the definition of TDM-treewidth

The definition of a TDM-tree-decomposition is rather technical and relies on specific properties of the adhesions. Perhaps it would be more natural to require all adhesions to have some bounded size outside of the protector set, or to not require some protectors to be strong. The reason for having an adhesion of size 1 outside of a hitting set is due to $\{0, 1\}$ -variables. Suppose all variables take values in $\{0, 1\}$, and we are doing dynamic programming on the tree decomposition. At some bag $\beta(t)$ for $t \in V(T)$, we guess the value of variables in $\alpha(t)$ and the value of the adhesion with the parent bag. However there may be an unbounded number of children bags, and for each child t' we must decide the value to take on $(\beta(t) \cap \beta(t')) \setminus \alpha(t)$. If this set has size 1, say it corresponds to variable x_v , then there's a nice trick: add a new variable $y_v \in \{0, 1\}$, add the constraint $x_v + y_v \leq 1$, set $w(x_v)$ equal to the optimal value of the subtree with root t' if $x_v = 1$, and set $w(y_v)$ equal to the optimal value of the subtree if $x_v = 0$. We may assume $w(x_v), w(y_v) \geq 0$ by adding a constant to the final weight. The optimal solution will then set exactly one of x_v or y_v to 1, and adding this constraint doesn't affect that the resulting problem is totally Δ -modular. Hence we can make this construction for an unbounded number of child bags. If instead x_v can take more than two values, or $|(\beta(t) \cap \beta(t')) \setminus \alpha(t)| > 1$, it is unclear how to encode this in the integer program without possibly increasing Δ .

The reason we can allow $|(\beta(t) \cap \beta(t')) \setminus \alpha(t)| = 1$ for some bags in our TDM-tree-decomposition is because we can reduce to $\{0, 1\}$ -variables for some of the subproblems during the dynamic programming. In the proof of [Theorem 1.3](#), we applied [Theorem 4.8](#) as a black box. What's really happening is that whenever we're in a subtree disjoint from J , we can guess on $\alpha(t)$ such that the remaining integer program corresponding to the subtree has coefficients in $\{-1, 0, 1\}$. We can then solve the LP-relaxation for the entire subtree and use proximity results to make all variables take values in $\{0, 1\}$ (see [\[CGK⁺25, Theorem 10.3\]](#) based on [\[FJWY25, Section 3\]](#)). Then we can use the above trick to fill the dynamic programming table even when t has an unbounded number of children with adhesions of size 1.

The above discussion implies, for instance, that if we know that all variables are in $\{0, 1\}$, we could solve the integer program even when no protectors are strong. This would correspond to a tame OCP-tree-decomposition (T, β, α) such that $K \cap \beta(t) \subseteq \alpha(t)$ for all $t \in V(T)$ because we may now take $J = T$. This leads to the following question, which asks for a grid theorem for such decompositions.

Question 7.1. What are the forbidden minors for the class of rooted signed graphs (G, γ, K) which have a tame OCP-tree-decomposition (T, β, α) of bounded width such that $K \cap \beta(t) \subseteq \alpha(t)$ for all $t \in V(T)$?

7.2 Bounded entries and unbounded variables

In [Theorem 1.3](#) we assume that $\|u - \ell\|_\infty$ is bounded, but we allow $\|A\|_\infty$ to be unbounded. Bounding $\|A\|_\infty$ is perhaps more natural from the perspective of totally Δ -modular matrices, as this would imply that the bags of the TDM-tree-decomposition are totally Δ -modular. This leads to the following question.

Question 7.2. Fix integers $d, k \geq 0$. Let A be a matrix with two nonzero entries per row such that the associated rooted signed graph has TDM-treewidth at most k . Let $w \in \mathbb{Z}^n, b \in \mathbb{Z}^m$, and $\ell, u \in (\mathbb{Z} \cup \{-\infty, \infty\})^n$. If $\|A\|_\infty \leq d$, can we solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

in polynomial time for constant d, k ?

In fact, something much simpler is not known.

Conjecture 7.3. Fix a positive integer d . Let A be a matrix with exactly two nonzero entries per row such that $G(A)$ is a tree and $\|A\|_\infty \leq d$. Then the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

can be solved in polynomial time for constant d .

If the tree $G(A)$ has constant depth, the [Conjecture 7.3](#) is implied by results on bounded treedepth (see, for instance, [\[EHK⁺19\]](#)). However [Conjecture 7.3](#) is not known even for $G(A)$ a path. The path case is perhaps particularly interesting due to the recent result of Briański et. al. [\[BLP⁺25\]](#) which shows that integer programming is NP-hard even when every non-zero coefficient appears in at most two consecutive constraints and $\|A\|_\infty \leq 8$. The resulting matrix has a “path-like” structure, but crucially may contain 3 nonzero entries per row. Due to the lack of hardness results for integer programs with at most two nonzero entries per row, it is not clear whether we must bound one of $\|u - \ell\|_\infty$ or $\|A\|_\infty$. It could be that [Conjecture 7.3](#) is true without the bound on $\|A\|_\infty$.

Another interesting question is whether the rooted grid in [Corollary 1.7](#) must be forbidden. In fact, it is unclear whether a grid with every vertex in the set of roots must be forbidden. That is, is there a hardness result for integer programs where large entries are everywhere, but the corresponding signed graph has no odd cycles?

Question 7.4. Is it NP-hard to solve the integer program

$$\max\{w^T x : Ax \leq b, x \in [\ell, u] \cap \mathbb{Z}^n\}$$

over instances where $G_\bullet^+(A) = (G, \gamma, K)$ is planar, has no odd cycles, but K is arbitrary?

The above question is unknown even for $G_\bullet^+(A)$ not necessarily planar. Any hardness result for integer programs with at most two nonzero entries per row that isn’t the standard reduction to MAXIMUM WEIGHTED INDEPENDENT SET would be interesting.

Acknowledgments. The author would like to thank Maximilian Gorsky, Rose McCarty, and Sebastian Wiederrecht for valuable discussions.

References

- [Ale04] Vladimir E Alekseev. Polynomial algorithm for finding the largest independent sets in graphs without forks. *Discrete Applied Mathematics*, 135(1-3):3–16, 2004.
- [AWZ17] Stephan Artmann, Robert Weismantel, and Rico Zenklusen. A strongly polynomial algorithm for bimodular integer linear programming. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pages 1206–1219, 2017.
- [BKP⁺24] Marcin Briański, Martin Koutecký, Kristýna Pekárková, Felix Schröder, et al. Characterization of matrices with bounded graver bases and depth parameters and applications to integer programming. *Mathematical Programming*, 208(1-2):497, 2024.

[BLP⁺25] Marcin Briański, Alexandra Lassota, Kristýna Pekárková, Michał Pilipczuk, and Janina Reuter. On integer programs that look like paths. *arXiv preprint arXiv:2510.22430*, 2025.

[CEH⁺21] Jana Cslovjecsek, Friedrich Eisenbrand, Christoph Hunkenschröder, Lars Rohwedder, and Robert Weismantel. Block-structured integer and linear programming in strongly polynomial and near linear time. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1666–1681. SIAM, 2021.

[CEP⁺20] Jana Cslovjecsek, Friedrich Eisenbrand, Michał Pilipczuk, Moritz Venzin, and Robert Weismantel. Efficient sequential and parallel algorithms for multistage stochastic integer programming using proximity. *arXiv preprint arXiv:2012.11742*, 2020.

[CG07] William H Cunningham and Jim Geelen. On integer programming and the branch-width of the constraint matrix. In *International conference on integer programming and combinatorial optimization*, pages 158–166. Springer, 2007.

[CGK⁺25] Mujin Choi, Maximilian Gorsky, Gunwoo Kim, Caleb McFarland, and Sebastian Wiederrecht. Odd-cycle-packing-treewidth: On the maximum independent set problem in odd-minor-free graph classes. *arXiv preprint arXiv:2511.10019*, 2025.

[CGST86] William Cook, Albertus MH Gerards, Alexander Schrijver, and Éva Tardos. Sensitivity theorems in integer linear programming. *Mathematical Programming*, 34(3):251–264, 1986.

[CKL⁺25] Jana Cslovjecsek, Martin Koutecký, Alexandra Lassota, Michał Pilipczuk, and Adam Polak. Parameterized algorithms for block-structured integer programs with large entries. *TheoretCS*, 4, 2025.

[CKPW22] Marcel Celaya, Stefan Kuhlmann, Joseph Paat, and Robert Weismantel. Improving the cook et al. proximity bound given integral valued constraints. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 84–97. Springer, 2022.

[CLB81] Derek G Corneil, Helmut Lerchs, and L Stewart Burlingham. Complement reducible graphs. *Discrete Applied Mathematics*, 3(3):163–174, 1981.

[Dad12] Daniel Dadush. *Integer programming, lattice algorithms, and deterministic volume estimation*. PhD thesis, Georgia Institute of Technology, 2012.

[DEG⁺21] Pavel Dvořák, Eduard Eiben, Robert Ganian, Dušan Knop, and Sebastian Ordyniak. The complexity landscape of decompositional parameters for ILP: Programs with few global variables and constraints. *Artificial Intelligence*, 300:103561, 2021.

[EGHK21] Eduard Eiben, Robert Ganian, Thekla Hamm, and O-joung Kwon. Measuring what matters: A hybrid approach to dynamic programming with treewidth. *Journal of Computer and System Sciences*, 121:57–75, 2021.

[EGK⁺19] Eduard Eiben, Robert Ganian, Dušan Knop, Sebastian Ordyniak, Michał Pilipczuk, and Marcin Wrochna. Integer programming and incidence treedepth. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 194–204. Springer, 2019.

[EHK⁺19] Friedrich Eisenbrand, Christoph Hunkenschröder, Kim-Manuel Klein, Martin Koutecký, Asaf Levin, and Shmuel Onn. An algorithmic theory of integer programming. *arXiv preprint arXiv:1904.01361*, 2019.

[EW19] Friedrich Eisenbrand and Robert Weismantel. Proximity results and faster algorithms for integer programming using the Steinitz lemma. *ACM Transactions on Algorithms (TALG)*, 16(1):1–14, 2019.

[FHT93] Martin Farber, Mihály Hujter, and Zsolt Tuza. An upper bound on the number of cliques in a graph. *Networks*, 23(3):207–210, 1993.

[FJWY25] Samuel Fiorini, Gwenaël Joret, Stefan Weltge, and Yelena Yuditsky. Integer programs with bounded subdeterminants and two nonzeros per row. *J. ACM*, 72(1):Art. 3, 50, 2025.

[Fre90] Eugene C. Freuder. Complexity of k -tree structured constraint satisfaction problems. In *Proceedings of the Eighth National Conference on Artificial Intelligence - Volume 1*, AAAI'90, page 4–9. AAAI Press, 1990.

[Ger89] AMH Gerards. A min-max relation for stable sets in graphs with no odd- K_4 . *Journal of Combinatorial Theory, Series B*, 47(3):330–348, 1989.

[GGR⁺09] Jim Geelen, Bert Gerards, Bruce Reed, Paul Seymour, and Adrian Vetta. On the odd-minor variant of Hadwiger’s conjecture. *Journal of Combinatorial Theory, Series B*, 99(1):20–29, 2009.

[GLS84] Martin Grötschel, László Lovász, and Alexander Schrijver. Polynomial algorithms for perfect graphs. In *North-Holland mathematics studies*, volume 88, pages 325–356. Elsevier, 1984.

[GO18] Robert Ganian and Sebastian Ordyniak. The complexity landscape of decompositional parameters for ILP. *Artificial Intelligence*, 257:61–71, 2018.

[GOR17] Robert Ganian, Sebastian Ordyniak, and M Ramanujan. Going beyond primal treewidth for (M) ILP. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 31, 2017.

[GW23] J Pascal Gollin and Sebastian Wiederrecht. Structure and algorithms for graphs excluding grids with small parity breaks as odd-minors. *arXiv preprint arXiv:2304.04504*, 2023.

[HLMR24] Jędrzej Hodor, Hoang La, Piotr Micek, and Clément Rambaud. Quickly excluding an apex-forest. *arXiv preprint arXiv:2404.17306*, 2024.

[JdK21] Bart MP Jansen and Jari JH de Kroon. FPT algorithms to compute the elimination distance to bipartite graphs and more. In *International Workshop on Graph-Theoretic Concepts in Computer Science*, pages 80–93. Springer, 2021.

[JDKW21] Bart MP Jansen, Jari JH De Kroon, and Michał Włodarczyk. Vertex deletion parameterized by elimination distance and even less. In *Proceedings of the 53rd annual ACM SIGACT symposium on theory of computing*, pages 1757–1769, 2021.

[JK15] Bart MP Jansen and Stefan Kratsch. A structural approach to kernels for ILPs: Treewidth and total unimodularity. In *Algorithms-ESA 2015: 23rd Annual European Symposium, Patras, Greece, September 14–16, 2015, Proceedings*, pages 779–791. Springer, 2015.

[JMST25] Lars Jaffke, Laure Morelle, Ignasi Sau, and Dimitrios M Thilikos. Dynamic programming on bipartite tree decompositions. *Journal of Computer and System Sciences*, page 103722, 2025.

[JS24] Bart MP Jansen and Céline MF Swennenhuis. Steiner tree parameterized by multiway cut and even less. *arXiv preprint arXiv:2406.19819*, 2024.

[Kan87] Ravi Kannan. Minkowski’s convex body theorem and integer programming. *Mathematics of operations research*, 12(3):415–440, 1987.

[Kob25] Stefan Kober. Totally δ -modular ips with two non-zeros in most rows. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 355–370. Springer, 2025.

[KPW20] Dušan Knop, Michał Pilipczuk, and Marcin Wrochna. Tight complexity lower bounds for integer linear programming with few constraints. *ACM Transactions on Computation Theory (TOCT)*, 12(3):1–19, 2020.

[KSSW25] Marcus Kühn, Lisa Sauermann, Raphael Steiner, and Yuval Wigderson. Disproof of the Odd Hadwiger Conjecture. *arXiv preprint arXiv:2512.20392*, 2025.

[LJ83] Hendrik W Lenstra Jr. Integer programming with a fixed number of variables. *Mathematics of operations research*, 8(4):538–548, 1983.

[LM08] Vadim V Lozin and Martin Milanič. A polynomial algorithm to find an independent set of maximum weight in a fork-free graph. *Journal of Discrete Algorithms*, 6(4):595–604, 2008.

[Min80] George J Minty. On maximal independent sets of vertices in claw-free graphs. *Journal of Combinatorial Theory, Series B*, 28(3):284–304, 1980.

[MSW17] Daniel Marx, Paul Seymour, and Paul Wollan. Rooted grid minors. *Journal of Combinatorial Theory, Series B*, 122:428–437, 2017.

[Pap81] Christos H Papadimitriou. On the complexity of integer programming. *Journal of the ACM (JACM)*, 28(4):765–768, 1981.

[PTW25] Evangelos Protopapas, Dimitrios M Thilikos, and Sebastian Wiederrecht. Colorful minors. *arXiv preprint arXiv:2507.10467*, 2025.

[PWW20] Joseph Paat, Robert Weismantel, and Stefan Weltge. Distances between optimal solutions of mixed-integer programs. *Mathematical Programming*, 179(1):455–468, 2020.

[RS86] Neil Robertson and Paul D Seymour. Graph minors. V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41(1):92–114, 1986.

[Sch03] A. Schrijver. *Combinatorial Optimization: Polyhedra and Efficiency*. Number 24 in Algorithms and Combinatorics. Springer, Berlin ; New York, 2003.

[She96] Valery N Shevchenko. *Qualitative topics in integer linear programming*. American Mathematical Soc., 1996.

[SST25] Ignasi Sau, Giannos Stamoulis, and Dimitrios M Thilikos. Parameterizing the quantification of CMSO: model checking on minor-closed graph classes. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 3728–3742. SIAM, 2025.

[Ste22] Raphael Steiner. Asymptotic equivalence of Hadwiger’s conjecture and its odd minor-variant. *Journal of Combinatorial Theory, Series B*, 155:45–51, 2022.

[Taz12] Siamak Tazari. Faster approximation schemes and parameterized algorithms on (odd-) H -minor-free graphs. *Theoretical Computer Science*, 417:95–107, 2012.

[Zas82] Thomas Zaslavsky. Signed graphs. *Discrete Applied Mathematics*, 4(1):47–74, 1982.

[Zas91] Thomas Zaslavsky. Orientation of signed graphs. *European Journal of Combinatorics*, 12(4):361–375, 1991.

[Zas13] Thomas Zaslavsky. Signed graphs and geometry. *arXiv preprint arXiv:1303.2770*, 2013.