

Unlocking the dynamics of Young Stellar Objects: Time-Domain Interferometry with six 4-m class telescopes

A. Soulain¹, B. Lopez¹, A. Matter¹, F. Lykou², P. Boley³, M. Scheuck³, R. van Boekel³, J.-C. Augereau⁴, M. leTessier⁴, J. Bouvier⁴, P. Berio¹, P. Ábrahám², N. Anugu¹³, J.-P. Berger⁴, R. Burn¹, W.-C Danchi⁵, W.J. de Wit⁶, F. Drewes¹, V. Fleury¹, V. Hocdé¹², W. Jaffe⁷, Á. Kóspál^{2,4}, E. Koumpia^{6,8}, J.-B. Lebouquin⁴, J. S. Martin¹⁰, H. Meheut¹, F. Millour¹, N. Nardetto¹, E. Pantin¹¹, K. Perraut⁴, R. Petrov¹, L.N.A van Haastere⁷, J. Varga², G. Weigelt⁹, et S. Wolf¹⁰

¹Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, Nice, France

²Konkoly Observatory, HUN-REN Research Centre for Astronomy and Earth Sciences, Konkoly Thege Miklós út 15-17, 1121 Budapest, Hungary

³Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany

⁴Université Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France

⁵NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁶ESO, Alonso de Córdova 3107, Vitacura, Casilla, 19001, Santiago, Chile

⁹Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121, Bonn, Germany

⁷Leiden Observatory, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

⁸ALMA Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile

¹⁰Institut für Theoretische Physik und Astrophysik, Christian-Albrechts-Universität zu Kiel, Leibnizstraße 15, 24118 Kiel, Germany

¹¹Université Paris-Saclay, Université Paris Cité CEA, CNRS, AIM, 91191, Gif-sur-Yvette, France

¹²Nicolaus Copernicus Astronomical Centre, Polish Academy of Sciences, Bartycka 18, 00-716 Warszawa, Poland

¹³The CHARA Array of Georgia State University, Mount Wilson Observatory, Mount Wilson, CA 91023, USA

Unlocking the Dynamics of Young Stellar Objects

Abstract

The dynamics of the inner regions of young stellar objects (YSOs) is driven by a variety of physical phenomena, from magnetospheres and accretion to the dust sublimation rim and inner disk flows. These inner environments evolve on timescales of hours to days, exactly when bursts, dips, and rapid structural changes carry the most valuable information about star and planet formations, but remain hardly reachable with current facilities. A better reactive infrastructure with six or more telescopes, combined with alerts from large time-domain surveys (e.g., at the era of LSST/Rubin type facilities), and equipped with instruments spanning from the *V*-band to the thermal infrared (*N*), would provide the instantaneous uv-coverage and spectral diagnostics needed to unambiguously interpret and image these events as they happen. Such a world's first time-domain interferometric observatory would enable qualitatively new science: directly linking optical and infrared variability to spatially resolved changes in magnetospheric accretion, inner-disk geometry, and dust and gas dynamics in the innermost astronomical unit. Crucially, connecting these processes to outer-scale unresolved information from JWST, ALMA, and the ELT would yield a complete tomography of the planet-forming region.

1 The dynamic inner au: Resolving the physics of star/planet formation in 4D

The innermost au scales of Young Stellar Objects (YSOs) is a complex interplay between magnetic fields, gas flows, and dust dynamics. While current optical interferometry has successfully resolved the static architecture of inner disks and magnetospheres, we possess limited spatially resolved information on the dynamic processes driving their evolution. Figure 1 illustrates the vast dynamic range of YSO variability in both timescales and amplitudes. We structure our science case around four key regimes where resolving these temporal signatures spatially is critical.

1.1 Unraveling the accretion/ejection mechanism

The magnetosphere governs the early evolution of stars by controlling angular-momentum loss and channeling accretion (Hartmann et al. 2016). However, the coupling between the star and the disk is highly unstable (Pantolmos et al. 2020; Zhu 2025). We observe accretion bursts, rapid photometric dips, jets, winds and stochastic variability, but we cannot currently see the structural changes that cause them.

- **The geometry of bursts:** Do accretion bursts result from large-scale disk instabilities or magnetic events?
- **Accretion in close binaries:** Resolving how binarity regulates accretion variability and shapes architecture is critical, as most stars form in multiple systems.
- **Magnetic funnels:** How do accretion funnels migrate and reconfigure on rotational timescales?
- **The “black box” of spectroscopy:** While it tells us *how fast* gas is moving, it cannot tell us *where* it is flowing.

1.2 The "Dipper" mystery: warps, clumps, or winds?

A large fraction of disk-bearing young stars exhibit day-to-week long quasi-periodic drops in optical brightness, suspected to be due to a dusty warp in the inner disk (dipper phenomenon, Roggero et al. 2021; Nagel et al. 2024). However, without time-resolved imaging, we cannot confirm the origin of these dips. Is the extinction caused by a coherent geometric warp, a transient dust lift, or a dusty disk wind? Understanding Dippers is crucial as they likely represent the standard geometry of the star-disk interaction, yet their dynamic inner structure remains spatially unresolved during the dimming events.

1.3 Transient signatures of planet formation

Beyond the inner rim, the planet-forming region is rich in transient phenomena that evolve on daily to monthly timescales.

- **Dynamics of disks:** Tracking the motion of vortices and dust traps triggered by the disk evolution or planet-disk interactions (Varga et al. 2021; Kuo et al. 2024).
- **Circumplanetary Disks (CPDs):** The formation of giant planets involves their own accretion disks. Detecting the moving variable thermal emission of a CPD against the stellar glare would provide a direct probe into the birth of gas giants (Benisty et al. 2021; Zhou et al. 2025).
- **Dust evolution and the ‘Cosmic Furnace’:** We observe sublimation cycles, re-condensation, and crystallization sites (e.g., in EX Lupi outbursts, Kóspál et al. 2023). Spatially resolving these ‘dust reactors’ is crucial to understand how the ISM is turned into planetary building blocks (CAIs, chondrules), ultimately setting the chemical budget inherited by planets (Krijger et al. 2020; Morbidelli et al. 2024).

1.4 Extreme variability: The FUor/EXor eruptions

Eruptive stars (FUors/EXors) represent the most violent manifestation of YSO dynamics which can occur for class I eruptions during a short time, with brightness increases up to 6 magnitudes (Contreras Peña et al. 2024), on timescales shorter than one year (Laznevoi et al. 2025). These events induce profound structural changes that only reactive interferometry can resolve:

- **Magnetospheric Crushing:** High accretion rates ($> 10^{-5} M_{\odot}/\text{yr}$) theoretically “crush” the magnetosphere, forcing a transition from funnel-flow to boundary-layer accretion (Liu et al. 2022).
- **Thermal Restructuring:** The outburst launches a thermal wave, rapidly pushing the water snowline outward (Cieza et al. 2016) and triggering in-situ dust crystallization (Ábrahám et al. 2009).

While surveys have tripled the known population (Contreras Peña et al. 2024), the triggering mechanism remains debated (Nayakshin et al. 2023). We need to image these targets *during* their brightness rise to distinguish between a propagating instability and a localized impact.




Figure 1: YSOs exhibit diverse, rapid, multi-scale variability modes that challenge the temporal resolution of existing arrays (adapted from Fischer et al. 2023). Unveiling their origin requires capturing the inner geometry simultaneously with the observed variability.

1.5 The missing dimension: time

The fundamental barrier to understanding these processes is not just spatial resolution or sensitivity, but temporal resolution. These environments evolve on timescales of hours to days – exactly the domain where bursts, dips, and structural changes carry the most information. To understand the physics of star and planet formation, we must be able to image the geometry of gas and dust as it changes, linking the variable flux directly to the evolving structure.

1.6 Broader impact: A facility for the transient sky

While time-domain photometry (Gaia, ZTF, LSST) provides the alerting trigger and the period (Hodgkin et al. 2021), only time-domain high angular resolution interferometry can capture the changing geometry. Only high-cadence monitoring at high spatial resolution can turn light curves into physical radii in the case of novae expansion (Schaefer et al. 2014) or disentangle static envelopes from dynamic mass-loss (Cepheids, Hocdé et al. 2025). This versatility ensures that a ≥ 6 telescope array becomes a cornerstone of ESO’s time-domain ecosystem, extending its legacy value far beyond YSOs.

2 Sensitivity is not the bottleneck

While the VLTI has achieved remarkable success in resolving the static architecture of bright objects, the transition to time-domain astronomy reveals fundamental structural limitations. The issue is no longer just seeing faint targets – a frontier being pushed by GRAVITY+ (GRAVITY+ Collaboration 2022) – but rather observing them fast enough and often enough to disentangle between the different models hypothesis and constrain their physics.

2.1 The “Time-Blurring” barrier

Standard interferometric imaging relies on Earth-rotation synthesis to fill the uv -plane. However, for dynamic YSOs, this assumption of a static target breaks down. Integrating over a night or months effectively “blurs” these dynamical signatures. To freeze these events, we need to secure robust constraints on the object’s geometry on timescales shorter than the variability itself. This requires **snapshot imaging capabilities** which is physically impossible with only 4 telescopes and movable configurations spread over several weeks.

2.2 Operational rigidity & sensitivity

Operationally, the 8-m Unit Telescopes (UTs) are too over-subscribed (1 week/month availability) to support monitoring campaigns or rapid response to alerts (Manara et al. 2021). Conversely, the available 1.8-m Auxiliary Telescopes (ATs) or the CHARA array lack the sensitivity ($\sim 12\%$ of close-in YSOs, Fig. 2). We have to chose between sensitive telescopes that are unavailable and available telescopes that are not sensitive enough.

3 Golden era of Time-Domain astronomy

Looking toward the 2035+ horizon, the project is timely positioned to harvest the legacy of the Vera C. Rubin Observatory and the Roman Space Telescope. By then, these surveys will have cataloged the entire *zoo* of YSO variability, delivering millions of light curves (Ivezić et al. 2019).

3.1 The discovery vs. characterization gap

However, such facilities are discovery engines, not a physical characterization engine. It identifies *that* an event has occurred, but it cannot resolve its origin. A sudden brightening could be a magnetic reconnection event or a clump accretion; a dip could be a warped disk or a dust cloud. **To understand these events, we must spatially resolve them.** Capturing the instantaneous geometry of the accretion flow and the dynamical inner disk structure mandates a reactive array capable of immediate response to external alerts.

3.2 The Keystone of the 2040 Landscape

In the 2040 ecosystem, this facility acts as the critical link between the thermal jets and cold disks monitored by SKA and the X-ray flares detected by Athena. By resolving the central core, it serves not merely as a follow-up machine, but as the indispensable spatially-resolved counterpart to the next generation of Great Observatories (ELT, HWO).

4 Six (or more) 4-m class telescopes: Agility, sensitivity, and instantaneous uv -coverage

To resolve the dynamical machinery of YSOs, we propose a paradigm shift: either (1) upgrading the VLTI auxiliary array with six or more 4-m class telescopes operating on baselines > 200 m with visible-to-mid-IR instrumentation; or (2) developing a new interferometric network.

Interferometry: Achievements & 2030 horizon

Outburst-driven dust evolution

- ↔ Scales: Fraction of au to > 10 au
- Impact of episodic outbursts on mineralogy,
- Observing crystallization and sublimation fronts.
- Key bands:** Mid-IR.

Dust vorticities & planet embryos

- ↔ Scales: Fraction of au to 10 au
- Monitoring warps, clumps & shadows
- Tracing dust traps and planet-disk interactions.
- Key bands:** NIR, mid-IR, mm.

Accretion & magnetospheric processes

- ↔ Scales: 0.01 to 1 au
- Resolving Bry emission size & position
- Distinguishing origin of spectroscopic lines
- Key Lines:** H α , Pa β , HeI, Bry, Br α .




Figure 2: Accessible parameter space (G_{RP} vs K) for YSOs. The upgrade to 4-m class (violet) bridges the gap between current ATs (gray) and UTs (dark gray), unlocking hundreds of targets for time-domain interferometry.

The first option proposes deploying fixed or relocatable telescopes on existing AT stations. While exact specifications (count ≥ 6 , agility, diameter) are outside this work’s scope, we assume a baseline of six 4-m telescopes on the AT infrastructure, though our concepts apply generally. This combination unlocks three critical capabilities.

4.1 Snapshot imaging & sensitivity

Moving from 4 to 6 telescopes increases the number of simultaneous baselines from 6 to 15, and closure phases from 4 to 20. This dense instantaneous *uv*-coverage enables snapshot characterization, allowing us to secure robust geometric constraints in a single epoch without relying on Earth rotation. This effectively “freezes” the target’s geometry, solving the time-blurring problem (Kraus et al. 2020).

Simultaneously, the shift to 4-m apertures bridges the sensitivity gap (reaching within 1.5 mag of the UTs). This is critical to unlock the vast population of Class I/II stars in nearby star formation region (~ 820 sources, Fig. 2).

4.2 True operational reactivity

Unlike the multi-purpose UTs, a dedicated array of 4-m class telescopes offers flexible scheduling that allows for:

- High-cadence monitoring to track rotational modulation,
- Immediate reaction to photometric alerts (within hours),
- Tailored observations during specific phases of a burst.

This transforms the VLTI from a static mapper into a dynamic response engine.

4.3 6D Tomography: From V to N Bands

To physically interpret the variability, the facility must cover a broad spectral range. Each band probes a distinct physical component:

- **Visible (V): A High-Sensitivity Option.** While core diagnostics lie in the NIR, extending coverage to the visible remains highly desirable. The sheer strength of $H\alpha$ (Alcalá et al. 2017) provides a unique lever to detect the faintest sources and probe high-velocity gas (>100 km/s), offering a powerful complement to infrared tracers.
- **Near-IR (J, H, K): The Inner Environment.** This primary window reveals the immediate star-disk interaction: the hot dust sublimation rim (K), magnetospheric funnel flows (Pa β , Bry, Tessore et al. 2023; GRAVITY Collaboration et al. 2023), and ionized winds, effectively linking the accretion shock to the reprocessing dust.
- **Mid-IR (L, M, N): The Disk Structure.** These bands capture the cool dust emission, chemical composition,

dynamical vortices/planet embryos, water snowlines and vertical disk structure (Lopez et al. 2022).

By coupling this spectral coverage with a **rapid-response** operational model, the facility enables true **6D tomography**. We can directly link a burst detected in the optical to its thermal echo in the inner disk (IR), causally connecting the central engine to planet-forming environments.

5 The missing piece of the puzzle

An interferometer built around six or more 4-m class telescopes represents more than an incremental upgrade; it is a paradigm shift. It will constitute the world’s first **high-sensitivity optical time-domain interferometer**.

In the landscape of the 2030s, the community will possess powerful large-scale spectro-imagers (ALMA, JWST, ELT) and high-cadence imagers (LSST/Rubin, Roman). Yet, a critical gap remains: the ability to **monitor** the variability processes on the spatial and temporal scales they occur, i.e., from less than 0.1 au on day timescales to a few au on weeks/months timescales. It will act as the **dynamical counterpart** to the ELT’s static resolution and the **physical interpreter** of large survey photometric alerts.

By unlocking the temporal dimension, this facility enables new science. It allows us to move beyond mapping *where* matter is, to understanding *how* it moves, falls, reacts and evolves to form stars and planets. Finally, by leveraging and upgrading existing ESO infrastructure, this project aligns with the European carbon footprint reduction strategy and minimises the need for new raw materials.

References

Ábrahám, P., Juhász, A., Dullemond, C. P., et al. 2009, *Nature*, 459, 224
 Alcalá, J. M., Manara, C. F., Natta, A., et al. 2017, *A&A*, 600, A20
 Benisty, M., Bae, J., Facchini, S., et al. 2021, *ApJ*, 916, L2
 Cieza, L. A., Casassus, S., Tobin, J., et al. 2016, *Nature*, 535, 258
 Contreras Peña, C., Lucas, P. W., Guo, Z., & Smith, L. 2024, *MNRAS*, 528, 1823
 Fischer, W. J. et al. 2023, *The Astrophysical Journal*
 GRAVITY+ Collaboration. 2022, GRAVITY+: The VLTI Facility Upgrade, eSO Messenger
 GRAVITY Collaboration, Soulain, A., Perraut, K., et al. 2023, *A&A*, 674, A203
 Hartmann, L., Herczeg, G., & Calvet, N. 2016, *ARA&A*, 54, 135
 Hocde, V., Matter, A., Nardetto, N., et al. 2025, *A&A*, 694, A101
 Hodgkin, S. T., Harrison, D. L., Breedt, E., et al. 2021, *A&A*, 652, A76
 Ivezic, Z., Kahn, S. M., Tyson, J. A., et al. 2019, *ApJ*, 873, 111
 Kóspál, Á., Ábrahám, P., Díezel, L., et al. 2023, *ApJ*, 945, L7
 Kraus, S., Keplin, A., Young, A. K., et al. 2020, *Science*, 369, 1233
 Kruijer, T. S., Kleine, T., & Borg, L. E. 2020, *Nature Astronomy*, 4, 32
 Kuo, I.-H. G., Yen, H.-W., & Gu, P.-G. 2024, *ApJ*, 975, L33
 Laznevoi, S. I., Akimkin, V. V., Pavlyuchenkov, Y. N., et al. 2025, *A&A*, 700, L24
 Liu, H., Herczeg, G. J., Johnstone, D., et al. 2022, *ApJ*, 936, 152
 Lopez, B., Lagarde, S., Petrov, R. G., et al. 2022, *A&A*, 659, A192
 Manara, C. F., Frasca, A., Venuti, L., et al. 2021, *A&A*, 650, A196
 Morbidelli, A., Marrocchi, Y., Ahmad, A. A., et al. 2024, *A&A*, 691, A147
 Nagel, E., Bouvier, J., & Duarte, A. E. 2024, *A&A*, 688, A61
 Nayakshin, S., Owen, J. E., & Elbakyany, V. 2023, *MNRAS*, 523, 385
 Pantolmos, G., Zanni, C., & Bouvier, J. 2020, *A&A*, 643, A129
 Roggero, N., Bouvier, J., Rebull, L. M., & Cody, A. M. 2021, *A&A*, 651, A44
 Schaefer, G. H., Brummelaar, T. T., Gies, D. R., et al. 2014, *Nature*, 515, 234
 Tessore, B., Soulain, A., Pantolmos, G., et al. 2023, *A&A*, 671, A129
 Varga, J., Hogerheijde, M., van Boekel, R., et al. 2021, *A&A*, 647, A56
 Zhou, Y., Bowler, B. P., Sanghi, A., et al. 2025, *ApJ*, 980, L39
 Zhu, Z. 2025, *MNRAS*, 537, 3701