

Non-perturbative renormalization for lattice massive QED₂: the ultraviolet problem

Simone Fabbri¹, Vieri Mastropietro², and Bruno Renzi¹

¹SISSA, Via Bonomea 265, 34136 Trieste, Italy

²University of Rome Sapienza, P.le Aldo Moro 2, 00185 Roma, Italy

February 4, 2026

Abstract

We consider a lattice regularization, preserving Ward Identities (WI) and with a Wilson term, of the Massive QED₂, describing a fermion with mass m and charge e interacting with a vector field with mass M , in the regime $m \ll M \ll a^{-1}$ (a being the lattice spacing) which is the suitable one to mimic a realistic 4d massive gauge theory like the Electroweak sector. The presence of the lattice and of the mass m breaks any solvability property. In this paper we prove that the effective action obtained after the integration of the ultraviolet degrees of freedom is expressed by expansions which are convergent for values of the coupling $|e| \leq e_0$, with e_0 independent on a and m , and with cut-off-independent bare parameters. By combining this result with the analysis of the infrared part in previous papers we get a complete construction of the model and a number of properties whose analogous are expected to hold in 4d. The analysis is done by integrating out the bosons and reducing to a fermionic theory; however, with respect to the case with momentum regularizations (which break essential features like the WI), the resulting effective fermionic action has not a simple form and this requires the developments of new methods to get the necessary bounds.

Contents

1	Introduction and main result	2
1.1	Previous results on the Massive QED ₂	4
1.2	The lattice massive QED ₂	5
1.3	Integration of the boson field	7
1.4	Ward-Identities and ξ -independence	9
1.5	The ultraviolet integration	11
1.6	Sketch of the proof	13
1.7	The infrared problem and construction of the lattice massive QED ₂	14
2	The fermionic multiscale	16
2.1	Multiscale decomposition	16
2.2	Tree expansion	18
2.3	Bounds for the tree expansion	20
3	Bounds for the relevant and marginal terms	22
3.1	The electron self-energy	23
3.2	The lowest-degree vertex function	31

3.3	The lowest-degree polarization bubble	33
3.4	The effective quartic interaction	38
3.5	The higher-degree marginal kernels	39
4	The complete generating functional	42
4.1	The chiral and fermion sources	43
4.2	The vector source	47
5	Proof of Theorem 1.4.	50
5.1	Bounds for the kernels	50
5.2	Correlation functions	51
A	Consequences of the local phase invariance	55
B	Identities for the effective potential	57
B.1	Preliminary identities	58
B.2	Identities for the kernels	62
C	The non-interacting bubbles	64
C.1	The finiteness of the non-interacting bubbles	64
C.2	Proof of Proposition 5.1	69
D	The tree expansion	71
E	Symmetries	74
F	Overview of the continuum limit	76

1 Introduction and main result

The *Massive Quantum Electro-Dynamics* (QED) describes Dirac fermions interacting with a $U(1)$ vector boson field. Its formal Euclidean action in dimension $d \geq 2$ is:

$$\int_{\mathbb{R}^d} d^d x \left(\frac{1}{4} \sum_{\mu, \nu=0}^{d-1} F_{\mu, \nu} F_{\mu, \nu} + \frac{M^2}{2} \sum_{\mu=0}^{d-1} A_{\mu}^2 + \sum_{\mu=0}^{d-1} \bar{\psi} \gamma_{\mu} \left(\partial_{\mu} + i e A_{\mu} \right) \psi + m \bar{\psi} \psi \right), \quad (1.1)$$

where: A_{μ} is the boson (real) field with mass M and $F_{\mu, \nu} = \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu}$; $\psi, \bar{\psi}$ is the fermionic (anti-commuting) field with mass m and charge e ; $\{\gamma_{\mu}\}_{\mu=0}^{d-1}$ are a representation of the Euclidean d -dimensional Clifford algebra. The model is expected to be well defined in $d = 4$ only as an effective theory with a finite ultraviolet cut-off, representing the maximal energy scale at which the theory is valid; it is indeed renormalizable but not asymptotically free. The cut-off must fulfill the following physical requirements:

- a) the Ward Identities (WI) must be verified, ensuring the validity of the conservation of the current coupled to A_{μ} ;
- b) the *longitudinal*, non-decaying term of the boson propagator (see comments after Eq. (1.4)) must not contribute to physical observables;
- c) for length scales much larger than the inverse cut-off scale the correlations must formally reduce to the continuum ones up to sub-leading corrections;

- d) the cut-off has to be at least exponentially high in the inverse coupling; for small e this ensures that the effect of the cut-off is undetectable at the energy scale of experiments.

A natural choice for the cut-off is the lattice one, obtained by replacing the continuum Euclidean space-time with a discrete lattice of step a and finite length L , and the derivative ∂_μ appearing in Eq. (1.1) with a discrete derivative¹; this however violates item **c**), as the lattice Dirac operator acquires spurious extra zero modes, a problem known as *fermion doubling*. Condition **c**) can be fulfilled adding a Wilson term to the action. In order to fulfill condition **a**) the interaction between boson and fermion has to be defined in a non-linear way, via the exponential of A_μ . As consequence of property **a**), property **b**) follows as well, which is crucial for ensuring the renormalizability of the theory and the validity of the property **d**). Note that the introduction of the Wilson term breaks the chiral symmetry which would be present at a formal level when the electron mass m is vanishing, a fact producing the chiral anomaly.

The regularization of QED, with a Wilson term and respecting WI is called *lattice Massive QED*. Other lattice definitions could be possible, but they are expected to be equivalent as far as the above requirements are true.

Massive QED is not physically realized in nature as a theory of elementary particles. However it is very much related to the Standard Model, the theory describing all elementary particles and forces except gravity, and in particular to the neutral Weak sector, where the force is mediated by an $U(1)$ massive boson, the Z^0 particle. Indeed both theories are expected to exist only as effective theories; moreover the Adler-Bardeen property [1] and the decrease of the divergence degree, due to the cancellation of the contribution of the longitudinal part of the propagator, are expected to hold. The lattice massive QED is therefore an ideal setting to test, in a simpler model, several features which are believed to be true on the basis of perturbative arguments, that is at the level of series which are not convergent. As the ratio of the boson and the electron mass is $\sim 10^6$, and the energies in accelerators are of the order of 100 times the boson mass, the range of interest of lattice massive QED is

$$m \ll M \ll a^{-1}. \quad (1.2)$$

In addition the charge is small as its physical value is, in dimensionless units, $e^2 \sim \frac{1}{137}$ (fixing the wave function renormalization to 1).

It should be recalled that other peculiarities of the Standard Model are instead absent in massive QED, like the fact that in the Standard Model the interaction has a *chiral* nature. This fact introduces a number of difficulties, in particular the fact that the Ward Identities are verified, even at a perturbative level, only if the chiral anomalies cancel out and if the particle charges verify certain conditions, a fact requiring the Adler-Bardeen property.

Even if the lattice massive QED in 4d is much simpler than the Standard Model, rigorous results in the regime of Eq. (1.2) are lacking, and the analysis has been limited either to the region $m \sim 1/a$ and any $M \geq 0$ [2] or with a momentum regularization with $M \sim 1/a$ and any $m \geq 0$ [3]. On the other hand massive QED in 2d is more tractable and at the same time the analogue of a number of features holding in 4d are true also in 2d.

In this paper we perform the construction of the lattice massive QED_2 in the regime of Eq. (1.2), and we prove a number of properties expected to be true in 4d or in the Standard Model. In particular we focus on the ultraviolet properties of the theory, obtaining results which, once combined with the infrared analysis in [4, 5, 6, 7], provide the construction of the model and the verification at a non-perturbative level of a number of properties which are expected to hold in 4d more realistic models.

¹Equivalently k_μ with $\frac{1}{a} \sin(ak_\mu)$ in momentum space.

1.1 Previous results on the Massive QED₂

The Massive QED₂ has been extensively studied in the past, see e.g. [8] for a review with more references, and we shortly summarize some of the main results.

Perturbative results. In the continuum, at a classical level, the formal action of QED₂ is invariant under the change of variables $\psi_x \rightarrow e^{i\alpha}\psi_x$ and this, by Noether theorem, implies the conservation of the current $\sum_\mu \partial_\mu j_\mu = 0$, with $j_\mu = e\bar{\psi}_x \gamma_\mu \psi_x$; in the same way if the electron mass is vanishing, $m = 0$, there is invariance under $\psi_x \rightarrow e^{i\alpha\gamma_5}\psi_x$ which implies the conservation of the axial current $\sum_\mu \partial_\mu j_\mu^5 = 0$ with $j_{\mu,x}^5 = e\bar{\psi}_x \gamma_\mu \gamma_5 \psi_x$.

At a quantum level, using Feynman graphs expansion with dimensional regularization, the following Ward Identities (WI) are found, to be understood as order by order identities among formal (non-convergent) power series in the coupling:

$$\sum_{\mu=0,1} p_\mu \langle \hat{j}_{\mu,p}; \hat{j}_{\nu,-p} \rangle = 0, \quad \sum_{\nu=0,1} p_\nu \langle \hat{j}_{\mu,p}; \hat{j}_{\nu,-p}^5 \rangle = \frac{ie^2}{\pi} \sum_{\nu=0,1} \varepsilon_{\mu,\nu} p_\nu, \quad (1.3)$$

with $\varepsilon_{\mu,\nu}$ being the antisymmetric tensor (namely $\varepsilon_{0,1} = -\varepsilon_{1,0} = 1, \varepsilon_{0,0} = \varepsilon_{1,1} = 0$). The first equation is the counterpart of the conservation of the current; the second, valid in the $m = 0$ case, shows that the axial current is not conserved at a quantum level. The coefficient $\frac{1}{\pi}$ in the r.h.s. is called the chiral anomaly. Such coefficient is in general a series in the coupling but the above statement says that only the lowest order contributes; this is the Adler-Bardeen's non-renormalization theorem in 2d [9]. The boson propagator has the form

$$\frac{1}{|k|^2 + M^2} \left(\delta_{\mu,\nu} + \frac{k_\mu k_\nu}{|k|^2 + M^2} \right). \quad (1.4)$$

The term $k_\mu k_\nu / (|k|^2 + M^2)$, called the *longitudinal* term, is not decaying for $|k| \gg M$ and the naive degree of divergence is $D = 2 - n_\psi/2 - n^A$ (n_ψ and n_A the number of ψ and A fields), corresponding to a renormalizable theory; however the WI corresponding to the current conservation implies that the contribution from the longitudinal term is vanishing and the real degree of divergence is $D = 2 - n - n_\psi/2$, if n is the perturbative order, corresponding to a super-renormalizable theory. A similar decrease of degree of divergence is at the basis of the perturbative renormalizability of the 4d gauge theories; indeed in 4d one passes from $D = 4 + n - 3n_\psi/2 - 2n_A$ to $D = 4 - 3n_\psi/2 - n^A$, that is there is a reduction from a non-renormalizable to a renormalizable behavior.

Exact solutions. There are exact solutions for the massive QED₂ in the $m = 0$ case in the continuum, which have been often used as a benchmark for perturbative analyses. In particular a solution was found in [10, 11, 12]; the $M = 0$ case, known as Schwinger model, was solved in [13]. Solutions are based either on a combination of Schwinger-Dyson equations and anomalous WI, obtained by a self-consistent argument previously used in [14] in the Thirring model, or on bosonization. Other exact solutions were found using functional integrals methods [15, 16]; see also [17] for related interesting applications.

The solutions rely on formal manipulations of diverging expressions, and the results appear to depend on a parameter introduced by the regularization. Main properties are:

- 1) a vanishing wave function renormalization $Z = 0^\eta$ with $\eta > 0$, see e.g. [11, Eq. (4.35)];
- 2) a value of the chiral anomaly given by

$$\sum_{\nu=0,1} p_\nu \langle \hat{j}_{\mu,p}; \hat{j}_{\nu,-p}^5 \rangle = \zeta \frac{ie^2}{\pi} \sum_{\nu=0,1} \varepsilon_{\mu,\nu} p_\nu, \quad (1.5)$$

where ζ is a parameter depending on the regularization; it was found $\zeta = 1/2$ in [10], $\zeta = 1$ in [11] and any ζ in [12]. The current was found to be non-conserved. Moreover a renormalized mass of the boson is found to be $M^2 + \zeta \frac{e^2}{\pi}$.

Note that the results found with exact solutions are somewhat different from the ones found in perturbation theory, due to the different regularizations adopted; in particular the conservation of the current is broken and there is no reduction of degree of divergence, as signaled by the singular Z ; moreover the value of the anomaly is different.

No solution is known when $m \neq 0$ or with a lattice regularization. The issue of lattice regularization is particularly relevant as anomalies appear in the continuum, but in a non-perturbative construction in 4d one needs a lattice cut-off. An intense activity in understanding the role of the lattice has been devoted in recent times, mostly in the context of the Schwinger model, see e.g. [18, 19, 20, 21, 22, 23].

Renormalization Group with momentum regularization. Non-perturbative Renormalization Group (RG) analysis have been performed in the continuum with momentum regularization. In [24, 7] it was considered a massive QED₂ model in the continuum, obtained starting from the formal action in Eq. (1.1), neglecting the longitudinal term of the propagator (so that the power counting is that of a super-renormalizable theory) and with a momentum fermionic cut-off $\chi_N(k)$, given by a smooth function vanishing for $|k| \geq 2^N$ (the boson can be assumed with or without cut-off [25]); this model has been also called *reference model*, and it was used to derive universality relations for several systems, see e.g. [26, 27]. The analysis is done by integrating out the boson field and obtaining a quartic interaction in the fermions; the analysis of the ultraviolet part is done using the non-locality of the boson propagator, see also [28], while the integration of the infrared scales is done by implementing cancellations due to Ward Identities at each step of the Renormalization Group, controlling the corrections due to the presence of the cut-offs, see [5], and proving the convergence to a line of fixed points. In this continuum massive QED₂ model, in which the ultraviolet limit $N \rightarrow \infty$ can be taken as well as the infinite volume and zero fermionic mass limit, the WI has the form:

$$\sum_{\mu=0,1} p_\mu \langle \hat{j}_{\mu,p} ; \hat{j}_{\nu,-p} \rangle = e \int \frac{d^2 k}{(2\pi)^2} \left(\not{k} \frac{1 - \chi_N(k)}{\chi_N(k)} - (\not{k} + \not{p}) \frac{1 - \chi_N(k+p)}{\chi_N(k+p)} \right) \langle \hat{\psi}_{k+p} \hat{\psi}_k ; \hat{j}_{\nu,-p} \rangle, \quad (1.6)$$

with $\not{k} \equiv \sum_{\mu=0,1} \gamma_\mu k_\mu$. The r.h.s. of Eq. (1.6) is not small, even for large N , and one can prove, see [24, 7], both the non-conservation of the current and of the axial current. The form of the limit appears to be strongly sensitive to the regularization adopted.

1.2 The lattice massive QED₂

In this paper we study a regularization of massive QED₂ fulfilling all the physical constraints **a)-d)** listed few lines below Eq. (1.1). We actually consider the lattice regularization of a generalized version of Eq. (1.1), obtained by adding an extra term $\frac{\xi}{2} \int d^2 x (\sum_{\mu=0,1} \partial_\mu A_{\mu,x})^2$ called *gauge-fixing term*, with $\xi \in [0, 1]$ called *gauge-fixing parameter*; the original theory is recovered by setting $\xi = 0$.² The model is defined on a lattice $\Lambda := a\mathbb{Z}^2 \cap (-\frac{L}{2}, \frac{L}{2}]^2$ with spacing $a \equiv 2^{-N} \in 2^{\mathbb{Z}}$. We impose periodic boundary conditions, so that $\Lambda = (a\mathbb{Z}^2)/(L\mathbb{Z}^2)$ and it comes equipped with the *torus metric* $\|x - y\|_L := \min_{n \in \mathbb{Z}^2} \sqrt{\sum_{\mu=0,1} (x_\mu - y_\mu + Ln_\mu)^2}$. The generating

²The addition of the gauge-fixing term does not modify the model as soon as gauge invariant observables are considered, see Lemma 1.3 and comments thereafter.

functional for correlation functions is

$$\mathcal{W}_\xi(\varphi; J; B) := -\log \frac{\int \mathcal{D}\psi \int \mathbb{P}_\xi(\mathcal{D}A) e^{-I(A+J,\psi) - (\varphi,\psi) - i(B,j^5(\psi))}}{\int \mathcal{D}\psi e^{-I(0,\psi)}}, \quad (1.7)$$

where:

1. ψ is the fermion field, identified with the Grassmann algebra generated by $\{\psi_{x,s}^\pm\}_{x \in \Lambda, s \in \{1,2\}}$, where, according to the Dirac notation, letting ψ_x^\pm be the row/column vector $(\psi_{x,1}^\pm, \psi_{x,2}^\pm)$, we identify $\psi_x^- \equiv \psi_x$ and $\psi_x^+ \equiv \bar{\psi}_x$. The symbol $\mathcal{D}\psi$ denotes the Grassmann-Berezin integration over the Grassmann algebra (namely the projector onto the highest degree monomial, cf. [29, Sect. 4.1]).
2. A is the vector-boson field, to be identified with $\{A_{\mu,x}\}_{x \in \Lambda, \mu \in \{0,1\}} \in \mathbb{R}^{\Lambda \times \{0,1\}}$, while $\mathbb{P}_\xi(\mathcal{D}A)$ is the real Gaussian integration with propagator

$$g_{\mu,\nu}^{A,\xi}(x-y) \equiv \mathbb{E}_\xi(A_{\mu,x} A_{\nu,y}) = \int_{\Lambda^*} dk \frac{e^{-ik \cdot (x-y)}}{|\sigma(k)|^2 + M^2} \left(\delta_{\mu,\nu} + (1-\xi) \frac{\sigma_\mu(k) \sigma_\nu^*(k)}{\xi |\sigma(k)|^2 + M^2} \right), \quad (1.8)$$

where: $\Lambda^* := \frac{2\pi}{L} \mathbb{Z}^2 \cap \left(-\frac{\pi}{a}, \frac{\pi}{a}\right]^2$ and

$$\int_{\Lambda^*} dk \cdot \equiv \frac{1}{L^2} \sum_{k \in \Lambda^*} \cdot; \quad (1.9)$$

$\sigma_\mu(k) \equiv i2^N(e^{-i2^{-N}k_\mu} - 1)$, while $\sigma_\mu^*(k)$ denotes its complex conjugate and $|\sigma(k)|^2 \equiv \sum_{\mu=0,1} |\sigma_\mu(k)|^2$; $M > 0$ is the bare boson mass; $\xi \in [0, 1]$ is the gauge-fixing parameter.

3. $J \equiv \{J_{\mu,x}\}_{x \in \Lambda, \mu \in \{0,1\}} \in \mathbb{R}^{\Lambda \times \{0,1\}}$ is a real external field, and

$$\begin{aligned} I(A, \psi) = & \frac{1}{2} 2^N \sum_{\mu=0,1} \int_{\Lambda} dx \psi_x^+ (\gamma_\mu - r) e^{i2^{-N} \mathbf{e} A_{\mu,x}} \psi_{x+2^{-N}\hat{e}_\mu}^- + \\ & - \frac{1}{2} 2^N \sum_{\mu=0,1} \int_{\Lambda} dx \psi_{x+2^{-N}\hat{e}_\mu}^+ (\gamma_\mu + r) e^{-i2^{-N} \mathbf{e} A_{\mu,x}} \psi_x^- + (m_N + 2r2^N) \int_{\Lambda} dx \psi_x^+ \psi_x^-, \end{aligned} \quad (1.10)$$

where:

$$\int_{\Lambda} dx \cdot \equiv a^2 \sum_{x \in \Lambda} \cdot; \quad (1.11)$$

$\hat{e}_0 = (1, 0)$ and $\hat{e}_1 = (0, 1)$; $\mathbf{e} \in \mathbb{R}$ is the electric charge; $r > 0$ is the Wilson mass, to be fixed once and for all (from now on we will assume $r = 1$); $m_N \in \mathbb{R}$ is the bare electron mass; γ_0, γ_1 are a representation of the Euclidean 2D Clifford algebra, i.e. they satisfy $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu,\nu}$; we chose them as

$$\gamma_0 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma_1 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}.$$

4. $\varphi \equiv \{\varphi_{x,s}^\pm\}_{x \in \Lambda, s \in \{1,2\}}$ is an external Grassmann field and: $(\varphi, \psi) \equiv \int_{\Lambda} dx (\varphi_x^+ \psi_x^- + \psi_x^+ \varphi_x^-)$.
5. $B \equiv \{B_{\mu,x}\}_{x \in \Lambda, \mu \in \{0,1\}}$ is an external real field coupled to the chiral current, namely $(B, j^5(\psi)) \equiv \int_{\Lambda} dx B_{\mu,x} j_{\mu,x}^5(\psi)$, where $j_{\mu,x}^5(\psi) := \mathbf{e} Z_N^5 \psi_x^+ \gamma_\mu \gamma_5 \psi_x^-$, with $\gamma_5 := i\gamma_0\gamma_1$ and Z_N^5 a suitable continuous function of \mathbf{e} , such that $Z_N^5|_{\mathbf{e}=0} = 1$ and, say, $|Z_N^5 - 1| \leq \frac{1}{2}$.

Remark 1.1. The Gaussian weight of the boson measure \mathbb{P}_ξ is explicit and is given by the exponential of:

$$\frac{1}{4} \sum_{\mu, \nu=0,1} \int_{\Lambda} dx \left(\nabla_\mu A_{\nu,x} - \nabla_\nu A_{\mu,x} \right)^2 + \frac{M^2}{2} \sum_{\mu=0,1} \int_{\Lambda} dx A_{\mu,x}^2 + \frac{\xi}{2} \int_{\Lambda} dx \left(\sum_{\mu=0,1} \nabla_\mu^T A_{\mu,x} \right)^2 \quad (1.12)$$

with a minus sign, where $\nabla_\mu A_{\nu,x} \equiv a^{-1} (A_{\nu,x+a\hat{e}_\mu} - A_{\nu,x})$ and $\nabla_\mu^T A_{\nu,x} \equiv a^{-1} (A_{\nu,x-a\hat{e}_\mu} - A_{\nu,x})$. Eq. (1.12) is nothing but the lattice counterpart of the free-boson action in Eq. (1.1) plus the gauge-fixing term.

Physical properties can be obtained by the correlations, for instance:

$$\hat{\Sigma}_{\mu,\nu}^\xi(p) := \int_{\Lambda} dx e^{-ip \cdot x} \frac{\delta^2 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta J_{\nu,0}}(0;0;0), \quad (1.13)$$

$$\hat{\Sigma}_{5;\mu,\nu}^\xi(p) := \int_{\Lambda} dx e^{-ip \cdot x} \frac{\delta^2 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta B_{\nu,0}}(0;0;0), \quad (1.14)$$

$$\hat{\Gamma}_\mu^\xi(p; k)_{s,s'} := \int_{\Lambda^2} dxdy e^{i(-p \cdot x + k \cdot y)} \frac{\delta^3 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta \varphi_{0,s}^- \delta \varphi_{y,s}^+}(0;0;0), \quad (1.15)$$

$$\hat{S}^\xi(k)_{s,s'} := \int_{\Lambda} dx e^{ik \cdot x} \frac{\delta^2 \mathcal{W}_\xi}{\delta \varphi_{x,s}^+ \delta \varphi_{0,s'}^-}(0;0;0), \quad (1.16)$$

with $\frac{\delta}{\delta J_{\mu,x}} := 2^{2N} \frac{\partial}{\partial J_{\mu,x}}$ and similarly for the other *functional derivatives*, with the understanding that derivatives w.r.t. φ are meant as anti-commuting derivatives.

Remark 1.2. The Wilson term, that is the term proportional to r , is introduced in order to avoid the fermion doubling problem. The mass m_N and Z_N^5 are *bare parameters* to be fixed in order to ensure the renormalization conditions. The Wilson term and the corresponding breaking of the chiral symmetry have the effect that the massless case, $m = 0$, requires a non-vanishing m_N , and that Z_N^5 is not equal to 1. In contrast with other related QFT in 2d, here we do not introduce in the model a bare wave function renormalization Z_N , which in several cases is singular in the limit of removed cut-off, as in the Thirring model; our aim is indeed to prove that one has finite observables uniformly in N by choosing $Z_N = 1$. With L, a, M, m_N finite, Eq. (1.7) is well defined (even if the boson field is unbounded) for ϵ small enough, as discussed in the following section.

1.3 Integration of the boson field

Our first step towards the construction of the theory is the integration of the boson field, which yields an effective fermion interaction. We consider the interacting part of the action,

$$\begin{aligned} V(A + J, \psi) &:= I(A + J, \psi) - I(0, \psi) \\ &= \frac{1}{2} \sum_{\mu=0,1} \int dx \psi_x^+ (\gamma_\mu - r) \psi_{x+2^{-N}\hat{e}_\mu}^- G_{x,\mu,+}(A + J) + \\ &\quad - \frac{1}{2} \sum_{\mu=0,1} \int dx \psi_{x+2^{-N}\hat{e}_\mu}^+ (\gamma_\mu + r) \psi_x^- G_{x,\mu,-}(A + J), \end{aligned} \quad (1.17)$$

where $G_{x,\mu,\pm}(A) := 2^N \left(e^{\pm i 2^{-N} \epsilon A_{\mu,x}} - 1 \right)$. We use the the following conventions.

1. Bosonic labels: $b \equiv (x, \mu, \epsilon) \in \Lambda_B \equiv \Lambda \times \{0, 1\} \times \{\pm\}$. Besides, $\int_{\Lambda_B} db \equiv \sum_{\mu \in \{0,1\}} \sum_{\epsilon \in \{\pm\}} \int_{\Lambda} dx$.

2. Fermionic labels: $\eta \equiv (x, s) \in \Lambda_F \equiv \Lambda \times \{1, 2\}$. Besides, $\int_{\Lambda_F} d\eta \equiv \sum_{s \in \{1, 2\}} \int_{\Lambda} dx$.

3. If $\underline{b} \equiv (b_1, \dots, b_p)$ we let $\int_{\Lambda_B}^p d\underline{b} := \int_{\Lambda_B} db_1 \cdots \int_{\Lambda_B} db_p$, and similarly for fermionic labels. Whenever the value of p and the nature of the label (bosonic or fermionic) will be clear from the context, we will simply write $\int d\underline{b}$ etc..

With this understanding, we can rewrite the action as

$$V(A + J, \psi) + (\varphi, \psi) + i(B, j^5) = \int_{\Lambda_B} db O_b(\psi) G_b(A + J) + \int_{\Lambda_F} d\eta (\varphi_\eta^+ \psi_\eta^- + \psi_\eta^+ \varphi_\eta^-) + \underbrace{\int_{\Lambda_B} db B_b O_{5;b}(\psi)}_{=: (B, O_5)}, \quad (1.18)$$

where $O_b(\psi) = \int d\eta d\eta' \psi_\eta^+ c_b(\eta, \eta') \psi_{\eta'}^-$, and, if $b = (y, \mu, \epsilon)$, $\eta = (x, s)$, $\eta' = (x', s')$:

$$c_b(\eta, \eta') = \begin{cases} \frac{1}{2} \delta(x - y) \delta(x' - y - 2^{-N} \hat{e}_\mu) (\gamma_\mu - r)_{s,s'} & \epsilon = + \\ -\frac{1}{2} \delta(x' - y) \delta(x - y - 2^{-N} \hat{e}_\mu) (\gamma_\mu + r)_{s,s'} & \epsilon = - \end{cases} \quad (1.19)$$

where hereafter $\delta(x) := \sum_{n \in \mathbb{Z}^2} 2^{2N} \delta_{x_0, n_0 L} \delta_{x_1, n_1 L}$.³ More explicitly:

$$\begin{cases} O_{x,\mu,+}(\psi) &= \frac{1}{2} \psi_x^+ (\gamma_\mu - r) \psi_{x+2^{-N} \hat{e}_\mu}^- \\ O_{x,\mu,-}(\psi) &= -\frac{1}{2} \psi_{x+2^{-N} \hat{e}_\mu}^+ (\gamma_\mu + r) \psi_x^- \end{cases}$$

Moreover, in the r.h.s. of Eq. (1.18), in order for the coupling with the chiral and the vector current to possess a similar, standard form (to be also compatible with the symmetries of the model, cf. Appendix E), we have set $B_{x,\mu,\epsilon} := i\epsilon e B_{\mu,x}$ and $O_{5;x,\mu,\epsilon}(\psi) := \int d\eta d\eta' \psi_\eta^+ c_{x,\mu,\epsilon}^5(\eta, \eta') \psi_{\eta'}^-$, where

$$c_{y,\mu,\epsilon}^5((x, s), (x', s')) := \frac{1}{2} \epsilon Z_N^5 \delta(x - y) \delta(x' - y) (\gamma_\mu \gamma_5)_{s,s'}. \quad (1.20)$$

The following identity holds:

$$\int \mathbb{P}_\xi(\mathcal{D}A) e^{-V(A+J,\psi)} = e^{-\mathcal{V}(\psi; G(J))},$$

where \mathcal{V} is a polynomial⁴ in $\{\psi_\eta^\pm\}_{\eta \in \Lambda_F}$ and $\{G_b(J)\}_{b \in \Lambda_B}$, given by

$$\begin{aligned} \mathcal{V}(\psi; G(J)) &= 2^N (\mu_{N,\lambda} - 1) \int db O_b(\psi) + \mu_{N,\lambda} \int_{\Lambda_B} db O_b(\psi) G_b(J) + \\ &+ \sum_{n \geq 2} \sum_{0 \leq m \leq n} 2^{N(2-n-m)} \int_{\Lambda_B^n} db \int_{\Lambda_B^m} db' w_{n,m}(\underline{b}; \underline{b}') O_{\underline{b}}^n(\psi) G_{\underline{b}'}^m(J), \end{aligned} \quad (1.21)$$

with $\mu_{N,\lambda} \equiv e^{-\frac{\lambda}{2} 2^{-2N} g_{\mu,\mu}^{A,\xi}(0)}$, $\lambda \equiv e^2$; given $\underline{b} = (b_1, \dots, b_n)$ we let $O_{\underline{b}}^n(\psi) := \prod_{j=1}^n O_{b_j}(\psi)$, similarly $G_{\underline{b}'}^m(J) := \prod_{i=1}^m G_{b'_i}(J)$. In complete analogy with [3, Sect. 2], as a straightforward

³For $\eta = (x, s)$ and $\eta' = (x', s')$ in Λ_F we also let $\delta(\eta - \eta') \equiv \delta_{s,s'} \delta(x - x')$, while for $b = (y, \mu, \epsilon) \in \Lambda_B$ and $b' = (y', \mu', \epsilon') \in \Lambda_B$, $\delta(b - b') \equiv \delta_{\mu,\mu'} \delta_{\epsilon,\epsilon'} \delta(x - y)$.

⁴Due to the finiteness of the fermionic Grassmann algebra, for any $L, N > 0$ fixed the sum in the r.h.s. of Eq. (1.21) has finitely many terms.

consequence of the Battle-Brydges-Federbush formula [30, Thm. 3.1], the kernels $w_{n,m}$ can be expressed via the following formulas:

$$w_{n,0}(\underline{b}) = \frac{1}{n!} \sum_{T \in \mathcal{T}(\{1, \dots, n\})} \prod_{\{i,j\} \in T} \lambda g_{b_i, b_j}^{A, \xi} \times \int dP_T(s) e^{-U(\underline{b}; s)} \quad (1.22)$$

$$w_{n,m}(\underline{b}; \underline{b}') = \frac{1}{m!} \sum_{\substack{i_1, \dots, i_m \in \{1, \dots, n\} \\ \text{distinct}}} \prod_{j=1}^m \delta(b'_j - b_{i_j}) \times w_{n,0}(\underline{b}), \quad m \geq 1, \quad (1.23)$$

where:

- $\mathcal{T}(\{1, \dots, n\})$ the set of all the spanning trees on the set $\{1, \dots, n\}$ where, given $X \subset \mathbb{N}$, a *spanning tree on X* is a set $T \subset X^2$ such that the graph $\mathcal{G} = (X, T)$ is a tree (connected, acyclic). If $|X| = 1$ then $\mathcal{T}(X) = \emptyset$ and in such case $w_{1,0} = \mu_{N,\lambda}$;
- $dP_T(s)$ is a probability measure supported on $[0, 1]^{\mathcal{P}(\{1, \dots, n\})}$, $\mathcal{P}(\{1, \dots, n\})$ being the set of pairs on $\{1, \dots, n\}$;
- $U(\underline{b}; s) \equiv \frac{1}{2} \lambda 2^{-2N} (\sum_{j=1}^n g_{b_j, b_j}^{A, \xi} + 2 \sum_{1 \leq i < j \leq n} s_{\{i,j\}} g_{b_i, b_j}^{A, \xi})$, and this is non-negative on the support of P_T for $\lambda \geq 0$.

It is an immediate consequence of the finiteness of both $\|g_{\mu, \nu}^{A, \xi}\|_1 := \int_{\Lambda} dx |g_{\mu, \nu}^{A, \xi}(x)|$ and $\|g_{\mu, \nu}^{A, \xi}\|_{\infty} := \sup_{x \in \Lambda} |g_{\mu, \nu}^{A, \xi}(x)|$ and of determinant bounds for fermionic expectations, the finiteness of the r.h.s. of Eq. (1.7) for any $\xi \in [0, 1]$ and $\mathbf{e} \in \mathbb{R}$.

The resulting fermionic theory has an interaction involving Grassmann monomials of all possible degree, multiplied by suitable non-local kernels; for every fixed tree T in the r.h.s. of Eq. (1.22) and given $\underline{b} = (b_1, \dots, b_n)$, we can graphically represent the quantity

$$2^{N(2-n)} O_{\underline{b}}^n(\psi) \prod_{\{i,j\} \in T} \lambda g_{b_i, b_j}^A$$

appearing in the expression for $\mathcal{V}(\psi; 0)$ as in Fig. 1.(i); there the nodes of the graph represent the points $\{1, \dots, n\}$: to each of them we assign a coordinate $b_j \in \Lambda_B$ and a dimensional factor $2^{N(1-d_j)}$, with d_j the number of lines of the tree incident in the node. The couple of half solid lines incident in the node j represents the fermionic bilinear $O_{b_j}(\psi)$; any wiggly line is a bond of the tree T , as in Fig. 3.(ii).

1.4 Ward-Identities and ξ -independence

Ward identities hold, generated by the following relation (to be called *local phase invariance*):

$$\mathcal{W}_{\xi}(\varphi; J; B) = \mathcal{W}_{\xi}(e^{i\mathbf{e}\alpha} \varphi; J + \nabla\alpha; B), \quad \forall \alpha \in \mathbb{R}^A, \quad (1.24)$$

with the understanding that $(e^{i\mathbf{e}\alpha} \varphi)_{x,s}^{\pm} \equiv e^{\pm i\mathbf{e}\alpha(x)} \varphi_{x,s}^{\pm}$, $(A + \nabla\alpha)_{\mu,x} \equiv A_{\mu,x} + \nabla_{\mu}\alpha(x)$, and ∇_{μ} the discrete-derivative operator as after Eq. (1.12). Note that \mathcal{W}_{ξ} is well defined for finite values of the cut-off and any $\xi \in [0, 1]$. Eq. (1.24) implies the *Ward identities*:

$$\sum_{\mu=0,1} \sigma_{\mu}(p) \hat{\Sigma}_{\mu,\nu}^{\xi}(p) = \sum_{\mu=0,1} \sigma_{\mu}(p) \hat{\Sigma}_{5;\mu,\nu}^{\xi}(p) = 0, \quad \forall p \in \Lambda^*, \quad (1.25)$$

$$\sum_{\mu=0,1} \sigma_{\mu}(p) \hat{\Gamma}_{\mu}^{\xi}(p; k) = \mathbf{e}(\hat{S}^{\xi}(k) - \hat{S}^{\xi}(k-p)), \quad \forall p, k \in \Lambda^*, \quad (1.26)$$

where $\hat{\Sigma}_{\mu,\nu}^{\xi}$, $\hat{\Sigma}_{5;\mu,\nu}^{\xi}$, $\hat{\Gamma}_{\mu}^{\xi}(p; k)$ and $\hat{S}^{\xi}(k)$ are as in Eqs. (1.13)-(1.16). As a consequence, see App. A, one has the ξ -independence for the generating functional.




Figure 1: (i) graphical representation of the contribution to the term $\mathcal{O}(\psi^{12})$ of the fermionic interaction, obtained by selecting in $w_{6,0}$ the tree with bonds $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{4, 5\}, \{4, 6\}$; (ii) graphical representation for the boson propagator.

Lemma 1.3 (ξ -independence). $W_\xi(0; J; B)$ is constant w.r.t. $\xi \in [0, 1]$.

Notice that the ξ -independence holds only for $\varphi = 0$; more generally it holds only for correlations of *gauge-invariant* observables, namely quantities $F(A, \psi)$ such that $F(A + \nabla\alpha, e^{i\alpha}\psi) = F(A, \psi)$ for every $\alpha \in \mathbb{R}^A$. Note in particular that $V(A, \psi) + i(B, j^5)$ is gauge-invariant.

Lemma 1.3 allows us to take $\xi = 1$, so that the longitudinal term of the boson propagator is not contributing. The identity with the original model with $\xi = 0$ holds only for the correlations of currents (correlations of fermions could be included if they respect the above invariance). From now on we will think of ξ as fixed once and for all at the value $\xi = 1$, and we will denote the boson propagator $g^{A,1}$ simply by g^A . The latter verifies the following bound:

$$|g_{\mu,\nu}^A(x - y)| \leq \delta_{\mu,\nu} \frac{K_p}{(N - h_M^*) + (2^{h_M^*} \|x - y\|_L)^{\frac{1}{p}}} \exp \left\{ -\kappa_1 \sqrt{2^{h_M^*} \|x - y\|_L} \right\} \quad (1.27)$$

with $h_M^* := \lfloor \log_2 M \rfloor$, for every $1 \leq p < \infty$ and suitable constants K_p, κ_1 ; as a consequence

$$\|g_{\mu,\nu}^A\|_1 = \mathcal{O}(2^{-2h_M^*}), \quad \|g_{\mu,\nu}^A\|_\infty = \mathcal{O}(N - h_M^*), \quad (1.28)$$

which has to be contrasted with the bounds in the $\xi = 0$ case, $\|g_{\mu,\nu}^{A,0}\|_1 = \mathcal{O}(2^{-2h_M^*}(N - h_M^*))$ and $\|g_{\mu,\nu}^{A,0}\|_\infty = \mathcal{O}(2^{2N-2h_M^*})$. The boundedness of the L^1 norm will play a crucial role in the following analysis.

Note also that it is convenient to rewrite $w_{n,0}$ as a dominant term plus a remainder, the latter carrying a dimensional improvement in N . To do so, in Eq. (1.23) we rewrite $e^{-U(\underline{b};s)} = e^{-U(\underline{b};0)} + (e^{-U(\underline{b};s)} - e^{-U(\underline{b};0)})$, which induces the splitting

$$w_{n,0}(\underline{b}) = v_n(\underline{b}) + 2^{2h_M^* - 2N} \tilde{v}_n(\underline{b}), \quad v_n(\underline{b}) := \frac{\mu_{N,\lambda}^n}{n!} \sum_{T(\{1, \dots, n\})} \prod_{\{i,j\} \in T} \lambda g_{b_i, b_j}^A, \quad (1.29)$$

and one can easily show, using Eq. (1.27), that there exists a constant $C > 0$ such that

$$\|v_n\|_1^w \leq C^n (2^{-2h_M^*} \lambda)^{n-1}, \quad \|\tilde{v}_n\|_1^w \leq C^n (2^{-2h_M^*} \lambda)^n, \quad (1.30)$$

where, for any function W (to be called *kernel* hereafter) over $\Lambda_F^{2q} \times \Lambda_B^m$ and $p > 0$, we let

$$\|W\|_p^w := \left(\frac{1}{L^2} \int_{\Lambda_F^{2q}} d\underline{\eta} \int_{\Lambda_B^m} d\underline{b} \exp\left(\frac{p}{2}\kappa\sqrt{2^{h_M^*} \delta^D(\underline{\eta}, \underline{b})}\right) |W(\underline{\eta}; \underline{b})|^p \right)^{\frac{1}{p}} \quad (1.31)$$

with: $\delta^D(\underline{\eta}, \underline{b})$ the diameter distance between the $2q + m$ space coordinates of $\eta_1, \dots, \eta_{2q}, b_1, \dots, b_m$; $\kappa := \min\{\kappa_1, \kappa_2, \kappa_3\}$, where $\kappa_1, \kappa_2, \kappa_3$ are suitable constants introduced in Eqs. (1.27), (2.4), (C.2) respectively.

1.5 The ultraviolet integration

Let $P^{(\leq N)}(\mathcal{D}\psi)$ be the Grassmann Gaussian integration with propagator:

$$g_{s,s'}^{(\leq N)}(x - y) \equiv \mathcal{E}_{(\leq N)}(\psi_{x,s}^- \psi_{y,s'}^+) = \int_{\Lambda^*} dk e^{-ik \cdot (x - y)} \left(-i\cancel{s}(k) + m_N + M_N(k) \right)_{s,s'}^{-1}, \quad (1.32)$$

where $\cancel{s}(k) := \sum_{\mu=0,1} \gamma_\mu s_\mu(k)$, with $s_\mu(k) := 2^N \sin(2^{-N} k_\mu)$, and $M_N(k) := 2r \sum_{\mu=0,1} 2^N \sin^2(2^{-N-1} k_\mu)$.

We consider $|m_N| \leq M \leq 2^N$ and, recalling that $h_M^* = \lfloor \log_2 M \rfloor$, we decompose the fermion propagator as $g^{(\leq N)}(x - y) = g^{(\leq h_M^*)}(x - y) + g^{(h_M^*, N]}(x - y)$, with

$$\hat{g}^{(\leq h_M^*)}(k) := \frac{\chi(2^{-h_M^*} |k|)}{-i\cancel{s}(k) + m_N + M_N(k)}, \quad \hat{g}^{(h_M^*, N]}(k) := \frac{1 - \chi(2^{-h_M^*} |k|)}{-i\cancel{s}(k) + m_N + M_N(k)}, \quad (1.33)$$

where χ is a non-negative, non-increasing Gevrey-2 function⁵, such that $\chi \equiv 1$ on $[0, \frac{1}{2}]$ and $\chi \equiv 0$ on $[1, \infty)$. By the addition principle for Gaussian integrations [29, Eq. (4.21)], we can rewrite:

$$\mathcal{W}(\varphi, J, B) = -\log \int P^{(\leq h_M^*)}(\mathcal{D}\psi) e^{-\mathcal{W}^{(u.v.)}(\psi; \varphi; J; B)} \quad (1.34)$$

with

$$\mathcal{W}^{(u.v.)}(\psi; \varphi; J; B) := -\log \int P^{(h_M^*, N]}(\mathcal{D}\zeta) \int \mathbb{P}(\mathcal{D}A) e^{-V(A + J, \zeta + \psi) - (\varphi, \zeta + \psi) - i(B, J^5(\zeta + \psi))}. \quad (1.35)$$

It is an immediate consequence of the above definitions that $\mathcal{W}^{(u.v.)}(\psi; \varphi; J; B)$ can be written in the form:

$$\begin{aligned} L^2 \mathcal{F}^{(u.v.)} + \sum_{\substack{q, q', \tilde{q}, \tilde{q}', p, p' \geq 0 \\ q+q'+\tilde{q}+\tilde{q}'+p+p' \geq 1 \\ q+\tilde{q}=q'+\tilde{q}'}} \int_{\Lambda_F^q} d\underline{\eta} \int_{\Lambda_F^{q'}} d\underline{\eta}' \int_{\Lambda_F^{\tilde{q}}} d\underline{\tilde{\eta}} \int_{\Lambda_F^{\tilde{q}'}} d\underline{\tilde{\eta}'} \int_{\Lambda_B^p} d\underline{b} \int_{\Lambda_B^{p'}} d\underline{b}' \times \\ \times W_{q, q'; \tilde{q}, \tilde{q}'; p; p'}^{(u.v.)}(\underline{\eta}, \underline{\eta}'; \underline{\tilde{\eta}}, \underline{\tilde{\eta}'}; \underline{b}; \underline{b}') \prod_{k=1}^q \psi_{\eta_k}^+ \prod_{k=1}^{q'} \psi_{\eta'_k}^- \prod_{k=1}^{\tilde{q}} \varphi_{\tilde{\eta}_k}^+ \prod_{k=1}^{\tilde{q}'} \varphi_{\tilde{\eta}'_k}^- \prod_{k=1}^p G_{b_k}(J) \prod_{k=1}^{p'} B_{b'_k}, \end{aligned} \quad (1.36)$$

for suitable complex kernels $W_{q, q'; \tilde{q}, \tilde{q}'; p; p'}^{(u.v.)}$. Our main result is the following.

⁵ $f \in \mathcal{C}^\infty(\mathbb{R})$ is said to be of Gevrey- s class if $\|f^{(n)}\|_{L^\infty} \leq C_1 C_2^n n!^s$ for every $n \geq 0$ and some $C_1, C_2 > 0$.

Theorem 1.4. *There exist $\lambda_*, C_*, N_* > 0$ such that for any $0 \leq |m_N| < 2^{h_M^*}$, $N \geq h_M^* + N_*$, $L \geq 2^{-h_M^*}$, $\xi = 1$, and $0 \leq \lambda \leq \lambda_* 2^{2h_M^*}$, the following are true.*

1. The kernels of $\mathcal{W}^{(u.v.)}$ in Eq. (1.36) satisfy the following bounds. If $q + q' + \tilde{q} + \tilde{q}' > 0$ or $p + p' > 2$,

$$\|W_{q,q';\tilde{q},\tilde{q}';p;p'}^{(u.v.)}\|_1^w \leq 2^{h_M^* D_{sc}} \times \begin{cases} C_* (2^{-2h_M^*} \lambda)^{\max\{1, \frac{2}{5}q\}}, & \tilde{q}, \tilde{q}', p, p' = 0 \\ C_*^{1+p+p'+\tilde{q}+\tilde{q}'} (2^{-2h_M^*} \lambda)^{\frac{1}{8} \max\{q+q'+\tilde{q}+\tilde{q}'-2, 0\}}, & \text{otherwise,} \end{cases} \quad (1.37)$$

with $D_{sc} \equiv D_{sc}(q, q'; \tilde{q}, \tilde{q}'; p, p') := 2 - \frac{1}{2}(q + q') - \frac{3}{2}(\tilde{q} + \tilde{q}') - p - p'$. Moreover:

$$|\mathcal{F}^{(u.v.)}| \leq C_* 2^{-2h_M^*} \lambda 2^{2N}; \quad W_{0,0;0,0;0,0;1}^{(u.v.)} = 0; \quad \|W_{0,0;0,0;1,0}^{(u.v.)}\|_1^w \leq C_* 2^N; \quad (1.38)$$

$$\|g^A * W_{0,0;0,0;2,0}^{(u.v.)}\|_1^w, \|g^A * W_{0,0;0,0;1,1}^{(u.v.)}\|_1^w \leq C_* 2^{-2h_M^*}; \quad \|W_{0,0;0,0;0,2}^{(u.v.)}\|_1^w \leq C_*(N - h_M^*). \quad (1.39)$$

2. Letting $\hat{\Sigma}_{\mu,\nu}^{(u.v.)}$, $\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}$ and $\hat{S}^{(u.v.)}$ be defined as the r.h.s. of Eqs. (1.13), (1.14) and (1.16) respectively, with \mathcal{W}_ξ in the r.h.s. replaced by $\mathcal{W}^{(u.v.)}$, the following relations hold:

$$|\hat{S}_{s,s'}^{(u.v.)}(k) - \hat{g}_{s,s'}^{(h_M^*, N)}(k)| \leq \frac{C_* 2^{-h_M^*} \lambda}{2^{2h_M^*} + |k|^2}, \quad (1.40)$$

$$|\hat{\Sigma}_{\mu,\nu}^{(u.v.)}(p) - \lambda \hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.)}(p)|, |\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}(p) - \lambda \hat{\mathfrak{B}}_{5;\mu,\nu}^{(u.v.)}(p)| \leq C_* 2^{-h_M^*} \lambda^{\frac{3}{2}}, \quad (1.41)$$

for every $p, k \in \Lambda^*$ such that $|p| \leq 2^{h_M^*}$, where $\hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.)}$ and $\hat{\mathfrak{B}}_{5;\mu,\nu}^{(u.v.)}$ are suitable functions over Λ^* , such that for $p \neq 0$

$$\lim_{M \rightarrow 0^+} \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.)}(p) = \frac{1}{\pi} \left(\delta_{\mu,\nu} - \frac{p_\mu p_\nu}{|p|^2} \right) + R_{\mu,\nu}(p), \quad (1.42)$$

$$\lim_{M \rightarrow 0^+} \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \hat{\mathfrak{B}}_{5;\mu,\nu}^{(u.v.)}(p) = \frac{i}{\pi} \left(\varepsilon_{\nu,\mu} - \sum_{\alpha=0,1} \varepsilon_{\nu,\alpha} \frac{p_\mu p_\alpha}{|p|^2} \right) + R_{5;\mu,\nu}(p), \quad (1.43)$$

with $|R_{\mu,\nu}(p)|, |R_{5;\mu,\nu}(p)| \leq C_*(2^{-N} |p|)^{\frac{1}{2}}$.

Remark 1.5.

- 1) The finiteness of the kernels obtained after the ultraviolet integration is established with a wave function renormalization independent of the lattice cut-off (cf. Remark 1.2). This is due to the ξ -independence implied by our lattice regularization, which is not true with other regularizations.
- 2) The result of the integration of the ultraviolet part is an effective interaction expressed by monomials which are bounded by the correct dimensional factor $2^{h_M^* D_{sc}}$ times constants independent on N , see Eq. (1.38). Our techniques can also be used for showing the existence of the limit $N \rightarrow \infty$ of the Fourier transform of the kernels (see Appendix F).
- 3) The behavior of the two-point Schwinger function $\hat{S}^{(u.v.)}(k)$ for large momenta is the same as the non-interacting one, up to sub-leading corrections; this is a difference with the Thirring model, where the behavior in the same regime is different from the non-interacting case, modified by the presence of an anomalous critical exponent.

For small momenta instead a critical exponent for $\hat{S}^{(u.v.)}(k)$ is found, see Theorem 1.6.

4) The current and axial current correlation functions, $\hat{\Sigma}_{\mu,\nu}^{(u.v.)}$ and $\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}$ respectively, are close to their non-interacting counterparts up to corrections. Note that the lattice regularization ensures the same expressions found with dimensional regularizations; with momentum regularizations different values would be obtained, for instance the non-interacting current correlation function would be given by $\frac{1}{2\pi}(\delta_{\mu,\nu} - 2\frac{p_\mu p_\nu}{|p|^2})$, violating the conservation of the current.

1.6 Sketch of the proof

As the bosons are integrated out, the power counting is the one of a fermionic theory in 2d, with scaling exponent D_{sc} as in Theorem 1.4. In the resulting effective fermionic theory, Eq. (1.21), there are marginal terms (i.e. with $D_{sc} = 0$) either quartic in ψ or quadratic in ψ and linear in $G(J)$, and relevant (i.e. with $D_{sc} > 0$) if quadratic in ψ ; all the others are irrelevant, namely with $D_{sc} < 0$.

By decomposing the fermionic integration (cf. Eq. (1.35)) in a multiscale fashion we obtain a series expansion for the kernels in terms of the relevant and marginal ones only. The expansion turns out to be convergent as long as the relevant and marginal kernels verify suitable bounds that are uniform w.r.t. N : this is the content of Proposition 2.1. The main problem is to show that the relevant and marginal kernels are indeed bounded (Theorem 3.1). To this end the idea is to perform, at each scale, a number of manipulations within the fermionic expectations in order to *extract* some fermionic or bosonic internal lines (i.e. propagators); this is done in order to gain from the non-locality of the boson propagator, $\hat{g}^A(k) \sim |k|^{-2}$, which decays faster than the fermionic one, $\hat{g}^{(\leq N)}(k) \sim |k|^{-1}$, for large k . In doing that one has to be careful and avoid losing the good combinatorial properties of the fermionic determinants. These manipulations lead to kernels identities for the relevant and marginal ones, which can be seen by differentiating a suitable generalized action with respect to new auxiliary sources.

However, while in the case with momentum regularization one introduces only one source, which is coupled to the current [28, 7, 25], in the present case the lattice interaction produces complicate kernels (cf. Eq. (1.21) and Fig. 1) so that one is forced to add source terms as in Eq. (1.44) below, by introducing an unbounded number of auxiliary variables coupled to new, complicated fermionic terms which hardly resemble standard observables.

$$\begin{aligned}
\mathcal{V}(\psi; G; \dot{G}; \ddot{G}) := & 2^N (\mu_{N,\lambda} - 1) \int_{\Lambda_B} d\underline{b} O_b(\psi) + \sum_{n \geq 2} 2^{N(2-n)} \int_{\Lambda_B^n} d\underline{b} w_{n,0}(\underline{b}) O_{\underline{b}}^n(\psi) \\
& + \sum_{n \geq 1} n^3 2^{N(1-n)} \int_{\Lambda_B^n} d\underline{b} v_n(\underline{b}) G_{b_1}^{(n)} O_{\underline{b}}^n(\psi) + \sum_{n \geq 2} n^3 2^{N(2-n)} \int_{\Lambda_B^n} d\underline{b} v_n(\underline{b}) \dot{G}_{b_1}^{(n)} O_{\underline{b}_*}^{n-1}(\psi) \quad (1.44) \\
& + \sum_{n \geq 2} n 2^{N(1-n)} \int_{\Lambda_B^n} d\underline{b} \tilde{v}_n(\underline{b}) \ddot{G}_{b_1}^{(n)} O_{\underline{b}_*}^{n-1}(\psi),
\end{aligned}$$

where $G \equiv \{G^{(n)}\}_{n \geq 1}$, $\dot{G} \equiv \{\dot{G}^{(n)}\}_{n \geq 2}$ and $\ddot{G} \equiv \{\ddot{G}^{(n)}\}_{n \geq 2}$ (n being called *degree* index) are the aforementioned auxiliary, complex fields⁶; moreover given $\underline{b} = (b_1, \dots, b_n)$, we let $\underline{b}_* := (b_2, \dots, b_n)$, $O_{\underline{b}}^n := \prod_{j=1}^n O_{b_j}$ and $O_{\underline{b}_*}^{n-1} := \prod_{j=2}^n O_{b_j}$. To motivate such definition (see Remark 3.3 below for details), we notice that by selecting one of the fermionic lines from the kernel v_n (see Fig. 2), there are two distinct situations: when the vertex of the extracted fermionic line is a leaf of the tree or not. In the first case (Fig. 2.(a)) the contribution from the remaining part of the kernel can be written as the expectation of the monomial in the r.h.s. of Eq. (1.44)

⁶ $G \equiv \{G^{(n)}\}_{n \geq 1}$ should not be confused with the function $G(J)$ in Eq. (1.21).




Figure 2: The two possibilities arising from the extraction of a fermionic line (black solid line attached to the vertex “1”), depending on whether the vertex it is attached to, “1”, is a leaf (a) or a not-leaf (b) of the spanning tree. The red sub-tree corresponds to a source term of type $G^{(n)}$ (with $n = 7$ for (a) and $n = 3$ for (b)) and the green one to a source term of type $\dot{G}^{(n)}$ (with $n = 5$).

multiplying $G^{(n-1)}$; in the second case (Fig. 2.(b)) it is the expectation of two monomials, the one multiplying $G^{(n_1)}$ and the one multiplying $G^{(n_2)}$, with $n_1 + n_2 = n$. All such new source fields $G^{(n)}$ and $\dot{G}^{(n)}$, with $n \geq 2$, have all scaling dimension $+1$.⁷ In fact the dimensional factors $2^{N(1-n)}, 2^{N(2-n)}, 2^{N(1-n)}$ in the r.h.s of Eq. (1.44) should be regarded as $2^{ND_{sc}}$, where the scaling dimension D_{sc} of a monomial $(\psi)^{2q}(G)^p(\dot{G})^{\dot{p}}(\ddot{G})^{\ddot{p}}$ is defined as $D_{sc}(2q; p; \dot{p}; \ddot{p}) := 2 - q - p - \dot{p} - 2\ddot{p}$. The dimensionless factors n^3 and n in front of the source terms are present for technical convenience, as part of the inductive structure of our proof (see Subsection 3.5, in particular the comments after Eq. (3.35)).

One has therefore to add in the multiscale expansion such an unbounded number of new source terms, and prove self-consistently that *all* the dimensionally *marginal* and *relevant* terms, namely those with $D_{sc} = 0$ and $D_{sc} > 0$ respectively, remain uniformly bounded at all the steps of the multiscale integration. The main challenge is to show that this whole amount of new terms can be still controlled via the same mechanism based on the non-locality of the boson propagator. This is the main content of Theorem 3.1.

1.7 The infrared problem and construction of the lattice massive QED₂

Once that the ultraviolet integration is performed and $\mathcal{W}^{(u.v.)}(\psi; \varphi; J; B)$ in Eq. (1.36) has been obtained, as expressed by Theorem 1.4, the full construction of the model requires the infrared integration, see Eq. (1.34). The analysis is done via a multiscale integration which is essentially identical, up to trivial adaptation, to the one performed (for instance) in the case of the Thirring model, see [6, Sect. 2]; in that case one starts from a quartic fermionic interaction, but after the integration of the first scale, an expression similar to Eq. (1.36) is obtained, with kernels verifying the bounds Eq. (1.37); hence the analysis in [6, Sect. 2] holds essentially identical. It should be noted that the integration of the infrared scales is completely different from the ultraviolet ones; there is no dimensional improvement due to the non-locality of the interaction, as briefly explained in the Subsection 1.6, hence one has to introduce a renormalized

⁷The dimension of $G^{(n)}$ and $\dot{G}^{(n)}$ is the same as in the last line of Eq. (1.44) the number of fermionic bilinears is $n - 1$.

expansion in terms of a set of running coupling constants, corresponding to the relevant and marginal terms. The running coupling constants correspond to: the quartic interaction, the wave function renormalization, the electron mass and the amplitude of the currents, see [6, Sect. 2.1]. The expansion in the running coupling constants is convergent if they are small enough, but one has to prove that the quartic running coupling constants remain small; this follows from the asymptotic vanishing of the beta function proved in [5, Theorem 3.1]. While the effective quartic coupling remains close to its initial value in the multiscale analysis, the wave function renormalization increases as a power law, with a critical exponent given by a non-trivial, analytic function of the quartic coupling. The parameters Z_N^5 and m_N in the r.h.s. of Eq. (1.7) have to be fixed to ensure the natural physical requirements, see [4]. The result can be summarized in the following theorem.

Theorem 1.6. *Under the same conditions of Theorem 1.4, the limit $L \rightarrow \infty$ of the two-point function and of the current and axial current correlations exists. In the massless case, for $|k| \leq M$ the two-point Schwinger function behaves like:*

$$\hat{S}_{s,s'}^1(k) = \mathcal{O}(|k|^{-1+\eta}), \quad (1.45)$$

with $\eta = \alpha\lambda^2 + \mathcal{O}(\lambda^3)$, for a suitable $\alpha > 0$, and the chiral current correlation verifies:

$$\sum_{\nu=0,1} p_\nu \hat{\Sigma}_{5;\mu,\nu}^\xi(p) = \sum_{\nu=0,1} \frac{i\lambda}{\pi} \varepsilon_{\nu,\mu} p_\nu + \mathcal{O}\left(\frac{\lambda|p|^{1+\theta}}{M^\theta}\right), \quad (1.46)$$

for some $\theta \in (0, 1)$.

The anomalous divergence of the two-point Schwinger function follows from [6, Theorem 1]. The fact that, instead, no anomalous exponent appears in the ultraviolet large-momentum behavior, as stated in Eq. (1.40) of Theorem 1.4, is a crucial expected property of the massive QED₂ [17].

Details for the existence of the $L \rightarrow \infty$ limit can be found e.g. in [31, App. D]. Finally the proof of Eq. (1.46) is done in [4, Sect. VI] (assuming the validity of Theorem 1.4 proved here). The integration of the ultraviolet degrees of freedom induces the decomposition $\hat{\Sigma}_{5;\mu,\nu}^\xi = \hat{\Sigma}_{5;\mu,\nu}^{(u.v.)} + \hat{\Sigma}_{5;\mu,\nu}^{(i.r.)}$, with $\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}$ as in Theorem 1.4, satisfying $\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}(p) - \hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}(0) = \mathcal{O}(|p|/M)$. The analysis of $\hat{\Sigma}_{5;\mu,\nu}^{(i.r.)}$ is based on [5, 6, Sect. 2] and references therein. As explained in [4, Sect. VI], one introduces a suitably tuned *reference model* (see also [5, Sect. 1.2]) such that its infrared fixed point coincides with that of the original model. This implies that, letting $\tilde{\Sigma}_{5;\mu,\nu}$ be the correlation function of the reference model, $\hat{\Sigma}_{5;\mu,\nu}(p) = Z_N^5 \tilde{\Sigma}_{5;\mu,\nu}(p) + \mathcal{R}_{\mu,\nu}(p)$ [4, Eq. 44], with $\mathcal{R}_{\mu,\nu}$ continuous (in particular $\mathcal{R}_{\mu,\nu}(p) - \mathcal{R}_{\mu,\nu}(0) = \mathcal{O}((|p|/M)^\theta)$) in which we can reabsorb also the contribution $\hat{\Sigma}_{5;\mu,\nu}^{(u.v.)}$. In this way $\hat{\Sigma}_{5;\mu,\nu}^\xi(p) = Z_N^5 \tilde{\Sigma}_{5;\mu,\nu}(p) + \mathcal{R}_{\mu,\nu}(0) + \mathcal{O}((|p|/M)^\theta)$. The contribution to the anomaly from $\tilde{\Sigma}_{5;\mu,\nu}$ is explicit [4, Eq. (46)], while the one from $\mathcal{R}_{\mu,\nu}(0)$ is fixed via the WI in Eq. (1.25) [4, Eq. 50]. Combining the two, the desired result, Eq. (1.46), follows.

Theorem 1.6 establishes the non-perturbative validity of the Adler-Bardeen theorem in the lattice QED₂; note that Eq. (1.45) ensures the validity of the anomaly non-renormalization with the coefficient in agreement with perturbation theory, Eq. (1.3) (and in contrast with exact solutions or momentum regularization, where a similar result is found with a different factor). This proves that the lattice regularization of QED₂, fulfilling the physical constraints **a)-d)** in Subsection 1.1 and in the regime of Eq. (1.2), verifies a number of properties at a non-perturbative level, like the decrease of divergence degree due to the absence of contribution of longitudinal part of boson propagator, the conservation of the current and the correct value

of the anomaly (namely the Adler-Bardeen non-renormalization), whose analogous are crucial ingredients for the consistency of realistic 4d QFT models.

It is an interesting open problem if a similar result could be achieved without integrating out the bosons but with a simultaneous decomposition of bosons and fermions, which seems necessary in four dimensions. Also, it would be interesting to consider a 2d chiral model, where finding a regularization preserving WI requires the cancellation of the anomalies.

Organization of the paper. The paper is organized as follows. In Section 2 we set up the fermionic multiscale analysis in absence of external sources φ, J, B . We rely on the *Gallavotti-Nicolò tree expansion*, which is reviewed in Appendix D, and we establish the well posedness of the multiscale expansion assuming the boundedness of the relevant and marginal terms. The latter property is the main focus of Section 3, and its proof relies on some exact identities for the kernels of the effective potential, which are proved in Appendix B. In Section 4 we reintroduce the external sources φ, J, B , thus recovering the complete generating functional, Eq. (1.35). In Section 5 we collect all the information from the preceding sections, thus proving the claims of Theorem 1.4. Appendix A is devoted to the proof of two main implications of *local phase invariance* (cf. Eq. (1.24)), namely the ξ -independence property, Lemma 1.3, and the Ward Identities, Eqs. (1.25)-(1.26). In Appendix C we collect some technical results about the non-interacting *bubble diagrams*, which are essential for the analysis in Sections 2-5. Finally, in Appendix E the symmetries of the lattice theory are discussed, whereas in Appendix F we sketch the proof for showing the existence of the removed-cut-off limit.

2 The fermionic multiscale

With no loss of generality it is sufficient to prove Theorem 1.4 with $M \in [1, 2)$, i.e. $h_M^* = 0$ (cf. below Eq. (1.27)). Indeed, as one can easily check, the rescaling

$$\begin{cases} \Lambda \ni x \mapsto y \in \Lambda^{(R)} := 2^{h_M^*} \Lambda; \quad \psi_x \mapsto \psi_y^{(R)} := 2^{-\frac{1}{2}h_M^*} \psi_{2^{-h_M^*}y}; \quad \varphi_x \mapsto \varphi_y^{(R)} := 2^{-\frac{3}{2}h_M^*} \varphi_{2^{-h_M^*}y}; \\ A_{\mu,x} \mapsto A_{\mu,y}^{(R)} := A_{\mu,2^{-h_M^*}y}; \quad J_{\mu,x} \mapsto J_{\mu,y}^{(R)} := J_{\mu,2^{-h_M^*}y}; \quad B_{\mu,x} \mapsto B_{\mu,y}^{(R)} := 2^{-h_M^*} B_{\mu,2^{-h_M^*}y} \end{cases} \quad (2.1)$$

maps the original theory into another one with parameters: $M \mapsto M^{(R)} := 2^{-h_M^*} M; m_N \mapsto m_N^{(R)} := 2^{-h_M^*} m_N; \mathbf{e} \mapsto \mathbf{e}^{(R)} := 2^{-h_M^*} \mathbf{e}$. In particular $M^{(R)} \in [1, 2)$. Then if Theorem 1.4 holds for the rescaled theory, it will hold for the original one as well.

2.1 Multiscale decomposition

As a first step towards the proof of Theorem 1.4, we restrict to the case of zero external sources, namely to the analysis of $\mathcal{W}^{(u.v.)}(\psi; 0; 0; 0)$, which can be expressed as

$$\mathcal{W}^{(u.v.)}(\psi; 0; 0; 0) = -\log \int P^{(0,N]}(\mathcal{D}\zeta) e^{-\mathcal{V}(\psi + \zeta; 0; 0; 0)}, \quad (2.2)$$

where $\mathcal{V}(\psi; G; \dot{G}; \ddot{G})$ is the generalized potential defined in Eq. (1.44).⁸ The external fields φ, J and B will be reintroduced in Section 4. We start by splitting the fermion propagator as

$$g^{(0,N]}(x - y) \equiv g^{[1,N]}(x - y) = \sum_{h=1}^N g^{(h)}(x - y),$$

⁸Note that for any $N, L < \infty$, due to the finiteness of the Grassmann algebra, all the sums in the r.h.s. of Eq. (1.44) are actually truncated at $n_* \equiv 2(2^N L)^2 + 1$, and \mathcal{V} depends only upon the auxiliary field variables with degree n less or equal than n_* .

where:

$$\begin{aligned} g^{(N)}(x-y) &:= \int_{\Lambda^*} dk e^{-ik \cdot (x-y)} \frac{1 - \chi_{N-1}(|k|)}{-i\sharp(k) + M_N(k) + m_N}, \\ g^{(h)}(x-y) &:= \int_{\Lambda^*} dk e^{-ik \cdot (x-y)} \frac{\chi_h(|k|) - \chi_{h-1}(|k|)}{-i\sharp(k) + M_N(k) + m_N}, \quad 1 \leq h \leq N-1 \end{aligned}$$

with $\chi_h(|k|) \equiv \chi(2^{-h}|k|)$, and χ defined after Eq. (1.33); recall also Eq. (1.32). It is possible to check that, due to the presence of the Wilson mass $M_N(k)$,

$$\left| (-i\sharp(k) + M_N(k) + m_N)^{-1} \right| \leq \frac{\sqrt{2\pi}}{\sqrt{|k|^2 + m_N^2}}, \quad \forall k \in \Lambda^*, \quad (2.3)$$

provided that $|m_N| \leq 1$ and $N \geq 2 \log_2 \pi + 3$. Moreover, from the properties of χ_h , i.e. compact support and Gevrey-2 regularity, it follows that [32, App. A]

$$\max_{s,s' \in \{1,2\}} |g_{s,s'}^{(h)}(x-y)| \leq K 2^h e^{-\kappa_2 \sqrt{2^h |x-y|}}, \quad \forall 0 \leq h \leq N, \quad (2.4)$$

for suitable $K, \kappa_2 > 0$, uniformly w.r.t. $m_N \in [-1, 1]$. Notice that by construction $\mathcal{W}^{(u.v.)}(\psi; 0; 0; 0) = \mathcal{V}^{(0)}(\psi; 0; 0; 0)$, where, for any $0 \leq h \leq N-1$,

$$\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}) := -\log \int P^{(h,N]}(\mathcal{D}\zeta) e^{-\mathcal{V}(\psi + \zeta; G; \dot{G}; \ddot{G})} \quad (2.5)$$

and $P^{(h,N]}$ is the Grassmann integration with propagator $g^{(h,N)}(x-y) \equiv \sum_{k=h+1}^N g^{(k)}(x-y)$. Letting also $P^{(h)}$ be the integration with propagator $g^{(h)}(x-y)$, by the Gaussian addition principle [29, Eq. (4.21)], we have that

$$\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}) = -\log \int P^{(h+1)}(\mathcal{D}\zeta) e^{-\mathcal{V}^{(h+1)}(\psi + \zeta; G; \dot{G}; \ddot{G})}. \quad (2.6)$$

As a standard fact [29, Eq. (4.19)], the effective potential $\mathcal{V}^{(h)}$ can be formally computed by the formula:

$$\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}) = \sum_{s \geq 1} \frac{(-1)^{s+1}}{s!} \mathcal{E}_{(h+1)}^T \left(\mathcal{V}^{(h+1)}(\psi + \cdot; G; \dot{G}; \ddot{G}) \cdot^s \right), \quad (2.7)$$

where, for Grassmann monomials $X_1(\psi, \zeta), \dots, X_s(\psi, \zeta)$, the *truncated expectation* \mathcal{E}^T is recursively defined as

$$\begin{aligned} \mathcal{E}_{(h+1)}^T (X_1(\psi, \cdot), \dots, X_s(\psi, \cdot)) &:= \mathcal{E}_{(h+1)} (X_1(\psi, \cdot), \dots, X_s(\psi, \cdot)) + \\ &- \mathbb{1}_{(s \geq 2)} \sum_{\Pi \in \text{Part}(\{1, \dots, s\})} \sigma_\Pi \prod_{j=1}^{|\Pi|} \mathcal{E}_{(h+1)}^T (X_{\pi_{j,1}}(\psi, \cdot), \dots, X_{\pi_{j,p(j)}}(\psi, \cdot)), \end{aligned} \quad (2.8)$$

with the following understanding. $\mathcal{E}_{(h+1)}(\cdot) \equiv \int P^{(h+1)}(\mathcal{D}\zeta) \cdot$ is the *simple expectation*; Π runs over all the partitions of $\{1, \dots, n\}$ into components $\{\pi_{j,1}, \dots, \pi_{j,p(j)}\}$ indexed by $j = 1, \dots, |\Pi|$, with $p(1) + \dots + p(|\Pi|) = s$; the ordering in the r.h.s. of Eq. (2.8) is such that $\pi_{j,1} < \pi_{j+1,1}$ and $\pi_{j,k} < \pi_{j,k+1}$ for every $j = 1, \dots, |\Pi| - 1$ and $k = 1, \dots, p(j) - 1$; σ_Π is the sign needed to order the Grassmann monomials as they appear in the r.h.s of Eq. (2.8) starting from the

ordering as in the l.h.s.. Finally, in Eq. (2.7), $\mathcal{E}_{(h+1)}^T(X(\psi, \cdot);^s) := \mathcal{E}_{(h+1)}^T(X(\psi, \cdot), \dots, X(\psi, \cdot))$ when $X_1, \dots, X_s = X$. Eq. (2.7) naturally yields an integral representation for $\mathcal{V}^{(h)}$:

$$\begin{aligned} \mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}) &= \sum_{q \geq 0} \sum_{\underline{n} \in \mathbb{N}_{\geq 1}^{\times}} \sum_{\dot{\underline{n}} \in \mathbb{N}_{\geq 2}^{\times}} \sum_{\ddot{\underline{n}} \in \mathbb{N}_{\geq 2}^{\times}} \int_{\Lambda_F} d\underline{\eta} d\underline{\eta}' \int_{\Lambda_B} d\underline{b} d\dot{\underline{b}} d\ddot{\underline{b}} \times \\ &\times W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}}^{(h)|q,q}(\underline{\eta}; \underline{\eta}'; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}) \prod_{k=1}^q \psi_{\eta_k}^+ \prod_{k=1}^q \psi_{\eta'_k}^- \prod_{k=1}^{\dim \underline{n}} G_{b_k}^{(n_k)} \prod_{k=1}^{\dim \dot{\underline{n}}} \dot{G}_{\dot{b}_k}^{(\dot{n}_k)} \prod_{k=1}^{\dim \ddot{\underline{n}}} \ddot{G}_{\ddot{b}_k}^{(\ddot{n}_k)}, \end{aligned} \quad (2.9)$$

where the notation is understood as follows:

- $\mathbb{N}_{\geq p}^{\times} \equiv \emptyset \sqcup \mathbb{N}_{\geq p} \sqcup (\mathbb{N}_{\geq p})^2 \sqcup (\mathbb{N}_{\geq p})^3 \sqcup \dots$ and $\mathbb{N}_{\geq p} = \{n \in \mathbb{N} : n \geq p\}$;
- if $\underline{n} = (n_1, \dots, n_p)$, then $|\underline{n}| \equiv n_1 + \dots + n_p$ and $\dim \underline{n} \equiv p$;
- $d\underline{b} \equiv \prod_{k=1}^{\dim \underline{n}} db_k$, $d\dot{\underline{b}} \equiv \prod_{k=1}^{\dim \dot{\underline{n}}} d\dot{b}_k$ and $d\ddot{\underline{b}} \equiv \prod_{k=1}^{\dim \ddot{\underline{n}}} d\ddot{b}_k$, and each of such variables runs in Λ_B ; similarly $d\underline{\eta} = d\eta_1 \dots d\eta_q$ and each variable runs in Λ_F ;
- the term of the sum in the r.h.s. of Eq. (2.9) with $q = 0$ and $\underline{n} = \dot{\underline{n}} = \ddot{\underline{n}} = \emptyset$ must be interpreted as $W_{\emptyset; \emptyset; \emptyset}^{(h)|0,0} \in \mathbb{C}$, and its L^p norm is by definition $\|W_{\emptyset; \emptyset; \emptyset}^{(h)|0,0}\|_p^w := L^{-\frac{2}{p}} |W_{\emptyset; \emptyset; \emptyset}^{(h)|0,0}|$ (compare with Eq. (1.31)).

The kernels $W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}}^{(h)|q,q}(\underline{\eta}; \underline{\eta}'; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})$ are assumed to be anti-symmetric under the exchange of any couple of ψ^+ or ψ^- labels, and symmetric under exchange of G, \dot{G} and \ddot{G} labels separately. Such symmetries can always be imposed and imply a representation for the kernels $W^{(h)}$ as functional derivatives of $\mathcal{V}^{(h)}$: letting $p = \dim \underline{n}, \dot{p} = \dim \dot{\underline{n}}, \ddot{p} = \dim \ddot{\underline{n}}$, if $q + p + \dot{p} + \ddot{p} > 0$, Eq. (2.9) implies that

$$\begin{aligned} W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}}^{(h)|q,q}(\underline{\eta}, \underline{\eta}'; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}) &= \frac{(-1)^q}{q!^2 p! \dot{p}! \ddot{p}!} \times \\ &\times \left(\frac{\delta^{2q}}{\delta \psi_{\eta_1}^+ \dots \delta \psi_{\eta_q}^+ \delta \psi_{\eta'_1}^- \dots \delta \psi_{\eta'_q}^-} \frac{\delta^p}{\delta G_{b_1}^{(n_1)} \dots \delta G_{b_p}^{(n_p)}} \frac{\delta^{\dot{p}}}{\delta \dot{G}_{\dot{b}_1}^{(\dot{n}_1)} \dots \delta \dot{G}_{\dot{b}_{\dot{p}}}^{(\dot{n}_{\dot{p}})}} \frac{\delta^{\ddot{p}}}{\delta \ddot{G}_{\ddot{b}_1}^{(\ddot{n}_1)} \dots \delta \ddot{G}_{\ddot{b}_{\ddot{p}}}^{(\ddot{n}_{\ddot{p}})}} \mathcal{V}^{(h)} \right) (\mathbf{0}), \end{aligned} \quad (2.10)$$

where $\mathbf{0} \equiv (0; 0; 0; 0)$; moreover for complex fields $G_b^{(n)} = \text{Re}G_b^{(n)} + i\text{Im}G_b^{(n)}$ and Grassmann monomials the action of functional derivatives is given respectively by

$$\frac{\delta}{\delta G_b^{(n)}} := 2^{2N} \left(\frac{\partial}{\partial \text{Re}G_b^{(n)}} - i \frac{\partial}{\partial \text{Im}G_b^{(n)}} \right), \quad \frac{\delta}{\delta \psi_{\eta}^{\varepsilon}} \prod_{j=1}^n \psi_{\eta_j}^{\varepsilon_j} := \sum_{k=1}^n (-1)^{k-1} \delta(\eta - \eta_k) \prod_{j \neq k} \psi_{\eta_j}^{\varepsilon_j}, \quad (2.11)$$

the second of which extends to general Grassmann polynomials by linearity.

2.2 Tree expansion

By Eq. (2.7), one has that Eq. (2.9) can be written as a recursion for the kernels $W^{(h)}$ in terms of $W^{(h+1)}$. The kernels $W^{(h+1)}$ can be either left implicit, or can be further expanded in terms of $W^{(h+2)}$. As it is customary in the renormalization group context, we expand only the irrelevant kernels, i.e. those with $D_{sc} < 0$, where for a generic kernel of type $(\psi)^{2q}(G)^p(\dot{G})^{\dot{p}}(\ddot{G})^{\ddot{p}}$,

$$D_{sc}(2q; p; \dot{p}; \ddot{p}) := 2 - q - p - \dot{p} - 2\ddot{p}. \quad (2.12)$$

By iterating this procedure, we find a representation for the kernels $W^{(h)}$ in terms of all the kernels at scale N and the relevant and marginal kernels at scales $h' > h$. The outcome of this expansion is conveniently represented as an infinite sum over trees for the *single-scale contributions* $\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q} := W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q} - W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h+1)|q,q}$:

$$\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q} = \mathcal{S} \sum_{\tau \in \mathcal{T}_{h,N}}^* \sum_{\mathbf{P} \in \mathcal{P}_\tau}^{**} \widetilde{W}[\tau, \mathbf{P}], \quad (2.13)$$

with the following understanding.

1. $\mathcal{T}_{h,N}$ is the set *Gallavotti-Nicolò trees* between scale h and N . A tree $\tau \in \mathcal{T}_{h,N}$ is such that the root is not a branching point and the nodes are partially ordered from the root towards the endpoints (we write $v \prec w$ if w comes after v on the tree). The nodes v that are not the root are called *vertices*, the set of which we denote by $V(\tau)$; each vertex v comes with a *scale label* h_v , an integer between $h+1$ and $N+1$; the scale label of the root equals h by definition. The $*$ over the sum over $\mathcal{T}_{h,N}$ is the constraint that there must always be one and one only vertex v_0 , which is not an endpoint, following the root, with scale label $h_{v_0} = h+1$ (see Fig. 3).
2. \mathcal{P}_τ is a set of decorations for the tree τ , each decoration $\mathbf{P} \equiv \{P_v, Q_v\}_{v \in V(\tau)}$ being called a collection of *field labels*. Any $f \in \mathbf{P}$ comes with a sign ε_f and with a variable $\eta_f \in \Lambda_F$, so that to any f we can associate a Grassmann variable $\psi_{\eta_f}^{\varepsilon_f}$.

At each vertex $v \in V(\tau)$, Q_v is the set of *internal fields*, i.e. those corresponding to the Grassmann variables that are integrated under the action of the truncated expectation $\mathcal{E}_{(h_v)}^T$; conversely P_v identifies the *external fields* at the vertex v , i.e. the Grassmann variables that are not integrated under the action of $\mathcal{E}_{(h_v)}^T$. The field labels are subject to the following constraints.

- 2.1. If v_0 is the vertex following the root, then P_{v_0} identifies the set of fields determined by the labels $q, \underline{n}, \dot{\underline{n}}, \ddot{\underline{n}}$ of the kernel $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}$ in the l.h.s. of Eq. (2.13).
- 2.2. If $v \in V(\tau)$ is not an endpoint (we will write $v \in V_0(\tau)$), then $|Q_v| \geq 2$, namely at least two fields are always integrated.
- 2.3. If v_1, \dots, v_s are the vertices following v on τ , then $P_v \dot{\cup} Q_v = \dot{\cup}_{j=1}^s P_{v_j}$.
- 2.4. If $v \in V(\tau)$ is an endpoint (we will write $v \in V_e(\tau)$), then $Q_v = \emptyset$; besides, if $h_v < N+1$, P_v identifies a set of fields (at most four, with at least two ψ variables) corresponding to a non-irrelevant kernel (i.e. such that $D_{sc} \equiv D_{sc}(P_v) \geq 0$, see Eq. (2.12)), $W_{P_v}^{(h_v-1)}$. If $h_v = N+1$ instead, P_v can identify either an irrelevant or non-irrelevant kernel (i.e. $D_{sc}(P_v) < 0$ or ≥ 0 respectively), $W_{P_v}^{(N)}$, related to the bare potential \mathcal{V} , Eq. (1.44).
- 2.5. If $v \in V_e(\tau)$ has P_v corresponding to a non-irrelevant kernel, then the vertex v' which precedes v has scale label $h_{v'} = h_v - 1$ (this reflects the prescription above that relevant and marginal kernels are not being expanded in terms of kernels at higher scales).

The sum $\sum_{\mathbf{P} \in \mathcal{P}_\tau}^{**}$ is simply the sum over all the choices of field labels coherent with the constraints above.

3. $\widetilde{W}[\tau, \mathbf{P}]$ has the following analytical expression:

$$\begin{aligned} \widetilde{W}[\tau, \mathbf{P}](\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}) = \\ \frac{\sigma_{\mathbf{P}}}{s_{v_0}!} \int_{\Lambda_F} d\underline{\eta}_{Q_{v_0}} \mathcal{E}_{(h_{v_0})}^T(\psi_{\tilde{P}_{w_1}}, \dots, \psi_{\tilde{P}_{w_{s_{v_0}}}}) \prod_{j=1}^{s_{v_0}} \widetilde{W}[\tau_{w_j}, \mathbf{P}_{w_j}]((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_{w_j}}), \end{aligned} \quad (2.14)$$

where:

- 3.1. $\sigma_{\mathbf{P}}$ is a sign;
- 3.2. for $v \in V_0(\tau)$, w_1, \dots, w_{s_v} are the vertices following v on τ and $\mathbf{P}_{w_j} \equiv \bigcup_{v \succeq w_j} P_v$;
- 3.3. $\tilde{P}_v := P_v \cap Q_{v'}$, with v' the vertex which precedes v on τ , and $\psi_{\tilde{P}_v} \equiv \prod_{f \in \tilde{P}_v} \psi_{\eta_f}^{\varepsilon_f}$, the product being performed in a suitable, prescribed order;
- 3.4. If $v \in V_0(\tau)$, τ_v is the sub-tree with $v_0 = v$ and $\widetilde{W}[\tau_v, \mathbf{P}_v](\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_v}$ is defined recursively via Eq. (2.14); if instead $v \in V_e(\tau)$, $\widetilde{W}[\tau_v, \mathbf{P}_v](\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_v}$ is the kernel associated with the endpoint v determined by the decoration P_v , to be denoted by $W_{P_v}^{(h_v-1)}((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_v})$.
4. \mathcal{S} is the *symmetrization operator*, which acts on every kernel taking the graded average over all the permutations of labels of identical Grassmann and complex fields, so that for the overall kernel Eq. (2.10) holds.




Figure 3: Example of a Gallavotti-Nicolò tree $\tau \in \mathcal{T}_{h,N}$, where the vertices $v \in V(\tau)$ are outlined as black dots.

2.3 Bounds for the tree expansion

Note that $\widetilde{W}[\tau, \mathbf{P}]$ can be regarded as a functional in the kernels $\{W_{P_v}^{(h_v-1)}\}_{v \in V_e(\tau)}$ associated with the endpoints. In fact, an iterative application of Eq. (2.14) yields:

$$\begin{aligned} \widetilde{W}_{n;\dot{\underline{n}};\ddot{\underline{n}}}[\tau, \mathbf{P}](\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}) = \\ \tilde{\sigma}_{\mathbf{P}} \prod_{v \in V_0(\tau)} \left\{ \frac{1}{s_v!} \int_{\Lambda_F} d\underline{\eta}_{Q_v} \mathcal{E}_{(h_v)}^T(\psi_{\tilde{P}_{w_1}}, \dots, \psi_{\tilde{P}_{w_{s_v}}}) \right\} \prod_{v \in V_e(\tau)} W_{P_v}^{(h_v-1)}((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_v}) \end{aligned} \quad (2.15)$$

with $\tilde{\sigma}_{\mathbf{P}} \equiv \prod_{v \in V_0(\tau)} \sigma_{\mathbf{P}_v} \in \{\pm\}$. The tree expansion turns out to be a convenient tool for controlling the size of the kernels $\tilde{W}^{(h)}$ as in Eq. (2.13). In fact, introducing

$$\Pi^{(h,N]}(b_1, b_2) := \frac{1}{2} \int d\eta_1 d\eta'_1 d\eta_2 d\eta'_2 c_{b_1}(\eta_1, \eta'_1) c_{b_2}(\eta_2, \eta'_2) g_{\eta'_1, \eta_2}^{(h,N]} g_{\eta'_2, \eta_1}^{(h,N]}, \quad (2.16)$$

$$Q^{(N)}(\eta_1, \eta_2, \eta'_1, \eta'_2) := -\frac{1}{4} \int db db' \lambda g_{b,b'}^A \left(c_b(\eta_1, \eta'_1) c_{b'}(\eta_2, \eta'_2) - c_b(\eta_2, \eta'_1) c_{b'}(\eta_1, \eta'_2) \right), \quad (2.17)$$

the lowest-order contributions to the kernels $W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}$ and $W_{\emptyset;\emptyset;\emptyset}^{(h)|2,2}$ respectively, we can prove the following Proposition.

Proposition 2.1. *For every $\vartheta, \theta \in (0, 1)$ fixed, there exists a constant $C_0 \geq 1$ such that for any $R \geq 1$ and $0 \leq h \leq N$, the following is true. Assume that at every scale $h < k \leq N$ the following bounds hold true for the non-irrelevant kernels:*

$$\|W_{\emptyset;\emptyset;\emptyset}^{(k)|1,1}\|_1^w \leq R\lambda 2^{-\vartheta k}; \quad \|(W_{\emptyset;\emptyset;\emptyset}^{(k)|2,2} - Q^{(N)})\|_1^w \leq R\lambda^{\frac{5}{4}}; \quad (2.18)$$

$$\|W_{1;\emptyset;\emptyset}^{(k)|1,1} - c\|_1^w \leq R\lambda; \quad \|W_{n;\emptyset;\emptyset}^{(k)|1,1}\|_1^w, \|W_{\emptyset;n;\emptyset}^{(k)|1,1}\|_1^w \leq \lambda^{\frac{n-1}{2}} \text{ for } n \geq 2; \quad (2.19)$$

$$\|g^A * (W_{(1,1);\emptyset;\emptyset}^{(k)|0,0} - \Pi^{(k,N]})\|_1^w \leq R\lambda; \quad (2.20)$$

$$\|W_{(n_1, n_2);\emptyset;\emptyset}^{(k)|0,0}\|_1^w, \|W_{\emptyset;(n_1, n_2);\emptyset}^{(k)|0,0}\|_1^w, \|W_{n_1, n_2;\emptyset}^{(k)|0,0}\|_1^w \leq \lambda^{\frac{n_1+n_2-2}{2}} \text{ for } n_1 + n_2 \geq 3. \quad (2.21)$$

Then the following bounds hold true at scale h , for $\lambda \leq (C_0 R^4)^{-1}$:

$$\|W_{n;\emptyset;\emptyset}^{(h)|0;0}\|_1^w, \|W_{\emptyset;n;\emptyset}^{(h)|0;0}\|_1^w \leq C_0 2^N \lambda^{\frac{n-1}{2}}; \quad \|W_{\emptyset;\emptyset;n}^{(h)|0;0}\|_1^w \leq C_0 \lambda^{\frac{5}{8}n}; \quad (2.22)$$

$$\|g^A * W_{(1,1);\emptyset;\emptyset}^{(h)|0;0}\|_1^w \leq C_0; \quad \|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq R C_0 \lambda 2^h \quad (2.23)$$

and for all the other kernels:

$$\|W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}\|_1^w \leq C_0^{1+\dim \underline{n}+\dim \dot{\underline{n}}+\dim \ddot{\underline{n}}} \lambda^d 2^{hD_{sc}} \times \begin{cases} 2^{\theta(h-N)} & \dim \ddot{\underline{n}} > 0 \\ 1 & \text{otherwise,} \end{cases} \quad (2.24)$$

with $D_{sc} \equiv D_{sc}(2q; \dim \underline{n}; \dim \dot{\underline{n}}; \dim \ddot{\underline{n}})$, see Eq. (2.12), and

$$d \equiv d(2q; \underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}) := \begin{cases} \max\{1, \frac{2}{5}q\}, & \dim \underline{n}, \dim \dot{\underline{n}}, \dim \ddot{\underline{n}} = 0 \\ \frac{1}{4} \max\{q-1, 0\} + \frac{1}{2}(|\underline{n}| + |\dot{\underline{n}}| - \dim \underline{n} - \dim \dot{\underline{n}}) + \frac{5}{8}|\ddot{\underline{n}}|, & \text{otherwise.} \end{cases} \quad (2.25)$$

Observe that Eqs. (2.18)-(2.21) are stronger than Eqs. (2.22)-(2.23) with $h = k$, for the non-irrelevant kernels. For this reason, Eqs. (2.22)-(2.23) will be called *standard dimensional bounds*, while we will refer to Eqs. (2.18)-(2.21) as *improved bounds*. Proposition 2.1 establishes the validity of the *standard dimensional bounds* taking as input the *improved bounds* at higher scales. The proof of the latter is the content of Section 3.

Proof. Proposition 2.1 is actually an immediate consequence of the following bounds for the kernels $\tilde{W}^{(h)}$ (cf. Eq. (2.13)):

$$\|\tilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}\|_1^w \leq (1 - 2^{\theta-1}) C_0^{1+\dim \underline{n}+\dim \dot{\underline{n}}+\dim \ddot{\underline{n}}} 2^{hD_{sc}} \lambda^d \times \begin{cases} R & |\underline{n}|, |\dot{\underline{n}}|, |\ddot{\underline{n}}| = 0, q = 1 \\ 2^{\theta(h-N)} & |\ddot{\underline{n}}| \geq 1 \\ 1 & \text{otherwise,} \end{cases} \quad (2.26)$$

under the same hypotheses of Proposition 2.1. Eq. (2.26) is proved in Appendix D.

For any irrelevant kernel with $|\underline{n}| \geq 1$, by construction we have:

$$W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q} = \sum_{k=h}^{N-1} \widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(k)|q,q} + W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(N)|q,q},$$

so that, combining Eq. (2.26) with the bounds for the kernels at scale N , we find:

$$\begin{aligned} \|W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}\|_1^w &\leq \sum_{k=h}^{N-1} 2^{kD_{sc}} \|\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(k)|q,q}\|_{(k)} + \|W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(N)|q,q}\|_1^w \\ &\leq \sum_{k=h}^N 2^{kD_{sc}} (1 - 2^{\theta-1}) C_0^{1+\dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}} \lambda^d 2^{\theta(k-N)} \\ &\leq C_0^{1+\dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}} \lambda^d 2^{hD_{sc}} 2^{\theta(h-N)}, \end{aligned} \quad (2.27)$$

where we used that $\|W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(N)|q,q}\|_1^w \leq 2^{ND_{sc}} (1 - 2^{\theta-1}) C_0^{1+\dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}} \lambda^d$ as well, and that for irrelevant kernels $D_{sc} \leq -1$. For $W_{\emptyset;\emptyset;n}^{(h)|0,0}$ and for the irrelevant kernels such that $|\underline{n}| = 0$, the estimate follows the same lines as in Eq. (2.27), except for the absence of the *short memory factor* $2^{\theta(h-N)}$. Following the same steps as Eq. (2.27), we also find the estimates in Eq. (2.22) for the relevant kernels $W_{n;\emptyset;\emptyset}^{(h)|0,0}$ and $W_{\emptyset;n;\emptyset}^{(h)|0,0}$, which are however unbounded in N , since $D_{sc} = +1$. In fact for the other non-irrelevant kernels we cannot proceed as in Eq. (2.27). Let us e.g. look at the kernel ψ^2 , which is relevant. Writing $W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1} = \widetilde{W}_{\emptyset;\emptyset;\emptyset}^{(h)|1,1} + W_{\emptyset;\emptyset;\emptyset}^{(h+1)|1,1}$, we use Eq. (2.26) for estimating the first term, while for the second one we exploit the first bound in Eq. (2.18):

$$\|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq \|\widetilde{W}_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w + \|W_{\emptyset;\emptyset;\emptyset}^{(h+1)|1,1}\|_1^w \leq (1 - 2^{\theta-1}) C_0 R \lambda 2^h + R \lambda 2^{-\vartheta(h+1)} \leq C_0 R \lambda 2^h,$$

if $C_0 \geq 2^{-2\theta+1}$. The estimates for the other non-irrelevant kernels qualitatively follow the very same lines, so they will be omitted for sake of brevity. All in all one recovers the bounds in Eqs. (2.22)- (2.23), with C_0 (the same in Proposition 2.1 and Eq. (2.26)) independent of R , and λ smaller than $(C_0 R^4)^{-1}$, thus proving the claim of Proposition 2.1. \square

Proposition 2.1 shows that the multiscale expansion is well posed and well behaved provided that the relevant and marginal kernels (i.e. those with $D_{sc} \leq -1$ according to Eq. (2.24)), satisfy suitable bounds at all the higher scales. This property, whose proof is actually a crucial point of the present work and will be discussed in the next section, is in general a crucial point in every renormalization group analysis.

3 Bounds for the relevant and marginal terms

In the present work, in the same way as [7, 24] and similarly to [28], the key underlying idea for controlling the non-irrelevant terms is to exploit the non-locality of the boson propagator (see Eq. (1.27) and lines thereafter).

Theorem 3.1. *For every $\vartheta, \theta \in (0, 1)$ fixed, there exist $N_0 \geq 1$, $R \geq 1$ and $\lambda_0 > 0$ such that, for any $N \geq N_0$ and $\lambda \leq \lambda_0$, the bounds in Eqs. (2.18)-(2.21) and Eqs. (2.22)-(2.24) for $k = h$ hold together at every scale $0 \leq h \leq N$.*

For the implications of Theorem 3.1 on the kernels bounds appearing in the main Theorem 1.4, see Subsection 5.1.

Proof of Theorem 3.1. The proof is by induction over the scale index h of the kernel: we will think of R entering the bounds in the inductive hypothesis as a free parameter and the idea is to show that the inductive step can be performed if $\lambda \leq \lambda_0$ with λ_0 small enough with respect to R , and R large but fixed. It turns out that a working choice for R and λ_0 is

$$R \geq \mathfrak{C} := \max_{j=0}^4 \mathfrak{c}_j C_0^8, \quad \lambda_0 = \min \left\{ (C_0 R^4)^{-1}, \min_{j=0}^7 (\mathfrak{c}'_j C_0^{12} R)^{-1} \right\}, \quad (3.1)$$

where $C_0 \geq 1$ is the same as in Proposition 2.1 while \mathfrak{c}_j and \mathfrak{c}'_j are constants independent of C_0, R that will appear along the proof in the present section.

To see the origin of Eq. (3.1), we start by observing that for the base of the induction, i.e. at scale $h = N$, since the potential has the explicit expression in Eq. (1.44), it is possible to find constants $\mathfrak{c}_0, \mathfrak{c}'_0 \geq 1$ such that bounds in Eqs. (2.22)-(2.24), (2.18)-(2.21), hold for $\lambda \leq (\mathfrak{c}'_0)^{-1}$ and $R \geq \mathfrak{c}_0$.

For the inductive step, we assume the claim true up to scale $h + 1$, and we prove its validity at scale h . As a first step, we can apply Proposition 2.1, which implies the validity of bounds in Eqs. (2.22)-(2.24) at scale h , for $\lambda \leq \lambda_0$ with R, λ_0 as above (note that by construction $\lambda_0 \leq (C_0 R^4)^{-1}$). The nontrivial point is now to show the validity of bounds in Eqs. (2.18)-(2.21) at scale h , taking for granted Eqs. (2.22)-(2.24) at the same scale. The remaining part of this section is dedicated to the proof of this fact. For sake of clarity, in order to avoid confusion with constants, throughout this section we will regard R and C_0 as generic parameters greater than 1 appearing in the bounds in Eqs. (2.22)-(2.24), so we will track explicitly their dependence in the upcoming estimates. Conversely, we will use the symbols $\mathfrak{c}, \mathfrak{c}', \mathfrak{c}'', \dots$ to denote quantities that are independent on R and C_0 .

3.1 The electron self-energy

We start by proving the bound regarding the kernel $W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}$ associated to the Grassmann monomial ψ^2 , also called *electron self-energy*. The goal of this section is to show:

$$\|W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}\|_1^w \leq \mathfrak{c}_1 C_0^5 \lambda 2^{-\vartheta h}, \quad (3.2)$$

with C_0 as in Proposition 2.1, $\lambda \leq (\mathfrak{c}'_1 C_0^2 R)^{-1}$, and suitable constants $\mathfrak{c}_1, \mathfrak{c}'_1 \geq 1$ independent on C_0, R . The starting point consists of deriving some convenient representation for the kernel.

Lemma 3.2. *The following identity holds:*

$$W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}(\eta_1, \eta_2) = \sum_{j=1}^4 \mathcal{A}_j(\eta_1, \eta_2) \quad (3.3)$$

where, recalling $\mu_{N,\lambda} = e^{-\frac{1}{2}\lambda 2^{-2N} g_{\mu,\mu}^A(0)}$, $\mathcal{A}_j(\eta_1, \eta_2)$ are given by

$$\mathcal{A}_1(\eta_1, \eta_2) = 2^N (\mu_{N,\lambda} - 1) \int db \left(c_b(\eta_1, \eta_2) - \int d\eta d\eta' c_b(\eta_1, \eta) g_{\eta, \eta'}^{(h,N]} W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2) \right), \quad (3.4)$$

$$\begin{aligned} \mathcal{A}_2(\eta_1, \eta_2) = & \sum_{n \geq 1} \frac{\mu_{N,\lambda}}{n^2} \int db_1 db \lambda g_{b_1, b}^A \left\{ c_{b_1}(\eta_1, \eta_2) W_{n; \emptyset; \emptyset}^{(h)|0,0}(b) + \int d\eta d\eta' c_{b_1}(\eta_1, \eta) g_{\eta, \eta'}^{(h,N]} \times \right. \\ & \left. \times \left[W_{n; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b) - W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2) W_{n; \emptyset; \emptyset}^{(h)|0,0}(b) \right] \right\}, \end{aligned} \quad (3.5)$$

$$\begin{aligned}
\mathcal{A}_3(\eta_1, \eta_2) &= \\
&= 2^{-N} \sum_{n_1 \geq 1, n_2 \geq 2} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \int db_1 db db' \lambda g_{b_1, b}^A \delta(b_1 - b') \times \\
&\times \left\{ c_{b_1}(\eta_1, \eta_2) \left[-W_{n_1; n_2; \emptyset}^{(h)|0,0}(b; b') + W_{n_1; \emptyset; \emptyset}^{(h)|0,0}(b) W_{\emptyset; n_2; \emptyset}^{(h)|0,0}(b') \right] + \int d\eta d\eta' c_{b_1}(\eta_1, \eta) g_{\eta, \eta'}^{(h, N)} \times \right. \\
&\times \left[-W_{n_1; n_2; \emptyset}^{(h)|1,1}(\eta', \eta_2; b; b') + W_{n_1; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b) W_{\emptyset; n_2; \emptyset}^{(h)|0,0}(b') + W_{n_1; \emptyset; \emptyset}^{(h)|0,0}(b) W_{\emptyset; n_2; \emptyset}^{(h)|1,1}(\eta', \eta_2; b') \right] \\
&\left. - \int d\eta d\eta' c_{b_1}(\eta_1, \eta) g_{\eta, \eta'}^{(h, N)} W_{\emptyset; \emptyset; n}^{(h)|1,1}(\eta', \eta_2) \left[-W_{n_1; n_2; \emptyset}^{(h)|0,0}(b; b') + W_{n_1; \emptyset; \emptyset}^{(h)|0,0}(b) W_{\emptyset; n_2; \emptyset}^{(h)|0,0}(b') \right] \right\}, \tag{3.6}
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_4(\eta_1, \eta_2) &= \sum_{n \geq 2} 2^{-N} \int db_1 db \delta(b_1 - b) \left\{ c_{b_1}(\eta_1, \eta_2) W_{\emptyset; \emptyset; n}^{(h)|0,0}(b) + \int d\eta d\eta' c_{b_1}(\eta_1, \eta) \times \right. \\
&\times g_{\eta, \eta'}^{(h, N)} \left[W_{\emptyset; \emptyset; n}^{(h)|1,1}(\eta', \eta_2; b) - W_{\emptyset; \emptyset; n}^{(h)|1,1}(\eta', \eta_2) W_{\emptyset; \emptyset; n}^{(h)|0,0}(b) \right] \left. \right\}, \tag{3.7}
\end{aligned}$$

where \mathcal{N}_{n_1, n_2} are suitable positive coefficients, such that $\mathcal{N}_{n_1, n_2} \leq \frac{(n_1 + n_2)^2}{n_1^2 n_2^2}$.

The proof of Lemma 3.2 given in Appendix B.

Notation. In order to show Eq. (3.2) we find convenient to exploit a graphical representation associated to the decomposition in Eq. (3.3), as shown in Fig. 4. For $\sharp = a, b, \dots, o$, we will denote by $\text{Val}(\sharp)(\eta, \eta')$, the analytical value of the diagram (\sharp) in Fig. 4, i.e. the corresponding term in the decomposition of Eq. (3.3) and by $\|\text{Val}(\sharp)\|_1^w$ its weighted norm, cf. Eq. (1.31). Even though there are 15 diagrams to discuss, we will see that only a fraction of them is independent, in a suitable sense (cf. Remark 3.4).

Remark 3.3. The decomposition in Fig. 4 is paradigmatic for understanding the combinatorial difficulty emerging from the lattice regularization of the theory. In the continuum version of this model [25], the only term in the r.h.s. would be diagram (d) with $n = 1$ (see [25, Fig.2]); similarly in [4], where only the terms ψ^2, ψ^4 and $\psi^2 G(J)$ are retained in the fermionic interaction (cf. Eq. (1.21)), one would get only diagrams $(a), (b), (d)$ and (e) with $n = 1$. In the present work, the multitude of terms in the expansion is due to the form of the “bare kernels” v_n , Eq. (1.29), which involve sums over trees of arbitrary order. If we adopt the representation in Fig. 5.(a) for the bare kernels v_n , we see that terms like the one in Fig. 5.(b) (corresponding to diagram (h) of Fig. 4) can emerge as contributions to the kernel $W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}$.

The blob in the l.h.s. of Fig 5.(b) should be regarded as

$$\frac{\delta^3 \mathcal{V}^{(h)}}{\delta G_{b_2}^{(1)} \delta G_{b_3}^{(1)} \delta G_{b_4}^{(3)}}(\mathbf{0}) \equiv 3! W_{(1, 1, 3); \emptyset; \emptyset}^{(h)|0,0}(b_2, b_3, b_4).$$

The factor 3 in the factorial corresponds to the number of G fields, which equals the number of wiggly lines attached to the blob in the diagram. Now, since in general we must expect an arbitrary number (say s) of wiggly lines in the l.h.s. of Fig. 5.(b), following this expansion we would end up with unbounded factorials (say $s!$), which would induce a combinatorial divergence of the expansion. The way we overcome this problem is by rewriting the diagram in the l.h.s. of Fig. 5.(b) as in the r.h.s., namely by “hiding” the bunch of wiggly lines at the end of the dashed line (which stands for a Dirac delta): in this way no combinatorial problem arises, being the number of points of the blob bounded. The role of the auxiliary source \tilde{G} is exactly to perform analytically what here is explained here by a graphical intuition.




Figure 4: Graphical representation of Eq. (3.3). The graphs are numbered as in the order they appear in the expansion of Eq. (3.3): graphs (a), (b) are relative to \mathcal{A}_1 , (c) – (e) to \mathcal{A}_2 , (f) – (l) to \mathcal{A}_3 and (m) – (o) to \mathcal{A}_4 . Solid and wiggly lines, with dotted endpoints, represent fermion and boson propagators respectively, while dashed and curly lines, with black dotted endpoints, represent delta functions. Blobs (big North East line patterned circles) represent kernels, associated to external fields distinguished by the type of attached lines to them: each solid, wiggly, dashed and curly line attached to a blob identifies a ψ , G , \dot{G} and \ddot{G} field respectively.

Diagram (a). This is simply the bare quartic kernel, whose weighted norm can be bounded by (cf. Eq. (1.27))

$$\|\text{Val}(a)\|_1^w \leq 2^N \|c\|_1^w |\mu_{N,\lambda} - 1| \leq 12K' \lambda N 2^{-N} \leq \mathfrak{c}_{1,1} \lambda 2^{-\vartheta N},$$

where we used definitions in Eqs. (1.19), (1.31) (recall $r = 1$, cf. below Eq. (1.10)) to get $\|c\|_1^w \leq 6e^{\hat{\kappa}2^{-\frac{N}{2}-1}} \leq 12$ for N large enough, the explicit expression of $\mu_{N,\lambda}$ (below Eq. (3.26)) and the bound on $|g_{\mu,\mu}^A(0)| \leq \|g^A\|_\infty \leq K'N$ (cf. Eq. (1.28)).

Diagram (b). The bound in this case is

$$\|\text{Val}(b)\|_1^w \leq 2^N \|c\|_1^w |\mu_{N,\lambda} - 1| \|g^{(h,N]}\|_1^w \|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq \tilde{K} C_0 R \lambda 2^N \|c\|_1^w |\mu_{N,\lambda} - 1|,$$

where we used that for some⁹ $\tilde{K} > 0$, $\|g^{(h,N]}\|_1^w \leq \tilde{K} 2^{-h}$ and that, by assumption, if $\lambda \leq (C_0 R^4)^{-1}$ we have $\|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq C_0 R \lambda 2^h$. Taking $\lambda \leq \min\{(C_0 R^4)^{-1}, (R C_0 \tilde{K})^{-1}\}$ we see that

⁹One can choose, e.g., $\tilde{K} = 320K\hat{\kappa}^{-2}$, cf. Eq. (2.4).




Figure 5: (a): synthetic representation for $v_n(\underline{b})O_b^n$, obtained from Fig. 1 by first collapsing each sub-tree rooted in b_2, b_3 and b_4 into one dark blob (the label (n) stands for the order of the sub-tree), and then by thinking of the sub-trees rooted at b_3 and b_4 as one only tree rooted at b' , with $b' = b$ (the dashed line stands for a Dirac delta). (b): contribution to the kernel $W_{\emptyset;0;\emptyset}^{(h)|1,1}$ arising from the bare kernel in (a).

$\text{Val}(b)$ can be bounded as $\text{Val}(a)$, i.e.

$$\|\text{Val}(b)\|_1^w \leq \mathfrak{c}_{1,1} \lambda 2^{-\vartheta N}. \quad (3.8)$$

The constraint on λ implies that $\mathfrak{c}'_1 \geq \tilde{K}$.

Remark 3.4. Diagram (b) is the first example that not all the graphs of Fig. 4 have independent bounds: each time that in a diagram we have a “dressed fermionic line” (first diagram in the first line of Fig. 4), in order to obtain an upper bound, we can graphically delete the dressed fermionic line at the price of shrinking λ , which, in the end, reflects in changing the values of the constant \mathfrak{c}'_1 in Eq. (3.2). In this way, we can say that graphs (b), (k) and (o) are apriori bounded by (a), (f) and (m) respectively.

Diagram (c). In this case $\text{Val}(c)(\eta_1, \eta_2) = 0$. This is actually a consequence of the *charge conjugation* symmetry (cf. Lemma E.1), which establishes the following cancellation:

$$\int_{\Lambda_B^p} db'_1 \dots db'_p W_{(n_1, \dots, n_p); \emptyset; \emptyset}^{(h)|0,0}(b'_1, \dots, b'_p) \prod_{j=1}^p g_{b_j, b'_j}^A = 0, \quad (3.9)$$

for every $p \geq 1$ odd, $n_1, \dots, n_p \geq 1$ and $b_1, \dots, b_p \in \Lambda_B$. In particular for $p = 1$ we find $\text{Val}(c)(\eta_1, \eta_2) = 0$. The proof of Eq. (3.9) follows immediately by noting that, in force of the *charge conjugation* symmetry (cf. Lemma E.1),

$$W_{(n_1, \dots, n_p); \emptyset; \emptyset}^{(h)|0,0}(b'_1, \dots, b'_p) = W_{(n_1, \dots, n_p); \emptyset; \emptyset}^{(h)|0,0}(\bar{b}'_1, \dots, \bar{b}'_p),$$

where recall that if $b = (x, \mu, \epsilon)$, $\bar{b} \equiv (x, \mu, -\epsilon)$. Using that $g_{b,b'}^A = -g_{b,\bar{b}'}^A$ and the fact that p is odd, we find:

$$\begin{aligned} \int d\bar{b}' \prod_{j=1}^p g_{b_j, b'_j}^A \times W_{\underline{n}; \emptyset; \emptyset}^{(h)|0;0}(\bar{b}') &= \int d\bar{b}' \prod_{j=1}^p g_{b_j, b'_j}^A \times W_{\underline{n}; \emptyset; \emptyset}^{(h)|0;0}(\bar{b}') = \\ &= \int d\bar{b}' \prod_{j=1}^p g_{b_j, \bar{b}'_j}^A \times W_{\underline{n}; \emptyset; \emptyset}^{(h)|0;0}(\bar{b}') = - \int d\bar{b}' \prod_{j=1}^p g_{b_j, b'_j}^A \times W_{\underline{n}; \emptyset; \emptyset}^{(h)|0;0}(\bar{b}'), \end{aligned}$$

as desired.

Diagram (d). This is the first case where we must use the non-locality of g^A . The analytical value of the diagram is

$$\text{Val}(d)(\eta_1, \eta_2) = \sum_{n \geq 1} \frac{\mu_{N,\lambda}}{n^2} \int db db' \int d\eta d\eta' c_b(\eta_1, \eta) \lambda g_{b,b'}^A g_{\eta,\eta'}^{(h,N]} W_{n; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b'),$$

which can be bounded as

$$\begin{aligned} \|\text{Val}(d)\|_1^w &\leq \\ &\leq \sum_{n \geq 1} \frac{\lambda}{n^2} \int \frac{db db'}{L^2} \int d\eta d\eta' d\eta_1 d\eta_2 e^{\frac{\kappa}{2} \sqrt{\delta^D(\eta, \eta', \eta_1, \eta_2, b, b')}} |c_b(\eta_1, \eta)| |g_{b,b'}^A| |g_{\eta,\eta'}^{(h,N]}| |W_{n; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b')| \\ &\leq \sum_{n \geq 1} \frac{\lambda}{n^2} \int \frac{db db'}{L^2} \int d\eta d\eta' d\eta_1 d\eta_2 \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(b, \eta, \eta_1)}} |c_b(\eta_1, \eta)| \right) \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(b-b')}} |g_{b,b'}^A| \right) |g_{\eta,\eta'}^{(h,N]}| \times \\ &\quad \times \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(\eta', \eta_2, b')}} |W_{n; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b')| \right) = \\ &= \sum_{n \geq 1} \frac{\lambda}{n^2} \int \frac{db db'}{L^2} \int d\eta' d\eta_2 \tilde{g}_{b,\eta'}^{(h,N]} \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(b, b')}} |g_{b,b'}^A| \right) \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(\eta', \eta_2, b')}} |W_{n; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b')| \right), \end{aligned}$$

with $\tilde{g}_{b,\eta'}^{(h,N]} := \int d\eta d\eta_1 e^{\frac{\kappa}{2} \sqrt{\delta^D(b, \eta, \eta_1)}} |c_b(\eta_1, \eta)| |g_{\eta,\eta'}^{(h,N]}|$. We proceed by applying the Hölder's inequality with weights $p = \frac{2}{1-\vartheta}$ and $p' = \frac{2}{1+\vartheta}$:

$$\|\text{Val}(d)\|_1^w \leq \sum_{n \geq 1} \frac{\lambda}{n^2} \|g^A\|_p^w \|\tilde{g}^{(h,N]}\|_{p'} \|W_{n; \emptyset; \emptyset}^{(h)|1,1}\|_1^w \leq \sum_{n \geq 1} \frac{\lambda}{n^2} \tilde{K}_\vartheta C_0^2 2^{-\vartheta h} \lambda^{\frac{n-1}{2}} \leq \lambda \mathfrak{c}_{1,2} C_0^2 2^{-\vartheta h},$$

where we used Eq. (2.24) with $\lambda \leq (C_0 R^4)^{-1}$, denoted with \tilde{K}_ϑ the constant coming from $\|\cdot\|_p, \|\cdot\|_{p'}$ (by using Eq. (1.27)) and used the fact that $\tilde{g}^{(h,N]}$ has the same dimensional bounds as $g^{(h,N]}$, being $c_b(\eta_1, \eta)$ local; finally $\mathfrak{c}_{1,2} = \tilde{K}_\vartheta \pi^2/6$ is independent of C_0, R .

Diagram (f). This is the first diagram with a dashed line, corresponding to a delta function (cf. the first term in \mathcal{A}_3 , Eq. (3.6)). In bounding such term, this forces us to take the L^∞ norm of the boson propagator associated with the wiggly line:

$$\begin{aligned} \|\text{Val}(f)\|_1^w &\leq \\ &\leq \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} 2^{-N} \lambda \|g^A\|_\infty \|W_{n_1; n_2; 0}^{(h)|0,0}\|_1^w \leq \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{\lambda}{n_1^2 n_2^2} 2^{-N} K' N C_0^3 \lambda^{\frac{n_1+n_2-2}{2}} \leq \mathfrak{c}_{1,3} C_0^3 \lambda^{3/2} 2^{-\vartheta N}, \end{aligned}$$

where we used Eqs. (1.27), (2.24), that $2^{-N} N \leq 2^{-\vartheta N}$ for any $\vartheta \in (0, 1)$ if N is large enough, and the fact that $\mathcal{N}_{n_1, n_2} \leq \frac{(n_1+n_2)^2}{n_1^2 n_2^2}$; $\mathfrak{c}_{1,3} = K' (\pi^2/6)^2$ does not depend on C_0, R .

Remark 3.5. Diagram (f) is the first nontrivial case (after diagram (a)) of a class of graphs which not only have the desired bound at any fixed scale h , but they are actually vanishing as $N \rightarrow \infty$. In view of proving Theorem 3.1 it is sufficient to replace the factor $2^{-\vartheta N}$ above with $2^{-\vartheta h}$; however, it is worth mentioning that tracking small terms in N would be crucial for the analysis of the $N \rightarrow \infty$ limit of the theory (see Appendix F for an overview).

Diagram (h). Again, due to the delta function associated with the dashed line, we are forced to take the L^∞ norm of both g^A and $g^{(h,N]}$. We find:

$$\|\text{Val}(h)\|_1^w \leq \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} 2^{-N} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \lambda \|g^A\|_\infty \|g^{(h,N)}\|_\infty \|W_{n_1; n_2; \emptyset}^{(h)|1,1}\|_1^w \leq C_0^3 K N \mathfrak{c}_{1,3} \lambda 2^{-h}, \quad (3.10)$$

with $\mathfrak{c}_{1,3}$ as in the previous item, and we used that $\|g^{(h,N)}\|_\infty \leq 2K2^N$ (cf. Eq. (2.4)) and that $\|W_{n_1; n_2; \emptyset}^{(h)|1,1}\|_1^w \leq C_0^3 \lambda^{\frac{n_1+n_2-2}{2}} 2^{-h}$, cf. Eq. (2.24). This apparent logarithmic divergence is related to the definition of the external source \dot{G} , which imposes to take the L^∞ norm of both g^A and $g^{(h,N)}$. We will show at the end of this paragraph (see Eq. (3.12) and below) how to solve this divergence; for the moment let us continue with the other graphs of Fig. 4.

Diagram (i). The analytical value in this case is:

$$\begin{aligned} \text{Val}(i)(\eta_1, \eta_2) = & \\ & \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} 2^{-N} \int d\eta d\eta' \int db db' c_b(\eta_1, \eta) \lambda g_{b, b'}^A g_{\eta, \eta'}^{(h,N)} W_{n_1; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b') W_{\emptyset; n_2; \emptyset}^{(h)|0,0}(b) \end{aligned} \quad (3.11)$$

and by noting that from translation invariance and Items 2, 3 of Lemma E.1, one has $W_{\emptyset; n_2; \emptyset}(b) = W_{\emptyset; n_2; \emptyset}((0, 0, +))$, the bound for this graph can be deduced from the one for graph (d):

$$\begin{aligned} \|\text{Val}(i)\|_1^w & \leq \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{2^{-N} \mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \|W_{\emptyset; n_2; \emptyset}^{(h)|0,0}\| \int \frac{d\eta_1 d\eta_2}{L^2} e^{\frac{\kappa}{2} \sqrt{\delta^D(\eta_1, \eta_2)}} \times \\ & \quad \times \left| \int d\eta d\eta' \int db db' c_b(\eta_1, \eta) \lambda g_{b, b'}^A g_{\eta, \eta'}^{(h,N)} W_{n_1; \emptyset; \emptyset}^{(h)|1,1}(\eta', \eta_2; b') \right| \leq \\ & \leq \sum_{n_2 \geq 2} \frac{1}{n_2^2} 2^{-N} \|W_{\emptyset; n_2; \emptyset}^{(h)|0,0}\|_1^w |\mu_{N, \lambda}^{-1}| \|\text{Val}(d)\|_1^w \leq C_0^3 (\pi^2/3) \mathfrak{c}_{1,2} \lambda^{3/2} 2^{-\vartheta h} \end{aligned}$$

where we used the bound of diagram (d), again that $\mathcal{N}_{n_1, n_2} \leq (n_1 + n_2)^2 (n_1 n_2)^{-2}$, that $|\mu_{N, \lambda}^{-1}| \leq 2$ for N large, and that $\|W_{\emptyset; n_2; \emptyset}^{(h)|0,0}\|_1^w \leq C_0 2^N \lambda^{\frac{n_2-1}{2}}$ by Eq. (2.22). Note that imposing also $\lambda \leq (C_0^2 \pi^4/9)^{-1}$ one has $\|\text{Val}(i)\|_1^w \leq C_0^2 \mathfrak{c}_{1,2} \lambda 2^{-\vartheta h}$ which is the same as case (d).

Remark 3.6. We observe a second reduction of the number of diagrams to bound related to the presence of a *dashed tadpole graph*, i.e. a blob with only one dashed external line, as in (g), (i), (l). By possibly shrinking λ , i.e. changing the value of \mathfrak{c}'_1 in Eq. (3.2), one has that $\text{Val}(g), \text{Val}(i), \text{Val}(\ell)$ can be bounded exactly as $\text{Val}(e), \text{Val}(d), \text{Val}(c)$ respectively (in particular $\text{Val}(g) = 0$).

Diagram (m). We have to bound the first of Eq. (3.7) which immediately gives

$$\|\text{Val}(m)\|_1^w \leq \sum_{n \geq 2} 2^{-N} \|c\|_1^w \|W_{\emptyset; \emptyset; n}^{(h)|0,0}\|_1^w \leq 12 \sum_{n \geq 2} 2^{-N} C_0 \lambda^{\frac{5}{8}n} \leq 12 C_0 \lambda^{5/4} 2^{-\vartheta h},$$

where we used $\|c\|_1^w \leq 12$ as in case (a), Eq. (2.22) and $\lambda \leq 1/2^{8/5}$ so that $(1 - \lambda^{5/8})^{-1} \leq 2$: this implies that $c'_1 \geq 2^{8/5}$.

Diagram (n). The curly line is again associated to a delta function, which forces us to take the L^∞ norm of the fermion propagator. The bound is:

$$\|\text{Val}(n)\|_1^w \leq \sum_{n \geq 2} 2^{-N} \|g^{(h,N)}\|_\infty \|W_{\emptyset; \emptyset; n}^{(h)|1,1}\|_1^w \leq \sum_{n \geq 2} 2 K C_0^2 \lambda^{\frac{5}{8}n} 2^{\theta(h-N)} 2^{-h} \leq 4 K C_0^2 \lambda^{5/4} 2^{-\vartheta h},$$

where we used Eq. (2.24) and the same hypothesis on λ of the previous item.

Improvement for (h). Finally, we return to diagram (h). The way to overcome the apparent divergence is to extract a second line, i.e. to perform an expansion for the kernel $W_{n_1; n_2; \emptyset}^{(h)|1,1}$: this allows to solve the apparent logarithmic divergence with an expansion in terms of a finite number of extra diagrams, each having good dimensional estimates. In analogy with Eq. (3.3), it is possible to obtain the following identity, proved in Appendix B:

$$\begin{aligned} W_{n_1; n_2; \emptyset}^{(h)|1,1}(\eta, \eta'; b_1, b_2) &= \\ &= \frac{n_2^2}{(n_2 - 1)^2} \mu_{N, \lambda} \int db \lambda g_{b_2, b}^A 2 W_{(n_1, n_2 - 1); \emptyset; \emptyset}^{(h)|1,1}(\eta, \eta'; b_1, b) + \\ &- \sum_{\substack{n'_1 \geq 1, n'_2 \geq 2 \\ n'_1 + n'_2 = n_2}} 2^{-N} \mathcal{N}_{n'_1, n'_2} \int db'_1 db'_2 \lambda g_{b_2, b'_1}^A \delta(b_2 - b'_2) \left[2 W_{(n_1, n'_1); n'_2; \emptyset}^{(h)|1,1}(\eta, \eta'; b_1, b'_1; b'_2) + \right. \\ &\left. - 2 W_{(n_1, n'_1); \emptyset; \emptyset}^{(h)|1,1}(\eta, \eta'; b_1, b'_1) W_{\emptyset; n'_2; \emptyset}^{(h)|0,0}(b'_2) - 2 W_{(n_1, n'_1); \emptyset; \emptyset}^{(h)|0,0}(b_1, b'_1) W_{\emptyset; n'_2; \emptyset}^{(h)|1,1}(\eta, \eta'; b'_2) + \right. \\ &\left. - W_{n'_1; \emptyset; \emptyset}^{(h)|1,1}(\eta, \eta'; b'_1) W_{n_1; n'_2; \emptyset}^{(h)|0,0}(b_1; b'_2) - W_{n'_1; \emptyset; \emptyset}^{(h)|0,0}(b'_1) W_{n_1; n'_2; \emptyset}^{(h)|1,1}(\eta, \eta'; b_1; b'_2) \right], \end{aligned} \quad (3.12)$$

which admits the graphical representation in Fig. 6.




Figure 6: Graphical representation of Eq. (3.12).




Figure 7: Diagram $(h1)$ obtained by plugging (1) in (h) of Fig. 4.

Let us denote by $(h1), (h2), \dots, (h6)$ the graphs obtained from (h) (Fig. 4) by replacing $W_{n_1, n_2; \emptyset}^{(h) | 1, 1}$ with each of the terms in its expansion $(1), \dots, (6)$ (as in Fig. 6). Similarly we denote by $\text{Val}(h1), \dots, \text{Val}(h6)$ their analytical value, as in the notation below Eq. (3.7).

Diagram $(h1)$. Explicitly,

$$\begin{aligned} -\text{Val}(h1) = & \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{2^{-N+1} \mu_{N, \lambda} n_2^2 \mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2 (n_2 - 1)^2} \int db_1 db d\tilde{b} d\eta d\eta' \lambda^2 g_{b_1, b}^A g_{b_1, \tilde{b}}^A c_{b_1}(\eta_1, \eta) g_{\eta_1, \eta'}^{(h, N]} W_{(n_1, n_2 - 1); \emptyset; \emptyset}^{(h) | (1, 1)}(\eta', \eta_2; b; b'), \end{aligned}$$

which can be bounded by taking the L^1 norm of the fermionic propagator and take the L^∞ norm of the two boson propagators:

$$\begin{aligned} \|\text{Val}(h1)\|_1^w \leq & \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{2^{-N+1} |\mu_{N, \lambda}| n_2^2 \mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2 (n_2 - 1)^2} \|c\|_1^w \lambda^2 \|g^A\|_\infty^2 \|g^{(h, N]}\|_1^w \|W_{(n_1, n_2 - 1); \emptyset; \emptyset}^{(h) | 1, 1}\|_1^w \leq \\ & \leq 24 K'^2 \tilde{K} C_0^3 \lambda^2 2^{-2h} N^2 2^{-N} \sum_{n_1, n_2 \geq 1} \frac{1}{n_1^2 n_2^2} \lambda^{\frac{n_1 + n_2 - 2}{2}} \leq C_0^3 \mathfrak{c}_{1,4} \lambda^2 2^{-\theta N} 2^{-2h}, \end{aligned}$$

where $\mathfrak{c}_{1,4} = (4/3)\pi^4 K'^2 \tilde{K}$ and the constants are the same appearing in the previous items.

Diagrams $(h2) - (h6)$. Observe that each of the remaining terms in the r.h.s. of Eq. (3.12) have a dimensional factor 2^{-N} , which graphically corresponds to the leftmost node with two dashed lines and a wiggly line in diagrams $(2) - (6)$ in Fig. 6. In this cases it is enough to note that each of such diagrams $(2) - (6)$ have good bounds for their analytical values:

$$\begin{aligned} \|\text{Val}(2)_{n_1, n_2}\|_1^w \leq & \sum_{\substack{n'_1 \geq 1, n'_2 \geq 2 \\ n'_1 + n'_2 = n_2}} 2^{-N} \mathcal{N}_{n'_1, n'_2} \lambda \|g^A\|_\infty \|W_{(n_1, n'_1); n'_2; \emptyset}^{(h) | 1, 1}\|_1^w \leq \\ & \sum_{\substack{n'_1 \geq 1, n'_2 \geq 2 \\ n'_1 + n'_2 = n_2}} K' N 2^{-N} \lambda \frac{n_2^2}{n_1^2 n_2^2} \lambda^{\frac{n_1 + n'_1 + n'_2 - 3}{2}} \|W_{(n_1, n'_1); n'_2; \emptyset}^{(h) | 1, 1}\|_1^w \leq C_0^4 (4\pi^2/3) K' 2^{-N-2h} N \lambda^{\frac{n_1 + n_2 - 1}{2}}, \end{aligned} \tag{3.13}$$

where we used Eq. (2.24) and the fact that $\sum_{\ell=1}^{n-2} \frac{n^2}{\ell^2 (n-\ell)^2} \leq 4\pi^2/3$. It is easy to check that also the other graphs $(3) - (6)$, in the same way, are bounded as

$$\sup_{j=3}^6 \|\text{Val}(j)_{n_1, n_2}\|_1^w \leq C_0^5 (4\pi^2/3) K' N 2^{-N} \lambda^{\frac{n_1 + n_2 - 1}{2}}. \tag{3.14}$$

Hence when inserted in graph (h) of Fig. 4, they produce a bound:

$$\begin{aligned} \sup_{j=2}^6 \|\text{Val}(hj)\|_1^w &\leq \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{2^{-N} \mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \lambda \|g^A\|_\infty \|g^{(h, N]}\|_\infty \sup_{j=2}^6 \|\text{Val}(j)_{n_1, n_2}\|_1^w \\ &\leq C_0^5 \mathfrak{c}_{1,5} \lambda^{3/2} N 2^{-N} \end{aligned} \quad (3.15)$$

with $\mathfrak{c}_{1,5} = K'^2 K (\pi^2/3)^3$. Collecting the bounds for diagrams (a) to (o) in Fig. 4 one realizes that all the graphs except (d) admit a bound with at a factor $\lambda^{1+\eta}$ with $\eta \geq 1/4$. The extra small factor $\lambda^{1/4}$ can be used to reduce, in each diagram but (d), the overall constant to $\mathfrak{c}_{1,2}/14$ at the price of choosing \mathfrak{c}'_1 depending on $\mathfrak{c}_{1,1}, \mathfrak{c}_{1,3}, \mathfrak{c}_{1,4}$ and such that $\mathfrak{c}'_1 \geq \max\{K', 2^{8/5}\}$ (see diagrams (b) and (m)). Summing all the contributions we obtain that if $\lambda \leq (\mathfrak{c}'_1 C_0^2 R)^{-1}$

$$\|W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}\|_1^w \leq 2\mathfrak{c}_{1,2} C_0^5 \lambda 2^{-\vartheta h}, \quad (3.16)$$

thus proving Eq. (3.2) with $\mathfrak{c}_1 \geq 2\mathfrak{c}_{1,2}$, with $\mathfrak{c}_{1,2}$ appearing in diagram (d).

3.2 The lowest-degree vertex function

Now we focus on the kernel $G^{(1)}\psi^2$. Recalling the definition of the bare vertex kernel c , in Eq. (1.19), the goal is to prove that

$$\|W_{1; \emptyset; \emptyset}^{(h)|1,1} - c\|_1^w \leq C_0^6 \mathfrak{c}_2 \lambda, \quad (3.17)$$

with C_0 as in Proposition 2.1, $\lambda \leq (\mathfrak{c}'_2 C_0^2 R)^{-1}$, and suitable constants $\mathfrak{c}_2, \mathfrak{c}'_2 \geq 1$ independent on C_0, R .

Again, the strategy is to derive a convenient identity for $W_{1; \emptyset; \emptyset}^{(h)|1,1}$. In analogy with Lemma 3.2, by applying $\delta/\delta\psi_{\eta_2}^-$ to Eq. (B.7), one can prove that

$$W_{1; \emptyset; \emptyset}^{(h)|1,1}(\eta_1, \eta_2; b) = \mu_{N,\lambda} c_b(\eta_1, \eta_2) + \mathfrak{D}W_{1; \emptyset; \emptyset}^{(h)|1,1}(\eta_1, \eta_2; b), \quad (3.18)$$

with $\mu_{N,\lambda}$ as below Eq. (3.3) and where $\mathfrak{D}W_{1; \emptyset; \emptyset}^{(h)|1,1}$ is a sum of terms, each interpretable as the analytical value $\text{Val}(\#)(\eta_1, \eta_2; b)$ of a graphical diagram $(\#)$ with the same understanding of Fig. 4 and the notation below it. Also in this case, using Remarks 3.4 and 3.6, one can obtain a bound for $\mathfrak{D}W_{1; \emptyset; \emptyset}^{(h)|1,1}$ by studying a *subclass* of terms contributing to it: by possibly shrinking λ , i.e. gauging the value of \mathfrak{c}'_2 above, it is sufficient to bound the diagrams labeled by $(\#) = (a), \dots, (k)$ appearing in Fig. 8, which, except (b)¹⁰, are obtained after the removal of a dashed tadpole or a dressed fermionic insertion from the general graph in $\mathfrak{D}W_{1; \emptyset; \emptyset}^{(h)|1,1}$.

The estimates follow the same lines as Subsection 3.1 and will not be repeated full detail: we only discuss the main novelties. In order to deduce Eq. (3.17) from the diagrams in Fig. 8, we start by observing the following.

Remark 3.7 (Bubble insertions). Differently from the analysis of the previous section, now some of the graphs in Fig. 8 involve also the ‘‘bubble kernel’’ $W_{(1,1); \emptyset; \emptyset}^{(h)|0,0}$ which, by direct inspection, appears always convoluted with a bosonic propagator, i.e. in the form $g^A * W_{(1,1); \emptyset; \emptyset}^{(h)|0,0}$. This

¹⁰Diagram (b) survives this procedure because, in order to recover the bound in Eq. (3.17), one does not have any extra power of λ to apply Remark 3.4 as it follows from the bound of $\|W_{\emptyset; \emptyset; \emptyset}^{(h)|1,1}\|_1^w$ being of order λ .




Figure 8: The diagram expansion for $\mathfrak{D}W_{1,\emptyset;\emptyset}^{(h)|1,1}(\eta_1, \eta_2; b)$. The symbol $\stackrel{!}{=}$ reminds that the identity holds up to diagrams that can be bounded in terms of the ones listed, i.e., only if one considers all the diagrams before the reduction procedure summarized in Remarks 3.4 and 3.6 by deleting dashed tadpoles and replacing dressed fermionic lines with bare ones.

implies that a bound on $\|g^A * W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w$, as provided in Eq. (2.24), is sufficient¹¹ for a bound of the whole graph: for concreteness, the simplest example is graph (c) which reads

$$\text{Val}(c)(\eta_1, \eta_2; b) = \mu_{N,\lambda} \sum_{n \geq 1} \frac{\lambda}{n^2} \int_{\Lambda_B^2} db' db'' c_{b'}(\eta_1, \eta_2) g_{b', b''}^A W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}(b'', b),$$

so that, using the bounds for the kernels $W_{(n,n');\emptyset;\emptyset}^{(h)|0,0}$ (Eqs. (2.23)-(2.24)) and the fact that according to Eq. (1.27), $\|g^A\|_1^w \leq \frac{32\pi}{\kappa^2}$, we find:

$$\begin{aligned} \|\text{Val}(c)\|_1^w &\leq \|c\|_1^w \lambda \left(\|g^A * W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w + \sum_{n \geq 2} \frac{1}{n^2} \|g^A\|_1^w \|W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w \right) \\ &\leq 12\lambda \left(C_0 + \sum_{n \geq 2} \frac{1}{n^2} \frac{32\pi}{\kappa^2} C_0^3 \lambda^{\frac{n-1}{2}} \right) \leq \mathfrak{c}_{2,1} C_0^3 \lambda, \end{aligned}$$

for some $\mathfrak{c}_{2,1}$ independent of C_0, R .

Let us stress that, in order to close the inductive procedure, in bounding the analytical value of diagram (b) one has to really use the just obtained bound Eq. (3.16) for the kernel $W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}$ instead of using the inductive hypothesis in Eq. (2.22).

¹¹By bounding $g^A * W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}$ as $\|g^A\|_1^w \|W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w$ one obtains, apriori, a logarithmic divergent factor $|h - N|$, as can be checked at first order in λ for the norm of the bubble diagram.

Also, a direct bound on graph (h) shows a logarithmic divergence, in complete analogy with graph (h) of the previous subsection (see Fig. 4). To solve such divergence, one has to extract another wiggly line, by expanding the kernel $W_{(1,n_1);n_2,\emptyset}^{(h)|1,1}$ that appears in graph (h) , following the same steps as in Eq. (3.12) and below.

Finally, in analogy with Remark 3.5, a careful analysis shows that the graphs of type $\sharp \in \{a, e, f, g, h, i, k, \ell\}$ actually admit a bound which is exponentially suppressed in N , namely $\|\text{Val}(\sharp)\|_1^w \leq \mathfrak{c}_{2,1} C_0^6 \lambda N^2 2^{-\theta N}$, for $\lambda \leq (\mathfrak{c}'_{2,1} C_0^2 R)^{-1}$, with some $\mathfrak{c}_{2,2}, \mathfrak{c}'_{2,1} \geq 1$.

Collecting all the bounds one finds that there exists $\mathfrak{c}_{2,3}, \mathfrak{c}'_2 \geq 1$ independent of C_0, R such that for $\lambda \leq (\mathfrak{c}'_2 C_0^2 R)^{-1}$,

$$\|\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq \mathfrak{c}_{2,3} C_0^6 \lambda,$$

so that from Eq. (3.18),

$$\begin{aligned} \|W_{1;\emptyset;\emptyset}^{(h)|1,1} - c\|_1^w &\leq \|\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)|1,1}\|_1^w + (1 - \mu_{N,\lambda}) \|c\|_1^w \\ &\leq \|\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)|1,1}\|_1^w + \lambda 2^{-2N} \|g^A\|_\infty \|c\|_1^w \leq \mathfrak{c}_2 \lambda C_0^6, \end{aligned} \quad (3.19)$$

for some $\mathfrak{c}_2 \geq \mathfrak{c}_{2,3}$.

3.3 The lowest-degree polarization bubble

We now analyze the kernel $W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}$ associated with the monomial $(G^{(1)})^2$, also called the *polarization bubble*. We will show that

$$\|g^A * (W_{(1,1);\emptyset;\emptyset}^{(h)|0,0} - \Pi^{(h,N]})\|_1^w \leq \mathfrak{c}_3 C_0^8 \lambda, \quad (3.20)$$

with $\Pi^{(h,N]}$ the *non-interacting bubble* defined in Eq. (2.16), C_0 as in Proposition 2.1, $\lambda \leq (\mathfrak{c}'_3 C_0^2 R)^{-1}$, and suitable constants $\mathfrak{c}_3, \mathfrak{c}'_3 \geq 1$ independent on C_0, R . In complete analogy with Lemma 3.2, by applying $\delta/\delta G_b^{(1)}$ to Eq. (B.12), for $n = 1$, one finds the following identity:

$$W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}(b, b') = \frac{1}{2} \int_{\Lambda_F^2} d\eta_1 d\eta_2 \mu_{N,\lambda} c_b(\eta_1, \eta_2) \int_{\Lambda_F^2} d\eta'_1 d\eta'_2 g_{\eta'_2, \eta'_1}^{(h,N]} g_{\eta_2, \eta_1}^{(h,N]} W_{1;\emptyset;\emptyset}^{(h)|1,1}(\eta'_1, \eta'_2; b'), \quad (3.21)$$

which can be graphically represented as in Fig. 9.




Figure 9: Expansion of the polarization bubble diagram.

Combining Eqs. (3.18), (3.21) we find that

$$W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}(b, b') = \mu_{N,\lambda} \Pi^{(h,N]}(b, b') + \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}(b, b') \quad (3.22)$$

where

$$\begin{aligned} \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}(b, b') &= \\ \frac{1}{2} \int_{\Lambda_F^2} d\eta_1 d\eta_2 \mu_{N,\lambda} c_b(\eta_1, \eta_2) \int_{\Lambda_F^2} d\eta'_1 d\eta'_2 g_{\eta'_2, \eta'_1}^{(h,N]} g_{\eta_2, \eta_1}^{(h,N]} &\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)|1,1}(\eta'_1, \eta'_2; b'). \end{aligned} \quad (3.23)$$

In this way we have:

$$\|g^A * (W_{(1,1);\emptyset;\emptyset}^{(h)0,0} - \Pi^{(h,N]})\|_1^w \leq |\mu_{N,\lambda} - 1| \|g^A * \Pi^{(h,N]}\|_1^w + \|g^A * \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w. \quad (3.24)$$

Using the finiteness of the non-interacting bubble, see Corollary C.2 in Appendix C, namely $\|g^A * \Pi^{(h,N]}\|_1^w \leq C'_\Pi$ for some constant $C'_\Pi \geq 1$, one finds that

$$\|g^A * (W_{(1,1);\emptyset;\emptyset}^{(h)0,0} - \Pi^{(h,N]})\|_1^w \leq C'_\Pi K' \lambda N 2^{-2N} + \|g^A * \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w, \quad (3.25)$$

where we also used that $|\mu_{N,\lambda} - 1| \leq K' N \lambda 2^{-2N-1}$ (see the analysis of diagram (a) in Subsection 3.1).

Now the idea is to control the quantity $\|g^A * \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w$ by expanding the kernel $\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)1,1}$ in the r.h.s. of Eq. (3.23) as we did in Subsection 3.2. The outcome can be expressed as a sum over graphs, obtained by replacing the blob in the r.h.s. of Fig. 9, by the sum over the diagrams in Fig. 8. For instance, recalling that the graphs of type $\sharp \in \{a, e, f, g, h, i, k, \ell\}$ are bounded as $\|\text{Val}(\sharp)\|_1^w \leq \mathfrak{c}_{2,1} C_0^6 \lambda N^2 2^{-\theta N}$ for $\lambda \leq (\mathfrak{c}'_{2,1} C_0^2 R)^{-1}$, their contribution to $\|\mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w$ can be easily bounded as follows:

$$\begin{aligned} & \frac{1}{L^2} \int_{\Lambda_B} db db' e^{\frac{\kappa}{2} \sqrt{\delta^D(b,b')}} \int_{\Lambda_F^2} d\eta_1 d\eta_2 d\eta'_1 d\eta'_2 \mu_{N,\lambda} |c_b(\eta_1, \eta_2)| \times \\ & \times \sum_{k,j=h+1}^N |g_{\eta'_2, \eta_1}^{(k)}| |g_{\eta_2, \eta'_1}^{(j)}| |\text{Val}(\sharp)(\eta'_1, \eta'_2; b')| \leq \\ & \sum_{h < k \leq j \leq N} \|c\|_1^w \|g^{(k)}\|_\infty \|g^{(j)}\|_1^w \|\text{Val}(\sharp)\|_1^w + \sum_{h < j \leq k \leq N} \|c\|_1^w \|g^{(j)}\|_\infty \|g^{(k)}\|_1^w \|\text{Val}(\sharp)\|_1^w \\ & \leq 2 \sum_{h < k \leq j \leq N} \|g^{(k)}\|_\infty \|g^{(j)}\|_1^w \mathfrak{c}_{2,1} C_0^6 \lambda N^2 2^{-\theta N} \leq \sum_{h < k \leq j \leq N} \mathfrak{c}_{3,1} C_0^6 2^{k-j} \lambda N^2 2^{-\theta N} \leq \\ & \leq 2 \mathfrak{c}_{3,1} C_0^6 \lambda N^3 2^{-\theta N}, \end{aligned} \quad (3.26)$$

for a suitable $\mathfrak{c}_{3,1} \geq 1$. The trick of expanding the product $g^{(h,N]} g^{(h,N]}$ in terms of single-scale contributions, namely $\sum_{h < k \leq N} g^{(k)} g^{(j)}$, and then to take the L^∞ (resp. L^1) norm of the propagator at the lowest (resp. highest) scale will be recurrently used throughout this paragraph.

We are left with bounding the contributions to $\|g^A * \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w$ in the r.h.s. of Eq. (3.25) which derive from diagrams (b), (c), (d), (j) of Fig. 8. The “composite” diagrams, obtained by combining (b), (c), (d), (j) of Fig. 8 with the r.h.s. of Fig. 9, are reported in Fig. 10. Note that differently from the previous section, where in Fig. 8 we could omit (according to Remark 3.4) the diagram (\tilde{c}) , obtained by attaching a dressed fermionic line to the bare vertex¹² in diagram (c), now one has to keep track of such diagram in the bound for $\|g^A * \mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)0,0}\|_1^w$. The reason is that a bound for $\|\text{Val}(\tilde{c})\|_1^w$ does not imply a finite bound on $\|\text{Val}(\tilde{c}')\|_1^w$ of Fig. 10 but one instead has to exploit the topological structure of the resulting composite diagram, as explained shortly.

Diagram (b'). Similarly to Eq. (3.26), we write each of the three propagators $g^{(h,N]}$ as $\sum_{h'=h+1}^N g^{(h')}$, and for any triple $g^{(h_1)}, g^{(h_2)}, g^{(h_3)}$ we take the L^∞ norm of the propagator at

¹²Exactly as (b) is obtained by (a) in Fig. 4.




Figure 10: Graphs with non-trivial bounds (i.e. after the reduction of Remarks 3.4, 3.6) obtained by combining (b), (c), (\tilde{c}), (d), (j) in Fig. 8 with Fig. 10 and contributing to $\mathfrak{D}W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}(b, b')$.

lowest scale, and L^1 norm for the other two. In this way the bound for diagram (b') becomes:

$$\begin{aligned} \|g^A * \text{Val}(b')\|_1^w &\leq \\ &\sum_{h < h_1 \leq h_2, h_3 \leq N} (\|c\|_1^w)^2 \|g^A\|_1^w \|g^{(h_1)}\|_\infty \|g^{(h_2)}\|_1^w \|g^{(h_3)}\|_1^w \|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq \\ &\sum_{h < h_1 \leq h_2, h_3 \leq N} \mathfrak{c}_{3,2} 2^{h_1 - h_2 - h_3} \|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq 4\mathfrak{c}_{3,2} 2^{-h} \|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq 4\mathfrak{c}_{3,2} \mathfrak{c}_1 C_0^5 \lambda 2^{-(1+\vartheta)h}, \end{aligned}$$

with a suitable $\mathfrak{c}_{3,2} \geq 1$ and for $\lambda \leq (\mathfrak{c}_1 C_0^2 R)^{-1}$, having used the improved bound for $W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}$, Eq. (3.2).

Diagram (c') . Explicitly we have that

$$\text{Val}(c')(b_1, b_2) = \sum_{n \geq 1} \frac{\lambda}{n^2} \mu_{N,\lambda}^2 \int_{\Lambda_B} db db' \Pi^{(h,N]}(b_1, b) g_{b,b'}^A W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}(b', b_2),$$

and exploiting the finiteness of the non-interacting bubble, namely $\|g^A * \Pi^{(h,N]}\|_1^w \leq C'_\Pi$ (cf. Corollary C.2), we find that

$$\begin{aligned} \|g^A * \text{Val}(c')\|_1^w &\leq \lambda \|g^A * \Pi^{(h,N]}\|_1^w \left(\|g^A * W_{(1,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w + \sum_{n \geq 2} \frac{1}{n^2} \|g^A\|_1^w \|W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w \right) \\ &\leq \lambda C'_\Pi \left(C_0 + \sum_{n \geq 1} \frac{1}{n^2} \frac{32\pi}{\kappa^2} C_0^3 \lambda^{\frac{n-1}{2}} \right) \leq \lambda \mathfrak{c}_{3,3} C_0^3, \end{aligned}$$

with some constant $\mathfrak{c}_{3,3} \geq 1$.

Diagram (\tilde{c}') . Analogously to diagram (b') , we use the trick of splitting the propagator $g^{(h,N]}$ into its single-scale counterparts:

$$\begin{aligned} \text{Val}(\tilde{c}')(b_1, b_2) &= \sum_{n \geq 1} \sum_{h < h_1, h_2, h_3 \leq N} \mu_{N,\lambda} \int_{\Lambda_F^6} d\eta_1 d\eta_2 d\eta'_1 d\eta'_2 d\eta''_1 d\eta''_2 \int_{\Lambda_B^2} db_3 db_4 \times \\ &\times c_{b_1}(\eta_1, \eta_2) c_{b_3}(\eta''_1, \eta''_2) g_{\eta'_2, \eta_1}^{(h_1)} g_{\eta_2, \eta'_1}^{(h_2)} g_{\eta''_2, \eta'_1}^{(h_3)} g_{b_3, b_4}^A W_{(1,n);\emptyset;\emptyset}^{(h)|1,1}(b_2, b_4). \end{aligned} \quad (3.27)$$

In the r.h.s. of Eq. (3.27) we must distinguish the three possible cases: $h_1 \leq h_2, h_3$ or $h_2 \leq h_1, h_3$ or $h_3 \leq h_1, h_2$. For instance, if $h_1 \leq h_2, h_3$, we can take the L^∞ norm of $g^{(h_1)}$, the L^1 norm of $g^{(h_2)}$ and, similarly to diagram (d) of Subsection 3.1, use the Hölder's inequality for the product $g^A \tilde{g}^{(h_3)}$ with weights $p = \frac{2}{1-\vartheta}$ and $p' = \frac{2}{1+\vartheta}$:

$$\begin{aligned} \|g^A * \text{Val}(\tilde{c}'|_{h_1 \leq h_2, h_3})\|_1^w &\leq \\ \sum_{h < h_1 \leq h_2, h_3 \leq N} \sum_{n \geq 1} \frac{\lambda}{n^2} \|g^A\|_1^w \|g^{(h_1)}\|_\infty \|g^{(h_2)}\|_1^w \|g^A\|_p^w \|\tilde{g}^{(h_3)}\|_{p'} \|W_{(1,n);\emptyset;\emptyset}^{(h)|1,1}\|_1^w &\leq \\ \sum_{h < h_1 \leq h_2, h_3 \leq N} \sum_{n \geq 1} \frac{1}{n^2} \mathfrak{c}_{3,4} C_0^3 \lambda \lambda^{\frac{n-1}{2}} 2^{h_1-h_2-\vartheta h_3} 2^{-h} &\leq \frac{\pi^2 \mathfrak{c}_{3,4}}{3(1-2^{-\vartheta})^2} C_0^3 \lambda 2^{-(1+\vartheta)h}, \end{aligned}$$

where $\tilde{g}_{b,\eta'}^{(h_3)} \equiv \int d\eta d\eta_1 e^{\frac{\kappa}{2} \sqrt{\delta^D(b,\eta,\eta_1)}} |c_b(\eta_1, \eta)| |\tilde{g}_{\eta,\eta'}^{(h_3)}|$ and $\mathfrak{c}_{3,4}$ is a suitable constant greater than 1. The regimes $h_2 \leq h_1, h_3$ and $h_3 \leq h_1, h_2$ can be worked out in the same way, taking the L^∞ norm of the propagator at the lowest scale, and they yield the same bound.

Diagram (d'). Here again we exploit the finiteness of the non-interacting bubble, namely $\|g^A * \Pi^{(h,N]}\|_1^w \leq C'_\Pi$ (cf. Corollary C.2), so that

$$\begin{aligned} \|g^A * \text{Val}(d')\|_1^w &\leq \\ \sum_{n \geq 1} \frac{\lambda}{n^2} \int_{\Lambda_B^3} \frac{db_1 db_2 db_3}{L^2} e^{\frac{\kappa}{2} \sqrt{\delta^D(b_1,b_2)} + \frac{\kappa}{2} \sqrt{\delta^D(b_2,b_3)}} &\left| (g^A * \Pi^{(h,N]})(b_1, b_2) (g^A * W_{n;\emptyset;\emptyset}^{(h)|0,0})(b_2, b_3) \right| \leq \\ \sum_{n \geq 1} \frac{\lambda}{n^2} \|g^A * \Pi^{(h,N]}\|_1^w \|g^A * W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w &\leq C'_\Pi \lambda \|g^A * W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w + \\ + C'_\Pi \lambda \sum_{n \geq 2} \frac{1}{n^2} \|g^A\|_1^w \|W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w &\leq \lambda C'_\Pi C_0 + \sum_{n \geq 2} \frac{1}{n^2} C'_\Pi \|g^A\|_1^w C_0^3 \lambda^{\frac{n+1}{2}} \leq \mathfrak{c}_{3,5} C_0^3 \lambda, \end{aligned}$$

for a suitable $\mathfrak{c}_{3,5} \geq 1$.

Diagram (d''). The bound for this diagram can be deduced from the bound for diagram (b'). Indeed:


$$\text{Val}(d'')(b_1, b_2) = \sum_{n \geq 1} \frac{\lambda}{n^2} \int_{\Lambda_B^2} db_3 db_4 (\text{Val}(b'))(b_1, b_3) g_{b_3, b_4}^A W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}(b_4, b_2)$$

from which:

$$\begin{aligned} \|g^A * \text{Val}(d'')\|_1^w &\leq \sum_{n \geq 1} \frac{\lambda}{n^2} \|g^A * \text{Val}(b')\|_1^w \|g^A * W_{(n,1);\emptyset;\emptyset}^{(h)|0,0}\|_1^w \leq \\ \sum_{n \geq 1} \frac{\lambda}{n^2} 4\mathfrak{c}_{3,2}\mathfrak{c}_1 C_0^5 \lambda 2^{-(1+\vartheta)h} C_0^3 \lambda^{\frac{n-1}{2}} &\leq \lambda \mathfrak{c}_{3,6} C_0^8 2^{-(1+\vartheta)h}, \end{aligned}$$

with $\mathfrak{c}_{3,6} = \frac{2\pi^2}{3} \mathfrak{c}_{3,2} \mathfrak{c}_1$.

Diagram (j'). Similarly to diagram (h) of Fig. 4, the present graph shows a superficial logarithmic divergence, which can be solved by a second extraction of line. Expanding the kernel $W_{\emptyset;n_2;\emptyset}^{(h)|1,1}$ which appears in the diagram, with a formula similar to Eq. (3.12), we find that diagram (j') can be graphically represented as a sum of two graphs (see Fig. below)



having analytical value, respectively:

$$\begin{aligned} \text{Val}(j'.1)(b, b') = & \sum_{n_1 \geq 1, n_2 \geq 2} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \mu_{N, \lambda}^2 2^{-N} \lambda^2 \int_{\Lambda_F^6} d\eta_1 d\eta_2 d\eta'_1 d\eta'_2 d\eta_3 d\eta_4 \int_{\Lambda_B^3} db_1 db_2 db_3 c_b(\eta_1, \eta_2) \times \\ & \times c_{b_1}(\eta'_1, \eta'_2) g_{b_1, b_2}^A g_{b_1, b_3}^A g_{\eta_2, \eta'_1}^{(h, N]} g_{\eta_4, \eta_1}^{(h, N]} g_{\eta'_2, \eta_3}^{(h, N]} W_{n_2-1; \emptyset; \emptyset}^{(h)|1,1}(\eta_3, \eta_4; b_2) W_{(n_1, 1); \emptyset; \emptyset}^{(h)|0,0}(b_3, b'); \end{aligned} \quad (3.28)$$

$$\begin{aligned} \text{Val}(j'.2)(b, b') = & \sum_{n_1, n'_2 \geq 1, n''_2 \geq 2} \frac{\mathcal{N}_{n_1, n'_2 + n''_2} \mathcal{N}_{n'_2, n''_2}}{(n_1 + n'_2 + n''_2)^2} \mu_{N, \lambda} 2^{-2N} \lambda^2 \int_{\Lambda_F^6} d\eta_1 d\eta_2 d\eta'_1 d\eta'_2 d\eta_3 d\eta_4 \times \\ & \times \int_{\Lambda_B^4} db_1 db_2 db_3 db_4 c_b(\eta_1, \eta_2) c_{b_1}(\eta'_1, \eta'_2) g_{b_1, b_2}^A g_{b_1, b_3}^A g_{\eta_2, \eta'_1}^{(h, N]} g_{\eta_4, \eta_1}^{(h, N]} g_{\eta'_2, \eta_3}^{(h, N]} \delta(b_1 - b_4) \times \\ & \times W_{n'_2; n''_2; \emptyset}^{(h)|1,1}(\eta_3, \eta_4; b_2; b_4) W_{(n_1, 1); \emptyset; \emptyset}^{(h)|0,0}(b_3, b'). \end{aligned} \quad (3.29)$$

Both these graphs can be bounded by writing $g_{\eta'_1, \eta'_2}^{(h, N]}$ and $g_{\eta_1, \eta'_2}^{(h, N]}$ as the sum of their single-scale contributions, and taking the L^∞ norm of the propagator at the lowest scale. In diagram $(j'.1)$ one exploits the extraction of the second line to take the L^1 norm of $g_{\eta'_2, \eta_3}^{(h, N]}$, which gives $\mathcal{O}(2^{-h})$ and the L^∞ norm of g_{b_1, b_3}^A , which gives $\mathcal{O}(N)$, so that the finiteness of the bound follows, due to the dimensional factor 2^{-N} in the r.h.s. of Eq. (3.28). In diagram $(j'.2)$ the dashed line, associated with a Dirac delta, forces one to bound in L^∞ norm both $g_{\eta'_2, \eta_3}^{(h, N]}$ and g_{b_1, b_3}^A , so obtaining a factor $\mathcal{O}(N2^N)$, which is however compensated by the dimensional factor $\mathcal{O}(2^{-2N})$ in the r.h.s. of Eq. (3.29). Summarizing,

$$\|g^A * \text{Val}(g')\|_1^w \leq \mathfrak{c}_{3,7} C_0^7 \lambda^2 2^{-N} N^2 2^{-h},$$

with a suitable $\mathfrak{c}_{3,7} \geq 1$. Recalling Eq. (3.25), and collecting the bounds obtained for the various contributions to $\mathfrak{D}W_{(1,1); \emptyset; \emptyset}^{(h)|0,0}$, the desired bound for the kernel $W_{(1,1); \emptyset; \emptyset}^{(h)|0,0}$ follows:

$$\begin{aligned} \|g^A * (W_{(1,1); \emptyset; \emptyset}^{(h)|0,0} - \Pi^{(h, N]})\|_1^w \leq & C'_\Pi K' \lambda N 2^{-2N} + 18 \|g^A\|_1^w \mathfrak{c}_{3,1} C_0^6 \lambda N^3 2^{-\theta N} + 2 \sum_{\sharp \in \{b', c', d', d'', g'\}} \|g^A * \text{Val}(\sharp)\|_1^w \leq \mathfrak{c}_3 C_0^8 \lambda, \end{aligned} \quad (3.30)$$

for a suitable \mathfrak{c}_3 bigger than all the constants $\mathfrak{c}_{3,1}, \mathfrak{c}_{3,2}, \dots$ appeared throughout the paragraph, and for $\lambda \leq \left(\max \{ \mathfrak{c}'_{1,2}, \mathfrak{c}'_1 \} C_0^2 R \right)^{-1}$. The factor 2 multiplying the sum over the graphs in the r.h.s. of Eq. (3.30) takes into account the fact that all the diagrams containing a dashed tadpole have not been considered in Fig. 10, coherently with the discussion in Remark 3.6 of Subsection 3.1.

3.4 The effective quartic interaction

We would like to prove the bound:

$$\|W_{\emptyset;0;\emptyset}^{(h)|2,2} - \mathcal{Q}^{(N)}\|_1^w \leq \mathfrak{c}_4 C_0^8 \lambda^{\frac{5}{4}}, \quad (3.31)$$

with $\mathcal{Q}^{(N)}$ defined in Eq. (2.17), C_0 as in Proposition 2.1, $\lambda \leq (\mathfrak{c}'_4 C_0^2 R)^{-1}$, and suitable constants $\mathfrak{c}_4, \mathfrak{c}'_4 \geq 1$ independent on C_0, R . In analogy with Lemma 3.2, one can prove that

$$W_{\emptyset;0;\emptyset}^{(h)|2,2}(\eta_1, \eta_2, \eta'_1, \eta'_2) = \mu_{N,\lambda} \mathcal{Q}^{(N)}(\eta_1, \eta_2, \eta'_1, \eta'_2) + \mathfrak{D}W_{\emptyset;0;\emptyset}^{(h)|2,2}(\eta_1, \eta_2, \eta'_1, \eta'_2),$$

where $\mathfrak{D}W_{\emptyset;0;\emptyset}^{(h)|2,2}$ can be graphically represented as a sum of graphs, whose analytical value is deduced in the same way as in Subsection 3.1. Such graphs are collected in Fig. 11 where, as before (see Remarks 3.4 and 3.6), we have omitted all those that after the elimination of a “dashed tadpole” or the replacement of a “dressed line” by a “bare line”, reduce to one of the graphs already present.




Figure 11: Graphs contributing to $\mathfrak{D}W_{\emptyset;0;\emptyset}^{(h)|2,2}(\eta_1, \eta_2, \eta'_1, \eta'_2)$. The darker blob in diagram (c) is the difference kernel $\mathfrak{D}W_{1;\emptyset;\emptyset}^{(h)|1,1}$. The exiting half-line attached to the bare vertex is always associated with the coordinate label η_1 , while the labels η'_1, η'_2 can be associated to the entering lines in all the two possible ways.

The estimates follow the same techniques used so far in this section, with no conceptual complication, and thus they will not be repeated in detail. Let us stress that in graphs (c) and

(e) one has to use the improved bounds for $\|W_{1;\emptyset;\emptyset}^{(h)|1,1} - c\|_1^w$ and $\|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\|_1^w$ respectively, namely Eqs. (3.17) and (3.16). Also, in analogy with diagrams (h) of Fig. 4 and Fig. 8, here we have that diagram (k) of Fig. 11 displays a superficial log-divergence, which is readily solved by a second extraction of line, i.e. by the expansion of the kernel $W_{n_1;n_2;\emptyset}^{(h)|2,2}$ (the analogous procedure was explained with all the details in Eq. (3.12) and below). Collecting the bounds from all the diagrams of Fig. 11, one finds:

$$\|\mathfrak{D}W_{\emptyset;\emptyset;\emptyset}^{(h)|2,2}\|_1^w \leq \mathfrak{c}_{4,1}C_0^8\lambda^{\frac{5}{4}} \quad (3.32)$$

for $\lambda \leq (\mathfrak{c}'_4 C_0^2 R)^{-1}$, with suitable constants $\mathfrak{c}_{4,1}, \mathfrak{c}'_4 \geq 1$. The exponent $5/4$ in λ takes into account the lowest power of λ among all the contributions to the l.h.s. of Eq. (3.32), namely diagram (b) with $n = 1$ and diagram (m) with $n = 2$. From Eq. (3.32) we get the desired bound:

$$\begin{aligned} & \|W_{\emptyset;\emptyset;\emptyset}^{(h)|2,2} - \mathcal{Q}^{(N)}\|_1^w \leq \\ & |\mu_{N,\lambda} - 1| \|\mathcal{Q}^{(N)}\|_1^w + \|\mathfrak{D}W_{\emptyset;\emptyset;\emptyset}^{(h)|2,2}\|_1^w \leq \mathfrak{c}_{4,2}\lambda^2 N 2^{-2N} + \mathfrak{c}_{4,1}C_0^8\lambda^{\frac{5}{4}} \leq (\mathfrak{c}_{4,1} + \mathfrak{c}_{4,2})C_0^8\lambda^{\frac{5}{4}} \end{aligned}$$

for $\lambda \leq (\mathfrak{c}'_4 C_0^2 R)^{-1}$, where we also used the fact that $\|\mathcal{Q}^{(N)}\|_1^w \leq \frac{\lambda}{2}(\|c\|_1^w)^2 \|g^A\|_1^w$ is bounded by constant times λ .

3.5 The higher-degree marginal kernels

So far we have been discussing the kernels which were already present in the quartic theory [7, 25, 4]. We are left with analyzing the marginal kernels due to the source fields $\{G^{(n)}, \dot{G}^{(n)}\}_{n \geq 2}$, namely $W_{n;\emptyset;\emptyset}^{(h)|1,1}, W_{\emptyset;n;\emptyset}^{(h)|1,1}$ with $n \geq 2$ and $W_{(n,n');\emptyset;\emptyset}^{(h)|0,0}, W_{\emptyset;(n,n');\emptyset}^{(h)|1,1}, W_{n;n';\emptyset}^{(h)|1,1}$ with $n + n' \geq 3$. Even though the aforementioned fields enter in the initial potential, Eq. (1.44), via irrelevant monomials, they all have scaling dimension 1, so they produce non-irrelevant terms at scales $h < N$. It turns out that the bounds for these extra marginal terms are somehow easier than those discussed so far in this section, the reason being that the fields $G^{(n)}, \dot{G}^{(n)}$ are coupled to kernels $\{v_n\}_{n \geq 2}$ (cf. Eq. (1.29)) which, differently from v_1 , include *at least* one wiggly line. As it already emerged throughout this section, the presence of wiggly lines generally yields dimensional gains, due to the non-locality of the boson propagator.

Since the analysis follows the same lines of the previous subsections, we will discuss with some more detail only the kernel $W_{n;\emptyset;\emptyset}^{(h)|1,1}$ and only give ideas on how to proceed for the other cases.

The higher-degree vertex. We consider the kernel $W_{n;\emptyset;\emptyset}^{(h)|1,1}$, with $n \geq 2$ and we aim to show that

$$\|W_{n;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \leq \mathfrak{c}_5 C_0^5 \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}}, \quad n \geq 2, \quad (3.33)$$

for $\lambda \leq (\mathfrak{c}''_5 C_0 R)^{-1}$, and suitable constants $\mathfrak{c}_5, \mathfrak{c}''_5 \geq 1$ independent on C_0, R . Notice that Eq. (3.33) directly implies the desired bound in Eq. (2.19) by requiring $\lambda \leq (\mathfrak{c}'_5 C_0^{10} R)^{-1}$, with $\mathfrak{c}'_5 = \max\{\mathfrak{c}_5^2, \mathfrak{c}''_5\}$. Let us recall that the choice of losing a fractional power of λ , so to gain in the r.h.s. of Eq. (3.34) a pre-factor which is exactly 1, is only technical and made so that the inductive structure of our proof results more tractable.

Again, the starting point is to derive an expansion for the kernel obtained from Lemma B.2 by applying $\delta^2/\delta\psi_{\eta_2}^- \delta\psi_{\eta_1}^+$ to Eq. (B.12). The output of the expansion is represented in Fig. 12 where, as before (cf. Remarks 3.4 and 3.6), we have omitted all those diagrams that after the elimination of a “dashed tadpole” or the replacement of a “dressed line” by a “bare line”, reduce to one of the graphs already present.

In order to read the value of a diagram, differently from the previous cases referred to Figures 4,8,10 and 11, now the nodes of the graphs in Fig. 12 are interpreted as follows. If the incoming and outgoing solid lines at the node have fermionic labels η_2, η_1 respectively, then:

- a node with two wiggly lines, with labels n and $n-1$, is associated to a factor $2^{-N} \frac{n^2}{(n-1)^2} \mu_{N,\lambda} c_b(\eta_1, \eta_2)$;
- a node with two wiggly lines and a dashed line, having labels n, n_1, n_2 , respectively, and satisfying $n_1 + n_2 = n$, is associated to a factor $2^{-2N} \mathcal{N}_{n_1, n_2} c_b(\eta_1, \eta_2)$ where we recall that $\mathcal{N}_{n_1, n_2} = \frac{(n_1+n_2)^2}{(n_1+n_2-1)n_1 n_2^2} \leq \frac{(n_1+n_2)^2}{n_1^2 n_2^2}$.




Figure 12: Graphs contributing to the kernel $W_{n;\emptyset;\emptyset}^{(h)|1,1}$, with $n \geq 2$. The n label of the leftmost wiggly line in each diagram reminds that the external field is of type $G^{(n)}$. The symbol $\stackrel{!}{=}$ reminds that the identity is up to diagrams which are not drawn, according to the reduction procedure explained below Eq. (3.33).

Diagram (a). In this case we get

$$\begin{aligned} \|\text{Val}(a)\|_1^w &\leq 2^{-N} n^2 (n-1)^{-2} \|c\|_1^w \lambda \|g^A\|_\infty \|g^{(h,N]}\|_1^w \|W_{n-1;\emptyset;\emptyset}^{(h)|1,1}\|_1^w \\ &\leq 24 \lambda K' \tilde{K} N 2^{-N} 2^{-h} C_0^2 \lambda^{\frac{n-2}{2}} \leq \mathfrak{c}_{5,1} C_0^2 \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}}, \end{aligned} \quad (3.34)$$

where $\mathfrak{c}_{5,1} \geq 1$ is a suitable constant depending on \tilde{K}, K' , already appearing along Section 3.1. Note that we neglected the vanishing factor $2^{-\theta N - h}$ as $N \rightarrow \infty$ (see Remark 3.5) Graphs (b) and (d) follow the same type of estimates as (a).

Diagram (c). Here we use the same idea of Eq. (3.26) and lines below by splitting in scales each $g^{(h,N]}$ and taking the L^∞ norm of g^A ,

$$\begin{aligned} \|\text{Val}(c)\|_1^w &\leq \lambda 2^{-N+1} n^2 (n-1)^{-2} \sum_{h \leq i \leq j \leq N} \|g^A\|_\infty \|g^{(i)}\|_\infty \|\tilde{g}^{(j)}\|_1^w \|W_{n-1; \emptyset; \emptyset}^{(h)|2,2}\|_1^w \\ &\leq \mathfrak{c}_{5,2} C_0^2 2^{-N} N^2 \lambda^{\frac{n-2}{2}} \leq \mathfrak{c}_{5,2} C_0^2 \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}}, \end{aligned}$$

for a suitable $\mathfrak{c}_{5,2} \geq 1$.

Diagram (e). The presence of the dashed line forces us to take the L^∞ norm of both $g^{(h,N]}$ and g^A :

$$\begin{aligned} \|\text{Val}(e)\|_1^w &\leq \sum_{\substack{n_1 \geq 1, n_2 \geq 2 \\ n_1 + n_2 = n}} 2^{-2N} \mathcal{N}_{n_1, n_2} \|g^{(h,N)}\|_\infty \lambda \|g^A\|_\infty \|W_{n_1; n_2; \emptyset}^{(h)|1,1}\|_1^w \\ &\leq \sum_{\substack{n_1 \geq 1, n_2 \geq 2 \\ n_1 + n_2 = n}} \frac{n^2}{n_1^2 n_2^2} \mathfrak{c}_{5,3} \lambda 2^{-N} N C_0^3 \lambda^{\frac{n_1+n_2-2}{2}} 2^{-h} \leq \frac{4}{3} \pi^2 \mathfrak{c}_{5,3} C_0^3 \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}} 2^{-h}, \end{aligned} \quad (3.35)$$

for a suitable $\mathfrak{c}_{5,3} \geq 1$, having also used the fact that $\sum_{n_1+n_2=n} \frac{n^2}{n_1^2 n_2^2} \leq \frac{4}{3} \pi^2$.

Remark 3.8. The origin of the quantity $\mathcal{N}_{n_1, n_2} \leq (n_1 + n_2)^2 n_1^{-2} n_2^{-2}$ is in the numerical factors n^3 in the definition of the source terms with $G^{(n)}$ and $\dot{G}^{(n)}$ in Eq. (1.44), as it can be explicitly derived from Lemmas B.1 and B.2. If in place of n^3 we had put n^k , in the r.h.s. of Eq. (3.35) we would have found:

$$\sum_{n_1+n_2=n} \left(\frac{n}{n_1 n_2} \right)^{k-1},$$

which is *bounded* in n for every $k \geq 3$, but for $k = 0, 1, 2$ it is actually *unbounded*. For instance, with the naive choice $k = 0$, the estimate for graph (d) would have been:

$$\|\text{Val}(d)\|_1^w \leq \mathcal{O}(n) C_0^3 N 2^{-N-h} \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}},$$

which has no hope to be consistent with the desired bound in Eq. (2.19). With our choice $k = 3$ instead, we manage to get $\mathcal{O}(1)$ in place of $\mathcal{O}(n)$.

The estimates for diagrams (f)-(o) can be performed similarly and we will not belabor the details. We mention that a direct bound on diagram (o) shows superficial divergences, which can be solved by expanding the kernel $W_{\emptyset; \emptyset; \emptyset}^{(h)|2,2}$ and exploiting the presence of an extra boson propagator (the analogous procedure for diagram (h) of Fig. 4 was explained in Eq. (3.12) and below).

Collecting all the contributions in Fig. 12 one finds the bound in Eq. (3.33), from which the desired bound in Eq. (2.19) follows.

Remaining higher-degree kernels. The analysis of the *higher order bubbles*, i.e. the kernels $W_{(n,n');\emptyset;\emptyset}^{(h)|0,0}$ with $n+n' \geq 3$, is almost identical to that for the higher-degree vertices and one can find that

$$\|W_{(n,n');\emptyset;\emptyset}^{(h)|0,0}\|_1^w \leq \mathfrak{c}_6 C_0^6 \lambda^{\frac{n+n'-2}{2}}, \quad n+n' \geq 3, \quad (3.36)$$

for $\lambda \leq (\mathfrak{c}_6'' C_0 R)^{-1}$ and with a suitable $\mathfrak{c}_6'' \geq 1$. As before, by requiring $\lambda \leq (\mathfrak{c}_6' C_0^{12} R)^{-1}$, with $\mathfrak{c}_6' = \max\{\mathfrak{c}_6'', \mathfrak{c}_6^2\}$, the desired bound in Eq. (2.21) follows.

Concerning instead the kernels $W_{\emptyset;n;\emptyset}^{(h)|1,1}$, $W_{\emptyset;(n,n');\emptyset}^{(h)|0,0}$ and $W_{n';n;\emptyset}^{(h)|0,0}$, their expansion is similar to Eq. (3.12) (depicted in Fig. 6) and is shown in Fig. 13.




Figure 13: Graphical expansions for the kernels $W_{\emptyset;n;\emptyset}^{(h)|0,0}$, $W_{\emptyset;(n,n');\emptyset}^{(h)|1,1}$, $W_{n';n;\emptyset}^{(h)|0,0}$ and $W_{\emptyset;(n',n);\emptyset}^{(h)|0,0}$ respectively.

Using the same tools as in the previous cases, one can check that each diagram in the r.h.s. of each line of Fig. 13 is bounded by $\mathcal{O}(C_0^4 \lambda^{\frac{1}{2}} \lambda^{\frac{n-1}{2}})$, without the need to extract any further boson line, so that:

$$\|W_{\emptyset;n;\emptyset}^{(h)|1,1}\|_1^w \leq \lambda^{\frac{n-1}{2}}, \quad \|W_{\emptyset;(n,n');\emptyset}^{(h)|1,1}\|_1^w \leq \lambda^{\frac{n+n'-2}{2}}, \quad \|W_{n';n;\emptyset}^{(h)|1,1}\|_1^w \leq \lambda^{\frac{n+n'-2}{2}}, \quad (3.37)$$

for any $n \geq 2$ and $\lambda \leq (\mathfrak{c}_7' C_0^8)^{-1}$, with a suitable $\mathfrak{c}_7' \geq 1$.

Summarizing, with Eqs. (3.2), (3.17), (3.20), (3.31), (3.33) and (3.37), we have shown the validity of the bounds in Eqs. (2.18)-(2.21) with $R = \max_{j=1}^4 \mathfrak{c}_j C_0^8$ and for λ such that $\lambda \leq \min\{(C_0 \hat{R}^4)^{-1}, \min_{j=1}^7 (\mathfrak{c}_j' C_0^{12} \hat{R})^{-1}\}$. This establishes the validity of the inductive step introduced at the beginning of Section 3, thus concluding the proof of Theorem 3.1.

4 The complete generating functional

In this section we reintroduce the external sources of the theory, i.e. the background vector field J , the chiral vector B and the Grassmann field φ with the aim of completing the proof of

Theorem 1.4 for the full generating functional:

$$\mathcal{W}^{(u.v.)}(\psi; \varphi; J; B) = -\log \int P^{(0,N]}(\mathcal{D}\zeta) e^{-\mathcal{V}(\psi+\zeta; G(J)) + (\varphi, \zeta + \psi) - (B, O_5(\zeta + \psi))} \quad (4.1)$$

(cf. Eqs. (1.35), (1.21)). We will follow two intermediate steps: we will first reintroduce the fields φ and B by keeping $J = 0$, and in a second moment we will discuss the case $J \neq 0$. The reason for this subdivision is that the multiscale analysis in presence of φ and B , at $J = 0$, requires a procedure which is very close to the one already explained in Sections 2 and 3 for the purely fermionic interaction. In particular, we will use a combination of new *standard dimensional bounds* and *improved bounds*, analogous to Eqs. (2.22)-(2.24) and Eqs. (2.18)-(2.21) which provide in the end the bounds for the new φ, B kernels appearing in Theorem 1.4. On the other hand the reintroduction of the field J and the bounds for the related kernels, appearing in Theorem 1.4, will require a simpler strategy based only on standard estimates with no *improved bounds* needed.

4.1 The chiral and fermion sources

At this stage we add to the fermionic potential $\mathcal{V}(\psi; 0)$, Eq. (1.21), the two source terms:

$$\int_{\Lambda_F} d\eta (\varphi_\eta^+ \psi_\eta^- + \psi_\eta^+ \varphi_\eta^-) + \int_{\Lambda_B} db B_b O_{5;b}(\psi).$$

The scaling dimension of these external fields is defined so that the monomials $\varphi\psi$ and $B\psi^2$ are both marginal. In this way, the scaling dimension of a monomial $(\psi)^q (\varphi)^{\tilde{q}} (B)^p$ is $2 - q - \frac{3}{2}\tilde{q} - p$. In analogy with Section 2, it is convenient to add to the potential some auxiliary source terms to help controlling the relevant and marginal terms. It turns out that it is enough to introduce exactly the same source fields G, \dot{G}, \ddot{G} of Section 2, with potential

$$\mathcal{V}(\psi; G; \dot{G}; \ddot{G}; \varphi; B) := \mathcal{V}(\psi; G; \dot{G}; \ddot{G}) + \int_{\Lambda_F} d\eta (\varphi_\eta^+ \psi_\eta^- + \psi_\eta^+ \varphi_\eta^-) + \int_{\Lambda_B} db B_b O_{5;b}(\psi), \quad (4.2)$$

where $\mathcal{V}(\psi; G; \dot{G}; \ddot{G})$ is the same as Eq. (1.44). In analogy with Section 2, we construct the effective potential at scale h :

$$\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}; \varphi; B) = -\log \int P^{(h,N]}(\mathcal{D}\zeta) e^{-\mathcal{V}(\psi+\zeta; G; \dot{G}; \ddot{G}; \varphi; B)}, \quad (4.3)$$

and we denote by $W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'}(\underline{\eta}, \underline{\eta}'; \tilde{\eta}, \tilde{\eta}'; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}; b')$ the generic kernel of $\mathcal{V}^{(h)}$ associated with the monomial

$$\prod_{k=1}^q \psi_{\eta_k}^+ \prod_{k=1}^{\tilde{q}} \varphi_{\tilde{\eta}_k}^+ \prod_{k=1}^{q'} \psi_{\eta'_k}^- \prod_{k=1}^{\tilde{q}'} \varphi_{\tilde{\eta}'_k}^- \prod_{k=1}^{\dim \underline{n}} G_{b_k}^{(n_k)} \prod_{k=1}^{\dim \dot{\underline{n}}} \dot{G}_{\dot{b}_k}^{(\dot{n}_k)} \prod_{k=1}^{\dim \ddot{\underline{n}}} \ddot{G}_{\ddot{b}_k}^{(\ddot{n}_k)} \prod_{k=1}^p B_{b'_k}, \quad (4.4)$$

where we are keeping the same notation of Eq. (2.9) and lines thereafter. The scaling dimension associated with the kernel $W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{n}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'}$ is

$$D_{sc} := 2 - \frac{1}{2}(q + q') - \frac{3}{2}(\tilde{q} + \tilde{q}') - \dim \underline{n} - \dim \dot{\underline{n}} - 2 \dim \ddot{\underline{n}} - p. \quad (4.5)$$

The new *relevant* and *marginal* (i.e. non-irrelevant) kernels that emerge are those satisfying $D_{sc} \geq 0$, which are graphically represented in Fig. 14.




Figure 14: The graphical representation of the marginal and relevant kernels associated to the fields φ, B , which are respectively represented as thick dashed and wiggly external lines. The thick dashed line is distinguishable from the dashed line associated to $\dot{G}^{(n)}$ thanks to the circled n label attached to the latter.

Now we state the analogue of Theorem 3.1, providing bounds for all the kernels $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}};p}^{(h)|q,q';\tilde{q},\tilde{q}'}$. Such bounds involve either the whole kernels, or their deviation from the value of their lowest-order counterparts. Specifically,

$$\Pi_5^{(h,N]}(b_1, b_2) := \frac{1}{2} \int d\eta_1 d\eta'_1 d\eta_2 d\eta'_2 c_{b_1}(\eta_1, \eta'_1) c_{b_2}^5(\eta_2, \eta'_2) g_{\eta'_1, \eta_2}^{(h,N]} g_{\eta'_2, \eta_1}^{(h,N]} \quad (4.6)$$

is the lowest-order counterpart of the kernel $W_{1;\emptyset;\emptyset;1}^{(h)|0,0;0,0}$, with c^5 the bare chiral vertex¹³ given by Eq. (1.20), which is also the lowest-order contribution to the kernel $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|1,1;0,0}$.

Proposition 4.1. *For every $\vartheta, \theta \in (0, 1)$ fixed, there exist $N'_0 \geq 1$, $R', C'_0 \geq 1$ and $\lambda'_0 > 0$ such that, for any $N \geq N'_0$, $\lambda \leq \lambda'_0$ and $0 \leq h \leq N$, the following is true.*

1. *The non-irrelevant kernels represented in Fig. 14 admit the following bounds:*

$$\|W_{\emptyset;\emptyset;\emptyset;0}^{(h)|0,1;1,0} - \delta\|_1^w \leq R' \lambda 2^{-(1+\vartheta)h}, \quad \|W_{\emptyset;\emptyset;\emptyset;0}^{(h)|1,0;0,1} - \delta\|_1^w \leq R' \lambda 2^{-(1+\vartheta)h}; \quad (4.7)$$

$$\|W_{\emptyset;\emptyset;\emptyset;1}^{(h)|1,1;0,0} - c^5\|_1^w \leq R' \lambda; \quad \|g^A * (W_{1;\emptyset;\emptyset;1}^{(h)|0,0;0,0} - \Pi_5^{(h,N]})\|_1^w \leq R' \lambda; \quad (4.8)$$

$$\|W_{n;\emptyset;\emptyset;1}^{(h)|0,0;0,0}\|_1^w \leq \lambda^{\frac{n-1}{2}} \quad \forall n \geq 2; \quad \|W_{\emptyset;n;\emptyset;1}^{(h)|0,0;0,0}\|_1^w \leq \lambda^{\frac{n-1}{2}} \quad \forall n \geq 2; \quad (4.9)$$

with δ the lattice Dirac delta $\delta(\eta - \eta')$ over Λ_F .

2. *All the irrelevant and non-irrelevant kernels not included in Eqs. (2.22)-(2.24), admit the following bounds:*

$$\|g^A * W_{1;\emptyset;\emptyset;1}^{(h)|0,0;0,0}\|_1^w \leq C'_0, \quad \|W_{\emptyset;\emptyset;\emptyset;2}^{(h)|0,0;0,0}\|_1^w \leq C'_0 N, \quad W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0} = 0 \quad (4.10)$$

and in the remaining cases

$$\|W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}};p}^{(h)|q,q';\tilde{q},\tilde{q}'}\|_1^w \leq C'_0^{1+\dim \underline{n}+\dim \dot{\underline{n}}+\dim \ddot{\underline{n}}+\tilde{q}+\tilde{q}'+p} \lambda^{\tilde{d}} 2^{hD_{sc}} \times \begin{cases} 2^{\theta(h-N)} & \dim \dot{\underline{n}} > 0 \\ 1 & \text{otherwise,} \end{cases} \quad (4.11)$$

with D_{sc} as in Eq. (4.5) and $\tilde{d} := \frac{1}{8} \max\{0, q + q' + \tilde{q} + \tilde{q}' - 2\} + \frac{1}{2}(|\underline{n}| + |\dot{\underline{n}}| - \dim \underline{n} - \dim \dot{\underline{n}}) + \frac{3}{4}|\ddot{\underline{n}}|$.

¹³Recall that by assumption $|Z_N^5| \leq \frac{3}{2}$, cf. Section 1.2.

As a direct consequence one can derive bounds for the kernels of the original generating functional $\mathcal{W}^{(u.v.)}(\psi; \varphi; 0; B)$ since $\mathcal{W}^{(u.v.)}(\psi; \varphi; 0; B) = \mathcal{V}^{(0)}(\psi; 0; 0; 0; \varphi; B)$, see Subsection 5.1.

Proposition 4.1 is the analogue of Theorem 3.1 for the kernels involving also the external fields φ and B , at $J = 0$, and its proof follows the same scheme as in Subsection 2 and Section 3, thus we only provide a sketch of it.

Sketch of the proof. The strategy is made of two steps.

1. First one derives the analogue of Proposition 2.1, namely that there exists $C'_0 \geq 1$ (the same C'_0 as in Proposition 4.1) such that, assuming the bounds in Eqs. (4.7)-(4.9) at scales higher than h , then at scale h the bounds in Eqs. (4.10)-(4.11) hold true, for λ small enough, say $\lambda \leq (C'_0 R')^4$. The arguments used to prove this fact are essentially the same as for Proposition 2.1, which are based on the expansion in Gallavotti-Nicolò trees (see Appendix D) and will not be rediscussed.
2. As a second step, one has to show the validity of the bounds in Eqs. (4.7)-(4.9) for some R' large enough and λ small enough, namely the analogue of Proposition 4.1. This is achieved by induction over the scale h , as explained at the beginning of Section 3. Again, the proof of this fact is ultimately based on the non-locality of the boson propagator, and can be exploiting crucial identities for the non-irrelevant kernels in the l.h.s. of Eqs. (4.7)-(4.9). Such identities are written as expansions in terms of all the kernels of $\mathcal{V}^{(h)}$, for which we shall use:


- the bounds in Eqs. (4.10)-(4.11), for the kernels involving at least a variable φ^\pm or B ;
- the bounds in Eqs. (2.22)-(2.23), with R defined at the beginning of Section 3, for the kernels not involving any φ^\pm or B variables.

Let us begin by discussing the bound for the kernels $\varphi^\pm \psi^\mp$, Eq. (4.7). In analogy with Lemma 3.2, one can obtain the following representation for these kernels by differentiating Eq. (4.3) with respect to ψ^\pm, φ^\mp :

$$W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|0,1;1,0}(\eta'; \tilde{\eta}) = \delta(\eta' - \tilde{\eta}) - \int d\eta'' g_{\tilde{\eta}, \eta''}^{(h,N)} W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|1,1;0,0}(\eta'', \eta'), \quad (4.12)$$

$$W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|1,0;0,1}(\eta; \tilde{\eta}') = \delta(\eta - \tilde{\eta}') - \int d\eta'' g_{\eta'', \tilde{\eta}'}^{(h,N)} W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|1,1;0,0}(\eta, \eta''), \quad (4.13)$$

which are graphically represented as in the figure below.



We therefore get the following estimate in L^1 norm:

$$\|W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|0,1;1,0} - \delta\|_1^w \leq \|g^{(h,N)}\|_1^w \|W_{\emptyset; \emptyset; \emptyset; 0}^{(h)|1,1;0,0}\|_1^w. \quad (4.14)$$

Recall that the kernel $W_{\emptyset;0;0;0}^{(h)|1,1;0,0}$ coincides with $W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}$ of the effective potential Eq. (2.9), which, in force of Theorem 3.1, admits the bound $\|W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}\| \leq R\lambda 2^{-\vartheta h}$ for $\lambda \leq \lambda_0$, where the quantities R and λ_0 are fixed by the analysis of Section 3. Recalling that $\|g^{(h,N]}\|_1^w \leq \tilde{K}2^{-h}$ for a suitable constant $\tilde{K} \geq 1$, from Eq. (4.14) we get

$$\|W_{\emptyset;\emptyset;\emptyset;0}^{(h)|0,1;1,0} - \delta\|_1^w \leq \tilde{K}R\lambda 2^{-(1+\vartheta)h}.$$

The same argument applies to the kernel $\psi^+ \varphi^-$, leading to the same bound.

The marginal kernels with one field B can be treated almost identically to the kernels where B is replaced by $G^{(1)}$, which have been discussed in Section 3. For instance, about the kernel $B\psi^2$, in complete analogy with the kernel $G^{(1)}\psi^2$ discussed in Subsection 3.2, one finds that the difference

$$W_{\emptyset;\emptyset;\emptyset;1}^{(h)|1,1;0,0}(\eta_1, \eta_2; b) - c_b^5(\eta_1, \eta_2)$$

admits an expansion which is graphically reported in Fig. 15, where the graphical rules are analogous to those of Section 3 (see also the notation of Fig. 14 for the new kernels), and the understanding that the node with a thick wiggly line and two solid lines is associated to the bare chiral vertex $c_b^5(\eta_1, \eta_2)$.




Figure 15: graphs contributing to the difference $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|1,1;0,0}(\eta_1, \eta_2; b) - c_b^5(\eta_1, \eta_2)$.

The graphs of Fig. 15 are in one to one correspondence with those in Fig. 8, and can be in fact bounded in the very same way. Each of them is indeed checked to be bounded by $\max\{C_0, C'_0\}^6 \mathfrak{c}_8 \lambda$ for $\lambda \leq (\mathfrak{c}'_8 \max\{C_0, C'_0\}^2 \max\{R', R\})^{-1}$, for suitable constants $\mathfrak{c}_8, \mathfrak{c}'_8 \geq 1$, independent of R' and C'_0 . Such a bound leads to the same qualitative estimate as Eq. (3.17), with $W_{1;\emptyset;\emptyset}^{(h)|1,1}$ and c in the l.h.s. replaced by $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|1,1;0,0}$ and c^5 respectively, and C_0 in the r.h.s. replaced by $\max\{C_0, C'_0\}$.

Similarly, one can study the kernels $G^{(n)}B$. For instance, in the case $n = 1$, following the same steps as in Subsection 3.3, one can expand the difference $W_{1;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(b; b') - \mu_{N,\lambda} \Pi_5^{(h,N]}(b, b')$ ending up with the same graphs as Fig. 10, where now the external wiggly line attached to one of the blobs is thicker, corresponding to the external field B , in place of $G^{(1)}$. Again, the estimates for the graphs are formally identical to those in Subsection 3.3, so they will not be repeated here. One crucial fact is that, as well as the *vector bubble* diagram $\Pi^{(h,N]}$, also its *chiral* counterpart $\Pi_5^{(h,N]}$ is uniformly bounded, in the sense that $\|g^A * \Pi_5^{(h,N)}\|_1^w \leq C'_\Pi$ (see Corollary C.2). \square

Remark 4.2. The property $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0} = 0$, namely the third of Eq. (4.10), is a simple consequence of the *charge conjugation* symmetry (cf. Eq. E.1). Under this transformation, $O_{5;x,\mu,\epsilon}(\psi) \mapsto O_{5;x,\mu,\epsilon}(\mathcal{Q}\psi) = O_{5;x,\mu,-\epsilon}(\psi)$. This immediately implies that $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(x, \mu, \epsilon) = W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(x, \mu, -\epsilon)$. On the other hand, since $O_{5;x,\mu,-\epsilon}(\psi) = -O_{5;x,\mu,\epsilon}(\psi)$, we also have that $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(x, \mu, \epsilon) = -W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(x, \mu, -\epsilon)$, which implies $W_{\emptyset;\emptyset;\emptyset;1}^{(h)|0,0;0,0}(x, \mu, \epsilon) = 0$.

4.2 The vector source

According to Eq. (1.21), in order to restore the original dependence upon J of the generating functional, we must consider as a starting potential the r.h.s. of Eq. (1.21):

$$\mathcal{V}(\psi; G(J)) = \mathcal{V}(\psi; 0) + \sum_{n \geq 1} \sum_{1 \leq p \leq n} 2^{N(2-n-p)} \int_{\Lambda_B^n} d\underline{b} \int_{\Lambda_B^p} d\underline{b}' w_{n,p}(\underline{b}; \underline{b}') O_{\underline{b}}^n(\psi) G_{\underline{b}'}^p(J), \quad (4.15)$$

with the kernels $w_{n,p}$ given by Eqs. (1.22), (1.23). The presence of infinitely many monomials in Eq. (4.15) is again a consequence of *local phase invariance*, Eq. (1.24). For dealing with such a combinatorial complication we adopt the following strategy.

1. All the terms in Eq. (4.15) that are linear in $G(J)$ will be recast in a form which is compatible with the r.h.s. of Eq. (1.44) after the replacement of $G^{(n)}$ by $G(J)$. In this way, for all the non-irrelevant terms produced by the multiscale integration involving only the variables $G^{(n)}$ and ψ , we will rely on the bounds provided by Theorem 3.1.
2. All the monomials in Eq. (4.15) involving at least two variables $G(J)$ will be checked not to produce any relevant or marginal term at lower scales (with a single exception which is however easily manageable), hence we will be able to control them by a standard multiscale analysis based on the tree-expansion techniques presented in Appendix D.

Practically, in the r.h.s. of Eq. (4.15) we isolate the terms that are linear in $G(J)$ and by splitting $w_{n,0}(\underline{b}) = v_n(\underline{b}) + 2^{-2N} \tilde{v}_n(\underline{b})$ (cf. Eq. (1.29)), we rewrite them as

$$\sum_{n \geq 2} n 2^{N(1-n)} \int d\underline{b} w_{n,0}(\underline{b}) G_{b_1} O_{\underline{b}}^n(\psi) \equiv \sum_{n \geq 2} n 2^{N(1-n)} \int d\underline{b} (v_n(\underline{b}) + 2^{-2N} \tilde{v}_n(\underline{b})) G_{b_1} O_{\underline{b}}^n(\psi),$$

so that we can rewrite Eq. (4.15) as

$$\begin{aligned} \mathcal{V}(\psi; G(J)) = & \mathcal{V}(\psi; 0) + \\ & + \sum_{n \geq 1} n 2^{N(1-n)} \int_{\Lambda_B^n} d\underline{b} v_n(\underline{b}) G_{b_1} O_{\underline{b}}^n(\psi) + \\ & + \sum_{n \geq 1} n 2^{-N(1+n)} \int_{\Lambda_B^n} d\underline{b} \tilde{v}_n(\underline{b}) G_{b_1} O_{\underline{b}}^n(\psi) + \\ & + \sum_{n \geq 2} \sum_{2 \leq p \leq n} 2^{N(2-n-p)} \int_{\Lambda_B^n} d\underline{b} \int_{\Lambda_B^p} d\underline{b}' w_{n,p}(\underline{b}; \underline{b}') O_{\underline{b}}^n(\psi) G_{\underline{b}'}^p(J). \end{aligned} \quad (4.16)$$

Now the idea is to treat differently the source field $G(J)$ in the three lines in the last r.h.s. of Eq. (4.16). To this purpose, we introduce three families of auxiliary fields:

$$H \equiv \{H_b^{(n)}\}_{n \geq 1, b \in \Lambda_B}, \dot{H} \equiv \{\dot{H}_b^{(n)}\}_{n \geq 2, b \in \Lambda_B}, \ddot{H} \equiv \{\ddot{H}_b\}_{b \in \Lambda_B},$$

and the generalized potential

$$\begin{aligned} \mathcal{U}(\psi; H; \dot{H}; \ddot{H}; \varphi; B) := & \mathcal{V}(\psi; 0) + \int_{\Lambda_F} d\eta (\varphi_\eta^+ \psi_\eta^- + \psi_\eta^+ \varphi_\eta^-) + \int_{\Lambda_B} db B_b O_{5;b}(\psi) + \\ & + \sum_{n \geq 1} n 2^{N(1-n)} \int_{\Lambda_B^n} d\underline{b} v_n(\underline{b}) H_{b_1}^{(n)} O_{\underline{b}}^n(\psi) + \\ & + \sum_{n \geq 2} n 2^{-(n+1)N} \int_{\Lambda_B^n} d\underline{b} \tilde{v}_n(\underline{b}) \dot{H}_{b_1}^{(n)} O_{\underline{b}}^n(\psi) + \\ & + \sum_{2 \leq p \leq n} 2^{N(2-n-p)} \int_{\Lambda_B^n} d\underline{b} \int_{\Lambda_B^p} d\underline{b}' w_{n,p}(\underline{b}; \underline{b}') O_{\underline{b}}^n(\psi) \ddot{H}_{\underline{b}'}^p. \end{aligned} \quad (4.17)$$

Note that by construction $\mathcal{V}(\psi; G(J); \varphi; B) = \mathcal{U}(\psi; G(J); G(J); G(J); \varphi; B)$ ¹⁴. We then consider, for every $0 \leq h \leq N$,

$$\mathcal{U}^{(h)}(\psi; H; \dot{H}; \ddot{H}; \varphi; B) := -\log \int P^{(h,N]}(\mathcal{D}\zeta) e^{-\mathcal{U}(\psi + \zeta; H; \dot{H}; \ddot{H}; \varphi; B)} \quad (4.18)$$

so that, in particular, the original generating functional $\mathcal{W}^{(u.v.)}(\psi; J; \varphi; B)$ can be recovered via the identification

$$\mathcal{W}^{(u.v.)}(\psi; J; \varphi; B) = \mathcal{U}^{(0)}(\psi; G(J); G(J); G(J); \varphi; B). \quad (4.19)$$

As usual $\mathcal{U}^{(h)}$ can be expressed in an integral form analogous to Eq. (2.9), where now we use the symbol $U_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{p}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'}(\underline{\eta}, \underline{\eta}'; \tilde{\eta}, \tilde{\eta}'; \underline{b}, \dot{\underline{b}}; \ddot{\underline{b}}; \underline{b}')$ to denote the generic kernel of $\mathcal{U}^{(h)}$ associated with the monomial

$$\prod_{k=1}^q \psi_{\eta_k}^+ \prod_{k=1}^{\tilde{q}} \varphi_{\tilde{\eta}_k}^+ \prod_{k=1}^{q'} \psi_{\eta'_k}^- \prod_{k=1}^{\tilde{q}'} \varphi_{\tilde{\eta}'_k}^- \prod_{k=1}^{\dim \underline{n}} H_{b_k}^{(n_k)} \prod_{k=1}^{\dim \dot{\underline{n}}} \dot{H}_{b_k}^{(\dot{n}_k)} \prod_{k=1}^{\ddot{p}} \ddot{H}_{b_k} \prod_{k=1}^p B_{b'_k}. \quad (4.20)$$

The scaling dimension associated with the kernels $U_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{p}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'}$ is defined, as usual, so to fit the dimensional factors in the r.h.s. of Eq. (4.17):

$$D_{sc} \equiv D_{sc} := 2 - \frac{1}{2}(q + q') - \frac{3}{2}(\tilde{q} + \tilde{q}') - \dim \underline{n} - 3 \dim \dot{\underline{n}} - \ddot{p} - p. \quad (4.21)$$

Remark 4.3. As anticipated in Item 1 below Eq. (4.15), the different labeling of the field $G(J)$ in terms of the auxiliary variables H, \dot{H}, \ddot{H} , has the purpose of providing an identification between $H^{(n)}$ and the auxiliary field $G^{(n)}$ introduced in Eq. (4.2), namely

$$\mathcal{U}(\psi; \{n^2 G^{(n)}\}_{n \geq 1}; 0; 0; \varphi; B) = \mathcal{V}(\psi; \{G^{(n)}\}_{n \geq 1}; 0; 0; \varphi; B).$$

As a consequence, the kernels of $\mathcal{U}^{(h)}$ are related to those of $\mathcal{V}^{(h)}$ (cf. Eq. (4.3) and below) via the relation:

$$U_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{p}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'} = \prod_{j=1}^s n_j^{-2} \times W_{\underline{n}; \dot{\underline{n}}; \ddot{\underline{p}}; p}^{(h)|q, q'; \tilde{q}, \tilde{q}'}, \quad \underline{n} \equiv (n_1, \dots, n_s). \quad (4.22)$$

¹⁴ $H = G(J)$ means that $H_b^{(n)}$ is replaced by $G_b(J)$ for every $n \geq 1$, and similarly for \dot{H}, \ddot{H} .

The kernels $W_{\underline{n};\emptyset;0;0}^{(h)|q,q';\tilde{q},\tilde{q}'}$ have already been studied in the previous analysis. In particular for those with $\tilde{q} + \tilde{q}' = 0$, we have the identification $W_{\underline{n};\emptyset;0;0}^{(h)|q,q;0,0} \equiv W_{\underline{n};\emptyset;0}^{(h)|q,q}$ and Theorem 3.1 applies; on the other hand, for the kernels with $\tilde{q} + \tilde{q}' \geq 1$ we can use Proposition 4.1.

We now state the result concerning the kernels not covered by the analysis carried out so far, namely those associated with at least one variable \dot{H} or \ddot{H} .

Proposition 4.4. *For every $\vartheta, \theta \in (0, 1)$ fixed, there exist $C_0'' \geq 1$ and $\lambda_0'' > 0$ such that for $\lambda \leq \lambda_0''$, the kernels $U^{(h)}$ admit the following bounds. $\|U_{\emptyset;\emptyset;2;0}^{(h)|0,0;0,0}\|_1^w \leq \frac{1-2^{\theta-1}}{1-2^{-\theta}} C_0''^3 \lambda^{\frac{1}{2}}$, while for any other kernel with $|\dot{n}| + \ddot{p} \geq 1$,*

$$\|U_{\underline{n};\dot{n};\ddot{p};p}^{(h)|q,q';\tilde{q},\tilde{q}'}\|_1^w \leq C_0''^{1+\dim \underline{n} + \dim \dot{n} + \ddot{p} + p} \lambda^d 2^{hD_{sc}} 2^{\theta(h-N)} \quad (4.23)$$

with D_{sc} as in Eq. (4.21) and $d := \frac{1}{2}(|\underline{n}| - \dim \underline{n}) + \frac{5}{8}|\dot{n}| + \frac{1}{4}\ddot{p} + \frac{1}{8} \max \{0, q + q' + \tilde{q} + \tilde{q}' - 2\}$.

In virtue of Eq. (4.19), Proposition 4.4 directly implies the bounds of the kernels appearing in Theorem 1.4, see Subsection 5.1 for more details.

Proof of Proposition 4.4. The proof goes along the same lines as that of Proposition 2.1. It is convenient to introduce the single-scale contributions to the kernels, $\tilde{U}_{\underline{n};\dot{n};\ddot{p};p}^{(h)|q,q';\tilde{q},\tilde{q}'} := U_{\underline{n};\dot{n};\ddot{p};p}^{(h)|q,q';\tilde{q},\tilde{q}'} - U_{\underline{n};\dot{n};\ddot{p};p}^{(h+1)|q,q';\tilde{q},\tilde{q}'}$, for $0 \leq h \leq N-1$. For these kernels, by using the tools of the tree expansion (see Appendix D), one gets the following bounds:

$$\|\tilde{U}_{\underline{n};\dot{n};\ddot{p};p;\underline{n}}^{(h)|q,q';\tilde{q},\tilde{q}'}\|_1^w \leq (1 - 2^{\theta-1}) C_0''^{1+\dim \underline{n} + \dim \dot{n} + \ddot{p} + p} 2^{hD_{sc}} \lambda^d 2^{\theta(h-N)}, \quad (4.24)$$

for $\lambda \leq \lambda_0''$. The proof of Eq. (4.24) is very similar to that of Eq. (2.26), and thus will be omitted. We just mention, as a main difference from Eq. (2.26), that in the r.h.s. of Eq. (4.24) the “short memory factor” $2^{\theta(h-N)}$ is always present. This is a trace of the fact that the monomials involving \dot{H} and \ddot{H} , in the r.h.s. of Eq. (4.17), are all irrelevant and do not produce, under the multiscale integration, any “running coupling function”, namely a relevant or marginal term involving ψ .

Let us consider a kernel $U^{(h)}$ which is irrelevant, namely has $D_{sc} \leq -1$. Using Eq. (4.24), we find:

$$\begin{aligned} \|U_{\underline{n};\dot{n};\ddot{p};p}^{(h)|q,q';\tilde{q},\tilde{q}'}\|_1^w &\leq \sum_{h'=h}^{N-1} \|\tilde{U}_{\underline{n};\dot{n};\ddot{p};p}^{(h')|q,q';\tilde{q},\tilde{q}'}\|_1^w + \|U_{\underline{n};\dot{n};\ddot{p};p}^{(N)|q,q';\tilde{q},\tilde{q}'}\|_1^w \\ &\leq \sum_{h'=h}^N (1 - 2^{\theta-1}) C_0''^{1+\dim \underline{n} + \dim \dot{n} + \ddot{p} + p} \lambda^d 2^{h'D_{sc}} 2^{\theta(h'-N)} \\ &\leq (1 - 2^{\theta-1}) C_0''^{1+\dim \underline{n} + \dim \dot{n} + \ddot{p} + p} \lambda^d 2^{hD_{sc}} 2^{\theta(h-N)} \sum_{h'=h}^N 2^{(h'-h)(\theta+D_{sc})} \\ &\leq C_0''^{1+\dim \underline{n} + \dim \dot{n} + \ddot{p} + p} \lambda^d 2^{hD_{sc}} 2^{\theta(h-N)}, \end{aligned}$$

where we used that $\|U_{\underline{n};\dot{n};\ddot{p};p}^{(N)|q,q';\tilde{q},\tilde{q}'}\|_1^w$ admits the same bound as the r.h.s. of Eq. (4.23) with $h = N$. Now we must analyze the non-irrelevant terms, namely those with $D_{sc} \geq 0$. Observe that since (by construction) there are no terms with a single \ddot{H} field in the r.h.s. of Eq. (4.17), nor they can be generated by the multiscale integration, the only non-irrelevant kernel with

$|\dot{n}| + \ddot{p} \geq 1$ is the one with $|\dot{n}| = 0, \ddot{p} = 2$. Even though this kernel is marginal, the “short memory factor” $2^{\theta(h-N)}$ in the r.h.s. of Eq. (4.24) is sufficient to provide a finite bound:

$$\|U_{\emptyset; \emptyset; 2; 0}^{(h)|0,0;0,0}\|_1^w \leq \sum_{h'=h}^{N-1} \|\tilde{U}_{\emptyset; \emptyset; 2; 0}^{(h')|0,0;0,0}\|_1^w \leq \sum_{h'=h}^{N-1} (1 - 2^{\theta-1}) C_0''^3 \lambda^{\frac{1}{2}} 2^{\theta(h'-N)} \leq \frac{1 - 2^{\theta-1}}{1 - 2^{-\theta}} C_0''^3 \lambda^{\frac{1}{2}}.$$

□

5 Proof of Theorem 1.4.

In this section we combine the partial results obtained along Sections 2-4 in order to prove the statements of Theorem 1.4. For the sake of clarity we stress the dependence on h_M^* of the generating functional in Eq. (1.35), by writing it as $\mathcal{W}^{(u.v.|h_M^*)}$, and similarly for its kernels appearing in Theorem 1.4. Recall that by Eq. (2.1) the kernels of $\mathcal{W}^{(u.v.|h_M^*)}$ and $\mathcal{W}^{(u.v.|0)}$ are related by a simple rescaling.

5.1 Bounds for the kernels

In order to prove Eqs. (1.37)-(1.39) we follow the structure of the paper by collecting the desired bounds first for $\varphi, J, B = 0$ (i.e. $\tilde{q}, \tilde{q}', p, p' = 0$, cf. Sections 2.2-3) then for $J = 0$ (i.e. $p = 0$, cf. Section 4.1) and finally in the most general case (i.e. $q, q', \tilde{q}, \tilde{q}', p, p' \neq 0$, cf. Section 4.2).

Case $\varphi, J, B = 0$. If also $\psi = 0$, by definition (cf. Eqs. (1.36), (2.9) for $h = 0$ and Eq. (2.13)):

$$\mathcal{F}^{(u.v.|0)} = L^{-2} W_{\emptyset; \emptyset; \emptyset}^{(0)|0,0} = \sum_{h=0}^{N-1} L^{-2} \tilde{W}_{\emptyset; \emptyset; \emptyset}^{(0)|0,0}, \quad (5.1)$$

which, after Eq. (2.26), implies that $|\mathcal{F}^{(u.v.|0)}| \leq \frac{1}{3}(1 - 2^{-\theta})C_0\lambda 2^{2N}$. Rescaling with Eq. (2.1), $\mathcal{F}^{(u.v.|h_M^*)} = 2^{2h_M^*} \mathcal{F}^{(u.v.|0)}$ and $\lambda \mapsto \lambda 2^{-2h_M^*}$, $N \mapsto N - h_M^*$ so that $|\mathcal{F}^{(u.v.|h_M^*)}| \leq C_\star(\lambda 2^{-2h_M^*}) 2^{2N}$, for $C_\star \geq \frac{1}{3}(1 - 2^{-\theta})C_0$, i.e. the first of Eq. (1.38).

If $\psi \neq 0$, combining Eqs. (2.2), (2.5) and Eq. (2.9), and recalling the rescaling in Eq. (2.1), the desired bounds in this case follow from Theorem 3.1 (applied with $h = 0$): we thus find the constraint that $N_\star \geq N_0$, $\lambda_\star \leq \lambda_0$ and $C_\star \geq C_0 \max\{R, \frac{1}{3}(1 + 2^{-\theta})\}$.

Case $J = 0, (\varphi, B) \neq 0$. In this case, since $\mathcal{W}^{(u.v.|0)}(\psi; \varphi; 0; B) = \mathcal{V}^{(0)}(\psi; 0; 0; 0; \varphi; B)$ (cf. Eqs. (4.1), (4.3)), Proposition 4.1 readily implies the claims of Theorem 1.4 concerning the kernels with zero J variables, namely $W_{q, q'; \tilde{q}, \tilde{q}'; 0; p}^{(u.v.|0)}$, with the constraint $N_\star \geq N'_0$, $\lambda_\star \leq \lambda'_0$ and $C_\star \geq C'_0$. For $W_{q, q'; \tilde{q}, \tilde{q}'; 0; p}^{(u.v.|h_M^*)}$ one simply uses the rescaling in Eq. (2.1).

Case $J \neq 0$. In this case the kernels $U_{n; \dot{n}; \ddot{p}; p'}^{(0)|q, q'; \tilde{q}, \tilde{q}'}$ of $\mathcal{U}^{(0)}$ (cf. Eq. (4.18)) collect all the information about $W_{q, q'; \tilde{q}, \tilde{q}'; p; p'}^{(u.v.|0)}$, since $\mathcal{W}^{(u.v.|0)}(\psi; J; \varphi; B) = \mathcal{U}^{(0)}(\psi; G(J); G(J); G(J); \varphi; B)$ (compare Eqs. (4.1), (4.18) and below). We discuss first the strategy in two cases of importance.

The tadpole $G(J)$. We have that

$$\begin{aligned} W_{0, 0; 0, 0; 1; 0}^{(u.v.|0)}(b) &= \sum_{n \geq 1} U_{n; \emptyset; \emptyset}^{(0)|0,0;0,0}(b) + \sum_{\dot{n} \geq 2} U_{\emptyset; \dot{n}; \emptyset}^{(0)|0,0;0,0}(b) \\ &= \sum_{n \geq 1} \frac{1}{n^2} W_{n; \emptyset; \emptyset}^{(0)|0,0}(b) + \sum_{\dot{n} \geq 2} U_{\emptyset; \dot{n}; \emptyset}^{(0)|0,0;0,0}(b). \end{aligned} \quad (5.2)$$

Note that the above kernels are actually constant w.r.t. $b \in \Lambda_B$ due to the *charge conjugation* and the *axes flip* symmetries (cf. Lemma E.1) and the translation invariance of the theory. Using the bounds for the kernels $W_{n;\emptyset;\emptyset}^{(0)|0,0,0}$ from Theorem 3.1, and the bounds for $U_{\emptyset;\dot{n};0;0}^{(0)|0,0,0,0}$ from Proposition 4.4, we find:

$$\begin{aligned} \|W_{0,0;0,0;1;0}^{(u.v.|0)}\|_1^w &\leq \sum_{n \geq 1} \frac{1}{n^2} \|W_{n;\emptyset;\emptyset}^{(0)|0,0,0}\|_1^w + \sum_{\dot{n} \geq 2} \|U_{\emptyset;\dot{n};0;0}^{(0)|0,0,0,0}\|_1^w \\ &\leq \sum_{n \geq 1} \frac{1}{n^2} C_0 2^N \lambda^{\frac{n-1}{2}} + \sum_{\dot{n} \geq 2} C_0''^2 \lambda^{\frac{5}{8}\dot{n}} 2^{-\theta N} \leq 2^N \left(\frac{\pi^2}{6} C_0^2 + \frac{\sqrt{2}}{\sqrt{2}-1} C_0''^2 \right), \end{aligned}$$

for $\lambda \leq \min\{\lambda_0, \lambda_0'', \frac{1}{2}\}$; by rescaling $\|W_{0,0;0,0;1;0}^{(u.v.|h_M^*)}\|_1^w = 2^{h_M^*} \|W_{0,0;0,0;1;0}^{(u.v.|0)}\|_1^w$ one obtains the third of Eq. (1.38).

The polarization bubble $G(J)G(J)$. We have that

$$\begin{aligned} W_{0,0;0,0;2;0}^{(u.v.|0)}(b, b') &= W_{(1,1);\emptyset;\emptyset}^{(0)|0,0}(b, b') + \sum_{\substack{n_1, n_2 \geq 1 \\ n_1 + n_2 \geq 3}} \frac{1}{n_1^2 n_2^2} W_{(n_1, n_2);\emptyset;\emptyset}^{(0)|0,0}(b, b') \\ &\quad + U_{\emptyset;\emptyset;2;0}^{(0)|0,0,0,0}(b, b') + \sum_{\substack{n \geq 1, \dot{n} \geq 2}} U_{n;\dot{n};0;0}^{(0)|0,0,0,0}(b, b'), \end{aligned} \quad (5.3)$$

which, again in force of Theorem 3.1 and Proposition 4.4, implies for $\lambda \leq \min\{\lambda_0, \lambda_0'', \frac{1}{2}\}$:

$$\begin{aligned} \|g^A * W_{0,0;0,0;2;0}^{(u.v.|0)}\|_1^w &\leq \\ C_0^3 + \sum_{\substack{n_1, n_2 \geq 1 \\ n_1 + n_2 \geq 3}} \frac{\lambda^{\frac{n_1+n_2-2}{2}}}{n_1^2 n_2^2} + C_0''^3 \lambda^{\frac{1}{2}} \sum_{n \geq 1, \dot{n} \geq 2} C_0''^3 \lambda^{\frac{1}{2}(n-1) + \frac{5}{8}\dot{n}} &\leq C_0^3 + C_0''^3 + \frac{\sqrt{2}}{\sqrt{2}-1} (1 + C_0''^3). \end{aligned} \quad (5.4)$$

By the rescaling, $g_{\mu,\nu}^A(x-y) \mapsto g_{\mu,\nu}^A(2^{h_M^*}(x-y))$ and $W_{0,0;0,0;2;0}^{(u.v.|0)} \mapsto W_{0,0;0,0;2;0}^{(u.v.|h_M^*)}$, with

$$W_{0,0;0,0;2;0}^{(u.v.|h_M^*)}((x, \mu, \epsilon), (y, \nu, \epsilon')) = 2^{2h_M^*} W_{0,0;0,0;2;0}^{(u.v.|0)}((2^{h_M^*}x, \mu, \epsilon), (2^{h_M^*}y, \nu, \epsilon')),$$

so that $\|g^A * W_{0,0;0,0;2;0}^{(u.v.|h_M^*)}\|_1^w$ is bounded by the r.h.s. of Eq. (5.4) times an extra $2^{-2h_M^*}$, proving the first bound in Eq. (1.39) for C_* greater than the r.h.s. of Eq. (5.4).

The desired bound for the kernel $G(J)B$, i.e. the second of Eq. (1.39), follows analogously.

Remaining kernels. Similar (though more cumbersome) steps yield finite bounds for all the remaining kernels of arbitrary order in the fields $G(J)$. As above one has to express the kernels of $W^{(u.v.|0)}$ in terms of those of $\mathcal{U}^{(0)}$, via the relation Eq. (4.19), and then exploit Theorem 3.1 and Propositions 4.1, 4.4 for bounding the various terms that appear. All in all one finds the validity of the whole collection of bounds in Item 1 of Theorem 1.4, for N_* large enough and suitable constants C_* large enough and λ_* small enough.

5.2 Correlation functions

Here we discuss the Item 2 of Theorem 1.4, concerning the two-point function $\hat{S}^{(u.v.)}$, the current-current function $\hat{\Sigma}_{\mu,\nu}^{(u.v.)}$ and its chiral counterpart. We keep using the same conventions introduced at the beginning of Section 5.

The two-point Schwinger function. By construction we have that $\hat{S}_{s,s'}^{(u.v.|0)}(k) = \int_{\Lambda} dx e^{ik \cdot x} W_{0,0;1,1;0;0}^{(u.v.|0)}((x, s), (0, s'))$. In analogy with Eqs. (4.12), (4.13), it is straightforward to check that the kernel $W_{0,0;1,1;0;0}^{(u.v.|0)}$ admits the expansion:

$$W_{0,0;1,1;0;0}^{(u.v.|0)}((x, s), (0, s')) = g_{s,s'}^{(0,N]}(x) - \sum_{s_1, s_2 \in \{1, 2\}} \int_{\Lambda} dy dz g_{s,s_1}^{(0,N]}(x - y) W_{1,1;0,0;0;0}^{(u.v.|0)}((y, s_1), (z, s_2)) g_{s_2,s'}^{(0,N]}(z). \quad (5.5)$$

It follows that

$$\hat{S}_{s,s'}^{(u.v.|0)}(k) = \hat{g}_{s,s'}^{(0,N]}(k) - \sum_{s_1, s_2 \in \{1, 2\}} \hat{g}_{s,s_1}^{(0,N]}(k) \hat{g}_{s_2,s'}^{(0,N]}(k) \int_{\Lambda} dy e^{ik \cdot y} W_{1,1;0,0;0;0}^{(u.v.|0)}((y, s_1), (0, s_2)).$$

Note that since $1 - \chi \equiv 0$ on $[0, \frac{1}{2}]$, the bound in Eq. (2.3) implies:

$$|\hat{g}_{s,s'}^{(0,N]}(k)| = (1 - \chi(|k|)) |(-i\cancel{s}(k) + m_N + M_N(k))_{s,s'}^{-1}| \leq \sqrt{2}\pi \frac{1 - \chi(|k|)}{\sqrt{|k|^2 + m_N^2}} \leq \frac{4\sqrt{2}\pi}{|k| + 1}.$$

Using also the bound for the kernel $W_{1,1;0,0;0;0}^{(u.v.|0)} \equiv W_{\emptyset;\emptyset;\emptyset}^{(0)|1,1}$ from Theorem 3.1, we find:

$$|\hat{S}_{s,s'}^{(u.v.|0)}(k) - \hat{g}_{s,s'}^{(0,N]}(k)| \leq \sum_{s_1, s_2 \in \{1, 2\}} \frac{32\pi^2}{1 + |k|^2} \|W_{1,1;0,0;0;0}^{(u.v.|0)}\|_1^w \leq \frac{32\pi^2 R \lambda}{1 + |k|^2},$$

which, after the rescaling in Eq. (2.1), yields the desired bound in Eq. (1.40) for $\hat{S}_{s,s'}^{(u.v.|h_M^*)}(k)$, if $C_{\star} \geq 32\pi^2 R$.

The non-interacting current functions. For the sake of concreteness we discuss only the *vector current function*, namely we prove the first bound of Eq. (1.41) and Eq. (1.42) since the second of Eq. (1.41) and Eq. (1.43), concerning the *chiral current function*, can be proved with the same strategy.

Proof of Eq. (1.41). From the definition $\hat{\Sigma}_{\mu,\nu}^{(u.v.|0)}(p) := \int_{\Lambda} dx e^{-ip \cdot x} \frac{\delta^2 \mathcal{W}^{(u.v.|0)}}{\delta J_{\mu,x} \delta J_{\nu,0}}(0)$, and the structure of $\mathcal{W}^{(u.v.|0)}$ (cf. Eq. (1.36)), it is straightforward to check that

$$\hat{\Sigma}_{\mu,\nu}^{(u.v.|0)}(p) = -2\lambda \left(2^{-N} \delta_{\mu,\nu} W_{0,0;0,0;1;0}^{(u.v.|0)} + \sum_{\epsilon, \epsilon'=\pm} \epsilon \epsilon' \int_{\Lambda} dx e^{-ip \cdot x} W_{0,0;0,0;2;0}^{(u.v.|0)}((x, \mu, \epsilon), (0, \nu, \epsilon')) \right), \quad (5.6)$$

where recall that the kernel $W_{0,0;0,0;1;0}^{(u.v.|0)}$ is actually a constant (see Eq. (5.2) and below). Now we explicitly characterize the dominant term in λ of the r.h.s. of Eq. (5.6). About the first term in the r.h.s., we start from the decomposition in Eq. (5.2) for the kernel $W_{0,0;0,0;1;0}^{(u.v.|0)}$ and we apply Lemma B.2 to further expand the kernel $W_{1;\emptyset;\emptyset}^{(0)|0,0}$:

$$W_{1;\emptyset;\emptyset}^{(0)|0,0}(b) = \mu_{N,\lambda} 2^N \mathcal{T}^{(0,N]} + \mu_{N,\lambda} \int d\eta_1 d\eta_2 c_b(\eta_1, \eta_2) \int d\eta d\eta' g_{\eta,\eta_1}^{(h,N]} g_{\eta_2,\eta'}^{(h,N]} W_{\emptyset;\emptyset;\emptyset}^{(0)|1,1}(\eta', \eta), \quad (5.7)$$

where

$$\mathcal{T}^{(h,N]} := 2^{-N} \mathcal{E}_{(h,N]}^T(O_b) \equiv -2^{-N} \int_{\Lambda_F^2} d\eta_1 d\eta_2 c_b(\eta_1, \eta_2) g_{\eta_2,\eta_1}^{(h,N]}. \quad (5.8)$$

Plugging the decomposition in Eq. (5.7) into Eq. (5.2), we find

$$\begin{aligned} & |2^{-N}(W_{0,0;0,0;1;0}^{(u.v.|0)} - \mathcal{T}^{(0,N]})| \leq \\ & (1 - \mu_{N,\lambda})2^{-N}\|c\|_1^w\|g^{(0,N]}\|_\infty + 2^{-N}\|c\|_1^w\|g^{(0,N]}\|_1^w\|g^{(0,N]}\|_\infty\|W_{\emptyset;\emptyset;\emptyset}^{(0)|1,1}\|_1^w + \\ & \sum_{n \geq 2} \frac{1}{n^2} \|W_{n;\emptyset;\emptyset}^{(0)|0,0}\|_1^w + \sum_{n \geq 2} \|U_{\emptyset;n;0;0}^{(0)|0,0;0,0}\|_1^w \leq C_1 \lambda^{\frac{1}{2}}, \end{aligned} \quad (5.9)$$

for a suitable constant $C_1 \geq 1$, where in the last step we used Theorem 3.1 for the bounds of $W_{\emptyset;\emptyset;\emptyset}^{(0)|1,1}$ and $W_{n;\emptyset;\emptyset}^{(0)|0,0}$, and Proposition 4.4 for the bound of $U_{\emptyset;n;0;0}^{(0)|0,0;0,0}$.

About the second summand in the r.h.s. of Eq. (5.6), by Eq. (5.3) we have that

$$\begin{aligned} & \|g^A * (W_{0,0;0,0;2;0}^{(u.v.|0)} - \Pi^{(0,N]})\|_1^w \leq \|g^A * (W_{(1,1);\emptyset;\emptyset}^{(0)|0,0} - \Pi^{(0,N]})\|_1^w + \\ & + \|g^A\|_1^w \left(\sum_{\substack{n_1, n_2 \geq 1 \\ n_1 + n_2 \geq 3}} \frac{1}{n_1^2 n_2^2} \|W_{(n_1, n_2);\emptyset;\emptyset}^{(0)|0,0}\|_1^w + \|U_{\emptyset;\emptyset;2;0}^{(0)|0,0;0,0}\|_1^w + \sum_{n \geq 1, n \geq 2} \|U_{n;n;0;0}^{(0)|0,0;0,0}\|_1^w \right) \leq C_2 \lambda^{\frac{1}{2}}, \end{aligned}$$

for a suitable constant $C_2 \geq 1$. In momentum space this bound reads:

$$\left| \sum_{\epsilon=\pm} \epsilon \int_{\Lambda} dx e^{-ip \cdot x} W_{0,0;0,0;2;0}^{(u.v.|0)}((x, \mu, \epsilon), (0, \nu, \epsilon')) - \epsilon' \hat{\Pi}_{\mu,\nu}^{(0,N]}(p) \right| \leq (|p|^2 + 4) C_2 \lambda^{\frac{1}{2}}, \quad (5.10)$$

where we used the fact that $|\sigma(p)|^2 + M^2 \leq |p|^2 + 4$ (recall that we are assuming $h_M^* = 0$) and we have introduced:

$$\begin{aligned} \hat{\Pi}_{\mu,\nu}^{(h,N]}(p) &:= \frac{1}{2} \sum_{\epsilon, \epsilon'=\pm} \epsilon \epsilon' \int_{\Lambda} \frac{dxdy}{L^2} e^{-ip \cdot (x-y)} \Pi^{(h,N]}((x, \mu, \epsilon), (y, \nu, \epsilon')) \\ &\equiv \sum_{\epsilon=\pm} \epsilon \epsilon' \int_{\Lambda} \frac{dxdy}{L^2} e^{-ip \cdot (x-y)} \Pi^{(h,N]}((x, \mu, \epsilon), (y, \nu, \epsilon')), \end{aligned} \quad (5.11)$$

with $\Pi^{(h,N]}$ defined in Eq. (2.16), and the last equality holds in force of the *charge conjugation* symmetry in Eq. (E.1). By simply reverting the rescaling in Eq. (2.1), Eqs. (5.7), (5.10) become:

$$|2^{-N}(W_{0,0;0,0;1;0}^{(u.v.|h_M^*)} - \mathcal{T}^{(h_M^*,N]})| \leq C_1 (2^{-2h_M^*} \lambda)^{\frac{1}{2}}, \quad (5.12)$$

$$\left| \sum_{\epsilon=\pm} \epsilon \int_{\Lambda} dx e^{-ip \cdot x} W_{0,0;0,0;2;0}^{(u.v.|h_M^*)}((x, \mu, \epsilon), (0, \nu, \epsilon')) - \epsilon' \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p) \right| \leq 5C_2 (2^{-2h_M^*} \lambda)^{\frac{1}{2}}, \quad (5.13)$$

for every $p \in \Lambda^*$ such that $|p| \leq 2^{h_M^*}$. Plugging these bounds back into Eq. (5.6), and letting:

$$\hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.|h_M^*)}(p, M, m_N, L) := -2 \left(\delta_{\mu,\nu} \mathcal{T}^{(h_M^*,N]}(m_N, L) + 2 \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, m_N, L) \right), \quad (5.14)$$

we find:

$$\begin{aligned} & \left| \hat{\Sigma}_{\mu,\nu}^{(u.v.|h_M^*)}(p) - \lambda \hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.|h_M^*)}(p, M, m_N, L) \right| \leq 2\lambda 2^{-N} \delta_{\mu,\nu} \left| W_{0,0;0,0;1;0}^{(u.v.|h_M^*)} - \mathcal{T}^{(h_M^*,N]} \right| + \\ & + 2\lambda \left| \sum_{\epsilon, \epsilon'=\pm} \epsilon \epsilon' \int_{\Lambda} dx e^{-ip \cdot x} W_{0,0;0,0;2;0}^{(u.v.|h_M^*)}((x, \mu, \epsilon), (0, \nu, \epsilon')) - 2 \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p) \right| \\ & \leq 2(C_1 + 5C_2) 2^{-h_M^*} \lambda^{\frac{3}{2}}, \end{aligned} \quad (5.15)$$

thus proving the first of Eq. (1.41), for $C_* \geq 2(C_1 + 5C_2)$.

For the incoming analysis it is also useful to introduce $\hat{\mathfrak{B}}_{\mu,\nu}(p, M, m_N, L) := \lim_{\lambda \rightarrow 0^+} \frac{1}{\lambda} \hat{\Sigma}_{\mu,\nu}(p)$, which, in force of Eq. (1.25), satisfies the following Ward Identity:

$$\sum_{\mu=0,1} \sigma_\mu(p) \hat{\mathfrak{B}}_{\mu,\nu}(p, M, m_N, L) = 0, \quad \forall p \in \frac{2\pi}{L} \mathbb{Z}^2 \cap (-\frac{\pi}{a}, \frac{\pi}{a}]. \quad (5.16)$$

It is straightforward to check that

$$\hat{\mathfrak{B}}_{\mu,\nu}(p, M, m_N, L) = -2 \left(\delta_{\mu,\nu} \mathcal{T}^{(\leq N)}(m_N, L) + 2 \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L) \right), \quad (5.17)$$

where $\mathcal{T}^{(\leq N)}$ and $\hat{\Pi}_{\mu,\nu}^{(\leq N)}$ are as in the r.h.s. of Eqs. (5.8) and (5.11) respectively, with the fermionic propagator $g^{(h,N)}$ in their definition replaced by $g^{(\leq N)}$. In order to proceed we make use of the following proposition for the characterization of $\hat{\mathfrak{B}}_{\mu,\nu}$ and $\hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.)}$.

Proposition 5.1. *For every M, m_N, L as in Theorem 1.4, with $m_N \neq 0$, the functions $\Lambda^* \ni p \mapsto \hat{\Pi}_{\mu,\nu}^{(h_M^*, N]}(p, m_N, L)$ and $\Lambda^* \ni p \mapsto \hat{\Pi}_{\mu,\nu}^{(h_M^*, N]}(p, m_N, L)$ admit a continuous extension over $(-\frac{\pi}{a}, \frac{\pi}{a}]^2$, which, with some abuse of notation, we will denote by the same symbol; their $L \rightarrow \infty$ limit exists and is reached uniformly w.r.t. $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$. Besides, for $p \neq 0$,*

$$\begin{aligned} \lim_{M \rightarrow 0^+} \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \hat{\Pi}_{\mu,\nu}^{(h_M^*, N]}(p, m_N, L) &= \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L) =: \hat{\Pi}_{\mu,\nu}(p), \\ \lim_{M \rightarrow 0^+} \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \mathcal{T}^{(h_M^*, N]}(m_N, L) &= \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \mathcal{T}^{(\leq N)}(m_N, L) =: \mathcal{T}, \end{aligned} \quad (5.18)$$

where

$$\hat{\Pi}_{\mu,\nu}(p) = \frac{1}{8\pi|p|^2} (2p_\mu p_\nu - |p|^2 \delta_{\mu,\nu}) + \tilde{R}_{\mu,\nu}(ap), \quad (5.19)$$

for a suitable function $\tilde{R}_{\mu,\nu}$ on $(-\pi, \pi]^2$, such that $|\tilde{R}_{\mu,\nu}(q) - \tilde{R}_{\mu,\nu}(0)| \leq C|q|^{\frac{1}{2}}$, for some constant $C > 0$ independent of N .

The proof is postponed to Appendix C.2. Let us see how Proposition 5.1 implies Eq. (1.42) (the derivation of Eq. (1.43) can be done analogously). First note that after taking the limit $L \rightarrow \infty$ at both sides of Eq. (5.16), in force of the uniform convergence established by Proposition 5.1, the equality above extends to every $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$. Taking also the limit $m_N \rightarrow 0$ and then $M \rightarrow 0^+$, we obtain:

$$\sum_{\mu=0,1} \frac{i}{a} (e^{-iap_\mu} - 1) \left(\frac{1}{4\pi|p|^2} (2p_\mu p_\nu - |p|^2 \delta_{\mu,\nu}) + 2\tilde{R}_{\mu,\nu}(ap) + \mathcal{T}\delta_{\mu,\nu} \right) = 0,$$

for every $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2 \setminus \{0\}$. Using the continuity in 0 of $\tilde{R}_{\mu,\nu}$, we find that

$$\sum_{\mu=0,1} p_\mu \left(\frac{1}{4\pi|p|^2} (2p_\mu p_\nu - |p|^2 \delta_{\mu,\nu}) + 2\tilde{R}_{\mu,\nu}(0) + \mathcal{T}\delta_{\mu,\nu} \right) + \mathcal{O}(a^{\frac{1}{2}}|p|^{\frac{3}{2}}) = 0,$$

and for arbitrariness of p , we must have that $2\tilde{R}_{\mu,\nu}(0) + \mathcal{T}\delta_{\mu,\nu} = -\frac{1}{4\pi} \delta_{\mu,\nu}$. Hence, going back to $\hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.|h_M^*)}$, we find that

$$\begin{aligned} \lim_{M \rightarrow 0^+} \lim_{m_N \rightarrow 0} \lim_{L \rightarrow \infty} \hat{\mathfrak{B}}_{\mu,\nu}^{(u.v.|h_M^*)}(p, M, m_N, L) &= -2 \left(\frac{1}{4\pi|p|^2} (2p_\mu p_\nu - |p|^2 \delta_{\mu,\nu}) + 2\tilde{R}_{\mu,\nu}(ap) + \mathcal{T}\delta_{\mu,\nu} \right) \\ &= \frac{1}{\pi} \left(\delta_{\mu,\nu} - \frac{p_\mu p_\nu}{|p|^2} \right) - 4 \left(\tilde{R}_{\mu,\nu}(ap) - \tilde{R}_{\mu,\nu}(0) \right), \end{aligned}$$

for every $p \neq 0$, which proves Eq. (1.42) with $R_{\mu,\nu}(p) = -4(\tilde{R}_{\mu,\nu}(ap) - \tilde{R}_{\mu,\nu}(0))$.

A Consequences of the local phase invariance

In this section we discuss two main consequences of the local phase invariance of the theory as in Eq. (1.24), namely the ξ -independence (Lemma 1.3) and the Ward identities (Eqs. (1.25) and (1.26)).

Proof of Lemma 1.3. Letting

$$Z_\xi(\varphi; J; B) := e^{-W_\xi(\varphi; J; B)} \equiv \frac{\int \mathbb{P}_\xi(\mathcal{D}A) \int \mathcal{D}\psi e^{-I(A+J,\psi) - (\varphi,\psi) - i(B,J^5(\psi))}}{\int \mathcal{D}\psi e^{-I(0,\psi)}}, \quad (\text{A.1})$$

we are going to show that $Z_\xi(0; J; B)$ is constant w.r.t. $\xi \in [0, 1]$. Observe that $Z_\xi(\varphi; J; B)$, as a function of ξ , is continuous in $[0, 1]$ and continuously differentiable in $(0, 1)$, with

$$\partial_\xi Z_\xi(\varphi; J; B) = \frac{1}{2} \sum_{\mu, \nu \in \{0, 1\}} \int_{\Lambda^2} dx dy \partial_\xi g_{\nu, \mu}^{A, \xi}(y - x) \frac{\delta^2 Z_\xi}{\delta J_{\mu, x} \delta J_{\nu, y}}(\varphi; J; B), \quad (\text{A.2})$$

as a standard formula for derivatives of Gaussian measures, where $\frac{\delta}{\delta J_{\mu, x}} \equiv 2^{2N} \frac{\partial}{\partial J_{\mu, x}}$. Hence it suffices to show that $\partial_\xi Z_\xi(0; J; B) = 0$ for every $\xi \in (0, 1)$. Eq. (A.2) can be rewritten in Fourier space:

$$\partial_\xi Z_\xi(\varphi; J; B) = \frac{1}{2} \sum_{\mu, \nu} \int_{\Lambda^*} dk \partial_\xi \left(\frac{1 - \xi}{\xi |\sigma(k)|^2 + M^2} \right) \frac{\sigma_\nu(k) \sigma_\mu^*(k)}{|\sigma(k)|^2 + M^2} \hat{K}_{\mu, \nu}^\xi(k, -k)_{\varphi; J; B}, \quad (\text{A.3})$$

where $\hat{K}_{\mu, \nu}^\xi(k, p)_{\varphi; J; B} := \int_{\Lambda^2} dx dy e^{ik \cdot x + ip \cdot y} \frac{\delta^2 Z_\xi}{\delta J_{\mu, x} \delta J_{\nu, y}}(\varphi; J; B)$ and $\sigma_\mu(k)$ defined after Eq. (1.8). Now one has the following equation:

$$\sum_{\mu=0,1} \sigma_\mu^*(k) \hat{K}_{\mu, \nu}^\xi(k, p)_{0; J; B} = 0, \quad \forall k, p \in \Lambda^*, \quad (\text{A.4})$$

which readily implies the vanishing of the r.h.s. of Eq. (A.3) at $\varphi = 0$, which in turn implies the independence of $Z_\xi(0; J; B)$ upon $\xi \in [0, 1]$. In order to prove Eq. (A.4), we must exploit the local phase invariance of the theory, namely

$$Z_\xi(e^{i\mathbf{e}\alpha} \varphi; J + \nabla \alpha; B) = Z_\xi(\varphi; J; B), \quad \forall \alpha \in \mathbb{R}^\Lambda \quad (\text{A.5})$$

with the understanding that $(e^{i\mathbf{e}\alpha} \varphi)_{x,s}^\pm \equiv e^{\pm i\mathbf{e}\alpha(x)} \varphi_{x,s}^\pm$, $(A + \nabla \alpha)_{\mu,x} \equiv A_{\mu,x} + \nabla_\mu \alpha(x)$, and where $\nabla_\mu \alpha(x) \equiv a^{-1}(\alpha(x + a\hat{e}_\mu) - \alpha(x))$. By differentiating both sides of Eq. (A.5) w.r.t. $\alpha(x)$ and setting $\alpha = 0$, we find:

$$\begin{aligned} a^{-1} \sum_{\mu=0,1} \left(\frac{\delta Z_\xi}{\delta J_{\mu, x - a\hat{e}_\mu}}(\varphi; J; B) - \frac{\delta Z_\xi}{\delta J_{\mu, x}}(\varphi; J; B) \right) = \\ -i\mathbf{e} \sum_{s=1,2} \left(\varphi_{x,s}^+ \frac{\delta Z_\xi}{\delta \varphi_{x,s}^+}(\varphi; J; B) - \varphi_{x,s}^- \frac{\delta Z_\xi}{\delta \varphi_{x,s}^-}(\varphi; J; B) \right). \end{aligned} \quad (\text{A.6})$$

If we further differentiate both sides of Eq. (A.6) w.r.t. $J_{\nu,y}$ and we set $\varphi = 0$, we get:

$$a^{-1} \sum_{\mu=0,1} \left(\frac{\delta^2 Z_\xi}{\delta J_{\mu, x - a\hat{e}_\mu} \delta J_{\nu, y}}(0; J; B) - \frac{\delta^2 Z_\xi}{\delta J_{\mu, x} \delta J_{\nu, y}}(0; J; B) \right) = 0. \quad (\text{A.7})$$

By taking the Fourier transform, we finally recover Eq. (A.4):

$$0 = a^{-1} \sum_{\mu=0,1} \int_{\Lambda^2} dx dy e^{i(k \cdot x + p \cdot y)} \left(\frac{\delta^2 Z_\xi}{\delta J_{\mu,x-a\hat{e}_\mu} \delta J_{\nu,y}}(0; J; B) - \frac{\delta^2 Z_\xi}{\delta J_{\mu,x} \delta J_{\nu,y}}(0; J; B) \right) = \\ \sum_{\mu=0,1} \frac{e^{iak_\mu} - 1}{a} \int_{\Lambda^2} dx dy e^{i(k \cdot x + p \cdot y)} \frac{\delta^2 Z_\xi}{\delta J_{\mu,x} \delta J_{\nu,y}}(0; J; B) = i \sum_{\mu=0,1} \sigma_\mu^*(k) \hat{K}_{\mu,\nu}^\xi(k, p)_{0;J;B}.$$

□

We now turn our focus to the generating functional, defined formally as $\mathcal{W}_\xi(\varphi; J; B) = -\log Z_\xi(\varphi; J; B)$. We are going to assume that \mathcal{W}_ξ is a well defined object, analytic w.r.t. J and B close enough to zero: note that this is apriori true only for $|\epsilon|$ very small depending on the cut-offs of the model, namely N, L, m_N ; however, one of the nontrivial outcomes of our analysis is the well posedness of \mathcal{W}_1 for $|\epsilon| \ll M$, uniformly in N, L, m_N .

Proof of the Ward Identities: Eqs. (1.25), (1.26). After taking the logarithm at both sides of Eq. (A.5), it follows that $\mathcal{W}_\xi(e^{i\epsilon\alpha}\varphi; J + \nabla\alpha; B) = \mathcal{W}_\xi(\varphi; J; B) \forall \alpha \in \mathbb{R}^\Lambda$. This implies the analogue of Eq. (A.6) with Z_ξ replaced by \mathcal{W}_ξ , which, after differentiation of both sides w.r.t. $J_{\nu,0}$, reads:

$$a^{-1} \sum_{\mu=0,1} \left(\frac{\delta}{\delta J_{\mu,x-a\hat{e}_\mu}} - \frac{\delta}{\delta J_{\mu,x}} \right) \left(\frac{\delta \mathcal{W}_\xi(\varphi; J; B)}{\delta J_{\nu,0}} \right) = \\ -i\epsilon \sum_{s=1,2} \left(\varphi_{x,s}^+ \frac{\delta}{\delta \varphi_{x,s}^+} - \varphi_{x,s}^- \frac{\delta}{\delta \varphi_{x,s}^-} \right) \left(\frac{\delta \mathcal{W}_\xi(\varphi; J; B)}{\delta J_{\nu,0}} \right). \quad (\text{A.8})$$

As a first step, we set φ, J, B to zero in both sides of Eq. (A.8); then if we also take the Fourier transform, we get:

$$a^{-1} \sum_{\mu=0,1} \int_{\Lambda} dx e^{-ip \cdot x} \left(\frac{\delta^2 \mathcal{W}_\xi}{\delta J_{\mu,x-a\hat{e}_\mu} \delta J_{\nu,0}}(0; 0; 0) - \frac{\delta^2 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta J_{\nu,0}}(0; 0; 0) \right) = 0$$

which readily implies the first of Eq. (1.25), namely $\sum_{\mu=0,1} \sigma_\mu(p) \hat{\Sigma}_{\mu,\nu}^\xi(p) = 0$ for every $p \in \Lambda^*$.

The second of Eq. (1.25), concerning $\hat{\Sigma}_{\nu,\mu,\nu}^\xi$, follows in full analogy. Let us now consider the analogue of Eq. (A.8) with the derivative w.r.t. $J_{\nu,0}$ replaced by the second derivative w.r.t. $\varphi_{0,s}'$ and $\varphi_{y,s''}^+$:

$$a^{-1} \sum_{\mu=0,1} \left(\frac{\delta^3 \mathcal{W}_\xi}{\delta J_{\mu,x-a\hat{e}_\mu} \delta \varphi_{0,s'}^- \delta \varphi_{y,s''}^+}(0; 0; 0) - \frac{\delta^3 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta \varphi_{0,s'}^- \delta \varphi_{y,s''}^+}(0; 0; 0) \right) = \\ i\epsilon \sum_{s=1,2} \left(\delta(y-x) \delta_{s,s''} \frac{\delta^2 \mathcal{W}_\xi}{\delta \varphi_{x,s}^+ \delta \varphi_{0,s'}^-}(0; 0; 0) - \delta(x) \delta_{s,s'} \frac{\delta^2 \mathcal{W}_\xi}{\delta \varphi_{y,s''}^+ \delta \varphi_{x,s}^-}(0; 0; 0) \right).$$

Then, taking the Fourier transform,

$$\sum_{\mu=0,1} \frac{e^{-iap_\mu} - 1}{a} \int_{\Lambda^2} dx dy e^{i(-p \cdot x + k \cdot y)} \frac{\delta^3 \mathcal{W}_\xi}{\delta J_{\mu,x} \delta \varphi_{0,s'}^- \delta \varphi_{y,s''}^+}(0; 0; 0) = \\ i\epsilon \left(\int_{\Lambda} dx e^{i(-p+k) \cdot x} \frac{\delta^2 \mathcal{W}_\xi}{\delta \varphi_{x,s''}^+ \delta \varphi_{0,s'}^-}(0; 0; 0) - \int_{\Lambda} dy e^{ik \cdot y} \frac{\delta^2 \mathcal{W}_\xi}{\delta \varphi_{y,s''}^+ \delta \varphi_{0,s'}^-}(0; 0; 0) \right),$$

which is exactly Eq. (1.26), namely

$$\sum_{\mu=0,1} \sigma_\mu(p) \hat{\Gamma}_\mu^\xi(p; k) = -\mathbf{e}(\hat{S}^\xi(k-p) - \hat{S}^\xi(k)), \quad \forall p, k \in \Lambda^*.$$

□

B Identities for the effective potential

Here we give the proof of the identities among the kernels of the effective potential, which have been used in Section 3. Since, as long as N, L are finite, $\mathcal{V}^{(h)}$ depends only upon field sources with degree n less or equal than $n_* \equiv 2(2^N L)^2 + 1$, throughout the section, functional derivatives and kernels related to $G^{(n)}, \dot{G}^{(n)}, \ddot{G}^{(n)}$ will be understood as zero for $n > n_*$. For completeness, recall that

$$\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G}) = -\log \mathcal{E}\left(e^{-\mathcal{V}(\psi + \cdot; G; \dot{G}; \ddot{G})}\right),$$

where in this section $\mathcal{E}(F) \equiv \mathcal{E}_{(h,N]}(F) := \int P^{(h,N]}(\mathcal{D}\zeta) F(\zeta)$ for any Grassmann polynomial $F(\zeta)$, while \mathcal{V} is as in Eq. (1.44).

The incoming analysis is based on the following two fundamental identities.

1. *Grassmann integration by parts.* Given any Grassmann polynomial $F(\zeta)$,

$$\int P^{(h,N]}(\mathcal{D}\zeta) \zeta_\eta^- F(\zeta) = \int_{\Lambda_F} d\eta' g_{\eta, \eta'}^{(h,N]} \int P^{(h,N]}(\mathcal{D}\zeta) \frac{\delta F}{\delta \zeta_{\eta'}^+}(\zeta) \quad (\text{B.1})$$

$$\int P^{(h,N]}(\mathcal{D}\zeta) \zeta_\eta^+ F(\zeta) = - \int_{\Lambda_F} d\eta' g_{\eta', \eta}^{(h,N]} \int P^{(h,N]}(\mathcal{D}\zeta) \frac{\delta F}{\delta \zeta_{\eta'}^-}(\zeta). \quad (\text{B.2})$$

2. *Tree property.* Let $I := \{1, \dots, n\}$, $I' := I \setminus \{1\}$, $g_T^A := \prod_{(i,j) \in T} \lambda g_{b_i, b_j}^A$ and recall that $\mathcal{T}(I)$ denotes set of all spanning trees on I (cf. below Eq. (1.23)). If $n = 1$ by definition $\mathcal{T}(I) = \emptyset$, otherwise:

$$\begin{aligned} \sum_{T \in \mathcal{T}(I)} g_T^A &= \\ &= \sum_{k \in I'} g_{b_1, b_k}^A \sum_{T \in \mathcal{T}(I')} g_T^A + \mathbb{1}_{(n \geq 3)} \sum_{\substack{X_1 \cup X_2 = I' \\ X_1 \ni 2}} \sum_{k \in X_1} g_{b_1, b_k}^A \sum_{T_1 \in \mathcal{T}(X_1)} g_{T_1}^A \sum_{T_2 \in \mathcal{T}(X_2 \cup \{1\})} g_{T_2}^A. \end{aligned} \quad (\text{B.3})$$

While the *Grassmann integration by parts* is a standard fact (see e.g. [25, Eq. (A4)]), the *tree property* is not completely straightforward, and since it is a crucial building block for the incoming analysis of this section, we are going to present its proof.

Proof of the tree property. There is a one-to-one correspondence between $\mathcal{T}(I)$ and

$$\mathcal{C}(I) := \left\{ (X_1, T_1, T_2, k) : X_1 \subseteq I', 2 \in X_1, T_1 \in \mathcal{T}(X_1), T_2 \in \mathcal{T}(I' \setminus X_1), k \in X_1 \right\}.$$

Indeed, we can define a map $\mathcal{T}(I) \mapsto \mathcal{C}(I)$ obtained by setting, for any $T \in \mathcal{T}(I)$,

- $X_1 \subseteq I'$ is the maximal connected set such that $2 \in X_1$.
- T_1 is the restriction of T to X_1 , namely $T_1 = \{(i, j) \in T : i, j \in X_1\}$.

- $k \in X_1$ is the unique vertex such that $(1, k) \in T$.
- Letting $X_2 := I' \setminus X_1$, T_2 is the restriction of T to $X_2 \cup \{1\}$.

See Fig. 16 for a graphical representation. On the other hand, we can consider the map $\mathcal{C}(I) \mapsto \mathcal{T}(I)$ defined as $T := T_1 \dot{\cup} T_2 \dot{\cup} (1, k)$. It is then straightforward to check that the maps $\mathcal{T}(I) \mapsto \mathcal{C}(I)$ and $\mathcal{C}(I) \mapsto \mathcal{T}(I)$ are one the inverse of the other.




Figure 16: Graphical representation of the correspondence $\mathcal{T}(I) \leftrightarrow \mathcal{C}(I)$. In the example (a), the vertex 1 is a leaf of T , $X_1 = I'$, $X_2 = \emptyset$, $T_1 = \{(2, 3), (3, 6), (6, 5), (6, 7), (6, 4), (4, 8)\}$ and $k = 3$. In the example (b), 1 is not a leaf, $X_1 = \{2, 4, 5\}$, $X_2 = \{3, 6, 7, 8\}$, $T_1 = \{(4, 2), (4, 5)\}$, $T_2 = \{(1, 3), (3, 6), (3, 7), (1, 8)\}$ and $k = 4$.

The bijection between $\mathcal{C}(I)$ and $\mathcal{T}(I)$ allows us to write:

$$\sum_{T \in \mathcal{T}(I)} g_T^A = \sum_{(X_1, T_1, T_2, k) \in \mathcal{C}(I)} g_{T_{(X_1, T_1, T_2, k)}}^A, \quad (\text{B.4})$$

where $T_{(X_1, T_1, T_2, k)}$ is the tree in $\mathcal{T}(I)$ uniquely identified by $(X_1, T_1, T_2, k) \in \mathcal{C}(I)$. Now it is convenient to distinguish whether $X_2 \equiv I' \setminus X_1$ is empty or not (i.e. the vertex 1 is a leaf or not, see Fig. 16). In the first case we simply have: $g_{T_{(X_1, T_1, T_2, k)}}^A = g_{(1, k)}^A g_{T_1}^A$, while, if $X_2 \neq \emptyset$, $g_{T_{(X_1, T_1, T_2, k)}}^A = g_{(1, k)}^A g_{T_1}^A g_{T_2}^A$. Therefore Eq. (B.4) becomes:

$$\sum_{T \in \mathcal{T}(I)} g_T^A = \sum_{\substack{(X_1, T_1, T_2, k) \in \mathcal{C}(I) \\ X_1 = I'}} g_{(1, k)}^A g_{T_1}^A + \sum_{\substack{(X_1, T_1, T_2, k) \in \mathcal{C}(I) \\ X_1 \subset I'}} g_{(1, k)}^A g_{T_1}^A g_{T_2}^A. \quad (\text{B.5})$$

Noting that the second sum in the r.h.s. of Eq. (B.5) is nonempty only if $n = 3$, the desired *tree property*, Eq. (B.3), follows. \square

B.1 Preliminary identities

We begin with two intermediate lemmas. The first one, Lemma B.1, characterizes the derivatives of $\mathcal{V}^{(h)}$ w.r.t. ψ and $G^{(1)}$. The next one, Lemma B.2, characterizes the derivatives of $\mathcal{V}^{(h)}$ w.r.t. $G^{(n)}$ and $\dot{G}^{(n)}$, with $n \geq 2$. We will use the notation $\mathcal{V}^{(h)}(\psi; \mathbf{0}) := \mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G})|_{G, \dot{G}, \ddot{G} \equiv 0}$ or $\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})$ when $G^{(n)}, \dot{G}^{(n)}, \ddot{G}^{(n)} \equiv 0$ for $n \geq 2$, etc.

Lemma B.1. *The following identities hold for the effective potential $\mathcal{V}^{(h)}$.*

$$\begin{aligned} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta_1}^+}(\psi; \mathbf{0}) &= e^{\mathcal{V}^{(h)}(\psi; \mathbf{0})} \int db_1 d\eta c_{b_1}(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \times \\ &\times \left\{ 2^N (\mu_{N, \lambda} - 1) - \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int db \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n)}} + \right. \\ &+ \left. \sum_{n_1 \geq 1, n_2 \geq 2} 2^{-N} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \int db \lambda g_{b_1, b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} - \sum_{n \geq 2} 2^{-N} \frac{\delta}{\delta \ddot{G}_{b_1}^{(n)}} \right\} e^{-\mathcal{V}^{(h)}} \Big|_{(\psi; \mathbf{0})}, \end{aligned} \quad (\text{B.6})$$

where $\mathcal{N}_{n_1, n_2} = \frac{(n_1 + n_2)^2}{(n_1 + n_2 - 1)n_1 n_2^2} \leq \frac{(n_1 + n_2)^2}{n_1^2 n_2^2}$. Moreover:

$$\begin{aligned} \frac{\delta^2 \mathcal{V}^{(h)}}{\delta G_b^{(1)} \delta \psi_{\eta_1}^+}(\psi; \mathbf{0}) &= \\ &\frac{\delta}{\delta G_b^{(1)}} \left\{ e^{\mathcal{V}^{(h)}} \int db_1 d\eta c_{b_1}(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \times \right. \\ &\times \left[2^N (\mu_{N, \lambda} - 1) - \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int db \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n)}} + \right. \\ &+ \left. \sum_{n_1 \geq 1, n_2 \geq 2} 2^{-N} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \int db \lambda g_{b_1, b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} - \sum_{n \geq 2} 2^{-N} \frac{\delta}{\delta \ddot{G}_{b_1}^{(n)}} \right] \left. \right\} e^{-\mathcal{V}^{(h)}} \Big|_{(\psi; \mathbf{0})} + \\ &+ \mu_{N, \lambda} \int d\eta c_b(\eta_1, \eta) \left(\psi_{\eta}^- - \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^+}(\psi; \mathbf{0}) \right). \end{aligned} \quad (\text{B.7})$$

Proof. We start from the identity:

$$\frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta_1}^+}(\psi; G; \dot{G}; \ddot{G}) = e^{\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G})} \mathcal{E} \left(\frac{\delta \mathcal{V}}{\delta \psi_{\eta_1}^+}(\psi + \cdot; G; \dot{G}; \ddot{G}) e^{-\mathcal{V}(\psi + \cdot; G; \dot{G}; \ddot{G})} \right)$$

and we set to zero all the fields except ψ and $G^{(1)}$. From Eq. (1.44) we find:

$$\begin{aligned} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta_1}^+}(\psi; G^{(1)}; \mathbf{0}) &= \\ &e^{\mathcal{V}^{(h)}} \left\{ \left[2^N (\mu_{N, \lambda} - 1) \int db_1 \mathcal{E} \left(\frac{\delta O_{b_1}}{\delta \psi_{\eta_1}^+}(\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right) + \right. \right. \\ &\sum_{n \geq 2} n 2^{N(2-n)} \int d\underline{b} v_n(\underline{b}) \mathcal{E} \left(\frac{\delta O_{b_1}}{\delta \psi_{\eta_1}^+}(\psi + \cdot) O_{\underline{b}'}(\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right) + \\ &\left. \left. \sum_{n \geq 2} n 2^{-nN} \int d\underline{b} \tilde{v}_n(\underline{b}) \mathcal{E} \left(\frac{\delta O_{b_1}}{\delta \psi_{\eta_1}^+}(\psi + \cdot) O_{\underline{b}'}(\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right) \right] \right\} + \\ &e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \int db G_b^{(1)} v_1(b) \mathcal{E} \left(\frac{\delta O_b}{\delta \psi_{\eta_1}^+}(\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right), \end{aligned} \quad (\text{B.8})$$

where we have used the decomposition $w_{n,0} = v_n + 2^{-2N}\tilde{v}_n$ according to Eq. (1.29). Noting that $\frac{\delta O_b}{\delta \psi_{\eta_1}^+}(\psi + \zeta) = \int d\eta c_b(\eta_1, \eta)(\psi_\eta^- + \zeta_\eta^-)$ and using Eq. (B.1), we find that the first line in the r.h.s. of Eq. (B.8) becomes

$$2^N (\mu_{N,\lambda} - 1) \int db \int d\eta c_b(\eta_1, \eta) \left(\psi_\eta^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})}.$$

Let us proceed with the second line in the r.h.s. of Eq. (B.8). Recalling that $v_n(\underline{b}) = \sum_{T \in \mathcal{T}(I)} \frac{\mu_{N,\lambda}^n}{n!} g_T^A$ (cf. Eq. (1.29)) and using Eq. (B.3) we find that

$$\begin{aligned} v_n(\underline{b}) &= \mu_{N,\lambda} \mathbf{1}_{(n=1)} + \mathbf{1}_{(n \geq 2)} \sum_{k \in I} \frac{\lambda \mu_{N,\lambda}}{n} g_{b_1, b_k}^A v_{n-1}(\underline{b}') + \\ &+ \mathbf{1}_{(n \geq 3)} \sum_{\substack{X_1 \cup X_2 = I' \\ 2 \in X_1}} \binom{n}{|X_1|}^{-1} \sum_{k \in X_1} \lambda g_{b_1, b_k}^A v_{|X_1|}(b_{X_1}) v_{|X_2|+1}(b_{X_2 \cup \{1\}}), \end{aligned} \quad (\text{B.9})$$

where $\underline{b}_{X_1} = \{b_i : i \in X_1\}$ and recall that v_n is symmetric under permutations of its variables. Thus we can rewrite the second line in the r.h.s. of Eq. (B.8) as

$$\begin{aligned} &\sum_{n \geq 2} \int d\underline{b} e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mathcal{E} \left(\frac{\delta O_{b_1}}{\delta \psi_{\eta_1}^+}(\psi + \cdot) \left\{ \mu_{N,\lambda}(n-1) \lambda g_{b_1, b_2}^A \left(2^{N(2-n)} v_{n-1}(\underline{b}') O_{\underline{b}'}(\psi + \cdot) \right) \right. \right. \\ &+ 2^{-N} n \mathbf{1}_{(n \geq 3)} \sum_{\substack{X_1 \cup X_2 = I' \\ X_1 \ni 2}} \binom{n}{|X_1|}^{-1} |X_1| \lambda g_{b_1, b_2}^A 2^{N(1-|X_1|)} v_{|X_1|}(b_{X_1}) O_{b_{X_1}}(\psi + \cdot) \times \\ &\left. \left. \times 2^{N(1-|X_2|)} v_{|X_2|+1}(b_{X_2 \cup \{1\}}) O_{b_{X_2}}(\psi + \cdot) \right\} e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right), \end{aligned} \quad (\text{B.10})$$

where we used the symmetry of v_n and $O_{b_{X_1}} := \prod_{i \in X_1} O_{b_i}$ etc.. Eq. (B.10) can be further rewritten by identifying the terms in parentheses as derivatives of \mathcal{V} w.r.t. G and \dot{G} (cf. Eq. (1.44)) as

$$\begin{aligned} &\int db_1 db e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mathcal{E} \left(\frac{\delta O_{b_1}}{\delta \psi_{\eta_1}^+}(\psi + \cdot) \left[- \sum_{n \geq 2} \frac{\mu_{N,\lambda}}{(n-1)^2} \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n-1)}} + \right. \right. \\ &+ \sum_{n \geq 3} 2^{-N} \sum_{\substack{n_1 \geq 1, n_2 \geq 2 \\ n_1 + n_2 = n}} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n_1)}} \frac{\delta}{\delta \dot{G}_{b_1}^{(n_2)}} \left. \right] e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right), \end{aligned}$$

where

$$\mathcal{N}_{n_1, n_2} = \frac{(n_1 + n_2)^3 n_1}{\binom{n_1 + n_2}{n_1} n_1^3 n_2^3} \sum_{\substack{X_1 \cup X_2 = I' \\ X_1 \ni 2 \\ |X_1| = n_1}} 1 = \frac{\binom{n_1 + n_2 - 2}{n_1 - 1} (n_1 + n_2)^3}{\binom{n_1 + n_2}{n_1} n_1^2 n_2^3} = \frac{(n_1 + n_2)^2}{(n_1 + n_2 - 1) n_1 n_2^2}. \quad (\text{B.11})$$

Expanding again $\frac{\delta O_b}{\delta \psi_{\eta_1}^+}(\psi + \zeta) = \int d\eta c_b(\eta_1, \eta)(\psi_\eta^- + \zeta_\eta^-)$ and using Eq. (B.1), we find that the second line of Eq. (B.8) can be written as

$$\begin{aligned} &e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \int db_1 db d\eta c_{b_1}(\eta_1, \eta) \left(\psi_\eta^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \times \\ &\times \left(- \sum_{n \geq 2} \frac{\mu_{N,\lambda} \lambda g_{b_1, b}^A}{(n-1)^2} \frac{\delta}{\delta G_b^{(n-1)}} + \sum_{n_1 \geq 1, n_2 \geq 2} 2^{-N} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \lambda g_{b_1, b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} \right) e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})}. \end{aligned}$$

Let us now discuss the third line in the r.h.s. of Eq. (B.8), which can be rewritten as

$$\begin{aligned} & \sum_{n \geq 2} 2^{-N} \int d\mathbf{b} e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mathcal{E} \left(\int d\eta c_{b_1}(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \times \right. \\ & \times \left. \left(n 2^{N(1-n)} \tilde{v}_n(\mathbf{b}) O_{\mathbf{b}'}(\psi + \cdot) \right) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right) = \\ & - \sum_{n \geq 2} 2^{-N} \int d\mathbf{b}_1 d\eta c_{b_1}(\eta_1, \eta) e^{\mathcal{V}^{(h)}} \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \frac{\delta}{\delta \dot{G}_{b_1}^{(n)}} e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})}. \end{aligned}$$

Finally, the fourth line in the r.h.s. of Eq. (B.8) can be easily handled using again Eq. (B.1) and recalling that $v_1(\mathbf{b}) = \mu_{N, \lambda}$:

$$\begin{aligned} & e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mu_{N, \lambda} \int d\mathbf{b} G_b^{(1)} \mathcal{E} \left(\frac{\delta O_b}{\delta \psi_{\eta_1}^+} (\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G^{(1)}; \mathbf{0})} \right) = \\ & e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mu_{N, \lambda} \int d\mathbf{b} d\eta G_b^{(1)} c_b(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})}. \end{aligned}$$

All in all we have rewritten Eq. (B.8) as

$$\begin{aligned} & \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta_1}^+} (\psi; G^{(1)}; \mathbf{0}) = \\ & e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \int d\mathbf{b}_1 d\eta c_{b_1}(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \left\{ 2^N (\mu_{N, \lambda} - 1) + \right. \\ & - \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int d\mathbf{b} \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n)}} + 2^{-N} \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2}} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \int d\mathbf{b} \lambda g_{b_1, b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} + \\ & \left. - 2^{-N} \sum_{n \geq 2} \frac{\delta}{\delta \dot{G}_{b_1}^{(n)}} \right\} e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} + \\ & + e^{\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})} \mu_{N, \lambda} \int d\mathbf{b} d\eta G_b^{(1)} c_b(\eta_1, \eta) \left(\psi_{\eta}^- + \int d\eta' g_{\eta, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} \right) e^{-\mathcal{V}^{(h)}(\psi; G^{(1)}; \mathbf{0})}. \end{aligned}$$

By setting also $G^{(1)} = 0$ we obtain Eq. (B.6), while, deriving both sides w.r.t. $G^{(1)}$ and then setting $G^{(1)} = 0$, we find Eq. (B.7). \square

Lemma B.2. *The following identities hold for the effective potential $\mathcal{V}^{(h)}$.*

$$\begin{aligned} & \frac{\delta \mathcal{V}^{(h)}}{\delta G_{b_1}^{(n)}} = \\ & e^{\mathcal{V}^{(h)}} \int d\eta_1 d\eta_2 c_{b_1}(\eta_1, \eta_2) \left(\psi_{\eta_1}^+ \psi_{\eta_2}^- + \psi_{\eta_2}^- \int d\eta g_{\eta, \eta_1}^{(h, N)} \frac{\delta}{\delta \psi_{\eta}^-} + \psi_{\eta_1}^+ \int d\eta' g_{\eta_2, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} + \right. \\ & \left. - g_{\eta_2, \eta_1}^{(h, N)} + \int d\eta d\eta' g_{\eta, \eta_1}^{(h, N)} g_{\eta_2, \eta'}^{(h, N)} \frac{\delta^2}{\delta \psi_{\eta'}^+ \delta \psi_{\eta}^-} \right) \times \\ & \times \left\{ \mathbb{1}_{(n=1)} \mu_{N, \lambda} - \mathbb{1}_{(n \geq 2)} \frac{n^2}{(n-1)^2} 2^{-N} \mu_{N, \lambda} \int d\mathbf{b} \lambda g_{b_1, b}^A \frac{\delta}{\delta G_b^{(n-1)}} + \right. \\ & + \mathbb{1}_{(n \geq 3)} 2^{-2N} \sum_{\substack{n_1 \geq 1, n_2 \geq 2 \\ n_1 + n_2 = n}} \mathcal{N}_{n_1, n_2} \int d\mathbf{b} \lambda g_{b_1, b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} \left. \right\} e^{-\mathcal{V}^{(h)}}, \end{aligned} \tag{B.12}$$

where \mathcal{N}_{n_1, n_2} are as in Lemma B.1. Moreover:

$$\begin{aligned} \frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b_1}^{(n)}} &= \\ e^{\mathcal{V}^{(h)}} \int db \left\{ -\frac{\mu_{N,\lambda} n^2}{(n-1)^2} \lambda g_{b_1,b}^A \frac{\delta}{\delta G_b^{(n-1)}} + 2^{-N} \mathbb{1}_{(n \geq 3)} \sum_{\substack{n_1 \geq 1 \\ n_2 \geq 2 \\ n_1 + n_2 = n}} \mathcal{N}_{n_1, n_2} \lambda g_{b_1,b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} \right\} e^{-\mathcal{V}^{(h)}}. \end{aligned} \quad (\text{B.13})$$

Proof. The proof follows the same ideas used for Lemma B.1 and will not be belabored in detail. The starting point is the identity:

$$\frac{\delta \mathcal{V}^{(h)}}{\delta G_{b_1}^{(n)}}(\psi; G; \dot{G}; \ddot{G}) = e^{\mathcal{V}^{(h)}(\psi; G; \dot{G}; \ddot{G})} \mathcal{E}\left(\frac{\delta \mathcal{V}}{\delta G_{b_1}^{(n)}}(\psi + \cdot; G; \dot{G}; \ddot{G}) e^{-\mathcal{V}(\psi + \cdot; G; \dot{G}; \ddot{G})}\right)$$

for proving Eq. (B.12), and the same one with the derivative w.r.t. $\dot{G}^{(n)}$ instead of $G^{(n)}$ for proving Eq. (B.13). Note that:

$$\frac{\delta \mathcal{V}}{\delta G_{b_1}^{(n)}}(\psi + \zeta; G; \dot{G}; \ddot{G}) = n^3 2^{N(1-n)} \int d\bar{b}' v_n(\bar{b}) O_{\bar{b}}(\psi + \zeta).$$

Using Eq. (B.9) for rewriting v_n and following the same steps as for the proof of Lemma B.1, we find:

$$\begin{aligned} \frac{\delta \mathcal{V}^{(h)}}{\delta G_{b_1}^{(n)}}(\psi; G; \dot{G}; \ddot{G}) &= e^{\mathcal{V}^{(h)}} \left\{ \mathbb{1}_{(n=1)} \mu_{N,\lambda} - 2^{-N} \mathbb{1}_{(n \geq 2)} \frac{\mu_{N,\lambda} n^2}{(n-1)^2} \int db \lambda g_{b_1,b}^A \frac{\delta}{\delta G_b^{(n-1)}} + \right. \\ &\quad \left. + 2^{-2N} \mathbb{1}_{(n \geq 3)} \int db \sum_{\substack{n_1 \geq 1, n_2 \geq 2 \\ n_1 + n_2 = n}} \mathcal{N}_{n_1, n_2} \lambda g_{b_1,b}^A \frac{\delta^2}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} \right\} \mathcal{E}\left(O_{b_1}(\psi + \cdot) e^{-\mathcal{V}(\psi + \cdot; G; \dot{G}; \ddot{G})}\right), \end{aligned} \quad (\text{B.14})$$

and the same expression holds for $\frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b_1}^{(n)}}$ without the factor $O_{b_1}(\psi + \zeta)$ within the Grassmann integral, yielding immediately Eq. (B.13). Letting $\mathcal{V} = \mathcal{V}(\psi + \zeta; G; \dot{G}; \ddot{G})$, we can rewrite:

$$\begin{aligned} \mathcal{E}\left(O_{b_1}(\psi + \cdot) e^{-\mathcal{V}}\right) &= \int d\eta_1 d\eta_2 c_{b_1}(\eta_1, \eta_2) \mathcal{E}\left((\psi_{\eta_1}^+ + (\cdot)_{\eta_1}^+) (\psi_{\eta_2}^- + (\cdot)_{\eta_2}^-) e^{-\mathcal{V}}\right) \\ &= \int d\eta_1 d\eta_2 c_{b_1}(\eta_1, \eta_2) \mathcal{E}\left(\left[-(\psi_{\eta_2}^- + (\cdot)_{\eta_2}^-) \left(\psi_{\eta_1}^+ - \int d\eta g_{\eta, \eta_1}^{(h, N)} \frac{\delta}{\delta (\cdot)_{\eta}^-} \right) - g_{\eta_2, \eta_1}^{(h, N)} \right] e^{-\mathcal{V}}\right) = \\ &= \int d\eta_1 d\eta_2 c_{b_1}(\eta_1, \eta_2) \left(\psi_{\eta_1}^+ \psi_{\eta_2}^- + \psi_{\eta_2}^- \int d\eta g_{\eta, \eta_1}^{(h, N)} \frac{\delta}{\delta \psi_{\eta}^-} + \psi_{\eta_1}^+ \int d\eta' g_{\eta_2, \eta'}^{(h, N)} \frac{\delta}{\delta \psi_{\eta'}^+} + \right. \\ &\quad \left. + \int d\eta d\eta' g_{\eta, \eta_1}^{(h, N)} g_{\eta_2, \eta'}^{(h, N)} \frac{\delta^2}{\delta \psi_{\eta'}^+ \delta \psi_{\eta}^-} - g_{\eta_2, \eta_1}^{(h, N)} \right) e^{-\mathcal{V}^{(h)}}, \end{aligned}$$

which, once plugged in the r.h.s. of Eq. (B.14), yields Eq. (B.12). \square

B.2 Identities for the kernels

Lemma B.1 and Lemma B.2 can be used to infer the identities among the kernels of $\mathcal{V}^{(h)}$, by means of Eq. (2.10).

Proof of Lemma 3.2. This is actually a corollary of Eq. (B.6):

$$\begin{aligned}
\frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta_1}^+}(\psi; \mathbf{0}) &= \int db_1 \int d\eta c_{b_1}(\eta_1, \eta) \left(\psi_{\eta}^- - \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^+} + \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \times \\
&\quad \times \left\{ 2^N (\mu_{N, \lambda} - 1) + \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int db \lambda g_{b_1, b}^A \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n)}} + \right. \\
&\quad - 2^{-N} \sum_{n_1 \geq 1, n_2 \geq 2} \frac{\mathcal{N}_{n_1, n_2}}{(n_1 + n_2)^2} \int db \lambda g_{b_1, b}^A \left(\frac{\delta^2 \mathcal{V}^{(h)}}{\delta G_b^{(n_1)} \delta \dot{G}_{b_1}^{(n_2)}} + \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n_1)}} \frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b_1}^{(n_2)}} \right) + \\
&\quad \left. + \sum_{n \geq 2} 2^{-N} \frac{\delta \mathcal{V}^{(h)}}{\delta \ddot{G}_{b_1}^{(n)}} \right\} (\psi; \mathbf{0}).
\end{aligned} \tag{B.15}$$

Eq. (3.3) follows by deriving both sides of Eq. (B.15) w.r.t. $\psi_{\eta_2}^-$ and then setting $\psi = 0$. $\mathcal{A}_1(\eta_1, \eta_2)$ comes from the first term in curly bracket in the r.h.s. of Eq. (B.15):

$$\begin{aligned}
\mathcal{A}_1(\eta_1, \eta_2) &= 2^N (\mu_{N, \lambda} - 1) \int db_1 \int d\eta c_{b_1}(\eta_1, \eta) \frac{\delta}{\delta \psi_{\eta_2}^-} \left[\psi_{\eta}^- - \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^+} \right] (\mathbf{0}) \\
&= 2^N (\mu_{N, \lambda} - 1) \int db_1 \int d\eta c_{b_1}(\eta_1, \eta) \left[\delta(\eta - \eta_2) - \int d\eta' g_{\eta, \eta'}^{(h, N]} W_{\emptyset; \emptyset; \emptyset}^{(h) | 1, 1}(\eta', \eta_2) \right].
\end{aligned}$$

$\mathcal{A}_2(\eta_1, \eta_2)$ comes from the second term in curly bracket in the r.h.s. of Eq. (B.15):

$$\begin{aligned}
\mathcal{A}_2(\eta_1, \eta_2) &= \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int db db_1 \int d\eta \lambda g_{b_1, b}^A c_{b_1}(\eta_1, \eta) \times \\
&\quad \times \left[\frac{\delta}{\delta \psi_{\eta_2}^-} \left(\psi_{\eta}^- - \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^+} + \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta}{\delta \psi_{\eta'}^+} \right) \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n)}} \right] (\mathbf{0}) = \\
&= \sum_{n \geq 1} \frac{\mu_{N, \lambda}}{n^2} \int db db_1 \int d\eta \lambda g_{b_1, b}^A c_{b_1}(\eta_1, \eta) \left[\delta(\eta - \eta_2) \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n)}} (\mathbf{0}) + \right. \\
&\quad \left. - \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta^2 \mathcal{V}^{(h)}}{\delta \psi_{\eta_2}^- \delta \psi_{\eta'}^+} (\mathbf{0}) \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n)}} (\mathbf{0}) + \int d\eta' g_{\eta, \eta'}^{(h, N]} \frac{\delta^3 \mathcal{V}^{(h)}}{\delta \psi_{\eta_2}^- \delta \psi_{\eta'}^+ \delta G_b^{(n)}} (\mathbf{0}) \right].
\end{aligned}$$

Similarly, $\mathcal{A}_3(\eta_1, \eta_2)$ and $\mathcal{A}_4(\eta_1, \eta_2)$ come from the third and fourth terms respectively in the curly bracket in the r.h.s. of Eq. (B.15). \square

Proof of Eq. (3.12). This is a corollary of Eq. (B.13):

$$\begin{aligned}
\frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b_2}^{(n_2)}} &= \frac{n_2^2}{(n_2 - 1)^2} \mu_{N, \lambda} \int db \lambda g_{b_2, b}^A \frac{\delta \mathcal{V}^{(h)}}{\delta G_b^{(n_2-1)}} + \\
&\quad - 2^{-N} \sum_{\substack{n'_1 \geq 1, n'_2 \geq 2 \\ n'_1 + n'_2 = n_2}} \mathcal{N}_{n'_1, n'_2} \int db'_1 db'_2 \lambda g_{b_2, b'_1}^A \delta(b_2 - b'_2) \left(\frac{\delta^2 \mathcal{V}^{(h)}}{\delta G_{b'_1}^{(n'_1)} \delta \dot{G}_{b'_2}^{(n'_2)}} - \frac{\delta \mathcal{V}^{(h)}}{\delta G_{b'_1}^{(n'_1)}} \frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b'_2}^{(n'_2)}} \right).
\end{aligned}$$

Deriving both sides w.r.t. ψ_η^+ , $\psi_{\eta'}^-$ and $G_{b_1}^{(n_1)}$, and setting the fields to zero, we find:

$$\begin{aligned}
& \frac{\delta^4 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b_1}^{(n_1)} \delta \dot{G}_{b_2}^{(n_2)}}(\mathbf{0}) = \\
& \frac{n_2^2}{(n_2 - 1)^2 \mu_{N,\lambda}} \int db \lambda g_{b_2,b}^A \frac{\delta^4 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b_1}^{(n_1)} \delta G_b^{(n_2-1)}}(\mathbf{0}) + \\
& - \sum_{\substack{n'_1 \geq 1, n'_2 \geq 2 \\ n'_1 + n'_2 = n_2}} 2^{-N} \mathcal{N}_{n'_1, n'_2} \int db'_1 db'_2 \lambda g_{b_2, b'_1}^A \delta(b_2 - b'_2) \left[\frac{\delta^5 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b_1}^{(n_1)} \delta G_{b'_1}^{(n'_1)} \delta \dot{G}_{b'_2}^{(n'_2)}}(\mathbf{0}) + \right. \\
& - \frac{\delta^4 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b_1}^{(n_1)} \delta G_{b'_1}^{(n'_1)}}(\mathbf{0}) \frac{\delta \mathcal{V}^{(h)}}{\delta \dot{G}_{b'_2}^{(n'_2)}}(\mathbf{0}) - \frac{\delta^2 \mathcal{V}^{(h)}}{\delta G_{b_1}^{(n_1)} \delta G_{b'_1}^{(n'_1)}}(\mathbf{0}) \frac{\delta^3 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta \dot{G}_{b'_2}^{(n'_2)}}(\mathbf{0}) + \\
& \left. - \frac{\delta^3 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b'_1}^{(n'_1)}}(\mathbf{0}) \frac{\delta^2 \mathcal{V}^{(h)}}{\delta G_{b_1}^{(n_1)} \delta \dot{G}_{b'_2}^{(n'_2)}}(\mathbf{0}) - \frac{\delta \mathcal{V}^{(h)}}{\delta G_{b'_1}^{(n'_1)}}(\mathbf{0}) \frac{\delta^4 \mathcal{V}^{(h)}}{\delta \psi_{\eta'}^- \delta \psi_\eta^+ \delta G_{b_1}^{(n_1)} \delta \dot{G}_{b'_2}^{(n'_2)}}(\mathbf{0}) \right],
\end{aligned}$$

which gives exactly Eq. (3.12), in force of. Eq. (2.10). \square

C The non-interacting bubbles

This appendix is dedicated to the characterization of the non-interacting bubbles $\Pi^{(h,N]}$ and $\Pi_5^{(h,N)}$:

$$\begin{aligned}
\Pi^{(h,N]}(b_1, b_2) &= -\frac{1}{2} \mathcal{E}_{(h,N]}^T(O_{b_1}, O_{b_2}) \equiv \frac{1}{2} \int d\eta_1 d\eta'_1 d\eta_2 d\eta'_2 c_{b_1}(\eta_1, \eta'_1) c_{b_2}(\eta_2, \eta'_2) g_{\eta'_1, \eta_2}^{(h,N]} g_{\eta'_2, \eta_1}^{(h,N]} \\
\Pi_5^{(h,N]}(b_1, b_2) &= -\frac{1}{2} \mathcal{E}_{(h,N]}^T(O_{b_1}, O_{5;b_2}) \equiv \frac{1}{2} \int d\eta_1 d\eta'_1 d\eta_2 d\eta'_2 c_{b_1}(\eta_1, \eta'_1) c_{b_2}^5(\eta_2, \eta'_2) g_{\eta'_1, \eta_2}^{(h,N]} g_{\eta'_2, \eta_1}^{(h,N]},
\end{aligned}$$

(cf. Eq. (2.16)) with c and c^5 defined in Eq. (1.19) and Eq. (1.19) respectively.

C.1 The finiteness of the non-interacting bubbles

In the first part of this appendix we establish the validity of the following result.

Lemma C.1. *There exists $C_\Pi \geq 1$ such that, for every $0 \leq h \leq N$, letting*

$$\hat{\Pi}_{\sharp; \mu, \nu}^{(h,N]}(p) := \frac{1}{2} \sum_{\epsilon, \epsilon' = \pm} \epsilon \epsilon' \int_{\Lambda} \frac{dxdy}{L^2} e^{-ip \cdot (x-y)} \Pi_{\sharp}^{(h,N]}((x, \mu, \epsilon), (y, \nu, \epsilon')), \quad (\text{C.1})$$

(cf. Eq. (5.11)) with \sharp either blank or 5, the following items are true.

1. $\hat{\Pi}_{\mu, \nu}^{(h,N]}(0) = \delta_{\mu, \nu} \hat{\Pi}_{0,0}^{(h,N]}(0)$; $\hat{\Pi}_{5; \mu, \nu}^{(h,N]}(0) = \varepsilon_{\mu, \nu} \hat{\Pi}_{5;0,1}^{(h,N]}(0)$.
2. $|\hat{\Pi}_{\sharp; \mu, \nu}^{(h,N]}(0)| \leq C_\Pi$.
3. For every $v : \Lambda_B \mapsto \mathbb{C}$ such that $v((x, \mu, \epsilon), (y, \nu, \epsilon')) = \epsilon \epsilon' \tilde{v}_{\mu, \nu}(x - y)$ for some $\tilde{v} : \Lambda \times \{0, 1\} \mapsto \mathbb{C}$, we have that

$$\|v * \Pi_{\sharp}^{(h,N]} - v \hat{\Pi}_{\sharp}^{(h,N]}(0)\|_1^w \leq C_\Pi 2^{-h} \|\nabla v\|_1^w,$$

where $\nabla_\alpha v((x, \mu, \epsilon), (y, \nu, \epsilon')) := 2^N \epsilon \epsilon' (\tilde{v}_{\mu, \nu}(x + 2^{-N} \hat{e}_\alpha) - \tilde{v}_{\mu, \nu}(x))$ and $\|\nabla v\|_1^w := \sum_{\alpha=0,1} \|\nabla_\alpha v\|_1^w$.

As a straightforward corollary of Lemma C.1 we have, as a crucial property of the theory, the finiteness of the quantity $\|g^A * \Pi_{\sharp}^{(h,N]}\|_1^w$, which is a main building block for the estimates in Sects. 3 and 4.

Corollary C.2. *There exists a constant $C'_{\Pi} \geq 1$ such that for every $0 \leq h \leq N$, one has: $\|g^A * \Pi_{\sharp}^{(h,N]}\|_1^w \leq C'_{\Pi}$.*

Proof. From Lemma C.1 we find:

$$\begin{aligned} \|g^A * \Pi_{\sharp}^{(h,N]}\|_1^w &\leq \|g^A \hat{\Pi}_{\sharp}^{(h,N]}(0)\|_1^w + \|g^A * \Pi_{\sharp}^{(h,N]} - g^A \hat{\Pi}_{\sharp}^{(h,N]}(0)\|_1^w \\ &\leq \left(2C_{\Pi} \|g^A\|_1^w + C_{\Pi} 2^{-h} \|\nabla g^A\|_1^w\right). \end{aligned}$$

We already know that $\|g^A\|_1^w$ is bounded by a constant (cf. Eq. (1.27) and comments thereafter); about ∇g^A , one can easily show that

$$|\nabla_{\alpha} g_{\mu,\nu}^A(x-y)| \leq \frac{K_{\nabla} \delta_{\mu,\nu}}{2^{-N} + |x-y|_L} e^{-\kappa_3 \sqrt{|x-y|_L}} \quad (\text{C.2})$$

for suitable $K_{\nabla}, \kappa_3 > 0$. Eq. (C.2) readily implies $\|\nabla g^A\|_1^w \leq K'_{\nabla}$, for a suitable $K'_{\nabla} > 0$, thus the claim of Corollary C.2. \square

Proof of Lemma C.1.

Item 1. This is a consequence of two symmetries of the theory, namely *axes flip* and *parity*, defined in Eqs. (E.2), (E.3) respectively. First, under the *axes flip* transformation, one has: $O_{x,\mu,\epsilon}(\psi) \mapsto O_{x,\mu,\epsilon}(\mathcal{F}\psi) = O_{Tx,1-\mu,\epsilon}(\psi)$, where, if $b = (x, \mu, \epsilon)$, $Tb \equiv (Tx, 1 - \mu, \epsilon)$. Moreover $O_{5;x,\mu,\epsilon}(\psi) \mapsto O_{5;x,\mu,\epsilon}(\mathcal{F}\psi) = -O_{5;Tx,1-\mu,\epsilon}(\psi)$. As a consequence:

$$\begin{aligned} \hat{\Pi}_{\mu,\nu}^{(h,N]}(0) &= -\frac{1}{4} \sum_{\epsilon,\epsilon'=\pm} \epsilon \epsilon' \int \frac{dxdy}{L^2} \mathcal{E}_{(h,N]}^T(O_{x,\mu,\epsilon}, O_{y,\nu,\epsilon'}) = \\ &= -\frac{1}{4} \sum_{\epsilon,\epsilon'=\pm} \epsilon \epsilon' \int \frac{dxdy}{L^2} \mathcal{E}_{(h,N]}^T(O_{Tx,1-\mu,\epsilon}, O_{Ty,1-\nu,\epsilon'}) = \hat{\Pi}_{1-\mu,1-\nu}^{(h,N]}(0). \end{aligned}$$

Similarly one finds that $\hat{\Pi}_{5;\mu,\nu}^{(h,N]}(0) = -\hat{\Pi}_{5;1-\mu,1-\nu}^{(h,N]}(0)$. On the other hand, under the *parity* transformation one has $O_b(\psi) \mapsto O_b(\mathcal{P}\psi) = O_{Pb}(\psi)$, where if $b = (x, \mu, \epsilon)$, then $Pb \equiv (Px - \delta_{\mu,1} 2^{-N} \hat{e}_1, \mu, (-1)^{\mu} \epsilon)$. Moreover $O_{5;x,\mu,\epsilon}(\psi) \mapsto O_{5;x,\mu,\epsilon}(\mathcal{P}\psi) = (-1)^{\mu-1} O_{5;Px,\mu,\epsilon}(\psi)$. It follows that

$$\begin{aligned} \hat{\Pi}_{0,1}^{(h,N]}(0) &= -\frac{1}{4} \sum_{\epsilon,\epsilon'=\pm} \epsilon \epsilon' \int \frac{dxdy}{L^2} \mathcal{E}_{(h,N]}^T(O_{x,0,\epsilon}, O_{y,1,\epsilon'}) = \\ &= -\frac{1}{4} \sum_{\epsilon,\epsilon'=\pm} \epsilon \epsilon' \int \frac{dxdy}{L^2} \mathcal{E}_{(h,N]}^T(O_{Px,0,\epsilon}, O_{Py-2^{-N} \hat{e}_1,1,-\epsilon'}) = \\ &= \frac{1}{4} \sum_{\epsilon,\epsilon'=\pm} \epsilon \epsilon' \int \frac{dxdy}{L^2} \mathcal{E}_{(h,N]}^T(O_{Px,0,\epsilon}, O_{Py-2^{-N} \hat{e}_1,1,\epsilon'}) = -\hat{\Pi}_{0,1}^{(h,N]}(0). \end{aligned}$$

Similarly, $\hat{\Pi}_{5;0,0}^{(h,N]}(0) = -\hat{\Pi}_{5;0,0}^{(h,N]}(0)$. All in all: $\hat{\Pi}_{\mu,\nu}^{(h,N]}(0) = \delta_{\mu,\nu} \hat{\Pi}_{0,0}^{(h,N]}(0)$ and $\hat{\Pi}_{5;\mu,\nu}^{(h,N]}(0) = \varepsilon_{\mu,\nu} \hat{\Pi}_{5;0,1}^{(h,N]}(0)$.

Item 2. First of all note that by *charge conjugation* symmetry, Eq. (E.3),¹⁵ we have (compare also with Eq. (5.11)):

$$\hat{\Pi}_{\sharp;\mu,\nu}^{(h,N]}(p) = \sum_{\epsilon=\pm} \epsilon \int_{\Lambda} \frac{dxdy}{L^2} e^{-ip \cdot (x-y)} \Pi_{\sharp}^{(h,N]}((x, \mu, \epsilon), (y, \nu, +)).$$

Let us begin with the *vector bubble*, $\Pi^{(h,N]}$. Recalling the explicit expression for $c_b(\eta_1, \eta_2)$, Eq. (1.19), we have that

$$\begin{aligned} \hat{\Pi}^{(h,N]}((\mu, +), (\nu, +))(0) &\equiv L^{-2} \int_{\Lambda} dxdy \Pi^{(h,N]}((x, \mu, +), (y, \nu, +)) = \\ &\frac{1}{8L^2} \int dz dy \text{Tr} \left\{ (\gamma_{\mu} - r) g^{(h,N]}(x + 2^{-N} \hat{e}_{\mu} - y) (\gamma_{\nu} - r) g^{(h,N]}(y + 2^{-N} \hat{e}_{\nu} - x) \right\} = \\ &\frac{1}{8} \int dz \text{Tr} \left\{ (\gamma_{\mu} - r) g^{(h,N]}(z) (\gamma_{\nu} - r) g^{(h,N]}(-z) \right\} + \\ &+ \frac{1}{8} \sum_{j,j'=h+1}^N \int dz \text{Tr} \left\{ (\gamma_{\mu} - r) g^{(j)}(z) (\gamma_{\nu} - r) \left[g^{(j')}(z) - g^{(j')}(z + 2^{-N}(\hat{e}_{\mu} + \hat{e}_{\nu})) \right] \right\}. \end{aligned} \quad (\text{C.3})$$

Note that the last line at the r.h.s. of Eq. (C.3) can be bounded by

$$\frac{1}{2} 2^{-N} \sum_{h+1 \leq j \leq j' \leq N} \|g^{(j)}\|_{\infty} \sum_{\mu=0,1} \|\nabla_{\mu} g^{(j')}\|_1 \leq \sum_{h+1 \leq j \leq j' \leq N} \mathcal{O}(2^{-N+j}) = \mathcal{O}(1),$$

where we used that $\|\nabla_{\mu} g^{(j')}\|_1 = \mathcal{O}(1)$, with $\mathcal{O}(1)$ denoting quantities which are bounded by universal constants. Therefore:

$$\hat{\Pi}^{(h,N]}((\mu, +), (\nu, +))(0) = \frac{1}{8} \int dz' \text{Tr} \left\{ (\gamma_{\mu} - r) g^{(h,N]}(z') (\gamma_{\nu} - r) g^{(h,N]}(-z') \right\} + \mathcal{O}(1). \quad (\text{C.4})$$

Similarly, one finds:

$$\hat{\Pi}^{(h,N]}((\mu, -), (\nu, +))(0) = -\frac{1}{8} \int dz' \text{Tr} \left\{ (\gamma_{\mu} + r) g^{(h,N]}(z') (\gamma_{\nu} - r) g^{(h,N]}(-z') \right\} + \mathcal{O}(1). \quad (\text{C.5})$$

Collecting Eqs. (C.4) and (C.5), we find:

$$\begin{aligned} \sum_{\epsilon=\pm} \epsilon \hat{\Pi}^{(h,N]}((\mu, \epsilon), (\nu, +))(0) &= \\ &\frac{1}{8} \sum_{\epsilon=\pm} \int dz' \text{Tr} \left\{ (\gamma_{\mu} - \epsilon r) g^{(h,N]}(z') (\gamma_{\nu} - r) g^{(h,N]}(-z') \right\} + \mathcal{O}(1) = \\ &\frac{1}{4} \int_{\Lambda^*} dk \frac{\chi_{(h,N]}^2(k)}{\left(|s(k)|^2 + (m_N + M_N(k))^2 \right)^2} \times \\ &\times \text{Tr} \left\{ \gamma_{\mu} (i\gamma(k) + m_N + M_N(k)) (\gamma_{\nu} - r) (i\gamma(k) + m_N + M_N(k)) \right\} + \mathcal{O}(1). \end{aligned} \quad (\text{C.6})$$

The trace in the r.h.s. of Eq. (C.6) can be computed using that $\text{Tr}\{\gamma_{\mu} \gamma_{\nu}\} = 2\delta_{\mu,\nu}$, $\text{Tr}\{\gamma_{\mu}\} =$

¹⁵Note that under the *charge conjugation* transformation, $O_{5;x,\mu,\epsilon}(\psi) \mapsto O_{5;x,\mu,\epsilon}(\Omega\psi) = O_{5;x,\mu,-\epsilon}(\psi)$.

$\text{Tr}\{\gamma_\mu\gamma_\nu\gamma_\alpha\} = 0$ and $\text{Tr}\{\gamma_\mu\gamma_\alpha\gamma_\nu\gamma_\beta\} = 2(\delta_{\mu,\alpha}\delta_{\nu,\beta} - \delta_{\mu,\nu}\delta_{\alpha,\beta} + \delta_{\mu,\beta}\delta_{\nu,\alpha})$. We find that

$$\begin{aligned} \sum_{\epsilon=\pm} \epsilon \hat{\Pi}^{(h,N]}((\mu, \epsilon), (\nu, +))(0) &= \\ &= -\frac{1}{2} \int_{\Lambda^*} dk \frac{\chi_{(h,N]}^2(k)}{(|s(k)|^2 + (m_N + M_N(k))^2)^2} \left(2s_\mu(k)s_\nu(k) - \delta_{\nu,\mu}|s(k)|^2 \right) + \\ &\quad - ir \int_{\Lambda^*} dk \chi_{(h,N]}^2(k) \frac{(m_N + M_N(k))s_\mu(k)}{(|s(k)|^2 + (m_N + M_N(k))^2)^2} + \\ &\quad + \frac{1}{2} \delta_{\mu,\nu} \int_{\Lambda^*} dk \chi_{(h,N]}^2(k) \frac{(m_N + M_N(k))^2}{(|s(k)|^2 + (m_N + M_N(k))^2)^2} + \mathcal{O}(1). \end{aligned}$$

Now, the first two terms in the r.h.s. above are checked to be zero after *axes flip*: $(k_0, k_1) \mapsto (k_1, k_0)$ and *axes inversion*: $(k_0, k_1) \mapsto -(k_0, k_1)$.¹⁶ The third term is non-zero, but still finite:

$$\int_{\Lambda^*} dk \chi_{(h,N]}^2(k) \frac{(m_N + M_N(k))^2}{(|s(k)|^2 + (m_N + M_N(k))^2)^2} \leq 4\pi^2 \int_{\Lambda^*} dk \chi_{(0,N]}^2(k) \frac{m_N^2 + r^2 2^{-2N}|k|^4}{(|k|^2 + m_N^2)^2} = \mathcal{O}(1),$$

where we used Eq. (2.3); it follows that $\hat{\Pi}_{\mu,\nu}^{(h,N]}(0) = \mathcal{O}(1)$.

Now let us discuss the *chiral bubble* $\hat{\Pi}_5^{(h,N]}$. In analogy with the *vector bubble*, we have:

$$\hat{\Pi}_5^{(h,N]}((\mu, \epsilon), (\nu, +))(0) = \frac{Z_N^5}{8} \int_{\Lambda} dz \text{Tr} \left\{ (\epsilon \gamma_\mu - r) g^{(h,N]}(z) \gamma_\nu \gamma_5 g^{(h,N]}(-z) \right\} + \mathcal{O}(1).$$

Note that $\gamma_\nu \gamma_5 = i \sum_{\sigma=0,1} \epsilon_{\nu,\sigma} \gamma_\sigma$, therefore:

$$\begin{aligned} \sum_{\epsilon=\pm} \epsilon \hat{\Pi}_5^{(h,N]}((\mu, \epsilon), (\nu, +))(0) &= \frac{iZ_N^5}{4} \sum_{\sigma=0,1} \epsilon_{\nu,\sigma} \int_{\Lambda^*} dk \frac{\chi_{(h,N]}^2(k)}{(|s(k)|^2 + (m_N + M_N(k))^2)^2} \times \\ &\quad \times \text{Tr} \left\{ \gamma_\mu (i\gamma(k) + m_N + M_N(k)) \gamma_\sigma (i\gamma(k) + m_N + M_N(k)) \right\} + \mathcal{O}(1) \end{aligned}$$

and the computation reduces exactly to the one for the simple bubble, so that:

$$\hat{\Pi}_{5;\mu,\nu}^{(h,N]}(0) \equiv \sum_{\epsilon=\pm} \epsilon \hat{\Pi}_5^{(h,N]}((\mu, \epsilon), (\nu, +))(0) = \mathcal{O}(1).$$

Item 3. Letting $P_{\sharp}^{(h)}(b, b') := (v * \hat{\Pi}_{\sharp}^{(h,N]}) (b, b') - (v \hat{\Pi}_{\sharp}^{(h,N]}(0)) (b, b')$, we have that

$$\begin{aligned} P_{\sharp}^{(h)}((x, \mu, \epsilon), (y, \nu, \epsilon')) &= \\ &\sum_{\alpha \in \{0,1\}} \sum_{\epsilon'' \in \{\pm\}} \epsilon \epsilon'' \int_{\Lambda} dz \left(v_{\mu,\alpha}(x-z) - v_{\mu,\alpha}(x-y) \right) \hat{\Pi}_{\sharp}^{(h,N]}((z, \alpha, \epsilon''), (y, \nu, \epsilon')). \end{aligned} \tag{C.7}$$

We rewrite: $v_{\mu,\alpha}(x-z) - v_{\mu,\alpha}(x-y) = \int_{\mathcal{C}_{y \rightarrow z}} d\ell \cdot \nabla v_{\mu,\alpha}(x-\ell)$, with the following understanding.

- $\mathcal{C}_{y \rightarrow z}$ is a path on the discrete torus Λ connecting y to z , with minimal length. More precisely, $\mathcal{C}_{y \rightarrow z}$ is a collection of ordered couples:

$$((x_1, x_2), (x_2, x_3), \dots, (x_{n-1}, x_n), (x'_1, x'_2), (x'_2, x'_3), \dots, (x'_{m-1}, x'_m))$$

with the constraint that $x_1 = y$, $x'_m = z$; $x_{i+1} - x_i = 2^{-N} \sigma_0 \hat{e}_0$ and $x'_{i+1} - x'_i = 2^{-N} \sigma_1 \hat{e}_1$, for some $0 \leq m, n \leq \frac{2^N L}{2} + 1$ and $\sigma_0, \sigma_1 \in \{\pm\}$.

¹⁶Note that the vanishing of the first term would not hold in dimension $d \neq 2$.

- If $\mathcal{C}_{y \rightarrow z} = \{(x_1, x_2), (x_2, x_3), \dots, (x_n, x_{n+1})\}$, then

$$\int_{\mathcal{C}_{y \rightarrow z}} d\ell \cdot \nabla v_{\mu, \alpha}(x - \ell) = \sum_{i=1}^n 2^{-N} \left(\frac{v_{\mu, \alpha}(x - x_{i+1}) - v_{\mu, \alpha}(x - x_i)}{2^{-N}} \right),$$

where note that the difference $\left(\frac{v_{\mu, \alpha}(x - x_{i+1}) - v_{\mu, \alpha}(x - x_i)}{2^{-N}} \right)$ equals, up to a sign, the discrete gradient $\nabla_\nu v_{\mu, \alpha}(x - u) \equiv \left(\frac{v_{\mu, \alpha}(x - u + 2^{-N} \hat{e}_\nu) - v_{\mu, \alpha}(x - u)}{2^{-N}} \right)$, with either $x_{i+1} + 2^{-N} \hat{e}_\nu = x_i = u$ or $x_i + 2^{-N} \hat{e}_\nu = x_{i+1} = u$.

Using this interpolated expansion for the difference of v 's, we can estimate:

$$\begin{aligned} \|P_\sharp^{(h)}\|_1^w &= \frac{1}{L^2} \sum_{\mu, \nu \in \{0, 1\}} \sum_{\epsilon, \epsilon' \in \{\pm\}} \int_{\Lambda^2} dx dy e^{\frac{\kappa}{2} \sqrt{|x - y|_L}} \left| P_\sharp^{(h)}((x, \mu, \epsilon), (y, \nu, \epsilon')) \right| \leq \\ &\leq \frac{1}{L^2} \sum_{\mu, \nu, \alpha \in \{0, 1\}} \sum_{\epsilon, \epsilon', \epsilon'' \in \{\pm\}} \int_{\Lambda^3} dx dy dz e^{\frac{\kappa}{2} \sqrt{|x - y|_L}} \left| \int_{\mathcal{C}_{y \rightarrow z}} d\ell \cdot \nabla v_{\mu, \alpha}(x - \ell) \right| \times \\ &\quad \times \left| \Pi_\sharp^{(h, N]}((z, \alpha, \epsilon''), (y, \nu, \epsilon')) \right| \end{aligned}$$

Note that for any $u \in \mathcal{C}_{y \rightarrow z}$, the quantity $\sqrt{|x - y|_L}$ is smaller than $\sqrt{|x - u|_L} + \sqrt{|z - y|_L}$. Therefore, using that the length of the path $\mathcal{C}_{y \rightarrow z}$ is smaller than $\sqrt{2}|z - y|_L$, we find that

$$\begin{aligned} \|P_\sharp^{(h)}\|_1^w &\leq C \|\nabla v\|_1^w \sum_{\nu, \alpha \in \{0, 1\}} \sum_{\epsilon', \epsilon'' \in \{\pm\}} \int_{\Lambda^2} \frac{dy dz}{L^2} \left| \Pi_\sharp^{(h, N]}((z, \alpha, \epsilon''), (y, \nu, \epsilon')) \right| e^{\frac{\kappa}{2} \sqrt{|y - z|_L}} |y - z|_L \\ &\equiv C \|\nabla v\|_1^w \int_{\Lambda_B^2} \frac{db db'}{L^2} \left| \Pi_\sharp^{(h, N]}(b, b') \right| e^{\frac{\kappa}{2} \sqrt{\delta^D(b, b')}} \delta^D(b, b'). \end{aligned} \tag{C.8}$$

for a suitable constant $C > 0$. Expanding the expression of $\Pi_\sharp^{(h, N]}$ and decomposing in scale each propagator in the same fashion as Eq. (3.26) and lines below, we can estimate the r.h.s. of Eq. (C.8) by

$$C' \|\nabla v\|_1^w \sum_{h+1 \leq j' \leq j \leq N} 2^{-j} \|g^{(j')}\|_\infty \|g^{(j)}(\cdot) e^{\frac{\kappa}{2} \sqrt{2^j | \cdot |_L}}\|_1 \leq \frac{C_\Pi}{4} \|\nabla v\|_1^w \sum_{h+1 \leq j' \leq j \leq N} 2^{j' - 2j},$$

for some $C_\Pi > 0$, where we used the fact that, according to Eq. (2.4),

$$\|g^{(j)}(\cdot) e^{\frac{\kappa}{2} \sqrt{2^j | \cdot |_L}}\|_1 \equiv \sum_{s, s' \in \{1, 2\}} \int_{\Lambda} dx |g_{s, s'}^{(j)}(x)| e^{\frac{\kappa}{2} \sqrt{2^j |x|_L}} \leq \tilde{K} 2^{-j},$$

for a suitable constant $\tilde{K} > 0$. Hence, as desired:

$$\|P_\sharp^{(h)}\|_1^w \leq \frac{C_\Pi}{4} \|\nabla v\|_1^w \sum_{h+1 \leq j' \leq j \leq N} 2^{j' - 2j} \leq C_\Pi 2^{-h} \|\nabla v\|_1^w.$$

□

C.2 Proof of Proposition 5.1

For definiteness we discuss only the claims concerning $\hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M, m_N, L)$ and $\hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L)$; the discussion for $\mathcal{T}^{(h_M^*,N]}$ and $\mathcal{T}^{(\leq N)}$ follows similarly.

Limits: proof of Eq. (5.18). By exploiting the *charge conjugation* symmetry, Eq. (E.1), combining the definition in Eq. (C.1) with the notations in Eq. (1.19), one explicitly obtains that, for every $p \in \frac{2\pi}{L}\mathbb{Z}^2 \cap (-\frac{\pi}{a}, \frac{\pi}{a}]^2$,

$$\begin{aligned} \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L) &= \\ \frac{1}{8} \int_{\Lambda^*} dk \sum_{\epsilon=\pm} e^{ia(\epsilon k_\mu - k_\nu - p_\nu + \frac{1-\epsilon}{2} p_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \hat{g}^{(\leq N)}(k) (\gamma_\nu - r) \hat{g}^{(\leq N)}(k + p) \right\}, \end{aligned} \quad (\text{C.9})$$

and $\hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M, m_N, L)$ having the same expression with an extra factor $(1 - \chi_{h_M^*}(|k|))(1 - \chi_{h_M^*}(|k + p|))$ under integral sign. Notice that the r.h.s. of Eq. (C.9) makes sense for every $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$, thus defining an extension of $\hat{\Pi}_{\mu,\nu}^{(\leq N)}$ which is continuous w.r.t. p , as long as $m_N \neq 0$. By rescaling $k \rightarrow ak = q \in a\Lambda^* \equiv \frac{2\pi a}{L}\mathbb{Z}^2 \cap (-\pi, \pi]^2$, one finds that

$$\begin{aligned} \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L) &= \\ \frac{1}{8} \int_{a\Lambda^*} dq \sum_{\epsilon=\pm} e^{i(\epsilon q_\mu - q_\nu - ap_\nu + \frac{1-\epsilon}{2} ap_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(k + ap) \right\}, \end{aligned} \quad (\text{C.10})$$

where $\tilde{g}(q) = 2^N \hat{g}^{(\leq N)}(2^N q) \equiv \left(-i \sum_{\mu=0,1} \gamma_\mu \sin(q_\mu) + am_N + 2r \sum_{\mu=0,1} \sin^2(q_\mu/2) \right)^{-1}$.

The limit $\lim_{L \rightarrow \infty} \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N, L) =: \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N)$ can be characterized via the Poisson summation formula [31, App. D], which establishes the convergence to the same expression as the r.h.s. of Eq. (C.10) where the Riemann sum over $a\Lambda^*$ is replaced by the integral over $(-\frac{\pi}{a}, \frac{\pi}{a}]^2$. Besides the convergence is uniform w.r.t. $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$ for any $a > 0$ and $m_N \neq 0$ fixed. Similar considerations hold for $\hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M, m_N) := \lim_{L \rightarrow \infty} \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M, m_N, L)$. The existence of $\hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M) := \lim_{m_N \rightarrow 0} \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M, m_N)$ instead trivially follows from the fact that $M > 0$ acts as an infrared cut-off. Finally, for $p \neq 0$, the Dominated Convergence Theorem¹⁷ implies that $\lim_{M \rightarrow 0^+} \hat{\Pi}_{\mu,\nu}^{(h_M^*,N]}(p, M)$ and $\lim_{m_N \rightarrow 0} \hat{\Pi}_{\mu,\nu}^{(\leq N)}(p, m_N)$ exist and are both given by

$$\hat{\Pi}_{\mu,\nu}(p) := \frac{1}{8} \int_{(-\pi, \pi]^2} \frac{d^2 q}{(2\pi)^2} \sum_{\epsilon=\pm} e^{i(\epsilon q_\mu - q_\nu - ap_\nu + \frac{1-\epsilon}{2} ap_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(q + ap) \right\}$$

with the understanding that $\tilde{g}(q)$ and $\tilde{g}(q + ap)$ are both computed at $m_N = 0$.

Computation of $\hat{\Pi}_{\mu,\nu}(p)$: proof of Eq. (5.19). We start by rewriting $\hat{\Pi}_{\mu,\nu}(p) = \hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p) + \hat{\Pi}_{\mu,\nu}^{(\text{reg})}(p)$, where

$$\begin{aligned} \hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p) &:= \frac{1}{8} \int_{(-\pi, \pi]^2} \frac{d^2 q}{(2\pi)^2} \sum_{\epsilon=\pm} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \frac{\chi(|q|)}{-iq} (\gamma_\nu - r) \frac{\chi(|q + ap|)}{-i(q + ap)} \right\} = \\ &- \sum_{\alpha, \beta=0,1} \int_{(-\pi, \pi]^2} \frac{d^2 q}{(2\pi)^2} \frac{\chi(|q|) \chi(|q + ap|) q_\alpha (q_\beta + ap_\beta)}{4|q|^2 |q + ap|^2} \text{Tr} \{ \gamma_\mu \gamma_\alpha \gamma_\nu \gamma_\beta \}, \quad \forall p \neq 0, \end{aligned} \quad (\text{C.11})$$

with $q \equiv \sum_{\mu=0,1} \gamma_\mu q_\mu$ and we used that $\text{Tr}\{\gamma_\mu \gamma_\nu \gamma_\alpha\} = 0$. Besides it is possible to show that

¹⁷It sufficient to observe that if $p \neq 0$ then $\text{Tr}\{(\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(q + ap)\}$ has two simple integrable singularities at $q = 0, q = -ap$.

$$\begin{aligned}\hat{\Pi}_{\mu,\nu}^{(\text{reg})}(p) &:= \frac{1}{8} \sum_{\epsilon=\pm} \int_{(-\pi,\pi]^2} \frac{d^2 q}{(2\pi)^2} \left(-\text{Tr} \left\{ (\gamma_\mu - \epsilon r) \frac{\chi(|q|)}{-i\cancel{q}} (\gamma_\nu - r) \frac{\chi(|q+ap|)}{-i(\cancel{q}+a\cancel{p})} \right\} \right. \\ &\quad \left. + e^{i(\epsilon q_\mu - q_\nu - ap_\nu + \frac{1-\epsilon}{2} ap_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(q+ap) \right\} \right)\end{aligned}\quad (\text{C.12})$$

is a well defined function of $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$, included the case $p = 0$, where the integrand is checked to be singular only at $q = 0$ with an integrable singularity, see Eqs. (C.16)-(C.17) and comments nearby. To conclude it is enough to show the two following properties:

$$\hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p) = \frac{1}{8\pi|p|^2} (2p_\mu p_\nu - |p|^2 \delta_{\mu,\nu}) + \tilde{R}'_{\mu,\nu}(ap), \quad p \neq 0, \quad (\text{C.13})$$

$$|\hat{\Pi}_{\mu,\nu}^{(\text{reg})}(p) - \hat{\Pi}_{\mu,\nu}^{(\text{reg})}(0)| \leq C(a|p|)^{1/2}, \quad (\text{C.14})$$

for some positive constant C and a function $\tilde{R}'_{\mu,\nu}$ such that $|\tilde{R}'_{\mu,\nu}(q)| \leq C|q|^{1/2}$. Notice that this concludes the proof of Eq. (5.19) with $\tilde{R}_{\mu,\nu} = \tilde{R}'_{\mu,\nu} + \hat{\Pi}_{\mu,\nu}^{(\text{reg})}$, with the understanding that $\tilde{R}'_{\mu,\nu}(0) \equiv 0$.

Proof of Eq. (C.13). Given $\omega = \pm$, we introduce $\gamma_\omega := \frac{1}{2}(-i\gamma_0 - \omega\gamma_1)$ so that $\gamma_\mu = \sum_{\omega=\pm} \Omega_{\mu,\omega} \gamma_\omega$, with $\Omega_{\mu,\omega} = i\delta_{\mu,0} - \omega\delta_{\mu,1}$. Using that $\text{Tr}\{\gamma_\mu \gamma_\alpha \gamma_\nu \gamma_\beta\} = 2(\delta_{\mu,\alpha} + \delta_{\mu,\beta} \delta_{\nu,\alpha} - \delta_{\mu,\nu} \delta_{\alpha,\beta})$, one can check that $\text{Tr}\{\gamma_\omega \gamma_\alpha \gamma_{\omega'} \gamma_\beta\} = \delta_{\omega,\omega'} (i\omega\delta_{\alpha,0} + \delta_{\alpha,1}) (i\omega\delta_{\beta,0} + \delta_{\beta,1})$, which means that we can rewrite Eq. (C.11) as

$$\hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p) = \sum_{\omega} \frac{\Omega_{\mu,\omega} \Omega_{\nu,\omega}}{4} \int_{\mathbb{R}^2} \frac{d^2 q}{(2\pi)^2} \frac{\chi(|q|) \chi(|q+ap|)}{D_\omega(q) D_\omega(q+ap)}, \quad D_\omega(q) := -iq_0 + \omega q_1. \quad (\text{C.15})$$

Now it is known [6, App. B.1] that the integral in Eq. (C.15) is given by $\frac{1}{4\pi} \frac{D_{-\omega}(p)}{D_\omega(p)} + \tilde{R}'_\omega(ap)$, with $|\tilde{R}'_\omega(q)| \leq C|q|^{1/2}$, so that letting $4\tilde{R}'_{\mu,\nu}(q) := \sum_{\omega} \Omega_{\mu,\omega} \Omega_{\nu,\omega} \tilde{R}'_\omega(q)$,

$$\begin{aligned}\hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p) - \tilde{R}'_{\mu,\nu}(ap) &= \frac{1}{16\pi|p|^2} \sum_{\omega=\pm} \Omega_{\mu,\omega} \Omega_{\nu,\omega} (p_1^2 - p_0^2 - 2i\omega p_0 p_1) \\ &= \frac{1}{8\pi|p|^2} \left((p_1^2 - p_0^2)(-\delta_{\mu,0}\delta_{\nu,0} + \delta_{\mu,1}\delta_{\nu,1}) - 2p_0 p_1 (\delta_{\mu,0}\delta_{\nu,1} + \delta_{\mu,1}\delta_{\nu,0}) \right) \\ &= \frac{1}{8\pi|p|^2} (2p_\mu p_\nu - \delta_{\mu,\nu} |p|^2).\end{aligned}$$

Proof of Eq. (C.14). For $p \in (-\frac{\pi}{a}, \frac{\pi}{a}]^2$ and $|p| \geq \frac{1}{2}a^{-1}$, by simple dimensional arguments one can easily check that $\hat{\Pi}_{\mu,\nu}(p)$ and $\hat{\Pi}_{\mu,\nu}^{(\text{sing})}(p)$ are both bounded by constant, so that Eq. (C.14) trivially holds for some constant $C > 0$. Hence we must analyze the case $|p| \leq \frac{1}{2}a^{-1}$. Explicitly, we have that the expression in the r.h.s. can be further expanded as $\hat{\Pi}_{\mu,\nu}^{(\text{reg})}(p) = \hat{\Pi}_{\mu,\nu}^{(\text{reg},1)}(p) + \hat{\Pi}_{\mu,\nu}^{(\text{reg},2)}(p)$, where:

$$\begin{aligned}\hat{\Pi}_{\mu,\nu}^{(\text{reg},1)}(p) &= \frac{1}{8} \sum_{\epsilon=\pm} \int_{(-\pi,\pi]^2} \frac{d^2 q}{(2\pi)^2} \left[(1 - \chi(|q+ap|)) + \chi(|q+ap|)(1 - \chi(|q|)) \right] \times \\ &\quad \times e^{i(\epsilon q_\mu - q_\nu - ap_\nu + \frac{1-\epsilon}{2} ap_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(q+ap) \right\},\end{aligned}\quad (\text{C.16})$$

$$\begin{aligned}\hat{\Pi}_{\mu,\nu}^{(\text{reg},2)}(p) &= \frac{1}{8} \sum_{\epsilon=\pm} \int_{(-\pi,\pi]^2} \frac{d^2 q}{(2\pi)^2} \left(e^{i(\epsilon q_\mu - q_\nu - ap_\nu + \frac{1-\epsilon}{2} ap_\mu)} \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \tilde{g}(q) (\gamma_\nu - r) \tilde{g}(q+ap) \right\} \right. \\ &\quad \left. - \text{Tr} \left\{ (\gamma_\mu - \epsilon r) \frac{1}{-i\cancel{q}} (\gamma_\nu - r) \frac{1}{-i(\cancel{q}+a\cancel{p})} \right\} \right) \chi(|q|) \chi(|q+ap|).\end{aligned}\quad (\text{C.17})$$

Now, since for $|p| \leq \frac{1}{2}a^{-1}$ we have that $\tilde{g}(\cdot + ap)$ is never singular on the support of $1 - \chi(|\cdot + ap|)$, it follows that $\hat{\Pi}_{\mu,\nu}^{(\text{reg},1)}$ is actually \mathcal{C}^∞ over the ball of radius $\frac{1}{2}a^{-1}$; ¹⁸ $|\hat{\Pi}_{\mu,\nu}^{(\text{reg},1)}(p) - \hat{\Pi}_{\mu,\nu}^{(\text{reg},1)}(0)| = \mathcal{O}(|ap|)$. In order to evaluate the continuity at zero of $\hat{\Pi}_{\mu,\nu}^{(\text{reg},2)}$ we further decompose, for $p' \in \{0, p\}$, $\hat{\Pi}_{\mu,\nu}^{(\text{reg},2)}(p') = \hat{\Pi}_{\mu,\nu}^{(\text{reg},2,1)}(p'; p) + \hat{\Pi}_{\mu,\nu}^{(\text{reg},2,2)}(p'; p)$, where the former (resp. the latter) is defined as the r.h.s. of Eq. (C.17) with the integral restricted to the ball (resp. the complement of the ball) of radius $\sqrt{a|p|}$. Since the integrand in the r.h.s. of Eq. (C.17) can be bounded, up to a constant, by $(|q||q + ap|)^{-1}$, we find that

$$\begin{aligned} & |\hat{\Pi}_{\mu,\nu}^{(\text{reg},2,1)}(p; p) - \hat{\Pi}_{\mu,\nu}^{(\text{reg},2,1)}(0; p)| \leq \\ & |\hat{\Pi}_{\mu,\nu}^{(\text{reg},2,1)}(p; p)| + |\hat{\Pi}_{\mu,\nu}^{(\text{reg},2,1)}(0; p)| \leq \int_{|q| \leq \sqrt{a|p|}} \frac{d^2q}{(2\pi)^2} \left[\mathcal{O}\left(\frac{1}{|q+ap|}\right) + \mathcal{O}\left(\frac{1}{|q|}\right) \right] = \mathcal{O}(\sqrt{a|p|}). \end{aligned}$$

About $\hat{\Pi}_{\mu,\nu}^{(\text{reg},2,2)}(p'; p)$, letting $F(q, ap)$ be the integrand in the r.h.s. of Eq. (C.17), after writing $F(q, ap) - F(q, 0) = \int_0^1 ds \partial_s F(q, asp)$, by dimensional considerations one finds that $|F(q, ap) - F(q, 0)| \leq \mathcal{O}\left(\frac{a|p|}{|q|^2}\right)$, for any $|q| \geq \sqrt{a|p|}$ and $|p| \leq \frac{1}{2}a^{-1}$. Therefore:

$$|\hat{\Pi}_{\mu,\nu}^{(\text{reg},2,2)}(p; p) - \hat{\Pi}_{\mu,\nu}^{(\text{reg},2,2)}(0; p)| \leq \int_{\sqrt{a|p|} \leq |q| \leq \sqrt{2}\pi} \frac{d^2q}{(2\pi)^2} \mathcal{O}\left(\frac{a|p|}{|q|^2}\right) = \mathcal{O}\left(a|p| \log\left(\frac{1}{a|p|}\right)\right) \leq \mathcal{O}(\sqrt{a|p|}),$$

which concludes the proof of Eq. (C.14). \square

D The tree expansion

In this appendix we prove Eq. (2.26). It is convenient to introduce the *dimensionless norm* $\|\widetilde{W}_{\underline{n};\dot{n};\ddot{n}}^{(h)|q,q}\|_{(h)} := 2^{-hD_{sc}} \|\widetilde{W}_{\underline{n};\dot{n};\ddot{n}}^{(h)|q,q}\|_1^w$. In terms of such norm one readily finds the following partial result.

Lemma D.1. *There exists $C \geq 1$ such that for every $\tau \in \mathcal{T}_{h,N}$ and $\mathbf{P} \in \mathcal{P}_\tau$,*

$$\|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} \leq \prod_{v \in V(\tau) \setminus \{v_0\}} 2^{(h_v - h_{v'})D_{sc}(P_v)} \prod_{v \in V_e(\tau)} C^{|P_v|} \|W_{P_v}^{(h_v-1)}\|_{(h_v-1)}, \quad (\text{D.1})$$

with the understanding that v' is the vertex which precedes v on τ .

We postpone the proof of Lemma D.1 to the end of this section. Using the bound of Eq. (D.1) inside Eq. (2.13), and recalling that only trees with $h_{v_0} = h + 1$ contribute, we have:¹⁹

$$\begin{aligned} \|\widetilde{W}_{\underline{n};\dot{n};\ddot{n}}^{(h)|q,q}\|_{(h)} & \leq \sum_{\tau \in \mathcal{T}_{h,N}}^* \sum_{\mathbf{P} \in \mathcal{P}_\tau}^{**} \|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} \\ & \leq 4^{|P_{v_0}|} \sum_{\tau \in \mathcal{T}_{h,N}}^* \sum_{\mathbf{P} \in \mathcal{P}_\tau}^{**} \prod_{v \in V(\tau)} 2^{(h_v - h_{v'})D_{sc}(P_v)} \prod_{v \in V_e(\tau)} C^{|P_v|} \|W_{P_v}^{(h_v-1)}\|_{(h_v-1)}, \end{aligned} \quad (\text{D.2})$$

where in order to simplify the expression, we have extended the product $\prod_v 2^{(h_v - h_{v'})D_{sc}(P_v)}$ also to $v = v_0$, at the cost of a factor $4^{|P_{v_0}|}$, which can be reabsorbed in the constant C . Now

¹⁸The smoothness is obvious for the contribution to the r.h.s. of Eq. (C.16) from the term $1 - \chi(|q + ap|)$ in square brackets. For the contribution from $\chi(|q + ap|)(1 - \chi(|q|))$, one can first extend the integral over \mathbb{R}^2 and then perform the change of variable $q \mapsto q - ap$, from which the smoothness of the integral follows.

¹⁹Note that the operator \mathcal{S} in the r.h.s. of Eq. (2.13) plays no role in view of an upper bound.

by construction $D_{sc}(P_v) \leq -1$ for every $v \in V(\tau)$ (here is why it is important to stop the expansion of the tree as soon as a non-irrelevant kernel, i.e. $D_{sc} \geq 0$, is encountered), except eventually for the vertex following the root and for the endpoints associated with non-irrelevant terms, for which, however, the quantity $h_v - h_{v'}$ is fixed to be 1. Moreover, if $D_{sc}(P_v) \leq -1$, $D_{sc}(P_v) \leq -\frac{1}{8}|P_v^\psi|$ too, with P_v^ψ the set of field labels associated with ψ variables. Hence one can write:

$$\prod_{v \in V(\tau)} 2^{(h_v - h_{v'})D_{sc}(P_v)} \leq 8^{|V_e|+1} \prod_{v \in V(\tau)} 2^{-\frac{1+\theta}{2}(h_v - h_{v'})} \prod_{v \in V(\tau)} 2^{-\frac{1-\theta}{16}|P_v^\psi|}.$$

The pre-factor $8^{|V_e|+1}$ can be absorbed into the constant C in the r.h.s. of Eq. (D.2), while the factor $\prod_{v \in V(\tau)} 2^{-\frac{1-\theta}{16}|P_v^\psi|}$ can be used to bound the sum over \mathbf{P} for any fixed configuration $\mathbf{P}_{V_e} \equiv \{P_v\}_{v \in V_e(\tau)}$. All in all [29, App. A.6]:

$$\|\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}\|_{(h)} \leq \sum_{\tau \in \mathcal{T}_{h,N}}^* \sum_{\mathbf{P}_{V_e}}^{**} \prod_{v \in V(\tau)} 2^{-\frac{1+\theta}{2}(h_v - h_{v'})} \prod_{v \in V_e(\tau)} C'^{|P_v|} \|W_{P_v}^{(h_v-1)}\|_{(h_v-1)}, \quad (\text{D.3})$$

with a suitable constant $C' \geq 1$. Now the quantity $\|W_{P_v}^{(h_v-1)}\|_{(h_v-1)}$ can be bounded as follows, for every $\epsilon > 0$:

$$\|W_{P_v}^{(h_v-1)}\|_{(h_v-1)} \leq \begin{cases} R\lambda 2^{-(1+\vartheta)(h_v-1)} \leq R\lambda, & P_v \sim \psi^2; \\ C_1^q \lambda^{q-1}, & P_v \sim \psi^{2q} \text{ with } q \geq 2; \\ C_1 \lambda^{\frac{n-1}{2}} \lambda^{\frac{1-\epsilon}{2}(q-1)}, & P_v \sim G^{(n)} \psi^{2q} \text{ with } q \in \{1, n\}; \\ C_1 \lambda^{\frac{n-1}{2}} \lambda^{\frac{1-\epsilon}{2}(q-1)}, & P_v \sim \dot{G}^{(n)} \psi^{2q} \text{ with } q \in \{1, n-1\}; \\ C_1 \lambda^{\frac{5}{8}n} \lambda^{\frac{3}{8}(1-\epsilon)q}, & P_v \sim \ddot{G}^{(n)} \psi^{2n-2}. \end{cases} \quad (\text{D.4})$$

with a suitable $C_1 \geq 1$, and for λ small enough, say²⁰ $\lambda \leq (C_1' R^4)^{-1}$. The convenience for losing a power ϵ in λ is to get in general a pre-factor C_1 instead of $C_1'^{|P_v|}$, in the r.h.s. above. The bounds in Eq. (D.4) can be checked to imply the following estimate for the r.h.s. of Eq. (D.3), after performing the sum $\sum_{\mathbf{P}_{V_e}}^{**}$:

$$\begin{aligned} \|\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)|q,q}\|_{(h)} &\leq C_2^{1+\dim \underline{n}+\dim \dot{\underline{n}}+\dim \ddot{\underline{n}}} \lambda^d \sum_{\tau \in \tilde{\mathcal{T}}_{h,N}} 2^{-\frac{1+\theta}{2}(h_v - h_{v'})} \lambda^{\epsilon'(|V_e| - \max\{1, \dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}\})} \times \\ &\times \begin{cases} R & q = 1, |\underline{n}| = |\dot{\underline{n}}| = |\ddot{\underline{n}}| = 0 \\ 1 & \text{otherwise,} \end{cases} \end{aligned} \quad (\text{D.5})$$

for some $C_2 \geq 1$ and ϵ' small enough, where $\tilde{\mathcal{T}}_{h,N}$ is the set of Gallavotti-Nicolò trees with number of endpoints $\geq \max\{1, \dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}\}$, and such that all the vertices except the endpoints are branching nodes. Note that the parameter R appears in the r.h.s. of Eq. (D.5) only if we are considering the kernel $W_{\emptyset;\emptyset;\emptyset}^{(h)|1,1}$: for every other kernel, indeed, every factor R can be checked to be always accompanied with some extra factor $\lambda^{\frac{1}{2}}$, and $R\lambda^{\frac{1}{2}} \leq 1$ for $\lambda \leq R^{-2}$. Now the sum in the r.h.s. of Eq. (D.5) can be bounded by

$$\sum_{\tau \in \tilde{\mathcal{T}}_{h,N}} 2^{-\theta(h_{\max}(\tau) - h)} 2^{-\frac{1-\theta}{2}(h_v - h_{v'})} \lambda^{\epsilon'(|V_e| - \max\{1, \dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}\})}, \quad (\text{D.6})$$

²⁰The dependence R^{-4} comes by requiring $\|W_{\emptyset;\emptyset;\emptyset}^{(h)|2,2}\|_1^w \leq C\lambda$, with C independent of R , cf. the second of Eq. (2.18).

with $h_{\max}(\tau)$ the highest scale over all the endpoints of τ . Note that in the case the kernel has labels $\underline{n} \neq \emptyset$, i.e. there is at least one field \tilde{G} , we must have $h_{\max}(\tau) = N + 1$, since by construction there are no endpoints with a field \tilde{G} at lower scales. Finally, the sum in the r.h.s. of Eq. (D.6) is readily checked to be bounded by $(8(2^{\frac{1-\theta}{2}} - 1)^{-1})^{2 \max\{1, \dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}\}}$ for λ small enough, therefore:

$$\|\widetilde{W}_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h)q,q}\|_{(h)} \leq C_3^{1+\dim \underline{n} + \dim \dot{\underline{n}} + \dim \ddot{\underline{n}}} \lambda^d \times \begin{cases} 2^{\theta(h-N)}, & |\ddot{\underline{n}}| \geq 1 \\ R, & q = 1, |\underline{n}| = |\dot{\underline{n}}| = |\ddot{\underline{n}}| = 0 \\ 1, & \text{otherwise,} \end{cases}$$

with $C_3 = C_2(8(2^{\frac{1-\theta}{2}} - 1)^{-1})^2$ and again λ small enough (say $\lambda \leq (C_4 R^4)^{-1}$ for a suitable constant $C_4 \geq 1$). The claim of Proposition 2.1 follows with $C_0 := (1 - 2^{\theta-1})^{-1} \max\{C_3, C_4\}$.

In order to complete the proof of Eq. (2.26), we need to prove Lemma D.1.

Proof of Lemma D.1. The starting point is Eq. (2.14):

$$\widetilde{W}[\tau, \mathbf{P}](\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}}) = \frac{\sigma_{\mathbf{P}}}{s_{v_0}!} \int_{\Lambda_F} d\underline{\eta}_{Q_{v_0}} \mathcal{E}_{(h_{v_0})}^T(\psi_{\tilde{P}_{w_1}}, \dots, \psi_{\tilde{P}_{w_{s_{v_0}}}}) \prod_{j=1}^{s_{v_0}} \widetilde{W}[\tau, \mathbf{P}_{w_j}]((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_{w_j}}),$$

from which we find:

$$\begin{aligned} \|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} &\leq \frac{2^{-(h_{v_0}-1)D_{sc}(P_{v_0})}}{L^2 s_{v_0}!} \int_{\Lambda_B} d\underline{b}_{v_0} \int_{\Lambda_F} d\underline{\eta}_{v_0} e^{\frac{\kappa}{2} \sqrt{\delta^D(\underline{b}_{v_0}, \underline{\eta}_{v_0})}} \times \\ &\quad \times |\mathcal{E}_{(h_{v_0})}^T(\psi_{\tilde{P}_{w_1}}, \dots, \psi_{\tilde{P}_{w_{s_{v_0}}}})| \prod_{j=1}^{s_{v_0}} |\widetilde{W}[\tau, \mathbf{P}_{w_j}]((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_{w_j}})|, \end{aligned} \quad (\text{D.7})$$

where $\underline{b}_{v_0} \equiv \underline{b}_{P_{v_0}}$ and $\underline{\eta}_{v_0} = (\underline{\eta}_{P_{v_0}}, \underline{\eta}_{Q_{v_0}})$. As a standard corollary of the Battle-Brydges-Federbush-Kennedy formula [29, Eq. (4.43)] and the Gram-Hadamard inequality [29, Sect. A.3.4], we have the following bound:

$$|\mathcal{E}_{(h_{v_0})}^T(\psi_{\tilde{P}_{w_1}}, \dots, \psi_{\tilde{P}_{w_{s_{v_0}}}})| \leq C_1^{|Q_{v_0}|} 2^{\frac{1}{2}h_{v_0}(\sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - |P_{v_0}^\psi| - 2(s_{v_0}-1))} \sum_T \prod_{(f, f') \in T} |g_{\eta_f, \eta_{f'}}^{(h_{v_0})}|, \quad (\text{D.8})$$

with a suitable $C_1 \geq 1$, where \sum_T is the sum over all the spanning trees over the *clusters* $\tilde{P}_{w_1}, \dots, \tilde{P}_{w_s}$, made of $s_{v_0} - 1$ edges, each of them connecting a field label $f \in \tilde{P}_{w_i}$, associated with a Grassmann variable $\psi_{\eta_f}^-$, with a field label $f' \in \tilde{P}_{w_j}$ (with $j \neq i$), associated with $\psi_{\eta_{f'}}^+$.

Plugging the r.h.s. of Eq. (D.8) into the r.h.s. of Eq. (D.7), and using that

$$\sqrt{\delta^D(\underline{b}_{v_0}, \underline{\eta}_{v_0})} \leq \sum_{j=1}^{s_{v_0}} \sqrt{\delta^D(\underline{b}_{P_{w_j}}, \underline{\eta}_{P_{w_j}})} + \sum_{(f, f') \in T} \sqrt{\delta^D(\eta_f, \eta_{f'})},$$

we find:

$$\begin{aligned} \|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} &\leq \\ &2^{-(h_{v_0}-1)D_{sc}(P_{v_0})} 2^{\frac{1}{2}h_{v_0}(\sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - |P_{v_0}^\psi| - 2(s_{v_0}-1))} \frac{C_1^{|Q_{v_0}|}}{L^2 s_{v_0}!} \sum_T \int_{\Lambda_B} d\underline{b}_{v_0} \int_{\Lambda_F} d\underline{\eta}_{v_0} \times \\ &\quad \times \prod_{(f, f') \in T} \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(\eta_f, \eta_{f'})}} |g_{\eta_f, \eta_{f'}}^{(h_{v_0})}| \right) \prod_{j=1}^{s_{v_0}} \left(e^{\frac{\kappa}{2} \sqrt{\delta^D(\underline{b}_{P_{w_j}}, \underline{\eta}_{P_{w_j}})}} |\widetilde{W}[\tau, \mathbf{P}_{w_j}]((\underline{\eta}; \underline{b}; \dot{\underline{b}}; \ddot{\underline{b}})_{P_{w_j}})| \right). \end{aligned} \quad (\text{D.9})$$

By performing standard *tree-stripping estimates* on T [29, Sect. 6.2], we get:

$$\begin{aligned} \|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} &\leq 2^{-(h_{v_0}-1)D_{sc}(P_{v_0})} 2^{\frac{1}{2}h_{v_0}(\sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - |P_{v_0}^\psi| - 2(s_{v_0}-1))} \frac{C_1^{|Q_{v_0}|}}{s_{v_0}!} \times \\ &\times \sum_T \prod_{(f, f') \in T} \|g^{(h_{v_0})}\|_1^w \prod_{j=1}^{s_{v_0}} \|\widetilde{W}[\tau, \mathbf{P}_{w_j}]\|_1^w. \end{aligned} \quad (\text{D.10})$$

Now, using the fact that $\|g^{(h)}\|_1^w = \mathcal{O}(2^{-h})$ (cf. Eq. (2.4)), that the number of spanning trees T is bounded by $s_{v_0}!e^{2|Q_{v_0}|}$ [29, Lemma A.5], and recalling the definition of the norm $\|\cdot\|_{(h)}$ at the beginning of this section, we find:

$$\begin{aligned} \|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} &\leq 2^{(h_{v_0}-1)[-D_{sc}(P_{v_0}) + \frac{1}{2}\sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - \frac{1}{2}|P_{v_0}^\psi| - 2(s_{v_0}-1)]} C_2^{|Q_{v_0}|} \times \\ &\times \prod_{j=1}^{s_{v_0}} 2^{(h_{w_j}-1)D_{sc}(P_{w_j})} \|\widetilde{W}[\tau, \mathbf{P}_{w_j}]\|_{(h_{w_j}-1)}, \end{aligned} \quad (\text{D.11})$$

with a suitable $C_2 \geq 1$. Now in the r.h.s. of Eq. (D.11) we write $h_{w_j} - 1 = (h_{w_j} - h_{v_0}) + (h_{v_0} - 1)$, and using the fact that $\sum_{j=1}^{s_{v_0}} (|P_{w_j}^G| + |P_{w_j}^{\dot{G}}| + 2|P_{w_j}^{\ddot{G}}|) = |P_{v_0}^G| + |P_{v_0}^{\dot{G}}| + 2|P_{v_0}^{\ddot{G}}|$ (no source fields can be integrated), we note that

$$\begin{aligned} -D_{sc}(P_{v_0}) + \frac{1}{2} \sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - \frac{1}{2}|P_{v_0}^\psi| - 2s_{v_0} + 2 + \sum_{j=1}^{s_{v_0}} D_{sc}(P_{w_j}) &= \\ = -2 + \frac{1}{2}|P_{v_0}^\psi| + |P_{v_0}^G| + |P_{v_0}^{\dot{G}}| + 2|P_{v_0}^{\ddot{G}}| + \frac{1}{2} \sum_{j=1}^{s_{v_0}} |P_{w_j}^\psi| - \frac{1}{2}|P_{v_0}^\psi| - 2s_{v_0} + 2 + \\ + \sum_{j=1}^{s_{v_0}} \left(2 - \frac{1}{2}|P_{w_j}^\psi| - |P_{w_j}^G| - |P_{w_j}^{\dot{G}}| - 2|P_{w_j}^{\ddot{G}}| \right) &= 0. \end{aligned}$$

Therefore Eq. (D.11) becomes

$$\|\widetilde{W}[\tau, \mathbf{P}]\|_{(h_{v_0}-1)} \leq C_2^{|Q_{v_0}|} \prod_{j=1}^{s_{v_0}} 2^{(h_{w_j}-h_{v_0})D_{sc}(P_{w_j})} \|\widetilde{W}[\tau, \mathbf{P}_{w_j}]\|_{(h_{w_j}-1)}, \quad (\text{D.12})$$

which, once iterated over the vertices of τ , yields the desired bound in Eq. (D.1). \square

E Symmetries

In this section we present the symmetries of the massive lattice QED₂: for definiteness we state them only for model with external sources $G(J), B, \varphi$, obtained right after the integration of the boson field A . However, the same symmetries hold for the model which includes the auxiliary external fields G, \dot{G}, \ddot{G} , Eqs. (1.44), (4.2), with the understanding that the transformation for the variables $G_b^{(n)}, \dot{G}_b^{(n)}, \ddot{G}_b^{(n)}$ is the same as the one for $G(J)$ prescribed in the following lemma.

Lemma E.1. *The potential $\mathcal{V}(\psi; G(J)) + (\varphi, \psi) + (B, O_5)$ and the Gaussian integrations $P^{(h)}$ (cf. Eq. (1.21) and comments after Eq. (2.5)) are separately invariant under the following symmetries, with the understanding that the transformation for φ^\pm is the same as for ψ^\pm .*

1. *Global $U(1)$:* $(\mathcal{U}_\alpha \psi)_{x,s}^\pm := e^{\pm i\alpha} \psi_{x,s}^\pm$, for every $\alpha \in [0, 2\pi)$.

2. Charge conjugation:

$$\begin{cases} (\mathcal{Q}\psi)_{x,s}^+ := \sum_{s'} \psi_{x,s'}^- (\gamma_1)_{s,s'} & , \\ (\mathcal{Q}\psi)_{x,s}^- := \sum_{s'} (\gamma_1)_{s,s'} \psi_{x,s'}^+ & , \end{cases} \quad \begin{cases} (\mathcal{Q}G)_b(J) := G_{\bar{b}}(J) & \\ (\mathcal{Q}B)_b := B_{\bar{b}}, & \end{cases} \quad (\text{E.1})$$

where $\bar{b} \equiv \overline{(x, \mu, \epsilon)} = (x, \mu, -\epsilon)$.

3. Axes flip:

$$\begin{cases} (\mathcal{F}\psi)_{x,s}^+ := \sum_{s'} \psi_{Tx,s'}^+ F_{s,s'}^* & , \\ (\mathcal{F}\psi)_{x,s}^- := \sum_{s'} F_{s,s'} \psi_{Tx,s'}^- & , \end{cases} \quad \begin{cases} (\mathcal{F}G)_b(J) := G_{Tb}(J) & \\ (\mathcal{F}B)_b := -B_{Tb}, & \end{cases} \quad (\text{E.2})$$

where $F \equiv \frac{1}{\sqrt{2}}(\gamma_0 - \gamma_1)\gamma_5 \equiv \frac{1}{\sqrt{2}} \begin{pmatrix} i & 1 \\ -1 & -i \end{pmatrix}$, $Tx \equiv (x_1, x_0)$ and $T(x, \mu, \epsilon) \equiv (Tx, 1 - \mu, \epsilon)$.

4. Parity:

$$\begin{cases} (\mathcal{P}\psi)_{x,s}^+ := \sum_{s'} \psi_{Px,s'}^+ (\gamma_5 \gamma_1)_{s',s} & , \\ (\mathcal{P}\psi)_{x,s}^- := \sum_{s'} (\gamma_1 \gamma_5)_{s,s'} \psi_{Px,s'}^- & , \end{cases} \quad \begin{cases} (\mathcal{P}G)_b(J) := G_{Pb}(J) & \\ (\mathcal{P}B)_{x,\mu,\epsilon} := (-1)^{\mu-1} B_{Px,\mu,\epsilon}, & \end{cases} \quad (\text{E.3})$$

where $Px \equiv (x_0, -x_1)$ and $P(x, \mu, \epsilon) \equiv (Px - \delta_{\mu,1} 2^{-N} \hat{e}_1, \mu, (-1)^\mu \epsilon)$.

Proof.

1. Global $U(1)$ is straightforward since every field ψ^+ or φ^+ is always accompanied with a field ψ^- or φ^- .
2. We note that $\gamma_1^T \gamma_\mu^T \gamma_1 = \gamma_\mu$ and $\gamma_1^T \gamma_1 = -1$, hence, recalling that $O_{x,\mu,+}(\psi) \equiv \frac{1}{2} \psi_x^+ (\gamma_\mu - r) \psi_{x+2^{-N} \hat{e}_\mu}^-$ and $O_{x,\mu,-}(\psi) = -\frac{1}{2} \psi_x^+ (\gamma_\mu + r) \psi_x^-$, we find:

$$\begin{cases} O_{x,\mu,+}(\mathcal{Q}\psi) = -\frac{1}{2} \psi_{x+2^{-N} \hat{e}_\mu}^+ \gamma_1^T (\gamma_\mu^T - r) \gamma_1 \psi_x^- = -\frac{1}{2} \psi_{x+2^{-N} \hat{e}_\mu}^+ (\gamma_\mu + r) \psi_x^- = O_{x,\mu,-}(\psi) \\ O_{x,\mu,-}(\mathcal{Q}\psi) = \frac{1}{2} \psi_x^+ \gamma_1^T (\gamma_\mu^T + r) \gamma_1 \psi_{x+2^{-N} \hat{e}_\mu}^- = \frac{1}{2} \psi_x^+ (\gamma_\mu - r) \psi_{x+2^{-N} \hat{e}_\mu}^- = O_{x,\mu,+}(\psi). \end{cases}$$

Besides, recalling Eqs. (1.22), (1.23), and noting that $g_{b,b'}^A = g_{\bar{b},\bar{b}'}^A$, we see that $w_{n,m}(\underline{b}, \underline{b}') = w_{n,m}(\bar{\underline{b}}, \bar{\underline{b}}')$, from which the invariance of $\mathcal{V}(\psi; G(J))$ follows. Moreover, since $\gamma_1^T (\gamma_\mu \gamma_5)^T \gamma_1 = -\gamma_1^T \gamma_5^T \gamma_1 \gamma_1^T \gamma_\mu^T \gamma_1 = -\gamma_5 \gamma_\mu = \gamma_\mu \gamma_5$, we see that $O_{5;x,\mu,-\epsilon}(\mathcal{Q}\psi) = O_{5;x,\mu,\epsilon}(\psi)$, which implies the invariance of the term (B, O_5) . The invariance of the Grassmann integration $P^{(h)}$ means $\mathcal{E}_{(h)}(f(\cdot)) = \mathcal{E}_{(h)}(f(\mathcal{Q}\cdot))$ for every Grassmann polynomial f . In force of the Grassmannian Wick rule [29, Eq. (4.20)], it is sufficient to focus on the case in which f is quadratic, so that the simple expectation reduces to the single-scale propagator.

$$\begin{aligned} \mathcal{E}_{(h)}((\mathcal{Q}\psi)_{x,s}^- (\mathcal{Q}\psi)_{y,s'}^+) &= -(\gamma_1 g^{(h)}(y-x) \gamma_1^T)_{s',s} = \\ &= - \int_{\Lambda^*} dk e^{-ik \cdot (y-x)} f_h(|k|) \left(\gamma_1 \left(-i \not{k} + m_N + M_N(k) \right)^{-1} \gamma_1^T \right)_{s',s} = \\ &= - \int_{\Lambda^*} dk e^{-ik \cdot (y-x)} f_h(|k|) \left(-i \not{k}^T - m_N - M_N(k) \right)_{s',s}^{-1} = \\ &= \int_{\Lambda^*} dk e^{-ik \cdot (x-y)} f_h(|k|) \left(-i \not{k} + m_N + M_N(k) \right)_{s,s'}^{-1} = \mathcal{E}_{(h)}(\psi_{x,s}^- \psi_{y,s'}^+), \end{aligned}$$

where we introduced $f_h(|k|) \equiv \chi_h(|k|) - \chi_{h-1}(|k|)$ for $0 \leq h < N$ and $f_N(|k|) \equiv 1 - \chi_{N-1}(|k|)$, and we used that $s_\mu(-k) = -s_\mu(k)$ and $M_N(-k) = M_N(k)$.

3. We observe that $F^\dagger \gamma_\mu F = \gamma_{1-\mu}$, with F^\dagger the Hermitian conjugate of F , therefore:

$$\begin{cases} O_{x,\mu,+}(\mathcal{F}\psi) = \frac{1}{2} \psi_{Tx}^+(\gamma_{1-\mu} - r) \psi_{Tx+2^{-N}\hat{e}_{1-\mu}}^- = O_{Tx,1-\mu,+}(\psi) \\ O_{x,\mu,-}(\mathcal{F}\psi) = -\frac{1}{2} \psi_{Tx+2^{-N}\hat{e}_{1-\mu}}^+(\gamma_{1-\mu} + r) \psi_{Tx}^- = O_{Tx,1-\mu,-}(\psi). \end{cases}$$

Recalling also that $g_{b,b'}^A = g_{Tb,Tb'}^A$, which implies that $w_{n,m}(b, b') = w_{n,m}(Tb, Tb')$, the invariance of \mathcal{V} follows. For the chiral current we use the fact that $F^\dagger \gamma_5 F = -\gamma_5$, and so $O_{5;Tx,\mu,\epsilon}(\mathcal{F}\psi) = -O_{5;x,1-\mu,\epsilon}(\psi)$. For the Grassmann integration, we have

$$\begin{aligned} \mathcal{E}_{(h)}((\mathcal{F}\psi)_{x,s}^-(\mathcal{F}\psi)_{y,s'}^+) &= \left(Fg^{(h)}(T(x-y))F^\dagger \right)_{s,s'} = \\ &\int_{\Lambda^*} dk e^{-iT\mathbf{k}\cdot(x-y)} f_h(|k|) \left(F(-i\cancel{s}(k) + m_N + M_N(k))^{-1} F^\dagger \right)_{s,s'} = \\ &\int_{\Lambda^*} dk e^{-ik\cdot(x-y)} f_h(|k|) \left(F(-i\cancel{s}(Tk) + m_N + M_N(Tk)) F^\dagger \right)_{s,s'}^{-1} = \\ &\int_{\Lambda^*} dk e^{-ik\cdot(x-y)} f_h(|k|) \left(-i\cancel{s}(k) + m_N + M_N(k) \right)^{-1} = \mathcal{E}_{(h)}(\psi_{x,s}^-\psi_{y,s'}^+), \end{aligned}$$

where we used $s_\mu(k) = s_{1-\mu}(Tk)$ and $M_N(k) = M_N(Tk)$.

4. We observe that $\gamma_5 \gamma_1 \gamma_\mu \gamma_1 \gamma_5 = (-1)^\mu \gamma_\mu$, which implies, by inspection, $O_b(\mathcal{P}\psi) = O_{Pb}(\psi)$. On the other hand we have $g_{b,b'}^A = g_{Pb,Pb'}^A$, so $w_{n,m}(b, b') = w_{n,m}(P\underline{b}, P\underline{b}')$, which implies the invariance of \mathcal{V} under *parity*. Moreover, we note that $(\gamma_5 \gamma_1) \gamma_\mu \gamma_5 (\gamma_1 \gamma_5) = (-1)^{\mu-1} \gamma_\mu \gamma_5$, hence $O_{5;Px,\mu,\epsilon}(\mathcal{P}\psi) = (-1)^{\mu-1} O_{5;x,\mu,\epsilon}(\psi)$. Finally:

$$\begin{aligned} \mathcal{E}_{(h)}((\mathcal{P}\psi)_{x,s}^-(\mathcal{P}\psi)_{y,s'}^+) &= \left(\gamma_1 \gamma_5 g^{(h)}(P(x-y)) \gamma_5 \gamma_1 \right)_{s,s'} = \\ &\int_{\Lambda^*} dk e^{-iPk\cdot(x-y)} f_h(|k|) \left((\gamma_5 \gamma_1)^{-1} (-i\cancel{s}(k) + m_N + M_N(k))^{-1} (\gamma_5 \gamma_1)^{-1} \right)_{s,s'} = \\ &\int_{\Lambda^*} dk e^{-iPk\cdot(x-y)} f_h(|k|) \left((\gamma_1 \gamma_5) (-i\cancel{s}(Pk) + m_N + M_N(Pk)) (\gamma_5 \gamma_1) \right)_{s,s'}^{-1} = \\ &\int_{\Lambda^*} dk e^{-ik\cdot(x-y)} f_h(|k|) \left(-i\cancel{s}(k) + m_N + M_N(k) \right)^{-1} = \mathcal{E}_{(h)}(\psi_{x,s}^-\psi_{y,s'}^+), \end{aligned}$$

where we used that $(\gamma_\mu \gamma_5)^{-1} = \gamma_5 \gamma_\mu$, $s_\mu(Pk) = (-1)^\mu s_\mu(k)$ and $M_N(Pk) = M_N(k)$.

□

F Overview of the continuum limit

We briefly present the scheme of the proof for showing the existence of the limit $N \rightarrow \infty$ of the kernels of the generating functional in Eq. (1.35).

One would like to show the existence of the $N \rightarrow \infty$ limit of the Fourier transforms of the kernels $W_{q,q';\tilde{q},\tilde{q}';p,p'}^{(u.v.;N)}$ of the generating functional, cf. Eqs. (1.35)-(1.36), given (for instance if $\tilde{q} = \tilde{q}' = p = p' = 0$) by

$$\hat{W}_{q,q;0;0;0}^{(u.v.;N)}((\underline{k}, \underline{s}), (\underline{k}', \underline{s}')) := \frac{1}{L^2} \int_{\Lambda_N^{2q}} dx dx' e^{i(\underline{k}\cdot\underline{x} + \underline{k}'\cdot\underline{x}')} W_{q,q;0;0;0}^{(u.v.;N)}(\underline{\eta}, \underline{\eta}'; \underline{b}), \quad \underline{k}, \underline{k}' \in (\Lambda_N^*)^q \quad (\text{F.1})$$

with $\Lambda_N \equiv \Lambda$ and $\Lambda_N^* \equiv \Lambda^*$ for which we have made the dependence on N explicit in every expression, recall the notations below Eq. (1.17).

It is then enough to show a *Cauchy property*, i.e. (assuming with no loss of generality $h_M^* = 0$) that for every $\underline{k}, \underline{k}' \in (\frac{2\pi}{L}\mathbb{Z}^2)^q$,

$$|\hat{W}_{q,q;0,0;0;0}^{(u.v.;N)}((\underline{k}, \underline{s}), (\underline{k}', \underline{s}')) - \hat{W}_{q,q;0,0;0;0}^{(u.v.;N')}((\underline{k}, \underline{s}), (\underline{k}', \underline{s}'))| \leq C^{1+q} \lambda^{\max\{1, \frac{2}{5}q\}} 2^{-\theta \min\{N, N'\}}, \quad (\text{F.2})$$

for any N, N' large enough, with a suitable $C > 0$ and $\theta \in (0, 1)$. Notice that the r.h.s of Eq. (F.2) is the same as the r.h.s. of Eq. (1.37) times a small factor $2^{-\theta \min\{N, N'\}}$.

Since any kernel in Fourier space is bounded in absolute value by the L^1 norm of its position-space counterpart (cf. Eq. (1.31)), and since for a non-perturbative multiscale analysis we are induced to work in position space, a natural way to derive Eq. (F.2) is by proving that the kernels $W^{(u.v.;N)}$ are *Cauchy sequences* in L^1 norm.

Comparing different lattices: smearing. The first issue is that, in presence of a lattice regularization, the kernels $W^{(u.v.;N)}$ and $W^{(u.v.;N')}$ are apriori defined on different lattices whenever, without loss of generality, $N < N'$. In order to give a meaning to $W^{(u.v.;N)} - W^{(u.v.;N')}$, we introduce a mollifier \mathfrak{d}_N given by

$$\mathfrak{d}_N(x) := \frac{1}{L^2} \int_{\Lambda_\infty^*} dk e^{-ik \cdot x} \chi(\frac{\gamma^{-N}}{2} |k|), \quad x \in \Lambda_\infty := \mathbb{R}^2 / (L\mathbb{Z}^2),$$

with $\Lambda_\infty^* \equiv \frac{2\pi}{L}\mathbb{Z}^2$, χ defined after Eq. (1.33) and $1 < \gamma \leq 2$ (e.g. $\gamma = \sqrt{2}$), and we define a smeared version of the kernels, denoted by $W^{(u.v.;N)}$, obtained via convolution with \mathfrak{d}_N , namely:

$$\begin{cases} W^{(u.v.;N)}(x_1, \dots, x_q) := \int_{\Lambda_N^q} dy_1 \dots dy_q W^{(u.v.;N)}(y_1, \dots, y_q) \prod_{j=1}^q \mathfrak{d}_N(x_j - y_j) \\ W^{(u.v.;N')}(x_1, \dots, x_q) := \int_{\Lambda_{N'}^q} dy_1 \dots dy_q W^{(u.v.;N')}(y_1, \dots, y_q) \prod_{j=1}^q \mathfrak{d}_N(x_j - y_j). \end{cases}$$

Note that Eq. (F.2) follows for $\max_{j=1}^q \{|k_j|, |k'_j|\} \leq \gamma^N$,²¹ if we show that

$$\|W_{q,q;0,0;0;0}^{(u.v.);N} - W_{q,q;0,0;0;0}^{(u.v.);N'}\|_{L^1(\Lambda_\infty)} \leq C^{1+q} \lambda^{\max\{1, \frac{2}{5}q\}} 2^{-\theta N}, \quad (\text{F.3})$$

where $\|\cdot\|_{L^1(\Lambda_\infty)}$ stands for Eq. (1.31) on Λ_∞ .

We stress that, since the smearing procedure cuts off all the modes $|k| \geq 2\gamma^N$ ($\chi = 0$ on $[1, \infty)$), the nested use of the smearing by \mathfrak{d}_N within the multiscale scheme would implicitly exclude all the contributions to correlation functions coming from modes at least greater than $2\gamma^N$, and this would be equivalent to saying that the lattice theory is well approximated by its continuum counterpart with a momentum cut-off. The latter property turns out to be actually incorrect, as one readily realizes that there are three irreducible diagrams, $\hat{\Pi}_{\mu,\nu}^{(0,N)}(p)$, $\hat{\Pi}_{5;\mu,\nu}^{(0,N)}(p)$ and $\mathcal{T}^{(0,N)}$, defined in Eqs. (C.1) and (5.8), which admit non-negligible contributions from momenta of size $\sim 2^N$. In fact it is possible to show that

$$\lim_{N \rightarrow \infty} \mathcal{T}^{(0,N)} = - \int_{(-\pi, \pi]^2} \frac{d^2 q}{(2\pi)^2} e^{-iq \cdot q} \frac{i\tilde{s}_0(q) - r\tilde{M}(q)}{|\tilde{s}(q)|^2 + \tilde{M}(q)^2}, \quad (\text{F.4})$$

where $\tilde{M}(q) \equiv 2r \sum_\mu \sin^2(\frac{q_\mu}{2})$ and $\tilde{s}_\mu(q) \equiv \sin(q_\mu)$. In the integral in the r.h.s. of Eq. (F.4), q has the interpretation of ‘‘rescaled momentum’’, namely $q = 2^{-N}k \in 2^{-N}\Lambda_\infty^*$: this shows that in the limit $N \rightarrow \infty$ all the contribution comes from momenta $k \in \Lambda_\infty^*$ such that $|k| \geq 2^{N(1-\epsilon)}$, for any $\epsilon > 0$. Similar considerations apply to the *bubble graphs* $\hat{\Pi}_{\mu,\nu}^{(0,N)}(0)$ and $\hat{\Pi}_{5;\mu,\nu}^{(0,N)}(0)$.

²¹Recall indeed that the cut-off χ equals 1 on $[0, \frac{1}{2}]$, hence the convolution by \mathfrak{d}_N does not affect the Fourier transform for momenta smaller than γ^N .

The Cauchy property. In virtue of the previous discussion one realizes that in order to prove Eq. (F.3), a refinement of Theorem 3.1 must be taken into account. In particular (see last paragraph for more details), it turns out that each non-irrelevant kernel can be decomposed as

$$W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} = w_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} + \mathcal{R}W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q},$$

where $w_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ is explicit, expressed in terms of the three diagrams above Eq. (F.4), while $\mathcal{R}W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ admits a bound with an extra improvement which is at least 2^{-h} smaller than the naive, dimensional one, $2^{hD_{sc}}$ (cf. Eq. (2.12)).²²

With this decomposition, fix e.g. $\gamma = \sqrt{2}$ and assume for simplicity N, N' to be multiples of 4, with $N' > N$; we can divide the range of scales into two different regimes as follows.

1. $N/4 \leq h \leq N$. In this regime, all the information about the theory is carried by the dominant part $w_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ of the marginal kernels, since the terms coming from $\mathcal{R}W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ contribute with an extra factor $2^{-h\vartheta} \leq 2^{-\frac{N\vartheta}{4}}$. Then in order to obtain Eq. (F.3), one uses the splitting of $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ and the fact that

$$\|w_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} - w_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N')|q,q}\|_{L^1(\Lambda_\infty)}^w = \mathcal{O}(2^{-N/4}), \quad (\text{F.5})$$

which follows from the explicit convergence of the graphs $\hat{\Pi}^{(h,N]}(0)$ and $\mathcal{T}^{(h,N]}$ to their continuum limit, namely

$$\begin{aligned} \hat{\Pi}^{(h,N]}(0) - \lim_{N \rightarrow \infty} \hat{\Pi}^{(h,N]}(0) &= \mathcal{O}(2^{-\frac{2}{3}N}) + \mathcal{O}(2^{h-N}), \\ \mathcal{T}^{(h,N]} - \lim_{N \rightarrow \infty} \mathcal{T}^{(h,N]} &= \mathcal{O}(2^{-\frac{2}{3}N}) + \mathcal{O}(2^{h-N}). \end{aligned}$$

2. $0 \leq h \leq N/4$. This regime is analyzed via the same inductive structure adopted in Sections 2, 3 for the proof of Theorem 3.1, with the difference that now the bounds in Eqs. (2.18)-(2.21) and Eqs. (2.22)-(2.24) involve, in their l.h.s., the differences $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} - W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N')|q,q}$ rather than the kernels $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q}$ themselves, and in the r.h.s. an extra smallness factor, say $2^{-N/8}$, is also present. The strategy is again based on a two-fold procedure.

As a first step one proves the counterpart of Proposition 2.1, establishing the smallness of the difference $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} - W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N')|q,q}$ for all the kernels, assuming it true for the relevant and marginal ones. This task is carried out by exploiting the tools of the tree expansion (cf. Section 2.2), combined with the dimensional bounds for the difference of fermionic propagators:

$$\left(\mathfrak{d}_N * g_N^{(h)} * \mathfrak{d}_N \right)(x, y) - \left(\mathfrak{d}_N * g_{N'}^{(h)} * \mathfrak{d}_N \right)(x, y) = \mathcal{O}\left(2^{2h-N} e^{-\kappa_4 \sqrt{2^h |x-y|_L}}\right) \quad (\text{F.6})$$

for a suitable $\kappa_4 > 0$. Note that the r.h.s. of Eq. (F.6) has a gain factor 2^{h-N} w.r.t. the dimensional bound for $g_N^{(h)}$ itself (cf. Eq. (2.4)), and for, say, $h \leq \frac{N}{2}$, such gain is better than $2^{-\frac{N}{2}}$.

Then, as a second step, we are left with proving the smallness of $W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N)|q,q} - W_{\underline{n};\dot{\underline{n}};\ddot{\underline{n}}}^{(h;N')|q,q}$ for the non-irrelevant kernels, and this can be done in the same spirit as Section 3, starting

²²For instance, in the first nontrivial case $\|\mathcal{R}W_{\emptyset;\emptyset;\emptyset}^{(h;N)|2,2}\|_1^w \leq R\lambda^{\frac{5}{4}} 2^{-h}$, which must be compared with Eq. (2.18)).

from the exact identities among the kernels (see e.g. Lemma 3.2), and exploiting inductively the bounds from the tree expansion, together with the bound for the difference of boson propagators:

$$\begin{aligned} & (\mathfrak{d}_N * g_N^A * \mathfrak{d}_N)(x, y) - (\mathfrak{d}_N * g_{N'}^A * \mathfrak{d}_N)(x, y) = \\ & \mathcal{O}\left(2^{-N} e^{-\kappa_4 \sqrt{|x-y|_L}}\right) + \mathcal{O}\left(\frac{2N'-N}{2} e^{-\kappa_4 \sqrt{2^{-\frac{N}{2}} |x-y|_L}}\right), \end{aligned}$$

and of fermion propagators (Eq. (F.6)).

Extracting the leading terms: refinement of Theorem 3.1. As explained above, in order for the previous strategy above to work, one needs to isolate the contribution from the diagrams $\hat{\Pi}_{\mu,\nu}^{(h,N)}$, $\hat{\Pi}_{5,\mu,\nu}^{(h,N)}$ and $\mathcal{T}^{(h,N)}$ in the multiscale analysis, and analyze their continuum limit separately. Such graphs typically appear in the kernels $W^{(h)}$ in the form of a geometrical series (corresponding to the sum of all the reducible diagrams), whose sum gives raise to

$$\Upsilon^{(h,N]}(b, b') := \delta_{\mu,\nu} \epsilon' \int_{\Lambda_N^*} dk \frac{e^{-ik \cdot (x-y)}}{|\sigma(k)|^2 + M^2 - 2\lambda(2\hat{\Pi}^{(h,N]}(0) + \mathcal{T}^{(h,N]})}, \quad (\text{F.7})$$

with the understanding that $b = (x, \mu, \epsilon)$, $b' = (y, \nu, \epsilon')$ and $\hat{\Pi}^{(h,N]}(0) \equiv \hat{\Pi}_{0,0}^{(h,N]}(0)$ (recall that by Lemma C.1, $\hat{\Pi}_{\mu,\nu}^{(h,N)}(0) = \delta_{\mu,\nu} \hat{\Pi}_{0,0}^{(h,N)}(0)$).²³ Let us show how to establish the dimensional improvement for a restricted class of kernels, namely $W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}$. It turns out that the dominant part of these kernels is given by

$$w_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0} = (nn')^2 \left(\lambda \mathcal{T}^{(h,N]} g^A \right)^{*n-1} * (1 + \lambda \Omega^{(h,N]}) * \Pi^{(h,N]} * \left(\lambda \mathcal{T}^{(h,N]} g^A \right)^{*n'-1}, \quad (\text{F.8})$$

so that

$$\|g^A * \mathcal{R}W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0}\|_1^w \leq R\lambda 2^{-h}; \quad \|\mathcal{R}W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}\|_1^w \leq \lambda^{\frac{n+n'-2}{2}} 2^{-h} \quad \forall n + n' \geq 3. \quad (\text{F.9})$$

The strategy for showing the dimensional improvement for $\mathcal{R}W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}$ is qualitatively the same as that discussed in Sections 2, 3. However, while the part related to Proposition 2.1 is proved almost identically, the inductive part (the part of Section 3) is way more involved, as we are going to show.

As a first step, using the same expansions introduced in Section 3 (see Fig. 10 for the kernel $W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0}$), one realizes that most of the terms have the right dimensional factor 2^{-h} as in the r.h.s of Eq. (F.9). Regarding the kernel $W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0}$, the only contributions without such dimensional improvement are the non-interacting bubble $\Pi^{(h,N]}$ and graph (c') of Fig. 10. Then one proceeds by isolating all the terms of the expansion which do not possess the dimensional improvement 2^{-h} ; in the case of the kernels $W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}$ we have:

$$W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} = \Pi^{(h,N]} + 2 \sum_{n \geq 1} \frac{1}{n^2} \lambda \Pi^{(h,N]} * g^A * W_{(n,1);\emptyset;\emptyset}^{(h;N)|0,0} + \mathcal{O}(\lambda 2^{-h}), \quad (\text{F.10})$$

$$W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0} = \frac{n^2}{(n-1)^2} \lambda \mathcal{T}^{(h,N]} g^A * W_{(n-1,n');\emptyset;\emptyset}^{(h;N)|0,0} + \mathcal{O}(\lambda^{\frac{n+n'-1}{2}} 2^{-h}), \quad n \geq 2, \quad (\text{F.11})$$

²³As established by Eqs. (5.14) and (5.15), if $h = 0$ the quantity $-2\lambda(2\hat{\Pi}^{(h,N]}(0) + \mathcal{T}^{(h,N]})$ appearing in the r.h.s. of Eq. (F.9) is exactly the lowest order contribution to the correlation function $\hat{\Sigma}_{\mu,\nu}^{(u,v)}(0)$.

where $\mathcal{O}(\cdot)$ is understood w.r.t. the weighted L^1 norm (cf. Eq. (1.31)). Eqs. (F.10), (F.11) can be regarded as a system of linear equations in the variables $\{W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}\}_{n \geq n' \geq 1}$. First, by an iterative use of Eq. (F.11), we find that

$$\begin{aligned} W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0} = \\ (nn')^2 (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1} * W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} * (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1} + \mathcal{O}(\lambda^{\frac{1}{2}} \lambda^{\frac{n+n'-2}{2}} 2^{-h}). \end{aligned} \quad (\text{F.12})$$

Now, combining Eq. (F.10) with Eq. (F.12), and letting $\Gamma^{(h,N]} := \sum_{n \geq 0} (\lambda \mathcal{T}^{(h,N]} g^A)^{*n}$, we find that

$$W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} = \Pi^{(h,N]} + 2\lambda \Pi^{(h,N]} * g^A * \Gamma^{(h,N]} * W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} + \mathcal{O}(\lambda 2^{-h}). \quad (\text{F.13})$$

Moreover, by convoluting with g^A both sides of Eq. (F.13), writing $g^A * \Pi^{(h,N]}$ as $\hat{\Pi}^{(h,N]}(0)g^A + (g^A * \Pi^{(h,N]} - \hat{\Pi}^{(h,N]}(0)g^A)$ and applying Item 3 of Lemma C.1, we further obtain:

$$g^A * (W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} - \Pi^{(h,N]} - 2\lambda \hat{\Pi}^{(h,N]}(0)g^A * \Gamma^{(h,N]} * W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0}) = \mathcal{O}(\lambda 2^{-h}). \quad (\text{F.14})$$

Finally, using that $\Omega^{(h,N]} = (\mathbb{1} - 2\lambda \hat{\Pi}^{(h,N]}(0)g^A * \Gamma^{(h,N]})^{-1}$, as well as $\Omega^{(h,N]} * g^A = g^A * \Omega^{(h,N]}$, and $\|\Omega^{(h,N]}\|_1^w = \mathcal{O}(1)$, we get the first of Eq. (F.9):

$$\left\| g^A * (W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} - \Omega^{(h,N]} * \Pi^{(h,N]}) \right\|_1^w \equiv \|g^A * \mathcal{R}W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0}\|_1^w \leq R\lambda 2^{-h}, \quad (\text{F.15})$$

for some R large enough. Besides, combining Eqs. (F.12), (F.15), one obtains the second of Eq. (F.9) concerning the kernels $W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}$ with $n + n' \geq 3$:

$$\begin{aligned} \|\mathcal{R}W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0}\|_1^w \equiv \\ \|W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0} - (nn')^2 (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1} * W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} * (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1}\|_1^w \leq \\ \|W_{(n,n');\emptyset;\emptyset}^{(h;N)|0,0} - (nn')^2 (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1} * W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} * (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1}\|_1^w + \\ + (nn')^2 \|(\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1} * \mathcal{R}W_{(1,1);\emptyset;\emptyset}^{(h;N)|0,0} * (\lambda \mathcal{T}^{(h,N]} g^A)^{*n-1}\|_1^w \leq \lambda^{\frac{n+n'-2}{2}} 2^{-h}, \end{aligned}$$

for λ small enough.

Acknowledgments. We thank A. Giuliani and M. Porta for useful comments, advices and discussions. We gratefully acknowledge financial support from the MUR, PRIN 2022 project MaIQuFi cod. 20223J85K3. This work has been carried out under the auspices of the GNFM of INdAM.

References

- [1] S. L. Adler and W. A. Bardeen, “Absence of higher-order corrections in the anomalous axial-vector divergence equation,” *Phys. Rev.*, vol. 182, pp. 1517–1536, Jun 1969.
- [2] J. Dimock and T. R. Hurd, “A renormalization group analysis of infrared qed,” *Journal of Mathematical Physics*, vol. 33, pp. 814–821, 02 1992.
- [3] V. Mastropietro, “Anomaly cancellation in the lattice effective electroweak theory,” *Journal of Mathematical Physics*, vol. 64, p. 032303, 03 2023.
- [4] V. Mastropietro, “Nonperturbative renormalization of the lattice sommerfield vector model,” *Physical Review D*, 2022.
- [5] G. Benfatto and V. Mastropietro, “Ward identities and chiral anomaly in the luttinger liquid,” *Communications in Mathematical Physics*, vol. 258, pp. 609–655, 06 2005.
- [6] G. Benfatto, P. Falco, and V. Mastropietro, “Functional integral construction of the massive thirring model: Verification of axioms and massless limit,” *Communications in Mathematical Physics*, vol. 258, pp. 67–118, 04 2007.
- [7] G. Benfatto, P. Falco, and V. Mastropietro, “Extended scaling relations for planar lattice models,” *Communications in Mathematical Physics*, 2009.
- [8] S. J. Summers, “A perspective on constructive quantum field theory,” *arXiv: Mathematical Physics*, 2012.
- [9] H. Georgi and J. M. Rawls, “Anomalies of the axial-vector current in two dimensions,” *Phys. Rev. D*, vol. 3, pp. 874–879, Feb 1971.
- [10] C. M. Sommerfield, “On the definition of currents and the action principle in field theories of one spatial dimension,” *Annals of Physics*, vol. 26, no. 1, pp. 1–43, 1964.
- [11] L. Brown, “Gauge invariance and mass in a two-dimensional model,” *Il Nuovo Cimento*, vol. 29, pp. 617—643, 1963.
- [12] C. R. Hagen, “Current definition and mass renormalization in a model field theory,” *Il Nuovo Cimento A*, 10 1967.
- [13] J. Schwinger, “Gauge invariance and mass. ii,” *Phys. Rev.*, vol. 128, pp. 2425–2429, Dec 1962.
- [14] K. Johnson, “Solution of the equations for the Green’s functions of a two-dimensional relativistic field theory,” *Nuovo Cim.*, vol. 20, pp. 773–790, 1961.
- [15] A. Das and V. S. Mathur, “Path-integral solubility of two-dimensional models,” *Phys. Rev. D*, vol. 33, pp. 489–495, Jan 1986.
- [16] J. Frohlich and E. Seiler, “The Massive Thirring-Schwinger Model (QED in Two-Dimensions): Convergence of Perturbation Theory and Particle Structure,” *Helv. Phys. Acta*, vol. 49, p. 889, 1976.
- [17] H. Georgi and Y. Kats, “Unparticle example in 2d,” *Phys. Rev. Lett.*, vol. 101, p. 131603, Sep 2008.

- [18] J. Kogut and L. Susskind, “Hamiltonian formulation of wilson’s lattice gauge theories,” *Phys. Rev. D*, vol. 11, pp. 395–408, Jan 1975.
- [19] F. Berruto, G. Grignani, G. W. Semenoff, and P. Sodano, “Chiral symmetry breaking on the lattice: A study of the strongly coupled lattice schwinger model,” *Phys. Rev. D*, vol. 57, pp. 5070–5083, Apr 1998.
- [20] R. Dempsey, I. R. Klebanov, S. S. Pufu, and B. Zan, “Discrete chiral symmetry and mass shift in the lattice hamiltonian approach to the schwinger model,” *Phys. Rev. Res.*, vol. 4, p. 043133, Nov 2022.
- [21] R. Dempsey, I. R. Klebanov, S. S. Pufu, B. T. Swgaard, and B. Zan, “Phase diagram of the two-flavor schwinger model at zero temperature,” *Phys. Rev. Lett.*, vol. 132, p. 031603, Jan 2024.
- [22] A. Chatterjee, S. D. Pace, and S.-H. Shao, “Quantized axial charge of staggered fermions and the chiral anomaly,” *Phys. Rev. Lett.*, vol. 134, p. 021601, Jan 2025.
- [23] N. Seiberg and S.-H. Shao, “Majorana chain and Ising model - (non-invertible) translations, anomalies, and emanant symmetries,” *SciPost Phys.*, vol. 16, p. 064, 2024.
- [24] V. Mastropietro, “Nonperturbative adler-bardeen theorem,” *J. Math. Phys.*, 2007.
- [25] P. Falco, “Vector and axial anomaly in the thirring-wess model,” *J. Math. Phys.*, 2010.
- [26] V. Mastropietro and M. Porta, “Multi-channel luttinger liquids at the edge of quantum hall systems,” *Communications in Mathematical Physics*, vol. 395, pp. 1097–1173, 09 2022.
- [27] A. Giuliani, V. Mastropietro, and F. Toninelli, “Non-integrable dimers: Universal fluctuations of tilted height profiles,” *Communications in Mathematical Physics*, vol. 377, pp. 1883–1959, 06 2020.
- [28] A. Lesniewski, “Effective action for the yukawa2 quantum field theory,” *Communications in Mathematical Physics*, 1987.
- [29] G. Gentile and V. Mastropietro, “Renormalization group for one-dimensional fermions. a review on mathematical results,” *Physics Reports*, 2001.
- [30] D. Brydges, “A short course on cluster expansions,” *Les Houches summer school*, 1984.
- [31] A. Giuliani and V. Mastropietro, “The two-dimensional hubbard model on the honeycomb lattice,” *Communications in Mathematical Physics*, 2010.
- [32] A. Giuliani, V. Mastropietro, and S. Rychkov, “Gentle introduction to rigorous renormalization group: a worked fermionic example,” *Journal of High Energy Physics*, 2021.