

AN ERDŐS–TROTTER PROBLEM ON ANTICHAINS WITH MULTIPLICITY r ON EACH OCCURRING LEVEL

YIXIN HE AND QUANYU TANG

ABSTRACT. Fix an integer $r \geq 2$. For each n we consider families $\mathcal{F} \subseteq 2^{[n]}$ that form an antichain and have the property that, for every t , if there exists $A \in \mathcal{F}$ with $|A| = t$ then there exist at least r members of \mathcal{F} of size t . A problem of Erdős and Trotter asserts that, for each fixed r , there exists a threshold $n_0(r)$ such that whenever $n > n_0(r)$ one can achieve $n - 3$ distinct set sizes in such a family, and asks for estimates on $n_0(r)$. We compute that $n_0(2) = 3$ and $n_0(3) = 8$. For all $r \geq 4$ we prove matching linear bounds up to lower-order terms, namely

$$2r + 2 \leq n_0(r) \leq 2r + 2 \log_2 r + O(\log_2 \log_2 r).$$

In particular, $n_0(r) = 2r + o(r)$.

1. INTRODUCTION

The study of antichains in the Boolean lattice $2^{[n]}$ (ordered by inclusion) is a cornerstone of extremal set theory, dating back to Sperner’s theorem [7], which determines the maximum possible size of an antichain. Since then, antichains and their generalizations have been extensively investigated; see, e.g., the monographs of Anderson [2] and Engel [4]. While classical results such as the LYM inequality focus on bounding the cardinality of a family, Paul Erdős often emphasized “profile” questions, asking which sets of cardinalities an antichain can realize under additional constraints.

In [5, Eq. (10.28)] (see also [6, p. 119]), Erdős and Trotter proposed the following problem, which also appears as Problem #776 on Bloom’s Erdős Problems website [3].

Problem 1.1 ([5, 6]). *Let S be a set with $|S| = n$. Consider a family (A_k) of subsets of S where no A_k contains any other and such that, for every t , if there is an A_k with $|A_k| = t$ then there are exactly r different A ’s of size t . If $r = 1$ and $n > 3$, one can always give $n - 2$ sets A_k and one cannot give $n - 1$. If $r > 1$ and $n >$ some number $n_0 = n_0(r)$, one can always give $r(n - 3)$ sets A_k but one cannot give $r(n - 2)$ such sets.*

Give estimates for $n_0(r)$.

In [6], Guy writes:

[...] We have no satisfactory estimate of $n_0(r)$. The content of the previous two paragraphs will be a small subset of a forthcoming paper of Erdős, Szemerédi and Trotter.

To the best of our knowledge, we have not been able to locate the announced paper of Erdős, Szemerédi and Trotter, nor have we found any subsequent work in the literature providing a satisfactory estimate for $n_0(r)$.

In this paper we obtain explicit upper and lower bounds for $n_0(r)$.

2020 *Mathematics Subject Classification.* 05D05, 06A07.

Key words and phrases. Erdős problem, extremal set theory, antichain.

1.1. A reformulation of Problem 1.1. Fix integers $n \geq 1$ and $r \geq 1$. For a family $\mathcal{F} \subseteq 2^{[n]}$ write

$$S(\mathcal{F}) := \{|A| : A \in \mathcal{F}\}, \quad \mathcal{F}_t := \{A \in \mathcal{F} : |A| = t\} \quad (t = 0, 1, \dots, n).$$

Call \mathcal{F} an r -multiplicity antichain if it satisfies:

- (i) (*Antichain*) $A \not\subseteq B$ for all distinct $A, B \in \mathcal{F}$.
- (ii) (r -multiplicity) For every $t \in S(\mathcal{F})$ one has $|\mathcal{F}_t| \geq r$.

Following Guy's statement [6], the underlying extremal quantity is

$$g(n, r) := \max\{|S(\mathcal{F})| : \mathcal{F} \subseteq 2^{[n]} \text{ is an } r\text{-multiplicity antichain}\},$$

i.e. the largest number of distinct cardinalities that can occur in an r -multiplicity antichain on $[n]$. Guy also reports that for $r > 1$ there exists a threshold $n_0(r)$ such that, for all sufficiently large n , one always has $g(n, r) = n - 3$ (equivalently, $n - 3$ distinct sizes are always achievable but $n - 2$ are not).

Remark 1.2. In the original formulation of Erdős–Trotter [5, Eq. (10.28)], one requires that for every occurring level $t \in S(\mathcal{F})$ there are *exactly* r sets of size t , i.e. $|\mathcal{F}_t| = r$. Guy [6] relaxes this to $|\mathcal{F}_t| \geq r$. For the extremal quantity $g(n, r)$ studied in this paper, these two conventions are equivalent. Indeed, given any antichain \mathcal{F} with $|\mathcal{F}_t| \geq r$ for all $t \in S(\mathcal{F})$, select an arbitrary subfamily $\mathcal{F}'_t \subseteq \mathcal{F}_t$ of size exactly r for each such t and set $\mathcal{F}' := \bigcup_{t \in S(\mathcal{F})} \mathcal{F}'_t$. Then \mathcal{F}' is still an antichain, satisfies $|\mathcal{F}'_t| = r$ for all $t \in S(\mathcal{F}')$, and preserves the set of occurring sizes: $S(\mathcal{F}') = S(\mathcal{F})$. Hence the maximum possible value of $|S(\mathcal{F})|$ is the same under either convention, and we adopt Guy's inequality formulation for convenience.

Accordingly, the threshold $n_0(r)$ appearing in Problem 1.1 can be equivalently defined as follows.

Definition 1.3. Fix an integer $r \geq 2$. We define $n_0(r)$ to be the smallest integer n_0 such that

$$g(n, r) = n - 3$$

for every integer $n > n_0$.

For small values of r one can determine $n_0(r)$ exactly by computation. In particular, $n_0(2) = 3$ and $n_0(3) = 8$; see Appendix A for details.

We now state our main results.

Theorem 1.4. *For every integer $r \geq 4$, one has*

$$n_0(r) \geq 2r + 2.$$

Theorem 1.5. *For every integer $r \geq 2$, one has*

$$n_0(r) \leq 2r + 2 \log_2 r + O(\log_2 \log_2 r).$$

1.2. Paper organization. The remainder of the paper is organized as follows. In Section 3 we prove Theorem 1.4, which provides a lower bound for $n_0(r)$. In Section 4 we prove Theorem 1.5, which gives an upper bound for $n_0(r)$. Section 5 collects further remarks and open problems. Finally, in Appendix A we determine the exact values $n_0(2)$ and $n_0(3)$.

2. PRELIMINARIES

We collect several auxiliary statements that will be used in the proofs of our main theorems.

Lemma 2.1. *For every integer $m \geq 1$,*

$$\binom{m}{\lfloor m/2 \rfloor} \geq \frac{2^m}{2\sqrt{m}}.$$

Proof. This follows from the standard estimate for the central binomial coefficient derived from Stirling's formula. We omit the details. \square

Corollary 2.2. *Let $K \geq 4$ be an integer and let m be the least integer such that*

$$\binom{m}{\lfloor m/2 \rfloor} \geq K.$$

Then

$$m \leq \left\lceil \log_2 K + \frac{1}{2} \log_2 \log_2 K + 2 \right\rceil. \quad (2.1)$$

Proof. Set $m_0 := \left\lceil \log_2 K + \frac{1}{2} \log_2 \log_2 K + 2 \right\rceil$. By Lemma 2.1,

$$\binom{m_0}{\lfloor m_0/2 \rfloor} \geq \frac{2^{m_0}}{2\sqrt{m_0}}.$$

Moreover $2^{m_0} \geq 2^{\log_2 K + \frac{1}{2} \log_2 \log_2 K + 2} = 4K \sqrt{\log_2 K}$, so

$$\frac{2^{m_0}}{2\sqrt{m_0}} \geq 2K \sqrt{\frac{\log_2 K}{m_0}}.$$

If $K \geq 16$ then $m_0 \leq 2 \log_2 K$, so the right-hand side is at least $\sqrt{2} K \geq K$. For $4 \leq K \leq 15$ the bound (2.1) is verified by a direct check. Thus $\binom{m_0}{\lfloor m_0/2 \rfloor} \geq K$, and by minimality of m we have $m \leq m_0$, proving (2.1). \square

We will use the following elementary classification of pairwise-intersecting families of 2-subsets.

Lemma 2.3. *Let X be a finite set and let $\mathcal{E} \subseteq \binom{X}{2}$ be a family of 2-subsets such that any two members of \mathcal{E} intersect. Then either*

- (i) \mathcal{E} is a star, i.e. there exists $a \in X$ such that $a \in E$ for every $E \in \mathcal{E}$, or
- (ii) \mathcal{E} is a triangle, i.e. there exist distinct $a, b, c \in X$ such that $\mathcal{E} = \{\{a, b\}, \{a, c\}, \{b, c\}\}$.

Proof. If $|\mathcal{E}| \leq 2$ then (i) holds trivially. Assume $|\mathcal{E}| \geq 3$ and pick two distinct edges $\{a, b\}, \{a, c\} \in \mathcal{E}$. If every edge in \mathcal{E} contains a , then (i) holds and we are done. Otherwise, there exists $E \in \mathcal{E}$ with $a \notin E$. Since E intersects both $\{a, b\}$ and $\{a, c\}$, we must have $b, c \in E$, so $E = \{b, c\}$. Now let $F \in \mathcal{E}$ be arbitrary. Then F intersects each of $\{a, b\}, \{a, c\}, \{b, c\}$. If $a \in F$, then F must contain b or c , hence $F \in \{\{a, b\}, \{a, c\}\}$. If $a \notin F$, then F must intersect both $\{a, b\}$ and $\{a, c\}$, so again $F = \{b, c\}$. Therefore $\mathcal{E} = \{\{a, b\}, \{a, c\}, \{b, c\}\}$ and (ii) holds. \square

We will also use the following simple ‘‘label’’ antichain gadget.

Lemma 2.4. *Let $m \geq 4$ and set $\ell := \lfloor m/2 \rfloor$. Let V_0 be a set of size m and let b be an element not in V_0 . Put $V := V_0 \cup \{b\}$. Let K be an integer such that*

$$K \geq \ell + 1 \quad \text{and} \quad K - (\ell + 1) \leq \binom{m}{\ell}. \quad (2.2)$$

Then there exist sets $L_t \subseteq V$ for all integers $t \in \{4, 5, \dots, K\}$ such that

- (i) $|L_t| = 2$ for every $t \in \{4, 5, \dots, \ell + 1\}$, and these L_t are pairwise distinct and all contain b ;
- (ii) $|L_t| = \ell$ for every $t \in \{\ell + 2, \ell + 3, \dots, K\}$, and these L_t are pairwise distinct and all lie in V_0 ;
- (iii) the family $\{L_t : 4 \leq t \leq K\}$ is an antichain under inclusion.

Proof. Since $\ell = \lfloor m/2 \rfloor$, we have $\ell - 2 \leq m$. Choose distinct $v_1, \dots, v_{\ell-2} \in V_0$ (possible because $m \geq 4$ implies $\ell \geq 2$), and define

$$L_t := \{b, v_{t-3}\} \quad (4 \leq t \leq \ell + 1).$$

For $t \geq \ell + 2$, we need $K - (\ell + 1)$ many ℓ -subsets of V_0 . By (2.2), we may choose distinct

$$L_t \in \binom{V_0}{\ell} \quad (\ell + 2 \leq t \leq K).$$

Finally, inclusion cannot occur between distinct L_t : within each of the two subfamilies all sets have the same size, so inclusion forces equality; and no L_t with $t \leq \ell + 1$ can be contained in an L_s with $s \geq \ell + 2$ because the former contains b while the latter does not. This proves (iii). \square

Finally, we record a universal obstruction: for $r \geq 2$ one cannot realize $n - 2$ distinct levels. This fact is taken for granted in the statement of Problem 1.1; we include a complete proof here for completeness.

Lemma 2.5. *Let $r \geq 2$ and $n \geq 4$. Then every r -multiplicity antichain $\mathcal{F} \subseteq 2^{[n]}$ satisfies $|S(\mathcal{F})| \leq n - 3$. Equivalently, for all $r \geq 2$ and $n \geq 4$,*

$$g(n, r) \leq n - 3.$$

Proof. Fix $r \geq 2$ and $n \geq 4$, and let $\mathcal{F} \subseteq 2^{[n]}$ be an r -multiplicity antichain.

Since $r \geq 2$, the levels $t = 0$ and $t = n$ cannot occur: indeed $|\mathcal{F}_0| \leq 1$ and $|\mathcal{F}_n| \leq 1$, contradicting the r -multiplicity condition whenever 0 or n lies in $S(\mathcal{F})$. Hence

$$S(\mathcal{F}) \subseteq \{1, 2, \dots, n - 1\}.$$

Claim 1. *If $1 \in S(\mathcal{F})$ then $n - 1 \notin S(\mathcal{F})$.*

Proof of Claim 1. Indeed, since $1 \in S(\mathcal{F})$ we have $|\mathcal{F}_1| \geq r \geq 2$, so choose two distinct singletons $\{x\}, \{y\} \in \mathcal{F}_1$. Any $(n - 1)$ -subset of $[n]$ misses exactly one element, hence contains at least one of x, y . Therefore every $(n - 1)$ -set H satisfies $\{x\} \subseteq H$ or $\{y\} \subseteq H$, contradicting the antichain property. So $\mathcal{F}_{n-1} = \emptyset$, i.e. $n - 1 \notin S(\mathcal{F})$. \square

Claim 2. *If $n - 1 \in S(\mathcal{F})$ then $1 \notin S(\mathcal{F})$.*

Proof of Claim 2. Indeed, if $n - 1 \in S(\mathcal{F})$ then $|\mathcal{F}_{n-1}| \geq r \geq 2$, so pick two distinct $(n - 1)$ -sets $H_1 = [n] \setminus \{x_1\}$ and $H_2 = [n] \setminus \{x_2\}$ with $x_1 \neq x_2$. For any singleton $\{z\}$, at least one of H_1, H_2 contains z (since z cannot equal both x_1 and x_2), so $\{z\} \subseteq H_i$ for some $i \in \{1, 2\}$, contradicting the antichain property. Hence $\mathcal{F}_1 = \emptyset$, i.e. $1 \notin S(\mathcal{F})$. \square

Suppose for contradiction that $|S(\mathcal{F})| \geq n - 2$. Then by Claims 1 and 2 necessarily $|S(\mathcal{F})| = n - 2$ and there exists a unique $t_0 \in \{1, \dots, n - 1\}$ such that

$$S(\mathcal{F}) = \{1, 2, \dots, n - 1\} \setminus \{t_0\}.$$

Now we split into cases.

Case 1: $1 \in S(\mathcal{F})$. By Claim 1 we have $n - 1 \notin S(\mathcal{F})$, so the unique missing level is $t_0 = n - 1$ and in particular $n - 2 \in S(\mathcal{F})$.

Choose r distinct singletons $\{x_1\}, \dots, \{x_r\} \in \mathcal{F}_1$ and set $X = \{x_1, \dots, x_r\}$. If $A \in \mathcal{F}$ has $|A| \geq 2$ then A cannot contain any x_i , otherwise $\{x_i\} \subseteq A$ contradicts the antichain property. Hence every $A \in \mathcal{F}$ with $|A| \geq 2$ satisfies $A \subseteq [n] \setminus X$, where $|[n] \setminus X| = n - r$. In particular, every $(n - 2)$ -set in \mathcal{F}_{n-2} must be an $(n - 2)$ -subset of an $(n - r)$ -set, so

$$|\mathcal{F}_{n-2}| \leq \binom{n - r}{n - 2}.$$

If $r \geq 3$ then $\binom{n-r}{n-2} = 0$, and if $r = 2$ then $\binom{n-r}{n-2} = \binom{n-2}{n-2} = 1$. In either case we get $|\mathcal{F}_{n-2}| < r$, contradicting $n-2 \in S(\mathcal{F})$ and r -multiplicity.

Case 2: $n-1 \in S(\mathcal{F})$. By Claim 2 we have $1 \notin S(\mathcal{F})$, so the unique missing level is $t_0 = 1$ and in particular $2 \in S(\mathcal{F})$.

Choose r distinct $(n-1)$ -sets $H_i \in \mathcal{F}_{n-1}$, say $H_i = [n] \setminus \{x_i\}$ with distinct x_1, \dots, x_r , and set $X = \{x_1, \dots, x_r\}$. Let $A \in \mathcal{F}$ with $|A| \leq n-2$. For each i , we must have $A \not\subseteq H_i$ by the antichain property. But if $x_i \notin A$ then $A \subseteq [n] \setminus \{x_i\} = H_i$, a contradiction. Hence $x_i \in A$ for every i , so $X \subseteq A$ and therefore $|A| \geq r$.

In particular, if $r \geq 3$ then $\mathcal{F}_2 = \emptyset$, contradicting $2 \in S(\mathcal{F})$. If $r = 2$ then every 2-set in \mathcal{F}_2 must equal $X = \{x_1, x_2\}$, so $|\mathcal{F}_2| \leq 1 < r$, again contradicting r -multiplicity. This finishes the contradiction.

Therefore $|S(\mathcal{F})| \leq n-3$ for every r -multiplicity antichain \mathcal{F} , i.e. $g(n, r) \leq n-3$. \square

3. LOWER BOUND

Proposition 3.1. *Let $r \geq 4$ and let n satisfy $r+3 \leq n \leq 2r+2$. Then every r -multiplicity antichain $\mathcal{F} \subseteq 2^{[n]}$ satisfies*

$$|S(\mathcal{F})| \leq n-4.$$

Proof. Assume for contradiction that there exists an r -multiplicity antichain $\mathcal{F} \subseteq 2^{[n]}$ with $|S(\mathcal{F})| \geq n-3$. By Lemma 2.5 every r -multiplicity antichain satisfies $|S(\mathcal{F})| \leq n-3$. Therefore \mathcal{F} must in fact satisfy

$$|S(\mathcal{F})| = n-3.$$

Since there are $n+1$ possible sizes in $\{0, 1, \dots, n\}$, the equality $|S(\mathcal{F})| = n-3$ means that exactly four sizes are missing.

Since $r \geq 4$, the levels 0 and n cannot occur in $S(\mathcal{F})$ (each contains only one set). If $1 \in S(\mathcal{F})$, then $|\mathcal{F}_1| \geq r$ and hence \mathcal{F}_1 contains r distinct singletons $\{x_1\}, \dots, \{x_r\}$. By the antichain property, no set in \mathcal{F} may contain any x_i ; thus every $A \in \mathcal{F}$ is contained in $[n] \setminus \{x_1, \dots, x_r\}$ and so $|A| \leq n-r$. In particular, none of the sizes $n-r+1, \dots, n$ occurs. Together with $0 \notin S(\mathcal{F})$, this forces at least $r+1 \geq 5$ missing sizes, contradicting that exactly four sizes are missing. Hence $1 \notin S(\mathcal{F})$.

Now consider the complement family $\mathcal{F}^c := \{[n] \setminus A : A \in \mathcal{F}\}$. It is again an r -multiplicity antichain, and $S(\mathcal{F}^c) = \{n-t : t \in S(\mathcal{F})\}$. Since $|S(\mathcal{F}^c)| = |S(\mathcal{F})| = n-3$, the same argument shows that $1 \notin S(\mathcal{F}^c)$. Thus $n-1 \notin S(\mathcal{F})$. Therefore the missing sizes are precisely $\{0, 1, n-1, n\}$ and hence

$$S(\mathcal{F}) = \{2, 3, \dots, n-2\}. \quad (3.1)$$

Let

$$\mathcal{M} := \mathcal{F}_{n-2}^c = \{[n] \setminus A : A \in \mathcal{F}_{n-2}\} \subseteq \binom{[n]}{2}.$$

Then $|\mathcal{F}_2| \geq r$ and $|\mathcal{M}| \geq r$ by r -multiplicity and (3.1). Moreover, for all $E \in \mathcal{F}_2$ and $M \in \mathcal{M}$ we have

$$E \cap M \neq \emptyset, \quad (3.2)$$

since otherwise $E \subseteq [n] \setminus M \in \mathcal{F}_{n-2}$ would contradict that \mathcal{F} is an antichain.

We claim that \mathcal{F}_2 is pairwise intersecting. Indeed, if \mathcal{F}_2 contained two disjoint edges $\{a, b\}$ and $\{c, d\}$, then any $M \in \mathcal{M}$ must intersect both, so necessarily

$$\mathcal{M} \subseteq \{\{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}\},$$

whence $|\mathcal{M}| \leq 4$. If $r > 4$ this contradicts $|\mathcal{M}| \geq r$. If $r = 4$, then $|\mathcal{M}| \geq 4$ forces equality, so $\mathcal{M} = \{\{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}\}$. But then the only 2-sets intersecting all members of \mathcal{M}

are $\{a, b\}$ and $\{c, d\}$, so $|\mathcal{F}_2| \leq 2$, contradicting $|\mathcal{F}_2| \geq r = 4$. This proves that \mathcal{F}_2 is pairwise intersecting.

We claim that \mathcal{F}_2 is a star, and all $(n-2)$ -sets avoid its centre. Indeed, by Lemma 2.3, a pairwise intersecting family of 2-sets is either a star or a triangle. Since $|\mathcal{F}_2| \geq r \geq 4$, it cannot be a triangle. Hence there exists a point $x \in [n]$ and a set $P \subseteq [n] \setminus \{x\}$ with $|P| = |\mathcal{F}_2| \geq r$ such that

$$\mathcal{F}_2 = \{\{x, p\} : p \in P\}. \quad (3.3)$$

Next we show that every $M \in \mathcal{M}$ contains x . Fix $M \in \mathcal{M}$. By (3.2) and (3.3), M must intersect $\{x, p\}$ for every $p \in P$. If $x \notin M$, then M is a 2-set and must contain every $p \in P$, impossible since $|P| \geq r \geq 4 > 2$. Therefore $x \in M$ for all $M \in \mathcal{M}$. Writing $M = \{x, q\}$, we obtain a set

$$Q := \{q \in [n] \setminus \{x\} : \{x, q\} \in \mathcal{M}\} \quad \text{with} \quad |Q| = |\mathcal{M}| \geq r,$$

and equivalently every $A \in \mathcal{F}_{n-2}$ has the form $A = [n] \setminus \{x, q\}$ for some $q \in Q$. In particular,

$$x \notin A \quad \text{for all } A \in \mathcal{F}_{n-2}.$$

Let $2 \leq t \leq r-1$ and take any $B \in \mathcal{F}_t$ (which exists by (3.1)). Suppose $x \notin B$. Since $|B| < |Q|$, we know that $Q \setminus B \neq \emptyset$. For any $q \in Q \setminus B$ we would have

$$B \subseteq [n] \setminus \{x, q\} \in \mathcal{F}_{n-2},$$

contradicting that \mathcal{F} is an antichain. This proves

$$x \in B \quad \text{for every } B \in \mathcal{F}_t \text{ and every } 2 \leq t \leq r-1. \quad (3.4)$$

Set

$$U := [n] \setminus (\{x\} \cup P).$$

Since $|P| \geq r$, we have

$$|U| = n - 1 - |P| \leq n - 1 - r \leq (2r + 2) - 1 - r = r + 1. \quad (3.5)$$

Moreover, any set $B \in \mathcal{F}$ with $x \in B$ and $|B| \geq 3$ cannot contain any $p \in P$, since otherwise $\{x, p\} \in \mathcal{F}_2$ would be contained in B , contradicting the antichain property. Thus every $B \in \mathcal{F}_t$ with $x \in B$ and $t \geq 3$ can be written as

$$B = \{x\} \cup B', \quad B' \subseteq U, \quad |B'| = t - 1.$$

Case 1: $|U| \leq r$. By (3.1) we have $r \in S(\mathcal{F})$, so $|\mathcal{F}_r| \geq r$.

We claim that there is at most one set $A \in \mathcal{F}_r$ with $x \notin A$. Indeed, let $A \in \mathcal{F}_r$ and suppose $x \notin A$. If there existed $q \in Q \setminus A$, then $A \subseteq [n] \setminus \{x, q\} \in \mathcal{F}_{n-2}$, contradicting that \mathcal{F} is an antichain. Hence $Q \subseteq A$, so $r = |A| \geq |Q|$. On the other hand, $|Q| = |\mathcal{M}| = |\mathcal{F}_{n-2}| \geq r$, and thus $|Q| = r$ and necessarily $A = Q$. This proves the claim.

Thus, we can choose $r-1$ distinct sets $A_1, \dots, A_{r-1} \in \mathcal{F}_r$ with $x \in A_i$. Write $A_i = \{x\} \cup X_i$ where $X_i \subseteq U$ and $|X_i| = r-1$. Since the A_i are distinct, the sets X_1, \dots, X_{r-1} are distinct. This forces $|U| \geq r$, and together with $|U| \leq r$ we get $|U| = r$.

Now $|\mathcal{F}_{r-1}| \geq r$ and by (3.4) every set in \mathcal{F}_{r-1} contains x , so pick r distinct sets $B_1, \dots, B_r \in \mathcal{F}_{r-1}$ and write $B_j = \{x\} \cup Y_j$ where $Y_j \subseteq U$ and $|Y_j| = r-2$. Since $|U| = r$, each Y_j has exactly two supersets of size $r-1$ in U . But U has exactly $\binom{r}{r-1} = r$ many $(r-1)$ -subsets, and the $r-1$ distinct sets X_1, \dots, X_{r-1} miss at most one of them; therefore, for each j at least one of the two $(r-1)$ -supersets of Y_j must equal X_i for some i . Hence for each j there exist i with $Y_j \subseteq X_i$, which implies

$$B_j = \{x\} \cup Y_j \subseteq \{x\} \cup X_i = A_i,$$

contradicting that \mathcal{F} is an antichain.

Case 2: $|U| = r + 1$. Then by (3.5) we know that $n = 2r + 2$. Since $r \geq 4$, we have $3 \leq r - 1$ and thus $\mathcal{F}_3 \neq \emptyset$ and every 3-set in \mathcal{F} contains x by (3.4). In particular, every $C \in \mathcal{F}_3$ has the form $C = \{x\} \cup e$ with $e \in \binom{U}{2}$. Define

$$\mathcal{H} := \left\{ e \in \binom{U}{2} : \{x\} \cup e \in \mathcal{F}_3 \right\}.$$

Then $|\mathcal{H}| = |\mathcal{F}_3| \geq r$.

Claim 3. *At most one set in \mathcal{F}_{r+1} contains x .*

Proof of Claim 3. Indeed, suppose $A \in \mathcal{F}_{r+1}$ with $x \in A$. Then $A = \{x\} \cup Z$ with $Z \subseteq U$ and $|Z| = r$. Since $|U| = r + 1$, we can write $Z = U \setminus \{u\}$ for a unique $u \in U$. If there exists $e \in \mathcal{H}$ with $e \subseteq Z$, then $\{x\} \cup e \subseteq A$, contradicting the antichain property. Thus for every $e \in \mathcal{H}$ we have $e \not\subseteq Z$, which is equivalent to $u \in e$. Hence $u \in \bigcap_{e \in \mathcal{H}} e$. But \mathcal{H} contains at least two distinct 2-sets (since $|\mathcal{H}| \geq r \geq 4$), so $\bigcap_{e \in \mathcal{H}} e$ has size at most 1. Therefore there is at most one possible u , and hence at most one such A . \square

By Claim 3 and $|\mathcal{F}_{r+1}| \geq r$, there are at least $r - 1$ distinct sets $\tilde{A}_1, \dots, \tilde{A}_{r-1} \in \mathcal{F}_{r+1}$ with $x \notin \tilde{A}_i$. Fix such an \tilde{A}_i . If there exists $q \in Q$ with $q \notin \tilde{A}_i$, then $\tilde{A}_i \subseteq [n] \setminus \{x, q\} \in \mathcal{F}_{n-2}$ and the inclusion is proper because $|\tilde{A}_i| = r + 1 < (2r + 2) - 2 = n - 2$, contradicting that \mathcal{F} is an antichain. Hence $Q \subseteq \tilde{A}_i$ for all i , and in particular $|Q| \leq r + 1$.

Since we have at least $r - 1$ distinct sets \tilde{A}_i of size $r + 1$ all containing Q , it follows that $|Q| \neq r + 1$. Therefore

$$|Q| = r. \tag{3.6}$$

Writing $R := [n] \setminus (\{x\} \cup Q)$, we have $|R| = n - 1 - |Q| = (2r + 2) - 1 - r = r + 1$ and each \tilde{A}_i is of the form

$$\tilde{A}_i = Q \cup \{a_i\} \quad (a_i \in R),$$

with a_1, \dots, a_{r-1} pairwise distinct.

Claim 4. *Every set in \mathcal{F}_{r+2} avoids x and contains Q .*

Proof of Claim 4. First, suppose $B \in \mathcal{F}_{r+2}$ contains x . Then $|B| \geq 3$ and hence $B = \{x\} \cup B'$ with $B' \subseteq U$ and $|B'| = r + 1$. Since $|U| = r + 1$, this forces $B' = U$ and hence $B = \{x\} \cup U$. But then every $C \in \mathcal{F}_3$ satisfies $C \subseteq \{x\} \cup U = B$, contradicting the antichain property. Thus $x \notin B$ for all $B \in \mathcal{F}_{r+2}$.

Now take any $B \in \mathcal{F}_{r+2}$. If there exists $q \in Q$ with $q \notin B$, then $B \subseteq [n] \setminus \{x, q\} \in \mathcal{F}_{n-2}$ and the inclusion is proper because $|B| = r + 2 < (2r + 2) - 2 = n - 2$, again contradicting that \mathcal{F} is an antichain. Hence $Q \subseteq B$ for all $B \in \mathcal{F}_{r+2}$, proving Claim 4. \square

By (3.6) and Claim 4, every $B \in \mathcal{F}_{r+2}$ has the form

$$B = Q \cup \{b, b'\} \quad (b, b' \in R, b \neq b').$$

Since $|\mathcal{F}_{r+2}| \geq r$, we can choose r distinct such pairs $\{b, b'\}$.

However, for each $i \in \{1, \dots, r - 1\}$ and each $B \in \mathcal{F}_{r+2}$ we must have $\tilde{A}_i \not\subseteq B$. Equivalently, none of the pairs $\{b, b'\}$ may contain any a_i . Thus every pair corresponding to a member of \mathcal{F}_{r+2} must lie in $R \setminus \{a_1, \dots, a_{r-1}\}$, which has size at most 2. Hence there is at most one possible pair, contradicting $|\mathcal{F}_{r+2}| \geq r \geq 4$.

This contradiction completes Case 2, and hence no r -multiplicity antichain on $[n]$ can satisfy $|S(\mathcal{F})| = n - 3$ when $r + 3 \leq n \leq 2r + 2$. Therefore $|S(\mathcal{F})| \leq n - 4$ for all such n . \square

We are now ready to present the following.

Proof of Theorem 1.4. By Proposition 3.1, in particular at $n = 2r + 2$ we have $g(2r + 2, r) \leq (2r + 2) - 4 < (2r + 2) - 3$. By Definition 1.3, this forces $n_0(r) \geq 2r + 2$. \square

4. UPPER BOUND

Proposition 4.1. *For every integer $r \geq 2$ and every integer n satisfying*

$$n \geq 2r + 2 \log_2 r + \log_2 \log_2 r + 15, \quad (4.1)$$

one has $g(n, r) = n - 3$.

Proof. Fix $r \geq 2$ and an integer n satisfying the displayed lower bound (4.1), and set $k := \lfloor n/2 \rfloor$. Let $m \geq 4$ be the least integer such that

$$\binom{m}{\lfloor m/2 \rfloor} \geq k, \quad (4.2)$$

and put $\ell := \lfloor m/2 \rfloor$. By Corollary 2.2 (applied with $K = k$) we have

$$m \leq \log_2 k + \frac{1}{2} \log_2 \log_2 k + 3. \quad (4.3)$$

We next verify that

$$k \geq r + m + 1. \quad (4.4)$$

Case 1: $n < 8r$. Then $k = \lfloor n/2 \rfloor \leq 4r$. Under the hypothesis (4.1) this case can only occur when $r \geq 4$ (since for $r = 2, 3$ the right-hand side of (4.1) is at least $8r$). Using (4.3) and $k \leq 4r$ we obtain

$$m + 1 \leq \log_2(4r) + \frac{1}{2} \log_2 \log_2(4r) + 4 = \log_2 r + 6 + \frac{1}{2} \log_2(\log_2 r + 2).$$

On the other hand,

$$k - r \geq \frac{n - 2r - 1}{2} \geq \log_2 r + \frac{1}{2} \log_2 \log_2 r + \frac{15 - 1}{2}.$$

Since $r \geq 4$ implies $\log_2 \log_2 r + 2 \geq \log_2(\log_2 r + 2)$, the last display gives $k - r \geq m + 1$, which is equivalent to (4.4).

Case 2: $n \geq 8r$. Then $k \geq 4r \geq 8$, so $k - r \geq \frac{3}{4}k$. By minimality of m we have $k > \binom{m-1}{\lfloor (m-1)/2 \rfloor} \geq \binom{m-1}{2}$, hence $k \geq (m-1)(m-2)/2 + 1$. If $m \leq 5$ then $m + 1 \leq 6 \leq \frac{3}{4}k$ since $k \geq 8$. If $m \geq 6$ then one checks that $\frac{3}{8}(m-1)(m-2) \geq m + 1$, so

$$k - r \geq \frac{3}{4}k \geq \frac{3}{8}(m-1)(m-2) \geq m + 1,$$

and again (4.4) follows.

Choose disjoint sets

$$[n] = \{a\} \dot{\cup} P \dot{\cup} \{u\} \dot{\cup} V \dot{\cup} R \dot{\cup} W$$

with $|P| = r$, $|V| = m + 1$, $|R| = k - 2$, and W the (possibly empty) remaining set (which exists by (4.4)). Write

$$P = \{p_1, \dots, p_r\}, \quad V = V_0 \cup \{b\}, \quad |V_0| = m,$$

and set $R' := R \cup W$, so that $|R'| \geq |R| = k - 2 \geq r$ (using (4.4) and $m \geq 4$).

Apply Lemma 2.4 with $K = k$. The hypothesis (4.2) implies $\binom{m}{\ell} \geq k$ and hence $k - (\ell + 1) \leq \binom{m}{\ell}$, while $k \geq \ell + 1$ holds since $m \geq 4$ and $k \geq r + m + 1$ by (4.4). Therefore we obtain sets $L_t \subseteq V$ for every $t \in \{4, 5, \dots, k\}$ such that:

- (i) $|L_t| = 2$ for $4 \leq t \leq \ell + 1$, and these L_t are distinct and all contain b ;
- (ii) $|L_t| = \ell$ for $\ell + 2 \leq t \leq k$, and these L_t are distinct and all lie in V_0 ;

(iii) $\{L_t : 4 \leq t \leq k\}$ is an antichain under inclusion.

Define $\tilde{\mathcal{F}}$ level-by-level as follows.

- *Size 2*: for $j \in [r]$, set $F_{2,j} := \{a, p_j\}$.
- *Size 3*: choose distinct $x_1, \dots, x_r \in R'$ and set $F_{3,j} := \{a, u, x_j\}$.
- *Size $t \in \{4, \dots, k\}$* : let $s(t) := t - 1 - |L_t|$. Note that $s(t) \geq 1$ and, since $\ell \geq 2$ and $|R'| \geq k - 2$,

$$s(t) \leq t - 1 - 2 \leq k - 3 \leq |R'| - 1.$$

Hence $1 \leq s(t) \leq |R'| - 1$ and therefore

$$\binom{|R'|}{s(t)} \geq |R'| \geq r,$$

so we can choose r distinct $s(t)$ -subsets $B_{t,1}, \dots, B_{t,r} \subseteq R'$. Define

$$F_{t,j} := \{a\} \cup L_t \cup B_{t,j} \quad (j \in [r]).$$

Let

$$\tilde{\mathcal{F}} := \{F_{t,j} : 2 \leq t \leq k, j \in [r]\}.$$

Then every $F \in \tilde{\mathcal{F}}$ contains a , and for each $t \in \{2, 3, \dots, k\}$ the family $\tilde{\mathcal{F}}$ contains at least r sets of size t .

We next check that $\tilde{\mathcal{F}}$ is an antichain. If $F_{t,j} \subseteq F_{t',j'}$ with $t, t' \geq 4$, then by disjointness of the parts

$$L_t \subseteq L_{t'} \quad \text{and} \quad B_{t,j} \subseteq B_{t',j'}.$$

If $t \neq t'$, this contradicts the label antichain property (iii). If $t = t'$, then $|B_{t,j}| = |B_{t',j'}| = s(t)$ and $B_{t,j} \subseteq B_{t',j'}$ forces $B_{t,j} = B_{t',j'}$, contradicting distinctness. Finally, the cases involving sizes 2 or 3 cannot create inclusion because P , $\{u\}$, and R' are disjoint and each of $F_{2,j}, F_{3,j}$ contains an element (p_j or u) that never appears in any $F_{t',j'}$ with $t' \geq 4$. Thus $\tilde{\mathcal{F}}$ is an antichain. Moreover, $|F| \leq k \leq n/2$ for all $F \in \tilde{\mathcal{F}}$.

Define

$$\mathcal{A} := \tilde{\mathcal{F}} \cup \{[n] \setminus F : F \in \tilde{\mathcal{F}}\}.$$

Since $\tilde{\mathcal{F}}$ is an antichain, so is its complement family $\{[n] \setminus F : F \in \tilde{\mathcal{F}}\}$. It remains to rule out inclusions between $\tilde{\mathcal{F}}$ and its complement family. Fix $F, G \in \tilde{\mathcal{F}}$. Because every set in $\tilde{\mathcal{F}}$ contains a , the complement $[n] \setminus G$ does not contain a , hence $F \not\subseteq [n] \setminus G$. For the reverse inclusion, note that

$$|[n] \setminus G| = n - |G| \geq n - k = \lceil n/2 \rceil \geq k, \quad \text{whereas} \quad |F| \leq k.$$

If n is odd then $\lceil n/2 \rceil \geq k+1 > |F|$, so $[n] \setminus G \not\subseteq F$. If n is even, then either $\lceil n/2 \rceil > k \geq |F|$, in which case $[n] \setminus G \not\subseteq F$ by cardinality, or else $\lceil n/2 \rceil = k$ (equivalently $|G| = k$). In the latter case, $[n] \setminus G \subseteq F$ would force $[n] \setminus G = F$, contradicting $a \in F$ and $a \notin [n] \setminus G$. Therefore \mathcal{A} is an antichain.

By construction, for each $t \in \{2, \dots, k\}$ there are at least r sets of size t in \mathcal{A} , and taking complements yields at least r sets of each size $t \in \{n-k, \dots, n-2\}$. Since $k = \lfloor n/2 \rfloor$, we have $n-k = \lceil n/2 \rceil \in \{k, k+1\}$ and hence

$$\{2, \dots, k\} \cup \{n-k, \dots, n-2\} = \{2, 3, \dots, n-2\}.$$

Therefore every size $t \in \{2, 3, \dots, n-2\}$ appears in \mathcal{A} at least r times, so $|S(\mathcal{A})| = n-3$ and \mathcal{A} is an r -multiplicity antichain.

Thus, we construct an r -multiplicity antichain $\mathcal{A} \subseteq 2^{[n]}$ with $|S(\mathcal{A})| = n-3$, hence $g(n, r) \geq n-3$. On the other hand, since $r \geq 2$ and the present hypothesis on n implies $n \geq 4$, Lemma 2.5 yields the universal upper bound $g(n, r) \leq n-3$. Therefore $g(n, r) = n-3$. \square

We are now ready to present the following.

Proof of Theorem 1.5. By Proposition 4.1, we know that $g(n, r) = n - 3$ for every integer $r \geq 2$ and every integer $n \geq 2r + 2 \log_2 r + \log_2 \log_2 r + 15$. By Definition 1.3, this implies

$$n_0(r) \leq 2r + 2 \log_2 r + \log_2 \log_2 r + 15. \quad (4.5)$$

In particular, one has $n_0(r) \leq 2r + 2 \log_2 r + O(\log_2 \log_2 r)$. \square

5. CONCLUDING REMARKS

In this paper we studied the threshold $n_0(r)$ in the Erdős–Trotter problem (Problem 1.1). For $r = 2$ and $r = 3$ one has $n_0(2) = 3$ and $n_0(3) = 8$. For all integers $r \geq 4$, Theorem 1.4 and Theorem 1.5 yield the two-sided estimate

$$2r + 2 \leq n_0(r) \leq 2r + 2 \log_2 r + O(\log_2 \log_2 r).$$

In particular, this determines the leading linear term of $n_0(r)$ as $2r$.

A natural next question is whether the error term beyond the linear main term is bounded.

Problem 5.1. *Does there exist an absolute constant $C > 0$ such that $n_0(r) \leq 2r + C$ for all $r \geq 4$?*

For several small values of r , namely $r = 4, 5, \dots, 10$, we found explicit r -multiplicity antichains on $[n]$ with $n = 2r + 5$ and

$$S(\mathcal{F}) = \{2, 3, \dots, n - 2\}.$$

Equivalently, $g(2r + 5, r) = (2r + 5) - 3$ for $4 \leq r \leq 10$. The corresponding constructions are recorded in [1, 2r_plus_5_r_4_to_10.txt]. These examples show that the extremal value $g(n, r) = n - 3$ is attained already at $n = 2r + 5$ for $4 \leq r \leq 10$, which provides some numerical evidence toward Problem 5.1. Using our current search implementation [1, antichain_solver_one_based.py], we did not obtain a construction for $r = 11$ within 24 hours of runtime.

APPENDIX A. DETERMINING $n_0(2)$ AND $n_0(3)$

A.1. **The case $r = 2$.** We begin with the extremal value in the smallest nontrivial ground set, which will serve as the base case for the threshold statement.

Proposition A.1. *One has $g(3, 2) = 1$.*

Proof. Let $\mathcal{F} \subseteq 2^{[3]}$ be a 2-multiplicity antichain. Clearly, levels 0 and 3 cannot occur, since $|\binom{[3]}{0}| = |\binom{[3]}{3}| = 1 < 2$. Assume for contradiction that both levels 1 and 2 occur, i.e. $\{1, 2\} \subseteq S(\mathcal{F})$. Then $|\mathcal{F}_1| \geq 2$, so we may choose two distinct singletons $\{x\}, \{y\} \in \mathcal{F}_1$. Every 2-subset of $[3]$ contains at least one of x, y , hence every $H \in \binom{[3]}{2}$ satisfies $\{x\} \subseteq H$ or $\{y\} \subseteq H$, contradicting the antichain property. Therefore at most one of the levels 1 and 2 can occur, and thus $|S(\mathcal{F})| \leq 1$. On the other hand, $\mathcal{F} = \{\{1\}, \{2\}\}$ is a 2-multiplicity antichain with $|S(\mathcal{F})| = 1$. Hence $g(3, 2) = 1$. \square

We can now determine the threshold $n_0(2)$ by combining Proposition A.1 with our general upper bound and an explicit verification for the remaining finite range.

Theorem A.2. *One has $n_0(2) = 3$.*

Proof. Proposition A.1 implies the lower bound $n_0(2) \geq 3$. On the other hand, (4.5) yields the upper bound $n_0(2) \leq 21$. By Definition 1.3, to prove $n_0(2) = 3$ it is enough to verify that $g(n, 2) = n - 3$ holds for every integer n with $4 \leq n \leq 21$. Moreover, Lemma 2.5 gives the general upper bound $g(n, 2) \leq n - 3$. Thus it suffices to construct, for each integer $4 \leq n \leq 21$, a 2-multiplicity antichain on $[n]$ whose occurring set sizes cover exactly $n - 3$ distinct levels. This certifies $g(n, 2) \geq n - 3$ for all $4 \leq n \leq 21$, and hence $g(n, 2) = n - 3$ in this range. The constructions are recorded in our GitHub repository [1, r2_n_4_to_21.txt]. \square

A.2. The case $r = 3$. In this subsection we determine $n_0(3)$. The key input is a small obstruction at $n = 8$, which we obtain by a short structural reduction followed by an exhaustive check.

Proposition A.3. *One has $g(8, 3) \leq 4$.*

Proof. Let $\mathcal{F} \subseteq 2^{[8]}$ be a 3-multiplicity antichain. First, levels 0 and 8 cannot occur: indeed, $\binom{[8]}{0} = \binom{[8]}{8} = 1$, so neither level can support multiplicity 3.

Claim A.1. *If level 1 occurs, then $|S(\mathcal{F})| \leq 4$.*

Proof of Claim A.1. Assume $1 \in S(\mathcal{F})$, so $|\mathcal{F}_1| \geq 3$. Choose three distinct singletons $\{x\}, \{y\}, \{z\} \in \mathcal{F}_1$. By the antichain property, no other set in \mathcal{F} may contain any of x, y, z . Hence every set in $\mathcal{F} \setminus \mathcal{F}_1$ is contained in

$$[8] \setminus \{x, y, z\},$$

which has size 5. In particular, every set in \mathcal{F} has size at most 5. Moreover, at level 5 there is only one possible set, namely $[8] \setminus \{x, y, z\}$, so $|\mathcal{F}_5| \leq 1 < 3$, and thus $5 \notin S(\mathcal{F})$. Therefore,

$$S(\mathcal{F}) \subseteq \{1, 2, 3, 4\},$$

and hence $|S(\mathcal{F})| \leq 4$. \square

Claim A.2. *If level 7 occurs, then $|S(\mathcal{F})| \leq 4$.*

Proof of Claim A.2. Assume $7 \in S(\mathcal{F})$, so $|\mathcal{F}_7| \geq 3$. Pick three distinct 7-sets $H_1, H_2, H_3 \in \mathcal{F}_7$. Each can be written as

$$H_i = [8] \setminus \{a_i\} \quad (i = 1, 2, 3),$$

with a_1, a_2, a_3 pairwise distinct. Let $A \in \mathcal{F}$ with $|A| \leq 6$. Since \mathcal{F} is an antichain, we must have $A \not\subseteq H_i$ for every i , i.e. $a_i \in A$ for every i . Thus

$$\{a_1, a_2, a_3\} \subseteq A \quad \text{for all } A \in \mathcal{F} \text{ with } |A| \leq 6.$$

In particular, no set of size 0, 1, 2 can belong to \mathcal{F} , and also the only possible 3-set is $\{a_1, a_2, a_3\}$, so $3 \notin S(\mathcal{F})$. Consequently,

$$S(\mathcal{F}) \subseteq \{4, 5, 6, 7\},$$

and hence $|S(\mathcal{F})| \leq 4$. \square

Suppose for contradiction that $|S(\mathcal{F})| \geq 5$. By Claims A.1 and A.2 we must have $1, 7 \notin S(\mathcal{F})$, and also $0, 8 \notin S(\mathcal{F})$. Therefore, the only way to have $|S(\mathcal{F})| = 5$ is $S(\mathcal{F}) = \{2, 3, 4, 5, 6\}$. In particular, $|\mathcal{F}_t| \geq 3$ for each $t \in \{2, 3, 4, 5, 6\}$. Choose an arbitrary subfamily $\mathcal{F}' \subseteq \mathcal{F}$ by selecting exactly 3 sets from each layer $t \in \{2, 3, 4, 5, 6\}$. Then \mathcal{F}' is still an antichain, and moreover

$$|\mathcal{F}'_t| = 3 \quad \text{for each } t = 2, 3, 4, 5, 6.$$

Hence it suffices to rule out the existence of an antichain $\mathcal{F}' \subseteq 2^{[8]}$ with exactly three sets on each of the five levels 2, 3, 4, 5, 6.

We performed an exhaustive backtracking search over all choices of three sets from each of the levels 2, 3, 4, 5, 6, pruning only by necessary antichain constraints. For the top two levels

we encode 6-sets (resp. 5-sets) by their complements, which are 2-sets (resp. 3-sets), so that all containment constraints reduce to explicit intersection/subset tests. The implementation is contained in [1, `check_g_8_3.py`]; it exhausts all candidates and returns `None`, meaning that no feasible antichain \mathcal{F}' exists.

Hence, we conclude that $|S(\mathcal{F})| \neq 5$ for every 3-multiplicity antichain $\mathcal{F} \subseteq 2^{[8]}$. Therefore $g(8, 3) \leq 4$, as claimed. \square

With Proposition A.3 in hand, we can now determine the exact threshold $n_0(3)$.

Theorem A.4. *One has $n_0(3) = 8$.*

Proof. Proposition A.3 implies the lower bound $n_0(3) \geq 8$. On the other hand, (4.5) yields the upper bound $n_0(3) \leq 24$. By Definition 1.3, to prove $n_0(3) = 8$ it is enough to verify that $g(n, 3) = n - 3$ holds for every integer n with $9 \leq n \leq 24$. Moreover, Lemma 2.5 gives the universal upper bound $g(n, 3) \leq n - 3$ for all $n \geq 4$. Thus it suffices to construct, for each integer $9 \leq n \leq 24$, a 3-multiplicity antichain on $[n]$ whose occurring set sizes cover exactly $n - 3$ distinct levels. This yields $g(n, 3) \geq n - 3$ for all $9 \leq n \leq 24$, and hence $g(n, 3) = n - 3$ in this range. The explicit constructions are recorded in our GitHub repository [1, `r3_n_9_to_24.txt`]. \square

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES IN THE MANUSCRIPT PREPARATION PROCESS

During the early stages of this project, an AI assistant was used for exploratory brainstorming and preliminary code drafting related to small- r computational checks. All mathematical statements, proofs, and the final presentation of this paper were independently verified and written by the authors.

REFERENCES

- [1] ep776-antichains-with-multiplicity.
<https://github.com/QuanyuTang/ep776-antichains-with-multiplicity>
- [2] I. Anderson, *Combinatorics of Finite Sets*, Courier Corporation, 2002.
- [3] T. F. Bloom, Erdős Problem #776, <https://www.erdosproblems.com/776>, accessed 2026-02-01.
- [4] K. Engel, *Sperner Theory*, Encyclopedia of Mathematics and its Applications, vol. 65, Cambridge University Press, Cambridge, 1997.
- [5] P. Erdős, *Problem sessions*, In: Ordered Sets (Proc. NATO Adv. Study), edited by I. Rival, Dordrecht: Reidel (1981), 860–861.
- [6] R. Guy, A Miscellany of Erdős Problems. *Amer. Math. Month.* **90** (1983), 118–120.
- [7] E. Sperner, Ein Satz über Untermengen einer endlichen Menge, *Math. Z.* **27** (1928), 544–548.

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200433, P. R. CHINA
Email address: hyx717math@163.com

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, P. R. CHINA
Email address: tang_quanyu@163.com