

# On thermal transpiration and thermomolecular pressure difference

I-Kun Chen  
*National Taiwan University*

Kai-Li Wang  
*National Taiwan University*

March 20, 2026

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Introduction . . . . .	2
1.2	Organization of the article . . . . .	7
<b>2</b>	<b>Preliminaries</b>	<b>7</b>
2.1	Estimates for $\nu$ and $K$ . . . . .	7
2.2	Proposition of $L$ . . . . .	10
2.3	Proposition of $S_\Omega$ and $J$ . . . . .	12
2.4	Estimate of $Q_+$ and $Q_-$ . . . . .	13
<b>3</b>	<b><math>L^2</math> estimate</b>	<b>14</b>
<b>4</b>	<b><math>L^\infty</math> estimate</b>	<b>18</b>
4.1	The existence of solution to (80) . . . . .	19
4.2	the proof of <b>Theorem 1.1</b> . . . . .	26
<b>5</b>	<b>Application to Thermal transpiration and thermomolecular pressure difference</b>	<b>28</b>
5.1	the model of the transpiration : the proof of <b>Theorem 1.2</b> and <b>Theorem 1.3</b> . . . . .	28
<b>6</b>	<b>Appendix A: The <math>H^2</math> estimate of Laplace equation with Robin boundary condition</b>	<b>33</b>
<b>7</b>	<b>Appendix B: The sufficient conditions for unsealed sets</b>	<b>36</b>
<b>8</b>	<b>Appendix C: The equivalent definition of solution</b>	<b>37</b>

## Abstract

In this article, we demonstrate the phenomenon of thermal transpiration in a bounded convex domain. We employ the stationary Boltzmann equation with a cutoff potential. For boundary condition, we partition the boundary into diffuse reflection and incoming regions. We establish the existence of solution in a weighted  $L^\infty$  space. Furthermore, we consider a convex domain with diffuse reflection boundary condition in the middle and incoming boundary condition at the two ends. We first consider Maxwellians with the same pressure but different temperatures at the two ends. We prove that the total flux  $U(x)$  is directed toward the hot end. Furthermore, we derive an estimate for the total flux:

$$U(x) \geq C \left( 1 - \frac{1}{\sqrt{T_2}} \right). \quad (1)$$

In addition, we show that when the pressures and temperatures on the two ends satisfy the relation

$$\frac{P_1}{P_2} = \sqrt{\frac{T_1}{T_2}}, \quad (2)$$

the total flux of the solution is of order  $\mathcal{O}(\frac{1}{\kappa})$ . This result is consistent with Knudsen's finding of thermomolecular pressure difference in 1909.

**Keyword** : Stationary Boltzmann equation, thermal transpiration, thermomolecular pressure difference, mixed boundary condition.

## Section 1. Introduction

### 1.1. Introduction

The **thermal transpiration** is a phenomenon that temperature gradient on the boundary induces a flow in a pipe [38]. This phenomenon has many engineering applications, including the Crookes radiometer [12], which can be used in energy harvesting [39]; the Knudsen compressor, which pumps gas without any moving mechanical part [46]; microelectromechanical systems [34], and MEMS-based multistage Knudsen compressor, where the gas moves from a cold chamber to a hot chamber creating a pressure difference across the compressor element [22]. Further applications and details can be found in [24], [38].

The thermal transpiration was first observed by Reynolds in 1879 [37]. He observed that gas moves along a solid surface due to temperature differences. In the same year, Maxwell developed a theory to explain this effect [30]. Moreover, when considering two chambers connected by a tube, Knudsen showed that in the limit of infinitely small tube, the pressure between the two chambers satisfies

$$\frac{P_1}{\sqrt{T_1}} = \frac{P_2}{\sqrt{T_2}} \quad (3)$$

in equilibrium. This phenomenon is called the thermomolecular pressure difference. This relation is discussed in [26], [27]. In an experiment, S. Chu Liang provided a more accurate expression of (3) in [28], with further applications given in [2], [42].

In kinetic theory, the stationary Boltzmann equation serves as a fundamental mathematical model for describing the behavior of rarefied gas. It reads

$$v \cdot \nabla_x F(x, v) = \frac{1}{\kappa} Q(F, F)(x, v), \quad \text{in } \Omega \times \mathbb{R}^3. \quad (4)$$

Here,  $\kappa$  is Knudsen number and  $F$  represents the velocity distribution function of the gas molecules. The velocity distribution function is used to describe the number of molecules  $dN$  in a six-dimensional phase-space element  $d\mathbf{X}d\boldsymbol{\xi}$ . That is,

$$dN = F(\mathbf{X}, \boldsymbol{\xi}) d\mathbf{X}d\boldsymbol{\xi}.$$

The collision operator  $Q$  that appears in (4) is defined as follows.

$$\begin{aligned} & Q(F, G)(x, v) \\ &= \int_{\mathbb{R}^3} \int_{S^2} [F(v')G(u') + F(u')G(v') - F(v)G(u) - F(u)G(v)] B(|v-u|, \omega) d\omega du, \end{aligned}$$

where

$$v' = v + ((u-v) \cdot \omega)\omega \quad \text{and} \quad u' = u - ((u-v) \cdot \omega)\omega. \quad (5)$$

Note that  $v', u'$  can also be expressed in the equivalent form

$$v' = \frac{v+u}{2} + \frac{|v-u|}{2}\sigma \quad \text{and} \quad u' = \frac{v+u}{2} - \frac{|v-u|}{2}\sigma, \quad (6)$$

where  $\sigma \in \mathbb{S}^2$ .

Consequently, stationary Boltzmann equation has been extensively applied to describe the gas distribution in the domains involving thermal transpiration. However, in engineering applications, obtaining numerical solutions requires substantial computational efforts, either deterministically ([40], [44], [48]) or stochastically ([3], [35]). Some numerical analysis of thermal transpiration employ the lattice Boltzmann model [41], the BGK model ([21], [29], [31]) or the linearized Boltzmann Equation model [43]. In particular, for the linearized Boltzmann equation model, Sone, Y. and Aoki, K. have made many significant contributions ([1], [32]). Moreover, Sone provided an explicit form of the solution when the free transport equation is used to describe the thermal transpiration [38].

Although there are many results on the numerical analysis of thermal transpiration using the linearized Boltzmann equation, there are only a few mathematical results. In 2007, Chiun-Chuan Chen, I-Kun Chen, Tai-Ping Liu and Yoshio Sone solved this problem for gas between two plates or within an infinite long pipe under diffuse reflection boundary conditions. They showed that when the Knudsen number  $\kappa$  is sufficiently large, the local flow between two plates was of the order  $\log \kappa$  and the flow in a pipe has finite positive velocity. Moreover, they proved that in both case, the local flux is proportional to the wall temperature gradient [7]. This result is used to study the boundary logarithmic singularity ([10],[43]).

In this paper, we address the problem in the case where the pipe has finite length, and we impose the incoming boundary condition at both ends with Maxwellians and diffusion boundary condition in the middle.

In this article, we formulate the mixed boundary problem for the Boltzmann equation as follows:

$$\begin{cases} v \cdot \nabla_x F = \frac{1}{\kappa} Q(F, F), & \text{in } \Omega \times \mathbb{R}^3, \\ F(x, v) = F_{in}(x, v), & \text{on } \partial\Omega_1^-, \\ F(x, v) = m_{T(x)} \int_{u \cdot n(x) > 0} F(x, u) u \cdot n(x) du, & \text{on } \partial\Omega_2^-, \end{cases} \quad (7)$$

where  $\Omega$  is a bounded convex domain in  $\mathbb{R}^3$ . Here,  $\partial\Omega_1$  and  $\partial\Omega_2$  are subsets of  $\partial\Omega$  and satisfy the following assumption

**Assumption 1.1.**

1.  $\partial\Omega_1 \cup \partial\Omega_2 = \partial\Omega$ .
2.  $\partial\Omega_1$  and  $\partial\Omega_2$  are of positive measure.
3.  $\partial\Omega_1$  and  $\partial\Omega_2$  are unions of connected components of  $\partial\Omega$ .
4.  $\partial\Omega_1$  is relative open to  $\partial\Omega$ .
5.  $\overline{\partial\Omega_1} \cap \overline{\partial\Omega_2}$  consists of finitely many  $C^2$  curves in  $\mathbb{R}^3$ .

Here  $m_T = \frac{1}{2\pi T^2} e^{-\frac{|v|^2}{2T}}$  and  $m_{T(x)} = \frac{1}{2\pi T^2(x)} e^{-\frac{|v|^2}{2T(x)}}$ . Also, we denote  $m = m_1$ . Let  $\partial E$  is a subset of  $\partial\Omega$ . We define  $\partial E^-$ ,  $\partial E^+$  and  $\partial E^0$  as follows.

$$\begin{aligned} \partial E^- &= \{(x, v) \in \partial E \times \mathbb{R}^3 : n(x) \cdot v < 0\}, \\ \partial E^+ &= \{(x, v) \in \partial E \times \mathbb{R}^3 : n(x) \cdot v > 0\}, \\ \partial E^0 &= \{(x, v) \in \partial E \times \mathbb{R}^3 : n(x) \cdot v = 0\}, \end{aligned}$$

where  $n(x)$  denotes the outward unit normal vector as  $x \in \partial\Omega$ . In this article, we always assume that  $T(x)$  is positive and belongs to  $L^\infty(\partial\Omega)$ . Furthermore, we assume that  $\mathbf{B}(|v - u|, \omega)$  satisfies

$$\mathbf{B}(|v - u|, \omega) = \mathfrak{B}|v - u|^\gamma \left( \frac{u - v}{|u - v|} \cdot \omega \right), \quad -3 < \gamma \leq 1, \quad (8)$$

for some constant  $\mathfrak{B} > 0$ . When we discuss the thermal transpiration, we define two parameters as

$$\begin{aligned} A &:= \inf\{x_3 \in \mathbb{R}^3 | (x_1, x_2, x_3) \in \Omega\}, \\ B &:= \sup\{x_3 \in \mathbb{R}^3 | (x_1, x_2, x_3) \in \Omega\}. \end{aligned}$$

Let  $A < a < b < B$ , and define

$$\partial\Omega_1 = \left(\partial\Omega \cap \{z < a\}\right) \cup \left(\partial\Omega \cap \{z > b\}\right). \quad (9)$$

Finally, we set the distribution of gas  $F(x, v)$  in  $\Omega$  which satisfies the following boundary value problem:

$$\left\{ \begin{array}{ll} v \cdot \nabla_x F = \frac{1}{\kappa} Q(F, F), & \text{in } \Omega \times \mathbb{R}^3, \\ F(x, v) = \frac{1}{(2\pi)^{3/2}} e^{-\frac{|v|^2}{2}}, & \text{on } \partial\Omega_{1,a}^- = \partial\Omega_1^- \cap \{x_3 < a\}, \\ F(x, v) = \frac{1}{(2\pi)^{3/2} T_2^{5/2}} e^{-\frac{|v|^2}{2T_2}}, & \text{on } \partial\Omega_{1,b}^- = \partial\Omega_1^- \cap \{x_3 > b\}, \\ F(x, v) = m_T(x, v) \int_{u \cdot n(x) > 0} F(x, u) u \cdot n(x) du, & \text{on } \partial\Omega_2^- = \partial\Omega^- \setminus \left(\partial\Omega_{1,a}^- \cup \partial\Omega_{1,b}^-\right). \end{array} \right. \quad (10)$$

Furthermore, we impose that  $T(x_1, x_2, x_3) = 1$  when  $x_3 \leq a$ , and that  $T(x_1, x_2, x_3) = T_2$  when  $x_3 \geq b$ . We write the solution as a perturbation near the standard Maxwellian  $M(v) = \frac{1}{\sqrt{2\pi}} m(v)$ . Let  $F = M + m^{1/2} f$  and  $F_{in} = M + m^{1/2} f_{in}$ . Then, the function  $f$  solves the following boundary value problem:

$$\left\{ \begin{array}{ll} v \cdot \nabla_x f = -\frac{1}{\kappa} L(f) + \frac{1}{\kappa} \Gamma(f, f), & \text{in } \Omega \times \mathbb{R}^3, \\ f(x, v) = f_{in}(x, v), & \text{on } \partial\Omega_1^-, \\ f(x, v) = m^{-1} m_{T(x)} P(f) + \frac{1}{\sqrt{2\pi}} m^{-1/2} (m_{T(x)} - m), & \text{on } \partial\Omega_2^-, \end{array} \right. \quad (11)$$

where

$$\begin{aligned} L(f) &= m^{-1/2} \left( Q(M, m^{1/2} f) + Q(m^{1/2} f, M) \right), \\ \Gamma &= m^{-1/2} Q(m^{1/2} f, m^{1/2} f), \\ P(f)(x, v) &= m^{1/2} \int_{u \cdot n(x, v) > 0} f(x, u) m^{1/2} u \cdot n(x) du. \end{aligned}$$

The operator  $L$  has two components: the multiplicative operator  $\nu$  and integral operator  $K$ , that is,  $L(f) = \nu(f)f - K(f)$ . Their definition can be found in **section 2** or [18]. In what follows, we discuss the well-posedness of the boundary value problem (11). The integral form of (11) is

$$f = \frac{1}{\kappa} S_\Omega^\kappa K f + J^\kappa \left( f_{in} \mathbf{1}_{\partial\Omega_1^-} + \left( m^{-1} m_{T(x)} P(f) + \frac{1}{\sqrt{2\pi}} m^{-1/2} (m_{T(x)} - m) \right) \mathbf{1}_{\partial\Omega_2^-} \right) + \frac{1}{\kappa} S_\Omega^\kappa \Gamma(f, f). \quad (12)$$

Here, we define

$$S_\Omega^\kappa f(x, v) := \int_0^{\tau_-(x, v)} e^{-\frac{1}{\kappa} \nu(v) s} f(x - sv, v) ds, \quad (13)$$

$$J^\kappa f := e^{-\frac{1}{\kappa} \nu(v) \tau_-(x, v)} f(q(x, v), v), \quad (14)$$

where

$$\tau_-(x, v) = \sup\{t | x - tv \in \bar{\Omega}\} \quad \text{and} \quad q(x, v) = x - \tau_-(x, v)v. \quad (15)$$

For convenience, we define

$$\langle f, g \rangle = \int_{\Omega \times \mathbb{R}^3} fg dx dv, \quad \langle f, g \rangle_v = \int_{\mathbb{R}^3} fg dv, \quad (16)$$

$$\langle f, g \rangle_{\partial_2^-} = \int_{\partial\Omega_2^-} fg |v \cdot n(x)| dx dv. \quad (17)$$

To state our main theorem, we introduce some function spaces. For  $\alpha \geq 0$ ,  $\beta \in \mathbb{R}$  and  $p \geq 1$ , we say  $f \in L_{\alpha, \beta}^p(\Omega \times \mathbb{R}^3)$  if

$$\begin{aligned} \|f\|_{L_{\alpha, \beta}^p} &= \|f\|_{L_{\alpha, \beta}^p(\Omega \times \mathbb{R}^3)} := \left( \int_{\Omega} \int_{\mathbb{R}^3} |f(x, v) e^{\alpha|v|^2} (1 + |v|)^{\beta}|^p dv dx \right)^{1/p} < \infty & \text{for } p < \infty, \\ \|f\|_{L_{\alpha, \beta}^{\infty}} &= \|f\|_{L_{\alpha, \beta}^{\infty}(\Omega \times \mathbb{R}^3)} := \operatorname{ess\,sup}_{(x, v) \in \Omega \times \mathbb{R}^3} |f(x, v) e^{\alpha|v|^2} (1 + |v|)^{\beta}| < \infty & \text{for } p = \infty. \end{aligned}$$

Moreover, we say  $f \in L_{\alpha, \beta}^p(\partial E^-)$  for some  $\partial E \subset \partial\Omega$ , if

$$\begin{aligned} \|f\|_{L_{\alpha, \beta}^p(\partial E^-)} &:= \left( \int_{\partial E^-} |f(x, v) e^{\alpha|v|^2} (1 + |v|)^{\beta}|^p |v \cdot n(x)| dS dv \right)^{1/p} < \infty & \text{for } p < \infty, \\ \|f\|_{L_{\alpha, \beta}^{\infty}(\partial E^-)} &:= \operatorname{ess\,sup}_{(x, v) \in \partial E^-} |f(x, v) e^{\alpha|v|^2} (1 + |v|)^{\beta}| < \infty & \text{for } p = \infty, \end{aligned}$$

where  $dS$  denotes the surface measure on  $\partial\Omega$ . For convenience, we write  $L^p(A) = L_{0,0}^p(A)$  when  $A = \Omega \times \mathbb{R}^3$  or  $\partial E^- \subseteq \partial\Omega \times \mathbb{R}^3$ . Throughout this article, we denote  $f(x) = \mathcal{O}(g(x))$  when  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$  is a constant.

We now introduce the concept of an unsealed set  $\Omega$  in  $\mathbb{R}^3$ . The term *unseal* describes the visibility of  $\partial\Omega_1$  from points on  $\partial\Omega$ . If every point  $x$  can directly see the  $\partial\Omega_1$ , then we call  $\Omega$  a **completely unsealed set**. If, for every  $x \in \partial\Omega$ , the  $\partial\Omega_1$  becomes visible after tracing back  $m$  times for some fixed integer  $m$ , then we call  $\Omega$  an **unsealed set**. The precise definition is in **section 4**.

**Theorem 1.1.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded convex domain with  $C^2$  boundary. Suppose that  $\Omega$  is unsealed with respect to  $\partial\Omega_1$  and satisfies the **assumption 1.1**. Assume that  $-3 < \gamma \leq 1$  and that (8) holds. Then, for any  $0 < \alpha < 1/4$  and  $\kappa_0 \in (0, \infty)$ , there exists  $\delta > 0$  which depends on  $\alpha$ ,  $\Omega$ ,  $\partial\Omega_1$ , and  $\kappa_0$  such that if*

$$\|m^{-1/2} (F_{in} - M)\|_{L_{\alpha, 0}^{\infty}(\partial\Omega_1^-)} \leq \delta \quad (18)$$

and

$$|T(x) - 1| \leq \delta, \quad \forall x \in \Omega_2 \quad (19)$$

then the equation (7) has a unique mild solution for  $\kappa \geq \kappa_0$ . Furthermore,

$$\begin{aligned} &\|m^{-1/2} (F - M)\|_{L_{\alpha, 0}^{\infty}} + \|m^{-1/2} (F - M)\|_{L_{\alpha, 0}^{\infty}(\partial\Omega_2^+)} \\ &\leq 2C_{\zeta} \left( \|m^{-1/2} (F_{in} - M)\|_{L_{\alpha, 0}^{\infty}(\partial\Omega_1^-)} + \|m^{-1/2} (m_{T(x)} - m)\|_{L_{\alpha, 0}^{\infty}(\partial\Omega_2^-)} \right). \end{aligned} \quad (20)$$

For  $x = (x_1, x_2, z) \in \Omega$ , the local flux  $u(x)$  and the total flux  $U(z)$  in the  $x_3$ -direction are defined as follows:

$$u(x) := \int_{\mathbb{R}^3} v_3 F(x, v) dv, \quad U(z) := \int_{\{(x_1, x_2, z) \in \Omega\}} u(x) dS. \quad (21)$$

By conservation law, the total flux is a constant with respect to  $z$  when  $z \in [a, b]$ .

**Theorem 1.2.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with  $C^1$  boundary such that  $\Omega \cap \{a \leq x_3 \leq b\}$  is convex. Suppose for every  $\kappa > \kappa_0 > 0$ , there exists a constant  $\mathcal{C}$  such that (10) admits a mild solution with the following estimate:*

$$\|F\|_{L_{0,\beta}^\infty(\Omega \times \mathbb{R}^3)} \leq \mathcal{C}, \quad (22)$$

where  $\beta > \max\{\gamma, 0\} + 5$ . Then, there exists  $\kappa_1 > \kappa_0$  such that the total flux  $U(x_3)$  of  $F$  is positive whenever  $\kappa \geq \kappa_1$ . Moreover,

$$U(x_3) \geq C \left(1 - \frac{1}{\sqrt{T_2}}\right). \quad (23)$$

In Knudsen's experiment ([26]), he connects two chambers with different temperature  $T_1, T_2$  with a narrow tube. He measures the pressures in two tanks  $P_1, P_2$  at equilibrium when there is no flux in the tube. He observes in the limit of infinitely small tubing,  $\frac{P_1}{\sqrt{T_1}} = \frac{P_2}{\sqrt{T_2}}$ . For the free transport equation case, the relation is exact ([38]). In our case, we obtain the same result by considering the following boundary value problem:

$$\begin{cases} v \cdot \nabla_x F = \frac{1}{\kappa} Q(F, F), & \text{in } \Omega \times \mathbb{R}^3, \\ F(x, v) = \frac{1}{(2\pi)^{3/2}} e^{-\frac{|v|^2}{2}}, & \text{on } \partial\Omega_{1,a}^- = \partial\Omega_1^- \cap \{x_3 < a\}, \\ F(x, v) = \frac{1}{(2\pi)^{3/2} T_2^2} e^{-\frac{|v|^2}{2T_2}}, & \text{on } \partial\Omega_{1,b}^- = \partial\Omega_1^- \cap \{x_3 > b\}, \\ F(x, v) = m_T(x, v) \int_{v \cdot n(x) > 0} F(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^- = \partial\Omega^- \setminus (\partial\Omega_{1,a}^- \cup \partial\Omega_{1,b}^-). \end{cases} \quad (24)$$

**Theorem 1.3.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with  $C^1$  boundary such that  $\Omega \cap \{a \leq x_3 \leq b\}$  is convex. Suppose that for every  $\kappa > \kappa_0$ , equation (10) admits a mild solution bounded in  $L_{0,\beta}^\infty$  by a constant independent of  $\kappa$ , where  $\beta > \max\{\gamma, 0\} + 5$ . Then, the total flux  $U(x_3)$  of  $F$  satisfies the following estimate:*

$$|U(x_3)| = \mathcal{O}\left(\frac{1}{\kappa}\right). \quad (25)$$

We can extend **Theorem 1.3** to the case when the incoming data is not localized at two places, i.e.,

$$\begin{cases} v \cdot \nabla_x F = \frac{1}{\kappa} Q(F, F), & \text{in } \Omega \times \mathbb{R}^3, \\ F(x, v) = \frac{1}{(2\pi)^{3/2} T_i^2} e^{-\frac{|v|^2}{2}}, & \text{on } \partial B_i^-, i = 1, \dots, n, \\ F(x, v) = m_T(x, v) \int_{v \cdot n(x) > 0} F(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^- = \partial\Omega^- \setminus (\cup_{i=1}^n \partial B_i^-), \end{cases} \quad (26)$$

where  $\partial B_i$  is an open subset of  $\partial\Omega$  and we assume that  $T(x) = T_i$  if  $x_i \in \partial B_i$ .

**Remark 1.1.** *Note that the condition  $\Omega \cap \{a \leq x_3 \leq b\}$  is convex ensures that  $\Omega$  is an unsealed set with respect to*

$$\left(\partial\Omega \cap \{z < a\}\right) \cup \left(\partial\Omega \cap \{z > b\}\right).$$

*The proof of **Theorem 1.2** and **Theorem 1.3** is valid for a domain in which  $\Omega \cap \{a \leq x_3 \leq b\}$  is not convex but is unsealed with respect to  $\left(\partial\Omega \cap \{z < a\}\right) \cup \left(\partial\Omega \cap \{z > b\}\right)$ .*

**Remark 1.2.** *Note that when  $\Omega$  is a bounded convex domain with  $C^2$  boundary. The **Theorem 1.1** ensures the condition of **Theorem 1.2**.*

## 1.2. Organization of the article

We briefly outline the proof in this article. We use the traditional  $L^2 - L^\infty$  estimate together with the bootstrap method to construct a solution of (7).

In **section 2**, we review some elementary properties of the linear operator  $L$ . In **section 3**, inspired by [14], [15], and [25], we introduce a simpler and more efficient approach to obtaining the  $L^2$  estimate. This approach is based on the operator  $S_\Omega^\kappa K$ , which is compact in  $L_{0,\gamma/2}^2$ .

In **section 4**, we use the unsealed condition to obtain the  $L^\infty$  estimate. This concept provides several advantages, such as reducing the weight requirement in  $L^\infty$ . Moreover, this is a key tool in proving **Theorem 1.2** and **Theorem 1.3**.

In **section 5**, we use original Boltzmann equation to address the problem by modeling the distribution as in (10), with incoming data at  $(\partial\Omega \cap \{z < a\})$  and  $(\partial\Omega \cap \{z > b\})$ . We cannot apply the powerful technique developed in [7]. However, we observe that the crucial mechanism for the appearance of flux lies in the different between the incoming data and the solution of the free transport model, which satisfies (8). By combining this observation with the concept of the unsealed set, we complete the proof of the main theorems in this article.

## Section 2. Preliminaries

In this section, we review some properties of the linear operator  $L$  and its components: the multiplicative operator  $\nu$  and the integral operator,  $K$ . We also present several propositions about the operator  $S_\Omega^\kappa$  and  $J^\kappa$ , which are useful in **section 3** and **section 4**. In addition, we recall some basic properties of  $Q_+$  and  $Q_-$ .

We emphasize that we linearize around Maxwellian  $M = \frac{1}{(2\pi)^{3/2}} e^{-\frac{|v|^2}{2}}$ . There is another popular convention to linearize around  $m = \frac{1}{\pi^{3/2}} e^{-|v|^2}$ , for example [25], [38]. Thus, the exponents differ by a factor of 2 and the constants are different.

### 2.1. Estimates for $\nu$ and $K$

Recall that  $L(f) = \nu(v)f - K(f)$  when the assumption (8) holds, and that  $\nu(v)$  and  $K(f)$  have explicit forms as follows

$$\begin{aligned}\nu(v) &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} B(|v-u|, \sigma) m(u) du d\sigma, \\ K(f)(x, v) &= \int_{\mathbb{R}^3} k(v, u) f(x, u) du = \int_{\mathbb{R}^3} (k_1(v, u) - k_2(v, u)) f(x, u) du.\end{aligned}$$

The explicit formulas for  $k_1$  and  $k_2$  are:

$$\begin{aligned}k_1(v, u) &= B_0 \pi |v-u|^\gamma e^{-\frac{1}{4}(|v|^2 + |u|^2)}, \\ k_2(v, u) &= \frac{2\mathfrak{B}}{\pi^{3/2}} \frac{1}{|v-u|} e^{-\frac{1}{8}|v-u|^2 - |V_1(v, u)|^2} \times \int_{W_{v-u}} e^{-|w+V_2(v, u)|^2} (|v-u|^2 + |w|^2)^{\frac{\gamma}{2}} dw,\end{aligned}$$

where

$$\begin{aligned}V_1(v, u) &:= \frac{1}{2\sqrt{2}} \frac{|v|^2 - |u|^2}{|v-u|^2} (v-u), \\ V_2(v, u) &:= \frac{(v-u) \times (v \times u)}{\sqrt{2}|v-u|^2}, \\ W_{v-u} &:= \{w \in \mathbb{R}^3 | w \perp (v-u)\}.\end{aligned}$$

Notice that we have the following identity

$$\begin{aligned}|V_1(v, u)|^2 + |V_2(v, u)|^2 &= \frac{1}{8}|v+u|^2, \\ V_2(v, u) \cdot (v-u) &= 0, \\ \frac{1}{8}|v-u|^2 + |V_1(v, u)|^2 + |V_2(v, u)|^2 &= \frac{1}{4}(|v|^2 + |u|^2).\end{aligned}$$

For their derivation, see [13]. We first give some estimates for the collision frequency  $\nu(v)$ .

**Proposition 2.1.** *Suppose that (8) holds and  $-3 < \gamma \leq 1$ . For the collision frequency  $\nu(v)$ , there exist positive constants  $\nu_0$  and  $\nu_1$  such that*

$$\nu_0(1 + |v|)^{\gamma/2} \leq \nu(v) \leq \nu_1(1 + |v|)^{\gamma/2}$$

for all  $v \in \mathbb{R}^3$ .

For the proof of **Proposition 2.1**, see [13], [18].

**Proposition 2.2.** *Let  $-3 < \gamma \leq 1$  and let  $\Omega$  be a bounded domain in  $\mathbb{R}^3$ . Suppose (8) holds. Then, we have*

$$\int_0^{\frac{L}{|v|}} e^{-\frac{1}{\kappa}\nu(|v|)t} dt \leq C_{\gamma,L} \frac{\kappa}{1 + |v|}$$

for any fix  $L > 0$  and  $(x, v) \in \Omega \times \mathbb{R}^3$ .

The proof of **Proposition 2.2** can be found in [47]. We next introduce some estimate for integral operator  $K(f)$ .

**Proposition 2.3.** *Let  $-3 < \gamma \leq 1$  and  $0 < \delta < 1$ . Suppose that (8) holds. Then, we have*

$$|k(v, u)| \leq C_{\gamma,\delta} \frac{w_\gamma(|v - u|)}{(1 + |v| + |u|)^{1-\gamma}} E_\delta(v, u) \quad (27)$$

for all  $v, u \in \mathbb{R}^3$ , where

$$E_\delta(v, u) := e^{-\frac{1-\delta}{8}(|v-u|^2 + 4|V_1(v,u)|^2)} = e^{-\frac{1-\delta}{8}\left(|v-u|^2 + \left(\frac{|v|^2 - |u|^2}{|v-u|}\right)^2\right)}$$

and

$$w_\gamma(|v - u|) := \begin{cases} 1, & -1 < \gamma \leq 1, \\ \frac{|\ln |v - u|| + 1}{|v - u|}, & \gamma = -1, \\ \frac{1}{|v - u|^{|\gamma|}}, & -3 < \gamma < -1. \end{cases} \quad (28)$$

The detail of proof can be found in [13] and the appendix of [25].

In the **section 4**, we estimate  $L_{\alpha,0}^\infty$  norm of the solution of Boltzmann equation. Hence, we need to discuss

$$K_\alpha(f) := w(v)K(w^{-1}f) = \int_{\mathbb{R}^3} w(v)k(v, u)w^{-1}(u)f(x, u)du := \int_{\mathbb{R}^3} k_\alpha(v, u)f(x, u)du$$

for  $w(v) = e^{\alpha|v|^2}$ ,  $\alpha > 0$ . It is known in [9] that

$$\begin{aligned} E_\delta(v, u) &= e^{a|v|^2} e^{-\alpha_{1,a,\delta}|v-u|^2} e^{-(1-\delta)\left(\frac{(v-u)\cdot v}{|v-u|} - \alpha_{2,a,\delta}|v-u|\right)^2} e^{-a|u|^2} \\ &= e^{-a|v|^2} e^{-\alpha_{1,a,\delta}|v-u|^2} e^{-(1-\delta)\left(\frac{(v-u)\cdot v}{|v-u|} + \alpha_{2,a,\delta}|v-u|\right)^2} e^{a|u|^2} \end{aligned}$$

for  $-1/4 < a < 1/4$ ,  $\delta > 0$  and  $v, u \in \mathbb{R}^3$ , where

$$\begin{aligned} \alpha_{1,a,\delta} &:= \frac{(1-\delta)^2 - 4a^2}{8(1-\delta)}, \\ \alpha_{2,a,\delta} &:= \frac{1-\delta-2a}{2\sqrt{2}(1-\delta)}. \end{aligned}$$

Plugging these identities into that estimate (27) with  $a = \alpha$  and  $\delta = 1/2 - \alpha$ , we obtain

$$|k_\alpha(v, u)| \leq C_{\gamma, \alpha} \frac{w(|v - u|)}{(1 + |v| + |u|)^{1-\gamma}} e^{-\alpha_1 |v-u|^2} e^{-\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2} \quad (29)$$

for some  $\alpha_1, \alpha_2, \alpha_3 > 0$  depending only on  $\alpha$ . Based on the above calculations, we obtain the following estimates, which are related to the kernel of the operator  $K_\alpha$ .

**Proposition 2.4.** [4] *Let  $\alpha_1, \alpha_3 > 0$  and  $\alpha_2 \in \mathbb{R}^3$ . Then, we have*

$$\int_{\mathbb{R}^3} \frac{1}{|v - u|^\mu} e^{-\alpha_1 |v-u|^2} e^{-\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2} \leq \frac{C_{\alpha, \mu}}{1 + |v|} \quad (30)$$

for any  $\mu < 3$  and  $v \in \mathbb{R}^3$ .

*Proof.* Let  $w = v - u$ . By using the sphere coordinate and setting  $\omega = |v| \cos \theta - \alpha_2 r$ , we have

$$\begin{aligned} \int_{\mathbb{R}^3} \frac{1}{|v - u|^\mu} e^{-\alpha_1 |v-u|^2} e^{-\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2} du &= \int_0^\infty \int_0^{2\pi} \int_0^\pi \frac{1}{r^{\mu-2}} e^{-\alpha_1 r^2} e^{-\alpha_3 (|v| \cos \theta - \alpha_2 r)^2} \sin \theta d\theta d\phi dr \\ &= \frac{2\pi}{|v|} \int_0^\infty \int_{-|v| - \alpha_2 r}^{|v| - \alpha_2 r} \frac{1}{r^{\mu-2}} e^{-\alpha_1 r^2} e^{-\alpha_3 w^2} d\omega \phi dr \\ &\leq C_{\alpha_1, \alpha_3} \frac{1}{|v|}. \end{aligned}$$

Moreover, we can deduce that

$$\int_{\mathbb{R}^3} \frac{1}{|v - u|^\mu} e^{-\alpha_1 |v-u|^2} e^{-\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2} \leq C_\alpha.$$

Here, the constant  $C_\alpha$  is independent to  $v$ . Combining these results, the proof is complete.  $\square$

**Proposition 2.5.** [25] *Let  $0 \leq \alpha < 1/4, \beta \in \mathbb{R}^3, -3 < \gamma \leq 1$  and  $1 \leq q < \min\{3, \frac{3}{|\gamma|}\}$ . Suppose that (8) holds. Then we have*

$$\int_{\mathbb{R}^3} |k_\alpha(v, u)|^q (1 + |u|)^\beta du \leq C_{\alpha, \gamma} (1 + |v|)^{\beta + q(\gamma-1) - 1} \quad (31)$$

for all  $v \in \mathbb{R}^3$ .

**Remark 2.1.** *We set  $\min\{3, \frac{3}{|\gamma|}\} = 3$  when  $\gamma = 0$*

The proof can be found in [25]. For convenience, we present the proof here.

*Proof. Case 1  $\beta \geq 0$ :*

Notice that  $(1 + |u|)^\beta \leq C_\beta ((1 + |v|)^\beta + |v - u|^\beta)$  for all  $v, u \in \mathbb{R}^3$ . By (29), we have

$$\begin{aligned} |k_\alpha(v, u)|^q (1 + |u|)^\beta &\leq C(1 + |v|)^{\beta + q(\gamma-1)} w_\gamma (|v - u|)^q e^{-q\alpha_1 |v-u|} e^{-q\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2} \\ &\quad + C(1 + |v|)^{q(\gamma-1)} w_\gamma (|v - u|)^q |v - u|^\beta e^{-q\alpha_1 |v-u|} e^{-q\alpha_3 \left( \frac{(v-u) \cdot v}{|v-u|} - \alpha_2 |v-u| \right)^2}. \end{aligned}$$

Observe that there exists  $\mu_{\gamma, q} < 3$  such that  $w_\gamma (|v - u|)^q \leq |v - u|^{-\mu_{\gamma, q}}$ . The constant  $\mu_{\gamma, q}$  only depend on  $\gamma$  and  $q$ . Also, we note

$$w_{-1} (|v - u|) e^{-\alpha_1 |v-u|^2} \leq C_\epsilon |v - u|^{-(1+\epsilon)} \quad (32)$$

for some small positive constant  $\epsilon$ . Thus, we can treat the case  $\gamma = -1$  in the same way as for  $-3 < \gamma < -1$ . In any case, by **proposition 2.4** we have

$$\begin{aligned} \int_{\mathbb{R}^3} |k_\alpha(v, u)|^q (1 + |u|)^\beta du &\leq C_{\alpha, \gamma} (1 + |v|)^{q(\gamma-1) - 1} + C_{\alpha, \gamma} (1 + |v|)^{\beta + q(\gamma-1) - 1} \\ &\leq C_{\alpha, \gamma} (1 + |v|)^{q(\gamma-1) - 1}. \end{aligned}$$

**Case 2**  $\beta < 0$ :  
Observe that

$$\begin{aligned} (1 + |u|)^\beta &= (1 + |v|)^\beta \frac{(1 + |v|)^{|\beta|}}{(1 + |u|)^{|\beta|}} \\ &\leq (1 + |v|)^\beta \frac{(1 + |u|)^{|\beta|} + |v - u|^{|\beta|}}{(1 + |u|)^{|\beta|}} \\ &\leq (1 + |v|)^\beta \left(1 + |v - u|^{|\beta|}\right). \end{aligned}$$

We apply the above argument to obtain the same estimate. Thus, the proof is complete.  $\square$

## 2.2. Proposition of $L$

Next, we review some basic properties of  $L$ , including the spectral gap and the projection of  $L$ . Recall that the normal orthogonal basis of the null space  $\text{Ker } L$  is given by

$$\begin{aligned} \Phi_0 &= (2\pi)^{-\frac{3}{4}} m^{\frac{1}{2}}(v), \\ \Phi_i &= (2\pi)^{-\frac{3}{4}} v_i m^{\frac{1}{2}}(v), \quad i = 1, 2, 3, \\ \Phi_4 &= \frac{(2\pi)^{-\frac{3}{4}}}{\sqrt{6}} (|v|^2 - 3) m^{\frac{1}{2}}(v). \end{aligned}$$

For each  $f = f(v) \in L^2(\mathbb{R}_v^3)$ , we denote the macroscopic part  $P_L(f)$  as the projection of  $f$  onto  $\text{Ker } L$ , that is,

$$P_L(f) = \sum_{i=0}^4 \langle f, \Phi_i \rangle_v \Phi_i,$$

and further denote  $(I - P_L)(f) = f - P_L(f)$  to be the microscopic part of  $f$ . It is well-known that there exists a constant  $C_L$  such that

$$\langle L(f), f \rangle_v \geq C_L \|(I - P_L)(f)\|_{L^2_{0,\gamma/2}(dv)}^2. \quad (33)$$

Here,  $L^2_{0,\gamma/2}(dv)$  denotes integration only with respect to the variable  $v$ . The proof can be found in [18], [21] and [23]. We present a key estimate concerning  $P_L$  and  $(I - P_L)$ .

**Proposition 2.6.** *Let  $\Omega$  be a bounded connected domain in  $\mathbb{R}^3$  with  $C^2$  boundary and  $-3 < \gamma \leq 1$ . Suppose that  $f \in L^2_{0,\gamma/2}$  is a solution to*

$$\begin{cases} v \cdot \nabla_x f + \frac{1}{\kappa} L(f) = \phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ f = f_{in}, & \text{on } \partial\Omega_1^-, \\ f = P(f) + \phi_2(x, v), & \text{on } \partial\Omega_2^- \end{cases} \quad (34)$$

in distribution sense. Then,

$$\|P_L(f)\|_{L^2_{0,\gamma/2}}^2 \leq C \left( \left[1 + \frac{1}{\kappa}\right] \|(I - P_L)(f)\|_{L^2_{0,\gamma/2}}^2 + \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|(I - P)f\|_{L^2(\partial\Omega_2^+)}^2 \right. \\ \left. + \|f\|_{L^2(\partial\Omega_1^+)}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (35)$$

*Proof.* Recall  $P_L(f) = \sum_{i=0}^4 \langle f, \Phi_i \rangle_v \Phi_i$ . Thus, it suffices to compute  $\langle f, \Phi_i \rangle_v$ , for  $i = 0, \dots, 4$ . The estimate of  $\langle f, \Phi_i \rangle_v$  are the same as those given in **lemma 3.3** in [15] for  $i = 1, \dots, 4$ . The standard elliptic estimates appearing in [15] can also be found in the section 2 of [19] or section 6 of [16]. For  $i = 0$ , due to the difference between the mixed and diffuse boundary conditions we need to find a

different test function to estimate  $\langle f, \Phi_0 \rangle$ . The Green's identity provides the following weak version of (34):

$$\int_{\partial\Omega \times \mathbb{R}^3} \psi f dSdv - \iint_{\Omega \times \mathbb{R}^3} v \cdot \nabla_x \psi f dx dv = - \iint_{\Omega \times \mathbb{R}^3} \psi \frac{1}{\kappa} L(\mathbf{I} - \mathbf{P})f dx dv + \iint_{\Omega \times \mathbb{R}^3} \psi \phi_1 dx dv. \quad (36)$$

We choose the test function

$$\psi = (|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v \cdot \nabla_x \phi_{\langle f, \Phi_0 \rangle_v} \Phi_0, \quad (37)$$

where  $\phi_{\langle f, \Phi_0 \rangle_v}$  is the solution of following elliptic problem

$$\begin{cases} -\Delta \phi_{\langle f, \Phi_0 \rangle_v} = \langle f, \Phi_0 \rangle_v, & \text{in } \Omega, \\ \chi(x) \phi_{\langle f, \Phi_0 \rangle_v} + \partial_n \phi_{\langle f, \Phi_0 \rangle_v} = 0, & \text{on } \partial\Omega. \end{cases} \quad (38)$$

Here  $\chi(x)$  is a non-negative  $C^2$  function. We require that  $\chi(x) = 0$  on  $\partial\Omega_2$  but  $\chi > 0$  on a positive measure subset of  $\partial\Omega$ . Notice that we have following estimate

$$\|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(\Omega)} \leq C \|\langle f, \Phi_0 \rangle_v\|_{L^2(\Omega)}. \quad (39)$$

This estimate can be found in [17] and [36]. We then bound the right hand side of (36) by

$$\frac{1}{\kappa} \|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(dx)} \|(\mathbf{I} - \mathbf{P}_L)f\|_{L^2_{0,\gamma/2}} + \|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(dx)} \|\phi_1\|_{L^2_{0,\gamma/2}}.$$

We choose  $\beta_{\langle f, \Phi_0 \rangle_v}$  such that

$$\int_{\mathbb{R}^3} (|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) \left( \frac{|v|^2}{2} - \frac{3}{2} \right) (v_i)^2 \mu(v) dv = 0. \quad (40)$$

Moreover, we decompose that

$$\begin{aligned} f &= \mathbf{1}_{\partial\Omega_1^+} f + \mathbf{1}_{\partial\Omega_1^-} f_{in} + \mathbf{1}_{\partial\Omega_2} P f + \mathbf{1}_{\partial\Omega_2^+} (I - P)f + \mathbf{1}_{\partial\Omega_2^-} \phi_2, & \text{on } \partial\Omega \times \mathbb{R}^3, \\ f &= \sum_{i=0}^4 \langle f, \Phi_i \rangle_v \Phi_i + (\mathbf{I} - \mathbf{P}_L)f, & \text{on } \Omega \times \mathbb{R}^3. \end{aligned} \quad (41)$$

By (41), since the contribution of  $\langle f, \Phi_4 \rangle_v$  vanishes in (36) due to our choice of  $\beta_{\langle f, \Phi_0 \rangle_v}$  and the contributions of  $\langle f, \Phi_1 \rangle_v$ ,  $\langle f, \Phi_2 \rangle_v$ , and  $\langle f, \Phi_3 \rangle_v$  vanish in (36) due to the oddness, the left-hand side of (36) takes the form of

$$\int_{\partial\Omega \times \mathbb{R}^3} n \cdot v (|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v \cdot \nabla_x \phi_{\langle f, \Phi_0 \rangle_v} \Phi_0 [\mathbf{1}_{\partial\Omega_1^+} f + \mathbf{1}_{\partial\Omega_1^-} f_{in} + \mathbf{1}_{\partial\Omega_2} P f + \mathbf{1}_{\partial\Omega_2^+} (I - P)f + \mathbf{1}_{\partial\Omega_2^-} \phi_2] \quad (42)$$

$$- \sum_{i,k=1}^d \iint_{\Omega \times \mathbb{R}^3} (|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v_i v_k \partial_{ik} \phi_{\langle f, \Phi_0 \rangle_v} \langle f, \Phi_0 \rangle_v \Phi_0^2 dx dv \quad (43)$$

$$- \sum_{i,k=1}^d \iint_{\Omega \times \mathbb{R}^3} (|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v_i v_k \partial_{ik} \phi_{\langle f, \Phi_0 \rangle_v} \Phi_0 (\mathbf{I} - \mathbf{P}_L)f. \quad (44)$$

We make an orthogonal decomposition at the boundary  $\partial\Omega_2$ ,

$$v = (v \cdot n)n + v - (v \cdot n)n = v_n n + v_\perp.$$

The contribution of  $Pf = z(x)m^{\frac{1}{2}} = z(x)\Phi_0$  is

$$\begin{aligned} & \int_{\partial\Omega_2 \times \mathbb{R}^3} v_n(|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v \cdot \nabla_x \phi_{\langle f, \Phi_0 \rangle_v}(x) z(x) \Phi_0^2 dSdv \\ &= \int_{\partial\Omega_2 \times \mathbb{R}^3} v_n(|v|^2 - \beta_{\langle f, \Phi_0 \rangle_v}) v_n z(x) \partial_n \phi_{\langle f, \Phi_0 \rangle_v} \Phi_0^2 dSdv + \int_{\partial\Omega_2 \times \mathbb{R}^3} v_n(|v|^2 - \beta_a) v_{\perp} \cdot \nabla_x \phi_{\langle f, \Phi_0 \rangle_v} z(x) \Phi_0^2 dSdv. \end{aligned}$$

Notice that the boundary condition (38) reduces to the Neumann boundary condition on  $\partial\Omega_2$ . Therefore, the first term vanish. While, the second term also vanishes due to the oddness of  $v_n v_{\perp}$ . Therefore, (42) and (44) are bounded by

$$\|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(dx)} \|(I - P_L)(f)\|_{L^2_{0,\gamma/2}}$$

and

$$\|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(dx)} \left( \|(I - P)f\|_{L^2(\partial\Omega_2^+)} + \|f\|_{L^2(\partial\Omega_1^+)} + \|\phi_2\|_{L^2(\partial\Omega_2^-)} + \|f_{in}\|_{L^2(\partial\Omega_1^-)} \right).$$

For (43), if  $k \neq i$ , it vanishes due to the oddness. Hence, the only contribution comes from  $k = i$ .

$$- \sum_{i=1}^d \iint_{\Omega \times \mathbb{R}^3} (|v|^2 - \beta_a) (v_i)^2 \mu \partial_{ii} \phi_a a.$$

Using  $-\Delta_x \phi_{\langle f, \Phi_0 \rangle_v} = \langle f, \Phi_0 \rangle_v$  and  $\|\phi_{\langle f, \Phi_0 \rangle_v}\|_{H^2(dx)} \leq C \|\langle f, \Phi_0 \rangle_v\|_{L^2(dx)}$ , we obtain

$$\|\langle f, \Phi_0 \rangle_v\|_{L^2(dx)} \lesssim \left( \begin{aligned} & [1 + \frac{1}{\kappa}] \|(I - P_L)(f)\|_{L^2_{0,\gamma/2}} + \|\phi_1\|_{L^2_{0,-\gamma/2}} + \|(I - P)f\|_{L^2(\partial\Omega_2^+)} \\ & + \|f\|_{L^2(\partial\Omega_1^+)} + \|f_{in}\|_{L^2(\partial\Omega_1^-)} + \|\phi_2\|_{L^2(\partial\Omega_2^-)} \end{aligned} \right).$$

We therefore deduce the proposition.  $\square$

### 2.3. Proposition of $S_{\Omega}$ and $J$

Here we introduce a proposition from [25].

**Proposition 2.7.** *Let  $\Omega$  be a bounded convex domain with  $C^1$  boundary,  $\alpha \geq 0, \beta \in \mathbb{R}^3, -3 < \gamma \leq 1$  and  $2 \leq p \leq \infty$ . Suppose that (8) holds. Then, the operator norm of  $S_{\Omega}^{\kappa} : L^p_{\alpha,\beta} \rightarrow L^p_{\alpha,\beta+1}$  is  $\mathcal{O}(\kappa)$ .*

We also have the following estimate.

**Proposition 2.8.** *Let  $\Omega$  be a bounded convex domain with  $C^1$  boundary and  $-3 < \gamma \leq 1$ , Suppose  $f \in L^2(\partial\Omega_2^+)$ . Then,*

$$\|J^{\kappa}(f_{in} \mathbf{1}_{\partial\Omega_1^-} + (P(f) + \phi_2) \mathbf{1}_{\partial\Omega_2^-})\|_{L^2_{0,\gamma/2}} \leq C\kappa \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)} + \|f\|_{L^2(\partial\Omega_2^+)} + \|\phi_2\|_{L^2(\partial\Omega_2^-)} \right). \quad (45)$$

*Proof.* By Lemma 2.1 ([8]), we have

$$\begin{aligned} & \|J^{\kappa}(f_{in} \mathbf{1}_{\partial\Omega_1^-} + (P(f) + \phi_2) \mathbf{1}_{\partial\Omega_2^-})\|_{L^2_{0,\gamma/2}}^2 \\ &= \int_{\Omega} \int_{\mathbb{R}^3} e^{-2\frac{1}{\kappa}\nu(|v|)\tau_-(x,v)} \left[ (f_{in} \mathbf{1}_{\partial\Omega_1^-} + (P(f) + \phi_2) \mathbf{1}_{\partial\Omega_2^-})(q(x,v),v) \right]^2 (1+|v|)^{\gamma} dv dx \\ &= \int_{\Omega^-} \int_0^{\tau_-(x,-v)} e^{-2\frac{1}{\kappa}\nu(|v|)t} \left[ (f_{in} \mathbf{1}_{\partial\Omega_1^-} + (P(f) + \phi_2) \mathbf{1}_{\partial\Omega_2^-})(z,v) \right]^2 (1+|v|)^{\gamma} |v \cdot n(x)| dt dSdv \\ &= \kappa \int_{\Omega^-} \left[ (f_{in} \mathbf{1}_{\partial\Omega_1^-} + (P(f) + \phi_2) \mathbf{1}_{\partial\Omega_2^-})(z,v) \right]^2 \frac{(1+|v|)^{\gamma}}{\nu(|v|)} |v \cdot n(x)| dSdv \\ &\leq C\kappa \int_{\Omega^-} (f_{in}^2 \mathbf{1}_{\partial\Omega_1^-} + P^2(f) \mathbf{1}_{\partial\Omega_2^-} + \phi_2^2 \mathbf{1}_{\partial\Omega_2^-})(z,v) |v \cdot n(x)| dSdv \\ &\leq C\kappa \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \|P(f)\|_{L^2(\partial\Omega_2^-)}^2 \right) \\ &= C\kappa \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \|P(f)\|_{L^2(\partial\Omega_2^+)}^2 \right) \\ &\leq C\kappa \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \|f\|_{L^2(\partial\Omega_2^+)}^2 \right). \end{aligned}$$

This finish the proof.  $\square$

## 2.4. Estimate of $Q_+$ and $Q_-$

**Lemma 2.1.** For any  $\beta \geq 1$ ,

$$\int_{\mathbb{S}^2} \langle v' \rangle^{-\beta} \langle u' \rangle^{-\beta} d\sigma \leq C_\beta \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{-(\beta+1)}, \quad (46)$$

where  $v' = \frac{v+u}{2} + \frac{|v-u|}{2}\sigma$  and  $u' = \frac{v+u}{2} - \frac{|v-u|}{2}\sigma$ .

This statement can be found in [45]. We provide a proof below.

*Proof.* For convenience, let  $A = 1 + \left| \frac{v+u}{2} \right|^2 + \left| \frac{v-u}{2} \right|^2$  and  $B = 2 \left| \frac{v+u}{2} \right| \left| \frac{v-u}{2} \right|$ . When  $\beta = 1$ ,

$$\int_{\mathbb{S}^2} \langle v' \rangle^{-1} \langle u' \rangle^{-1} d\sigma = \int_0^{2\pi} \int_0^\pi \frac{\sin \theta}{\sqrt{A^2 - B^2 \cos^2 \theta}} d\theta d\phi.$$

Here we use sphere coordinate with  $e_1 = \frac{v+u}{|v+u|}$ . Let  $z = \cos \theta$  and  $y = \sin^{-1}(Bz/A)$ . We obtain

$$\begin{aligned} \int_0^{2\pi} \int_0^\pi \frac{\sin \theta}{\sqrt{A^2 - B^2 \cos^2 \theta}} d\theta d\phi &= 4\pi \int_0^1 \frac{1}{\sqrt{A^2 - B^2 z^2}} dz \\ &= 4\pi \int_0^{\sin^{-1}(B/A)} \frac{\cos(y)}{B|\cos(y)|} dy \\ &\leq 8\pi A^{-1} \\ &\leq \frac{16\pi}{1 + |v|^2 + |u|^2}. \end{aligned}$$

The penultimate inequality follows from  $\sin^{-1}(x) \leq 2x$  for any  $0 \leq x \leq 1$ . For  $\beta > 1$ , we first consider the case of  $\beta \in \mathbb{N}$ . We use induction to establish the result. Since  $\beta = 1$  holds by the above calculation. Assume that  $\beta = n$  holds. When  $\beta = n + 1$ , we have

$$\begin{aligned} \int_{\mathbb{S}^2} \langle v' \rangle^{-(n+1)} \langle u' \rangle^{-(n+1)} d\sigma &= 4\pi \int_0^{B/A} BA^{-n} \frac{1}{(\sqrt{1-y^2})^{n+1}} dy \\ &\leq \frac{4\pi}{A\sqrt{1-(B/A)^2}} \int_0^{B/A} BA^{1-n} \frac{1}{(\sqrt{1-y^2})^n} dy \\ &\leq \frac{C}{\sqrt{A^2 - B^2}} \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{n+1} \\ &\leq \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{n+2}. \end{aligned}$$

The second-to-last line follows from the induction hypothesis, so (46) holds when  $\beta \in \mathbb{N}$ . For  $\beta \notin \mathbb{N}$ , suppose that  $\beta = n + d$  for some  $n \in \mathbb{N}$  and  $d \in (0, 1)$ . Then, we have

$$\begin{aligned} 4\pi \int_0^{B/A} BA^{1-\beta} \frac{1}{(\sqrt{1-y^2})^\beta} dy &\leq \frac{4\pi}{(\sqrt{A^2 - B^2})^d} \int_0^{B/A} BA^{1-n} \frac{1}{(\sqrt{1-y^2})^n} dy \\ &\leq \frac{C}{(\sqrt{A^2 - B^2})^d} \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{-n-1} \\ &= C \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{-(\beta+1)}. \end{aligned}$$

□

Now, we provide estimates of  $Q_+, Q_-$ , which will be used in **section 5**.

**Proposition 2.9.** *Suppose that (8) holds. Then,*

$$\left| \int_{\mathbb{R}^3} |v| S_{\Omega}^{\kappa}(Q_+[f, f])(x, v) dv \right| \leq C_{\gamma, \beta} \|f\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}^2 \quad (47)$$

for any  $\beta > 5 + \max\{0, \gamma\}$  and

$$\left| \int_{\mathbb{R}^3} |v| S_{\Omega}^{\kappa}(Q_-[f, f])(x, v) dv \right| \leq C_{\gamma, \beta} \|f\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}^2 \quad (48)$$

for any  $\beta > 3 + \max\{0, \gamma\}$ .

*Proof.* The proof of (48) follows directly from

$$\|S^{\kappa}(f)\|_{L_{\alpha, \beta}^{\infty}(\Omega \times \mathbb{R}^3)} \leq \frac{C}{|v|} \|f\|_{L_{\alpha, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}. \quad (49)$$

The proof of (47) proceeds as follows:

$$\begin{aligned} \left| \int_{\mathbb{R}^3} |v| S_{\Omega}^{\kappa}(Q_+[f, f])(x, v) dv \right| &\leq C \left| \int_{\mathbb{R}^3} \|Q_+[f, f]\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)} dv \right| \\ &\leq \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - u|^{\gamma} \langle v' \rangle^{-\beta} \langle u' \rangle^{-\beta} d\sigma dudv \|f\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}^2 \\ &\leq C \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - u|^{\gamma} \left( \frac{1}{\sqrt{1 + |v|^2 + |u|^2}} \right)^{-(\beta+1)} dudv \|f\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}^2 \\ &\leq C_{\gamma, \beta} \|f\|_{L_{0, \beta}^{\infty}(\Omega \times \mathbb{R}^3)}^2. \end{aligned}$$

The second-to-last inequality follows from **Lemma (2.1)**. □

### Section 3. $L^2$ estimate

Before we begin the  $L^2$  estimate, we recall a lemma concerning boundary data. This lemma ensures that the trace of  $f$  is well-defined locally for a certain class of functions. The proof can be found in [15].

**Lemma 3.1** (Ukai Trace Theorem). Define

$$\gamma^{\epsilon} \equiv \{(x, v) \in \partial\Omega \times \mathbb{R}^3 : |n(x) \cdot v| \geq \epsilon, \epsilon \leq |v| \leq \frac{1}{\epsilon}\}. \quad (50)$$

Then

$$\|f 1_{\gamma^{\epsilon}}\|_{L^1(\partial\Omega \times \mathbb{R}^3)} \leq C_{\epsilon, \Omega} \left( \|f\|_{L^1(\Omega \times \mathbb{R}^3)} + \|v \cdot \nabla_x f\|_{L^1(\Omega \times \mathbb{R}^3)} \right). \quad (51)$$

Also, we recall Green's identity. Let  $\gamma$  denote the trace operator.

**Lemma 3.2.** *Assume that  $f(x, v), g(x, v) \in L^2(\Omega \times \mathbb{R}^3)$ ,  $v \cdot \nabla_x f, v \cdot \nabla_x g \in L^2(\Omega \times \mathbb{R}^3)$ , and  $\gamma f, \gamma g \in L^2(\partial\Omega \times \mathbb{R}^3)$ . Then*

$$\iint_{\Omega \times \mathbb{R}^3} \{(v \cdot \nabla_x f) g + (v \cdot \nabla_x g) f\} dv dx = \int_{\partial\Omega \times \mathbb{R}^3} \gamma f \gamma g dS dv. \quad (2.3)$$

The proof can be found in [6]. We now state the main theorem of this section. The statement and strategy are similar to [14] and [15]. However, we need to modify the proof due to the difference between the mixed and diffuse boundary conditions. Moreover, we use **Lemma 1.2** ([25]) to simplify the argument. We present the details below.

**Theorem 3.1.** *Let  $\Omega \in \mathbb{R}^3$  be a bounded convex domain with  $C^2$  boundary. Assume that  $-3 < \gamma \leq 1$  and that (8) holds. Then, for any  $\phi_1 \in L^2_{0, \max\{0, -\gamma/2\}}(\Omega \times \mathbb{R}^3)$ ,  $\phi_2 \in L^2_{0, \gamma/2}(\partial\Omega_1^-)$ , and  $f_{in} \in L^2_{0, \gamma/2}(\partial\Omega_2^-)$ . There exists a unique solution  $f \in L^2_{0, \gamma/2}$  to the following boundary value problem.*

$$\begin{cases} v \cdot \nabla_x f + \frac{1}{\kappa} L(f) = \phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ f = f_{in}, & \text{on } \partial\Omega_1^-, \\ f = P(f) + \phi_2(x, v), & \text{on } \partial\Omega_2^-. \end{cases} \quad (52)$$

Moreover,

$$\|f\|_{L^2_{0, \gamma/2}}^2 + \|f\|_{L^2(\partial\Omega_1^+)}^2 \leq C \max\{1, \kappa^2\} \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right), \quad (53)$$

*Proof.* We consider the following iteration with  $f_1 = 0$ ,

$$\begin{cases} \varpi f_{k+1} + v \cdot \nabla_x f_{k+1} + \frac{1}{\kappa} L(f_{k+1}) = \phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ f_{k+1} = f_{in}, & \text{on } \partial\Omega_1^-, \\ f_{k+1} = \eta P(f_k) + \phi_2(x, v), & \text{on } \partial\Omega_2^-, \end{cases} \quad (54)$$

where  $0 < \varpi, \eta < 1$ . Note that (54) differs slightly (52). More precisely, the boundary value problem (54) contains a penalization term  $\varpi f_{k+1}$  and the factor  $\eta$  inserted in front of  $P(f)$ . We first construct a solution to (54) in  $L^2_{0, \gamma/2}$  for arbitrary  $\varpi$  and  $\eta$  and denote this solution by  $f^{\varpi, \eta}$ . We then show that, for fixed  $\eta$ , the family  $\{f^{\varpi, \eta}\}$  converges in  $L^2_{0, \gamma/2}$ . Finally, we remove  $\eta$  by proving that  $\{f^\eta\}$  is a convergent sequence in  $L^2_{0, \gamma/2}$ , where  $f^\eta = \lim_{\varpi \rightarrow 0} f^{\varpi, \eta}$ . We emphasize that the penalization term is introduced in order to justify the application of Green's formula to (54) for all  $-3 < \gamma \leq 1$ . Consequently, when  $\gamma < 0$ , the procedure for removing  $\varpi$  and  $\eta$  are different from that in the case of  $\gamma \geq 0$ . Since the argument for the negative  $\gamma$  case relies on an inequality established in **section 4**, for the sake of clarity we present only the case  $\gamma \geq 0$  here. The case  $\gamma < 0$  can be treated by applying the same argument as in [14]. To construct the solution of (54), by Green formula, we have

$$\begin{aligned} \varpi \|f_{k+1}\|_{L^2}^2 + \frac{1}{2} \|f_{k+1}\|_{L^2(\partial\Omega_1^+)}^2 + \frac{1}{\kappa} \langle L(f_{k+1}), f_{k+1} \rangle \\ = \langle \phi_1, f_{k+1} \rangle + \frac{1}{2} \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \frac{1}{2} \|\eta P(f_k) + \phi_2\|_{L^2(\partial\Omega_2^-)}^2. \end{aligned} \quad (55)$$

In particular, we have

$$\|f_{k+1}\|_{L^2(\partial\Omega_1^+)}^2 \leq C \varpi \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|P(f_k)\|_{L^2(\partial\Omega_2^+)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (56)$$

Using **Proposition 2.8**, we deduce that

$$J^\kappa(f_{in}\chi_{\partial\Omega_1^-} + (P(f_k) + \phi_2)\chi_{\partial\Omega_2^-}) \in L^2_{0, \gamma/2} \quad \text{if } f_k \in L^2_{\Omega_2^+}. \quad (57)$$

We can then rewrite (54) in integral form as:

$$\left( I - \frac{1}{\kappa} S_\Omega^\kappa K \right) f_{k+1} = J^\kappa(f_{in}\chi_{\partial\Omega_1^-} + (P(f_k) + \phi_2)\chi_{\partial\Omega_2^-}) + S_\Omega \phi_1. \quad (58)$$

Applying **Lemma 1.2** ([25]), we establish the existence of  $f_{k+1}$ . By induction, we can then prove that  $f_{k+1}$  exists and belongs to  $L^2_{0, \gamma/2}$  for any  $k \in \mathbb{N}$ . Applying Yong's inequality, we have

$$\begin{aligned} \|\eta P(f_k) + \phi_2\|_{L^2(\partial\Omega_2^-)}^2 &= \eta^2 \|P(f_k)\|_{L^2(\partial\Omega_2^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + 2\eta \langle P(f_k), \phi_2 \rangle_{\partial\Omega_2^-} \\ &\leq \eta \|P(f_k)\|_{L^2(\partial\Omega_2^-)}^2 + \frac{1}{1-\eta} \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \\ &\leq \eta \|f_k\|_{L^2(\partial\Omega_2^+)}^2 + \frac{1}{1-\eta} \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \end{aligned} \quad (59)$$

Then, we combine the (55) and (59) to obtain

$$\begin{aligned} \|f_{k+1}\|_{L^2}^2 + (2\varpi)^{-1}\|f_{k+1}\|_{L^2(\partial\Omega^+)}^2 &\leq \eta \left( \|f_k\|_{L^2}^2 + (2\varpi)^{-1}\|f_k\|_{L^2(\partial\Omega^+)}^2 \right) \\ &\quad + C_{\varpi,\eta} \left( \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 \right) + C_{\varpi,\eta}\|\phi_2\|_{L^2(\partial\Omega_2^-)}^2. \end{aligned} \quad (60)$$

By same process, we have

$$\begin{aligned} \|f_{n+1} - f_{m+1}\|_{L^2}^2 + (2\varpi)^{-1}\|f_{n+1} - f_{m+1}\|_{L^2(\partial\Omega^+)}^2 \\ \leq \eta \left( \|f_n - f_m\|_{L^2}^2 + (2\varpi)^{-1}\|f_n - f_m\|_{L^2(\partial\Omega^+)}^2 \right). \end{aligned} \quad (61)$$

This implies that  $\{f_k\}$  is a Cauchy sequence. For any fixed  $\eta$  and  $\varpi$ , let  $\lim_{k \rightarrow \infty} f_k := f^{\varpi,\eta}$ . By applying Green's formula once more, we have

$$\begin{aligned} \varpi\|f^{\varpi,\eta}\|_{L^2}^2 + \frac{1}{2}\|f^{\varpi,\eta}\|_{L^2(\partial\Omega^+)}^2 + \frac{1}{\kappa}\langle L(f^{\varpi,\eta}), f^{\varpi,\eta} \rangle \\ = \langle \phi_1, f^{\varpi,\eta} \rangle + \frac{1}{2}\|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \frac{1}{2}\|\eta P(f^{\varpi,\eta}) + \phi_2\|_{L^2(\partial\Omega_2^-)}^2. \end{aligned} \quad (62)$$

Applying (35) we have

$$\|P_L(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 \leq C \left( \begin{aligned} &\varpi\|f^{\varpi,\eta}\|_{L^2}^2 + \left[1 + \frac{1}{\kappa}\right] \|(I - P_L)(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 + \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 \\ &+ \|f^{\varpi,\eta}\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)f^{\varpi,\eta}\|_{L^2(\partial\Omega_2^+)}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \end{aligned} \right). \quad (63)$$

Multiplying it with a constant  $a > 0$  and combining it with equation (62), we derive that

$$\begin{aligned} &\varpi(1 - aC)\|f^{\varpi,\eta}\|_{L^2}^2 + a\|P_L(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 + C_L\frac{1}{\kappa}\|(I - P_L)(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 + \frac{1}{2}\|f^{\varpi,\eta}\|_{L^2(\partial\Omega^+)}^2 \\ &\leq aC \left( \left[1 + \frac{1}{\kappa}\right] \|(I - P_L)(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 + \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|f^{\varpi,\eta}\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)f^{\varpi,\eta}\|_{L^2(\partial\Omega_2^+)}^2 \right) \\ &+ aC \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right) \\ &+ \frac{1}{2}\|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \frac{1}{2}\|\eta P(f^{\varpi,\eta}) + \phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \langle \phi_1, f^{\varpi,\eta} \rangle. \end{aligned}$$

Finally, by choosing  $a$  sufficiently small, we conclude that

$$\begin{aligned} &\|f^{\varpi,\eta}\|_{L^2_{0,\gamma/2}}^2 + \|f^{\varpi,\eta}\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)f^{\varpi,\eta}\|_{L^2(\partial\Omega_2^+)}^2 + C_L\|(I - P_L)(f^{\varpi,\eta})\|_{L^2_{0,\gamma/2}}^2 \\ &\leq C_\eta \cdot \kappa \left( \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \end{aligned} \quad (64)$$

Hence,  $f^{\varpi,\eta}$  is a bounded sequence in  $L^2_{0,\gamma/2}$  for any fixed  $\eta$ . Observe that  $f^{\eta,\varpi_1} - f^{\eta,\varpi_2}$  satisfies

$$\begin{cases} v \cdot \nabla_x (f^{\eta,\varpi_1} - f^{\eta,\varpi_2}) + \frac{1}{\kappa}L(f^{\eta,\varpi_1} - f^{\eta,\varpi_2}) = \varpi_1 f^{\eta,\varpi_1} - \varpi_2 f^{\eta,\varpi_2}, & \text{in } \Omega \times \mathbb{R}^3, \\ f^{\eta,\varpi_1} - f^{\eta,\varpi_2} = 0, & \text{on } \partial\Omega_1^-, \\ f^{\eta,\varpi_1} - f^{\eta,\varpi_2} = \eta P(f^{\eta,\varpi_1} - f^{\eta,\varpi_2}), & \text{on } \partial\Omega_2^-, \end{cases}$$

for any  $0 < \varpi_1, \varpi_2 < 1$ . After taking  $\varpi_1 f^{\eta,\varpi_1} - \varpi_2 f^{\eta,\varpi_2}$  as  $\phi_1$  and setting  $f_{in} = \phi_2 = 0$ , by Hölder's inequality, we have

$$\begin{aligned} &\|f^{\eta,\varpi_1} - f^{\eta,\varpi_2}\|_{L^2_{0,\gamma/2}}^2 + \|f^{\eta,\varpi_1} - f^{\eta,\varpi_2}\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)(f^{\eta,\varpi_1} - f^{\eta,\varpi_2})\|_{L^2(\partial\Omega_2^+)}^2 \\ &\leq C_\eta \cdot \kappa \left( \varpi_1\|f^{\eta,\varpi_1}\|_{L^2_{0,-\gamma/2}}^2 + \varpi_2\|f^{\eta,\varpi_2}\|_{L^2_{0,-\gamma/2}}^2 \right) \\ &\leq C_\eta \cdot \kappa \max\{\varpi_1, \varpi_2\} \left( \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \end{aligned} \quad (65)$$

This implies that  $f^\eta = \lim_{\varpi \rightarrow 0} f^{\varpi, \eta}$  exists in  $L^2_{0, \gamma/2}$  for any  $0 < \eta < 1$ . Moreover, applying the same procedure as in (64), we obtain

$$\begin{aligned} & \|f^\eta\|_{L^2_{0, \gamma/2}}^2 + \|f^\eta\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)f^\eta\|_{L^2(\partial\Omega_2^+)}^2 + C_L \|(I - P_L)(f)\|_{L^2_{0, \gamma/2}}^2 \\ & \leq C \cdot \kappa \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + C_\beta \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \beta \|P(f^\eta)\|_{L^2(\partial\Omega_2^-)}^2 \right), \end{aligned} \quad (66)$$

where  $\beta$  will be determined later. Furthermore, we have

$$v \cdot \nabla_x (f^\eta)^2 = -\frac{2}{\kappa} L(f^\eta) f^\eta + 2\phi_1 f^\eta. \quad (67)$$

Taking absolute values and integrating over  $\Omega \times \mathbb{R}^3$  in (67), and using (66), we deduce that

$$\|v \cdot \nabla_x (f^\eta)^2\|_{L^1} \leq C \max\{1, \kappa\} \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + C_\beta \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \beta \|P(f^\eta)\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (68)$$

Using **Lemma 3.1**, for any  $\gamma^\epsilon$  in (50) away from  $\partial\Omega_0$ , we obtain

$$\|f^\eta \mathbf{1}_{\gamma^\epsilon}\|_{L^2(\partial\Omega \times \mathbb{R}^3)}^2 \leq C_\epsilon \max\{1, \kappa\} \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + C_\beta \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \beta \|P(f^\eta)\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (69)$$

By the definition of  $P(f)$ , we can write  $P(f)(x, v) = g(x)m^{1/2}(v)$  for some  $g(x)$ . When  $\epsilon$  is sufficiently small,

$$\begin{aligned} \|P(f^\eta) \mathbf{1}_{\gamma^\epsilon}\|_{L^2(\partial\Omega_2^+)}^2 &= \int_{\partial\Omega_2} |g(x)|^2 \int_{n(x) \cdot v \geq \epsilon, |v| \leq \frac{1}{\epsilon}} m(v) |v \cdot n(x)| dv dx \\ &\geq \int_{\partial\Omega_2} |g(x)|^2 dx \times \frac{1}{2} \int_{v \cdot n(x) > 0} m(v) |v \cdot n(x)| dv \\ &= \frac{1}{2} \|P(f^\eta)\|_{L^2(\partial\Omega_2^+)}^2. \end{aligned} \quad (70)$$

Here, we have used the fact that

$$\begin{aligned} \int_{|n(x) \cdot v| \leq \epsilon} m(v) |n(x) \cdot v| dv &\leq \int_{-\epsilon'}^{\epsilon} e^{-v^2} |v| dv \int_{\mathbb{R}^2} e^{-|v_\perp|^2/2} dv_\perp \leq C\epsilon, \\ \int_{|v| \geq 1/\epsilon} m(v) |n(x) \cdot v| dv &\leq C\epsilon. \end{aligned} \quad (71)$$

We observe that

$$\|P(f^\eta)\|_{L^2(\partial\Omega_2^-)}^2 = \|P(f^\eta)\|_{L^2(\partial\Omega_2^+)}^2 \quad (72)$$

$$f^\eta = P(f^\eta) + (I - P)(f^\eta) \quad (73)$$

which yields

$$\|P(f^\eta) \mathbf{1}_{\gamma^\epsilon}\|_{L^2(\partial\Omega_2^+)}^2 \leq 2\|f^\eta \mathbf{1}_{\gamma^\epsilon}\|_{L^2(\partial\Omega_2^+)}^2 + 2\|(I - P)(f^\eta) \mathbf{1}_{\gamma^\epsilon}\|_{L^2(\partial\Omega_2^+)}^2. \quad (74)$$

Combining this estimate with (66), (69), (70) and (74), we conclude that

$$\|P(f^\eta)\|_{L^2(\partial\Omega_2^+)}^2 \leq C \max\{1, \kappa\} \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + C_\beta \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 + \beta \|P(f^\eta)\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (75)$$

Taking  $\beta$  sufficiently small and combining (66) with (75), we obtain

$$\|f^\eta\|_{L^2_{0, \gamma/2}}^2 + \|f^\eta\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)f^\eta\|_{L^2(\partial\Omega_2^+)}^2 \quad (76)$$

$$\leq C \max\{1, \kappa^2\} \left( \|\phi_1\|_{L^2_{0, -\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \quad (77)$$

We observe that

$$\begin{cases} v \cdot \nabla_x (f^{\eta_1} - f^{\eta_2}) + \frac{1}{\kappa} L(f^{\eta_1} - f^{\eta_2}) = 0, & \text{in } \Omega \times \mathbb{R}^3, \\ f^{\eta_1} - f^{\eta_2} = 0, & \text{on } \partial\Omega_1^-, \\ f^{\eta_1} - f^{\eta_2} = \eta_1 P(f^{\eta_1} - f^{\eta_2}) + (\eta_1 - \eta_2) P(f^{\eta_2}), & \text{on } \partial\Omega_2^-, \end{cases}$$

for any  $0 < \eta_1, \eta_2 < 1$ . By regarding  $(\eta_1 - \eta_2)P(f^{\eta_2})$  as  $\phi_2$ , we obtain

$$\|f^{\eta_1} - f^{\eta_2}\|_{L^2_{0,\gamma/2}}^2 + \|f^{\eta_1} - f^{\eta_2}\|_{L^2(\partial\Omega_1^+)}^2 + \|(I - P)(f^{\eta_1} - f^{\eta_2})\|_{L^2(\partial\Omega_2^+)}^2 \quad (78)$$

$$\begin{aligned} &\leq C \max\{1, \kappa^2\} (\eta_1 - \eta_2)^2 \|P(f^{\eta_2})\|_{L^2(\partial\Omega_2^-)}^2 \\ &\leq C \max\{1, \kappa^2\} \max\{\kappa^{-2}, \kappa^2\} (\eta_1 - \eta_2)^2 \left( \|\phi_1\|_{L^2_{0,-\gamma/2}}^2 + \|f_{in}\|_{L^2(\partial\Omega_1^-)}^2 + \|\phi_2\|_{L^2(\partial\Omega_2^-)}^2 \right). \end{aligned} \quad (79)$$

This means that  $\{f^\eta\}_\eta$  is a Cauchy sequence in  $L^2_{0,\gamma/2}$ . Therefore  $f := \lim_{\eta \rightarrow 1} f^\eta$  is the solution of (52).  $\square$

#### Section 4. $L^\infty$ estimate

In this section, we denote  $w(v) = e^{\alpha|v|^2}$ . Let  $h(x, v) = w(v)f(x, v)$ . Then,  $h$  satisfies

$$\begin{cases} v \cdot \nabla_x h + \frac{1}{\kappa} L_w(h) = w\phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ h = wf_{in}, & \text{on } \partial\Omega_1^-, \\ h = P_w(h) + w\phi_2(x, v), & \text{on } \partial\Omega_2^-, \end{cases} \quad (80)$$

where  $P_w(\cdot) := wP(w^{-1}\cdot)$  and  $L_w(\cdot) := wL(w^{-1}\cdot)$ . Furthermore, we denote

$$L_w(f) = \nu(v)f - K_\alpha(f)$$

by defining  $K_\alpha(\cdot) := wK(w^{-1}\cdot)$ . Note that  $K_\alpha$  and  $|K_\alpha|(f) := w \int_{\mathbb{R}^3} |k(v, u)| w^{-1} f(x, u) du$  have the same properties, where  $k(v, u)$  denotes the kernel of  $K$ . In this section, we prove that problem (80) admits a unique solution in  $L^\infty$  when  $\Omega$  is unsealed with respect to  $\partial\Omega_1$ , and therefore **Theorem 1.1** in **subsection 4.2**. First, we consider the approximating problem with the parameters  $\varpi$  and  $\eta$ , and we define a recursive sequence  $h^l$  with  $h^1 = 0$ .

$$\begin{cases} \varpi h^{l+1} + v \cdot \nabla_x h^{l+1} + \frac{1}{\kappa} L_w(h^{l+1}) = w\phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ h^{l+1} = wf_{in}, & \text{on } \partial\Omega_1^-, \\ h^{l+1} = \eta P_w(h^l) + w\phi_2, & \text{on } \partial\Omega_2^-. \end{cases} \quad (81)$$

Second, we establish the following proposition to relate the  $L^\infty$  norm of  $h^l$  to the weighted  $L^2$  norm of  $h^l/w$ . This relation allows us to apply the results from previous section to construct a solution to (81) in  $L^\infty$  and then remove the parameters  $\varpi$  and  $\eta$ . This approach is known as the  $L^2 - L^\infty$  estimate, which is a useful tool ([14], [15]) for constructing solutions of Boltzmann equation. Finally we note that **proposition 4.1** remains valid when  $\varpi = 0$  and  $\eta = 1$ . The statement of the proposition is

**Proposition 4.1.** *Let  $\Omega$  be a bounded set in  $\mathbb{R}^3$  with  $C^2$  boundary, and  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ . Suppose that  $h^{l+1}$  is the solution of (81) and (8) holds. Then, there exists  $n$  depending only on the constant of unsealed set  $\zeta$  such that*

$$\begin{aligned} &\|h^{l+1}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\ &\leq C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + \frac{1}{8} \max_{0 \leq m \leq 2(n-1)} \|h^{l-m}\|_{L^\infty(\Omega \times \mathbb{R}^3)} + C_\zeta \min\{1, \kappa^{-1}\} \max_{0 \leq m \leq 2(n-1)} \|h^{l+1-m} w^{-1}\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)}. \end{aligned} \quad (82)$$

for any  $l \geq 2n$ .

To prove this proposition we need the following estimate for  $h^l$ .

$$|h^{l+1}|(x, v) \leq \frac{1}{\kappa} S_{\Omega}^{\kappa} |K_{\alpha}| (|h^{l+1}|) + S_{\Omega}^{\kappa} (|w\phi_1|) + J^{\kappa} \mathbf{1}_{\partial\Omega_1^-} |w f_{in}| + J^{\kappa} \mathbf{1}_{\partial\Omega_2^-} |w\phi_2| + J^{\kappa} \mathbf{1}_{\partial\Omega_2^-} P_w (|h^l|). \quad (83)$$

After iterating for  $n$  times, we obtain

$$|h^{l+1}|(x, v) \leq \sum_{k=1}^4 A_k(x, v) + \sum_{j=1}^{n-1} \hat{P}^j(A_1^j)(x, v) + \sum_{k=2}^4 \sum_{j=1}^{n-1} \hat{P}^j(A_k)(x, v) + \hat{P}^n(|h^{l+1-n}|)(x, v). \quad (84)$$

Here the definitions of  $A_k^j$  and  $\hat{P}^j$  are given as follows.

1.  $A_1^j(x, v) = \frac{1}{\kappa} S_{\Omega}^{\kappa} |K_{\alpha}| (|h^{l+1-j}|)$ ,  $A_1(x, v) = A_1^0$ .
2.  $A_2(x, v) = S_{\Omega}^{\kappa} (|w\phi_1|)$ .
3.  $A_3(x, v) = J_1^{\kappa} |w f_{in}|(x, v) = \mathbf{1}_{\partial\Omega_1^-} (q(x, v)) e^{-\frac{1}{\kappa} \nu(v) \tau_-(x, v)} |w(v) f_{in}|(q(x, v), v)$ .
4.  $A_4(x, v) = J_2^{\kappa} |w\phi_2|(x, v) = \mathbf{1}_{\partial\Omega_2^-} (q(x, v)) e^{-\frac{1}{\kappa} \nu(v) \tau_-(x, v)} |w(v) \phi_2|(q(x, v), v)$ .
5.  $\hat{P}(h)(x, v) = J_2^{\kappa} w P(w^{-1}h) = J_2^{\kappa} \left( W \int h w^{-1} m^{-1/2} d\sigma_{0,1} \right) (x, v)$ .
6.  $\hat{P}^j(h)(x, v) = J_2^{\kappa} \left( W \int \cdots \int h w^{-1} m^{-1/2} d\sigma_{j-1,j} d\sigma_{j-2,j-1} \cdots d\sigma_{0,1}^J \right) (x, v) \quad \forall j \geq 2$ .

The definitions of the notations appearing above are as follows

1.  $W := w(v) m^{1/2}(v)$ ,  $x_0 = q(x, v)$ ,  $x_j := q(x_{j-1}, v_j)$  for all  $j \in \mathbb{N}$ .
2.  $J_i^{\kappa} := \mathbf{1}_{\partial\Omega_i} J^{\kappa}$  for  $i = 1, 2$ .
3.  $\left( \int h d\sigma_{j-1,j} \right) (x_{j-1}, v_{j-1}) := \int_{v_j \cdot n(x_{j-1}) > 0} h(x_{j-1}, v_j) m(v_j) v_j \cdot n(x_{j-1}) dv_j \quad \forall j \geq 1$ .
4.  $\left( \int h d\sigma_{j-1,j}^1 \right) (x_{j-1}, v_{j-1}) := \int_{v_j \cdot n(x_{j-1}) > 0} h(x_j, v_j) \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_j) m(v_j) v_j \cdot n(x_{j-1}) dv_j \quad \forall j \geq 1$ .
5.  $\left( \int h d\sigma_{j-1,j}^J \right) (x_{j-1}, v_{j-1}) := \int_{v_j \cdot n(x_{j-1}) > 0} J_2^{\kappa} h(x_{j-1}, v_j) m(v_j) v_j \cdot n(x_{j-1}) dv_j \quad \forall j \geq 1$ .

#### 4.1. The existence of solution to (80)

We now introduce the definition of an unsealed set:

**Definition 4.1.** Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with  $C^2$  boundary, and let  $\partial\Omega_1$  be an open subset of  $\partial\Omega$ . We say that  $\Omega$  is **unsealed** with respect to  $\partial\Omega_1$  if there exist  $0 < \zeta < 1$  and  $m \in \mathbb{N}$  such that

$$\int \cdots \int \mathbf{1} d\sigma_{m-1,m}^1 \cdots d\sigma_{0,1}^1 \leq \zeta \quad \forall x \in \partial\Omega \setminus \partial\Omega_1, \quad (85)$$

In particular,  $\Omega$  is said to be **completely unsealed** with respect to  $\partial\Omega_1$  when  $m = 1$ .

**Remark 4.1.** The  $m$  can be replaced by  $m_T$  in **Definition 4.1**. Additional information about unsealed set is given in **Appendix 7**.

Then, we prove **proposition 4.1** and state our main goal of this section. The statement is as follows:

**Theorem 4.1.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded convex domain with  $C^2$  boundary and suppose  $\Omega$  is unsealed with respect to  $\partial\Omega_1$ . Assume that  $-3 < \gamma \leq 1$ ,  $0 < \alpha < \frac{1}{4}$  and (8) holds. Then, for any  $\phi_1 \in L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)$ ,  $f_{in} \in L_{\alpha,0}^\infty(\partial\Omega_1)$ , and  $\phi_2 \in L_{\alpha,0}^\infty(\partial\Omega_2)$ , there exists a unique solution  $h \in L_{0,0}^\infty(\Omega \times \mathbb{R}^3)$  to (80). Furthermore, the solution  $h$  satisfies following estimate*

$$\|h\|_{L^\infty(\Omega \times \mathbb{R}^3)} + \|h\|_{L^\infty(\partial\Omega_2^+)} \leq C_\zeta \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \max\{1, \kappa\} \|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \right). \quad (86)$$

For clarity, we present the proof for the case of  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ . The proof for unsealed case is similar. In order to achieve our goal, we need the following lemmas to establish **proposition 4.1**.

**Lemma 4.1.** *Let  $-3 < \gamma \leq 1$  and assume that (8) holds. Then*

$$\begin{aligned} |h^{l+1}|(x, v) &\leq C_\zeta \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + \frac{\zeta^{n-1}}{1-\zeta} \max_{0 \leq j \leq n-1} \|h^{l+1-j-n}\|_{L^\infty(\Omega \times \mathbb{R}^3)} + \left( \frac{1}{\kappa} \right)^{2n-1} \sum_{j=0}^{n-1} \sum_{i=0}^{n-1} \hat{P}^j S_\Omega^\kappa |K_\alpha| \hat{P}^i S_\Omega^\kappa |K_\alpha| (|h^{l+1-j-i}|)(x, v) \end{aligned} \quad (87)$$

for any  $(x, v) \in \bar{\Omega} \times \mathbb{R}^3$ .

*Proof.* For  $A_3$  and  $A_4$ , because  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ , we have

$$\begin{cases} \|A_3\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq C \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)}, \\ \|A_4\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq C \|w(v)\phi_2\|_{L^\infty(\partial\Omega_2^-)}. \end{cases} \quad (88)$$

Also, similar to (88), we obtain

$$\begin{cases} \left\| \sum_{j=1}^{n-1} \hat{P}^j(A_3) \right\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq C \frac{1}{1-\zeta} \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)}, \\ \left\| \sum_{j=1}^{n-1} \hat{P}^j(A_4) \right\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq C \frac{1}{1-\zeta} \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)}. \end{cases} \quad (89)$$

Using the **Proposition 2.2**, we obtain

$$\|A_2\|_{L^\infty(\Omega \times \mathbb{R}^3)} \leq C \kappa \| \langle v \rangle^{-1} w(v) \phi_1 \|_{L^\infty(\partial\Omega_1^-)} \quad (90)$$

and

$$\left\| \sum_{j=1}^{n-1} \hat{P}^j(A_2) \right\|_{L^\infty(\Omega \times \mathbb{R}^3)} \leq C \frac{\kappa}{1-\zeta} \|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)}. \quad (91)$$

Applying that  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ , we again yield

$$\|\hat{P}^n(h^{l+1-n})\|_{L^\infty(\Omega \times \mathbb{R}^3)} \leq C \zeta^{n-1} \|h^{l+1-n}\|_{L^\infty(\Omega \times \mathbb{R}^3)}. \quad (92)$$

Combine the above result, we conclude that

$$\begin{aligned} |h^{l+1}| &\leq C \frac{1}{1-\zeta} \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + \frac{1}{\kappa} \sum_{j=0}^{n-1} \hat{P}^j S_\Omega^\kappa |K_\alpha| (|h^{l+1-j}|) + \zeta^{n-1} \|h^{l+1-n}\|_{L^\infty(\Omega \times \mathbb{R}^3)}. \end{aligned} \quad (93)$$

For each  $0 < j < n-1$ , replace  $l+1$  to  $l+1-j$ , we have

$$\begin{aligned} |h^{l+1-j}| &\leq C \frac{1}{1-\zeta} \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + \frac{1}{\kappa} \sum_{i=0}^{n-1} \hat{P}^i S_\Omega^\kappa |K_\alpha| (|h^{l+1-j-i}|) + \zeta^{n-1} \|h^{l+1-j-n}\|_{L^\infty(\Omega \times \mathbb{R}^3)}. \end{aligned} \quad (94)$$

Substitute  $h^{l+1-j}$  by 94, we obtain

$$\begin{aligned}
|h^{l+1}| &\leq C_\zeta \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \kappa\|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega\times\mathbb{R}^3)} \right) \\
&\quad + \frac{1}{\kappa} \sum_{j=0}^{n-1} \zeta^j \left\| S_\Omega^\kappa |K_\alpha| \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} \right) \right\|_{L^\infty(\Omega\times\mathbb{R}^3)} \\
&\quad + \frac{1}{\kappa} \sum_{j=0}^{n-1} \zeta^j \left\| S_\Omega^\kappa |K_\alpha| \left( \kappa\|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega\times\mathbb{R}^3)} + \zeta^{n-1}\|h^{l+1-j-n}\|_{L^\infty(\Omega\times\mathbb{R}^3)} \right) \right\|_{L^\infty(\Omega\times\mathbb{R}^3)} \\
&\quad + \left( \frac{1}{\kappa} \right)^2 \sum_{j=0}^{n-1} \sum_{i=0}^{n-1} \hat{P}^j S_\Omega^\kappa |K_\alpha| \hat{P}^i S_\Omega^\kappa |K_\alpha| (|h^{l+1-j-i}|) \\
&\leq C_\zeta \left( \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} + \|\phi_2\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} + \kappa\|\phi_1\|_{L_{\alpha,-1}^\infty(\Omega\times\mathbb{R}^3)} \right) \\
&\quad + \frac{\zeta^{n-1}}{1-\zeta} \max_{0\leq j\leq n-1} \|h^{l+1-j-n}\|_{L^\infty(\Omega\times\mathbb{R}^3)} + \left( \frac{1}{\kappa} \right)^2 \sum_{j=0}^{n-1} \sum_{i=0}^{n-1} \hat{P}^j S_\Omega^\kappa |K_\alpha| \hat{P}^i S_\Omega^\kappa |K_\alpha| (|h^{l+1-j-i}|). \quad (95)
\end{aligned}$$

□

For convenience, we define

$$A_{ij}(f) := \left( \frac{1}{\kappa} \right)^2 \hat{P}^j S_\Omega^\kappa |K_\alpha| \hat{P}^i S_\Omega^\kappa |K_\alpha|(f) \quad (97)$$

and denote  $\mathcal{D} = \text{diam}(\Omega)$ . Now, we prove the following lemma

**Lemma 4.2.** *Suppose  $\Omega$  is a bounded set in  $\mathbb{R}^3$  with  $C^2$  boundary and  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ . Given  $\epsilon > 0$  and  $i, j \in \mathbb{N}$ . Suppose  $-3 < \gamma \leq 1$  and (8) holds. Then*

$$\|A_{ij}(f)\|_{L^\infty(\Omega\times\mathbb{R}^3)} \leq C\zeta^{j+i-2} \left( \epsilon\|f\|_{L^\infty(\Omega\times\mathbb{R}^3)} + C_\epsilon \min \left\{ 1, \frac{1}{\kappa} \right\} \|f/w\|_{L_{0,\gamma/2}^2(\Omega\times\mathbb{R}^3)} \right) \quad (98)$$

*Proof.* Without loss of generality, we assume  $f \geq 0$ . For  $i = j = 0$ , we use properties of  $|k_\alpha|$  like 29, proposition 2.6, and proposition 2.5. By definition,

$$A_{00} = \left( \frac{1}{\kappa} \right)^2 \int_0^{\tau_-(x,v)} \int_{\mathbb{R}^3} \int_0^{\tau_-(x-tv,u)} \int_{\mathbb{R}^3} e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds dudt. \quad (99)$$

Fixed some  $\epsilon_1, \epsilon_2 > 0$  and  $N, M \in \mathbb{N}$ , we define

$$A = \{|v-u| > \epsilon_2, |u| < N\} \text{ and } B = \{|u-w| > \epsilon_1\}. \quad (100)$$

Moreover, we assume that  $f \geq 0$ . Then, we have

$$\begin{aligned}
& A_{00} \tag{101} \\
& \leq C\epsilon_1^{3-|\gamma|} \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + \left(\frac{1}{\kappa}\right)^2 \int_0^{\tau_-(x,v)} \int_{\mathbb{R}^3} \int_0^{\tau_-(x-tv,u)} \int_B e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds du dt \\
& \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + \left(\frac{1}{\kappa}\right)^2 \int_0^{\tau_-(x,v)} \int_{\{|v-u|>\epsilon_2\}} \int_0^{\tau_-(x-tv,u)} \int_B e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds du dt \\
& \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|} + \frac{1}{1+N}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + \left(\frac{1}{\kappa}\right)^2 \int_0^{\tau_-(x,v)} \int_A \int_0^{\tau_-(x-tv,u)} \int_B e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds du dt \\
& \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|} + \frac{1}{1+N} + \frac{1}{N}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + \left(\frac{1}{\kappa}\right)^2 \int_0^{\tau_-(x,v)} \int_A \int_{1/N}^{\tau_-(x-tv,u)} \int_B e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds du dt. \tag{102}
\end{aligned}$$

Using Hölder inequality and noticing that  $k(v,u), k(u,w)$  are bounded on  $A, B$  respectively. Let  $y = x - tv - su$ , we obtain

$$\begin{aligned}
A_{00} & \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|} + \frac{1}{1+N} + \frac{1}{N} + \frac{1}{1+M}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + C\mathbf{1}_{|v|<M} \left(\frac{1}{\kappa}\right)^2 \int_{\frac{D}{5M}}^{\frac{D}{|v|}} \int_A \int_{1/N}^\infty \int_B e^{-\frac{1}{\kappa}\nu(v)t} |k_\alpha|(v,u) e^{-\frac{1}{\kappa}\nu(u)s} |k_\alpha|(u,w) f(x-tv-su,w) dw ds du dt \\
& \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|} + \frac{1}{1+N} + \frac{1}{N} + \frac{1}{M}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\
& + C\left(\frac{1}{\kappa}\right)^2 \mathbf{1}_{|v|<M} \left(\int_{\frac{D}{5M}}^{\frac{D}{|v|}} \int_{\mathbb{R}^3} \int_{1/N}^\infty \int_{\mathbb{R}^3} e^{-\frac{C_M}{\kappa}t} \frac{1}{s^3} \nu(w) f^2(y,w) w^{-2}(w) dw ds dy dt\right)^{1/2} \\
& \leq C\left(\epsilon_1^{3-|\gamma|} + \epsilon_2^{3-|\gamma|} + \frac{1}{1+N} + \frac{1}{N} + \frac{1}{M}\right) \|f\|_{L^\infty(\Omega \times \mathbb{R}^3)} + C \min\left\{1, \frac{1}{\kappa}\right\} \|f/w\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)}. \tag{103}
\end{aligned}$$

Hence, if  $\epsilon_1, \epsilon_2$  are sufficiently small and  $N, M$  are chosen large enough, the proof for  $A_{00}$  is complete. For  $i$  or  $j \neq 0$ , since  $wm^{1/2} \leq C$  and  $\Omega$  is completely unsealed with respect to  $\partial\Omega_1$ , it suffice to estimate  $S_\Omega^\kappa |K_\alpha| \hat{P}^i S_\Omega^\kappa |K_\alpha|(f)$ . Furthermore, we can observe that the character of  $|K_\alpha|$  can be replace by  $\hat{P}^i$ . Therefore, the estimate  $A_{ij}$  is similar to  $A_{00}$  and it is sufficient to estimate  $\frac{1}{\kappa} \hat{P}^i S_\Omega^\kappa |K_\alpha|(f)$ . For instance, let  $i = 1$  and denote  $q = q(x, v)$ ,

$$\begin{aligned}
& \hat{P} S_\Omega^\kappa |K_\alpha|(f)(x, v) \tag{104} \\
& = \frac{1}{\kappa} \int W(v) J_2^-(q(x, v), v) S_\Omega^\kappa |K_\alpha|(f)(q(x, v), v_1) w^{-1}(v_1) d\sigma_{0,1} \\
& \leq \frac{1}{\kappa} \int_{v_1 \cdot n(q) > 0} \int_0^{\tau_-(q, v_1)} \int_{\mathbb{R}^3} e^{-\frac{1}{\kappa}\nu(v_1)t} |k_\alpha|(v_1, u) f(q - tv_1, u) w^{-1}(v_1) m^{1/2}(v_1) v_1 \cdot n(q) du dt dv_1 \\
& = \frac{1}{\kappa} \int_{v_1 \cdot n(q) > 0} \int_0^{\tau_-(q, v_1)} \int_{\mathbb{R}^3} e^{-\frac{1}{\kappa}\nu(v_1)t} |k|(v_1, u) f(q - tv_1, u) w^{-1}(u) m^{1/2}(v_1) v_1 \cdot n(q) du dt dv_1.
\end{aligned}$$

Similar to  $A_{00}$ , fix some  $\epsilon_1 > 0$  and  $N \in \mathbb{N}$ , we define

$$A = \{|v_1| < N\}, \quad B = \{|v_1 - u| > \epsilon_1\}. \tag{105}$$

Then,

$$\begin{aligned}
& \hat{P}S_{\Omega}^{\kappa}|K_{\alpha}|(f)(x, v) \tag{106} \\
& \leq C\epsilon_1^{3-|\gamma|}\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \\
& \quad + \frac{1}{\kappa}\int_{v_1\cdot n(q)>0}\int_0^{\frac{D}{|v_1|}}\int_B e^{-\frac{1}{\kappa}\nu(v_1)t}|k|(v_1, u)f(q-tv_1, u)w^{-1}(u)m^{1/2}(v_1)v_1\cdot n(q)dudtdv_1 \\
& \leq C\left(\epsilon_1^{3-|\gamma|}+\frac{1}{1+N}\right)\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \\
& \quad + \frac{1}{\kappa}\int_{\{v_1\cdot n(q)>0\}\cap A}\int_0^{\frac{D}{|v_1|}}\int_B e^{-\frac{1}{\kappa}\nu(v_1)t}|k|(v_1, u)f(q-tv_1, u)w^{-1}(u)m^{1/2}(v_1)v_1\cdot n(q)dudtdv_1. \tag{107}
\end{aligned}$$

Using Hölder inequality and notice that  $k(v_1, u)$  is bounded on  $B$ . Let  $y = q(x, v) - tv_1$ . We obtain

$$\begin{aligned}
& \leq C\left(\epsilon_1^{3-|\gamma|}+\frac{1}{1+N}\right)\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \tag{108} \\
& \quad + \frac{1}{\kappa}\int_{\{v_1\cdot n(q)>0\}\cap A}\int_{\frac{D}{N}}^{\frac{D}{|v_1|}}\int_B e^{-\frac{1}{\kappa}\nu(v_1)t}|k|(v_1, u)f(q-tv_1, u)w^{-1}(u)m^{1/2}(v_1)v_1\cdot n(q)dudtdv_1 \\
& \leq C\left(\epsilon_1^{3-|\gamma|}+\frac{1}{1+N}\right)\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \\
& \quad + \frac{C}{\kappa}\left(\int_{\{v_1\cdot n(q)>0\}\cap A}\int_{\frac{D}{N}}^{\frac{D}{|v_1|}}\int_B e^{-\frac{C_N}{\kappa}t}|k|(v_1, u)f^2(q-tv_1, u)w^{-2}(u)dudtdv_1\right)^{1/2} \tag{109}
\end{aligned}$$

$$\begin{aligned}
& \leq C\left(\epsilon_1^{3-|\gamma|}+\frac{1}{1+N}\right)\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \\
& \quad + C\min\left\{1, \frac{1}{\kappa}\right\}\left(\int_{\Omega}\int_{\frac{D}{N}}^{\infty}\int_{\mathbb{R}^3}\frac{1}{t^3}\nu(u)f^2(y, u)w^{-2}(u)dudtdy\right)^{1/2} \\
& \leq C\left(\epsilon_1^{3-|\gamma|}+\frac{1}{1+N}\right)\|f\|_{L^{\infty}(\Omega\times\mathbb{R}^3)}+C\min\left\{1, \frac{1}{\kappa}\right\}\|f/w\|_{L^2_{0, \gamma/2}(\Omega\times\mathbb{R}^3)}. \tag{110}
\end{aligned}$$

Hence, when  $\epsilon_1$  is sufficiently small and  $N$  is sufficiently large, the proof for  $A_{ij}$  is complete.  $\square$

*Proof. The proof of property 4.1*

We use **Lemma 4.1** and **Lemma 4.2** to obtain

$$\begin{aligned}
& |h^{l+1}|(x, v) \tag{111} \\
& \leq C_{\zeta}\left(\|f_{in}\|_{L^{\infty}_{\alpha, 0}(\partial\Omega_1^-)}+\|\phi_2\|_{L^{\infty}_{\alpha, 0}(\partial\Omega_2^-)}+\kappa\|\phi_1\|_{L^{\infty}_{\alpha, -1}(\Omega\times\mathbb{R}^3)}\right) \\
& \quad + \frac{\zeta^{n-1}}{1-\zeta}\max_{0\leq j\leq n-1}\|h^{l+1-j-n}\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \\
& \quad + \frac{C}{\zeta^2(1-\zeta)^2}\left(\epsilon\max_{0\leq i, j\leq n-1}\|h^{l+1-i-j}\|_{L^{\infty}(\Omega\times\mathbb{R}^3)}+C_{\epsilon}\min\left\{1, \frac{1}{\kappa}\right\}\max_{0\leq i, j\leq n-1}\|h^{l+1-i-j}/w\|_{L^2_{0, \gamma/2}(\Omega\times\mathbb{R}^3)}\right).
\end{aligned}$$

Taking  $n$  is enough large and  $\epsilon$  is sufficiently small, we have

$$\begin{aligned}
& \|h^{l+1}\|_{L^{\infty}(\Omega\times\mathbb{R}^3)} \tag{112} \\
& \leq C_{\zeta}\left(\|f_{in}\|_{L^{\infty}_{\alpha, 0}(\partial\Omega_1^-)}+\|\phi_2\|_{L^{\infty}_{\alpha, 0}(\partial\Omega_2^-)}+\kappa\|\phi_1\|_{L^{\infty}_{\alpha, -1}(\Omega\times\mathbb{R}^3)}\right) \\
& \quad + \frac{1}{8}\max_{0\leq m\leq 2(n-1)}\|h^{l-m}\|_{L^{\infty}(\Omega\times\mathbb{R}^3)}+C_{\zeta}\min\left\{1, \frac{1}{\kappa}\right\}\max_{0\leq m\leq 2(n-1)}\|h^{l+1-m}/w\|_{L^2_{0, \gamma/2}(\Omega\times\mathbb{R}^3)}.
\end{aligned}$$

The proof is complete.  $\square$

*Proof. The proof of Theorem 4.1*

Now, we construct the solution of (80) by starting from (81). It is clear that **Property 4.1** still holds when  $\varpi = 0$  or  $\eta = 1$ . We rewrite (112) as

$$\begin{aligned} & \|h^{i+2(n-1)+1}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\ & \leq C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + \frac{1}{8} \max_{0 \leq m \leq 2(n-1)} \|h^{i+m}\|_{L^\infty(\Omega \times \mathbb{R}^3)} + C_\zeta \min \left\{ 1, \frac{1}{\kappa} \right\} \max_{0 \leq m \leq 2(n-1)} \|h^{i+1+m}/w\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)}, \end{aligned} \quad (113)$$

for any  $i = 0, 1, \dots$ . By **Lemma 4.3** with  $a_i = \|h^i\|_{L^\infty(\Omega \times \mathbb{R}^3)}$  and  $k = 2(n-1)$  and notice that

$$\begin{aligned} \|h^i/w\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} & = \|f^i\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \\ & \leq C_\eta \max\{1, \kappa\} \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)} + \|\phi_2\|_{L^2(\partial\Omega_1^-)} + \|\phi_1\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \right), \end{aligned} \quad (114)$$

for any  $i \in \mathbb{N}$ . Then

$$\begin{aligned} \max_{0 \leq m \leq 2(n-1)} \|h^{i+m}\|_{L^\infty(\Omega \times \mathbb{R}^3)} & \leq \left( \frac{1}{8} \right)^{\lfloor \frac{i}{2n-1} \rfloor} \max_{0 \leq m \leq 4(n-1)} \|h^m\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\ & \quad + C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + C_{\varpi,\eta} \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)} + \|\phi_2\|_{L^2(\partial\Omega_1^-)} + \|\phi_1\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \right). \end{aligned} \quad (115)$$

The statement of **Lemma 4.3** is

**Lemma 4.3** ([14](lemma 6.1)). *Let  $\{a_i\}_{i=0} \in \mathbb{R}_+$ . For any fixed  $k \in \mathbb{N}$ , we denote  $A_i^k = \max\{a_i, a_{i+1}, \dots, a_{i+k}\}$ .*

1. *Assume  $D \geq 0$ . If  $a_{i+1+k} \leq \frac{1}{8}A_i^k + D$  for  $i = 0, 1, \dots$ , then it holds that*

$$A_i^k \leq \left( \frac{1}{8} \right)^{\lfloor \frac{i}{k+1} \rfloor} \max\{A_0^k, A_1^k, \dots, A_k^k\} + \frac{8+k}{7}D, \quad \text{for } i \leq k+1 \quad (116)$$

2. *Let  $0 \leq \eta < 1$  with  $\eta^{k+1} \geq \frac{1}{4}$ . If  $a_{i+1+k} \leq \frac{1}{8}A_i^k + C_k \eta^{i+k+1}$  for  $i = 0, 1, \dots$ , then it holds that*

$$A_i^k \leq \left( \frac{1}{8} \right)^{\lfloor \frac{i}{k+1} \rfloor} \max\{A_0^k, A_1^k, \dots, A_k^k\} + 2C_k \frac{8+k}{7}D, \quad \text{for } i \leq k+1 \quad (117)$$

The proof of **Lemma 4.3** is omitted as it appears in the appendix of [14]. Now, we should control  $\max_{0 \leq m \leq 4(n-1)} \|h^m\|_{L^\infty(\Omega \times \mathbb{R}^3)}$ . We can use same techniques in the proof of **Property 4.1** and induction to show that

$$\begin{aligned} \|h^m\|_{L^\infty(\Omega \times \mathbb{R}^3)} & \leq C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + C_{\zeta,\varpi,\eta} \left( \|f_{in}\|_{L^2(\partial\Omega_1^-)} + \|\phi_2\|_{L^2(\partial\Omega_1^-)} + \|\phi_1\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \right), \end{aligned} \quad (118)$$

for  $0 \leq m \leq 4(n-1)$ . Here,  $C_{n,\eta}, C_{n,\zeta}$  are increasing with respect to  $n$ . In particular, combining (115) and (118), we have

$$\begin{aligned} \|h^l\|_{L^\infty(\Omega \times \mathbb{R}^3)} & \leq \left( \frac{1}{8} \right)^{\lfloor \frac{l}{2n-1} \rfloor} C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + C_{\zeta,\varpi,\eta} \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right), \end{aligned} \quad (119)$$

since  $L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3) \subset L^\infty_{\alpha,0}(\Omega \times \mathbb{R}^3)$ .

Now, we show that  $\{h^l\}$  is a Cauchy sequence in  $L^\infty(\Omega \times \mathbb{R}^3)$ . Consider  $2n \ll m_1, m_2$ , we notice that  $h^{m_1+1} - h^{m_2+1}$  satisfies

$$\begin{cases} \varpi\nu(v)(h^{m_1+1} - h^{m_2+1}) + v \cdot \nabla_x (h^{m_1+1} - h^{m_2+1}) + \frac{1}{\kappa} L_w (h^{m_1+1} - h^{m_2+1}) = 0, & \text{in } \Omega \times \mathbb{R}^3, \\ (h^{m_1+1} - h^{m_2+1}) = 0, & \text{on } \partial\Omega_1^-, \\ (h^{m_1+1} - h^{m_2+1}) = \eta P_w (h^{m_1} - h^{m_2}), & \text{on } \partial\Omega_2^-. \end{cases}$$

By the same procedure as in **Property 4.1**, we obtain

$$\begin{aligned} \|h^{m_1+1} - h^{m_2+1}\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq \frac{1}{8} \max_{0 \leq m \leq 2(n-1)} \|h^{m_1-m} - h^{m_2-m}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\ &\quad + C_\zeta \min\{1, \kappa^{-1}\} \max_{0 \leq m \leq 2(n-1)} \|f^{m_1+1-m} - f^{m_2+1-m}\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)}. \end{aligned} \quad (120)$$

Note that  $\{f_m\}$  converges in  $L^2_{0,\gamma/2}$  (see the proof of **Theorem 3.1**). Given  $\epsilon > 0$ , by definition, there exists  $M$  such that for any  $m_1, m_2 > M$ , we have

$$C_\zeta \max_{0 \leq m \leq 2(n-1)} \|f^{m_1+1-m} - f^{m_2+1-m}\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} < \frac{\epsilon}{2(2n+7)/7}. \quad (121)$$

we then obtain

$$\|h^{m_1+1} - h^{m_2+1}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \leq \frac{1}{8} \max_{0 \leq m \leq 2(n-1)} \|h^{m_1-m} - h^{m_2-m}\|_{L^\infty(\Omega \times \mathbb{R}^3)} + \epsilon. \quad (122)$$

By (119), we find that

$$\|h^{m_1} - h^{m_2}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \leq C_{f_{in}, \phi_1, \phi_2} C_{n, \eta, \zeta} \left(\frac{1}{8}\right)^{\lfloor \frac{\min\{m_1, m_2\}}{2n-1} \rfloor} + \epsilon/2. \quad (123)$$

Hence, if  $m_1, m_2$  satisfy

$$C_{f_{in}, \phi_1, \phi_2} C_{n, \eta, \zeta} \left(\frac{1}{8}\right)^{\lfloor \frac{\min\{m_1, m_2\}}{2n-1} \rfloor} < \epsilon/2,$$

then it follows that  $\|h^{m_1} - h^{m_2}\|_{L^\infty(\Omega \times \mathbb{R}^3)} < \epsilon$ . From the discussion above,  $\{h^l\}$  is a Cauchy sequence. Therefore, the limit  $\lim_{l \rightarrow \infty} h^l = h^{\varpi, \eta}$  exists for any  $\alpha, \eta$ . Moreover,  $h^{\varpi, \eta}$  satisfies

$$\begin{cases} \varpi\nu(v)h^{\varpi, \eta} + v \cdot \nabla_x h^{\varpi, \eta} + \frac{1}{\kappa} L_w (h^{\varpi, \eta}) = w\phi_1, & \text{in } \Omega \times \mathbb{R}^3, \\ h^{\varpi, \eta} = wf_{in}, & \text{on } \partial\Omega_1^-, \\ h^{\varpi, \eta} = \eta P_w (h^{\varpi, \eta}) + w\phi_2, & \text{on } \partial\Omega_2^-. \end{cases}$$

Moreover, we observe that the argument used in **Property 4.1** also applies to  $h^{\varpi, \eta}$  for any fixed  $0 < \eta < 1$ . Then we obtain

$$\begin{aligned} \|h^{\varpi, \eta}\|_{L^\infty(\Omega \times \mathbb{R}^3)} &\leq C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + \frac{C_\zeta}{\kappa} \|h^{\varpi, \eta}/w\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \\ &= C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\varpi, \eta}\|_{L^2_{0,\gamma/2}(\Omega \times \mathbb{R}^3)} \\ &\leq C_\zeta \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \max\{1, \kappa\} \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \\ &\quad + C_{\zeta, \varpi, \eta} \left( \|f_{in}\|_{L^\infty_{\alpha,0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha,0}(\partial\Omega_2^-)} + \|\phi_1\|_{L^\infty_{\alpha,-1}(\Omega \times \mathbb{R}^3)} \right) \end{aligned} \quad (124)$$

Let  $0 < \varpi_1, \varpi_2 < 1$ . We recall that

$$\begin{cases} v \cdot \nabla_x (h^{\varpi_1, \eta} - h^{\varpi_2, \eta}) + \frac{1}{\kappa} L_w (h^{\varpi_1, \eta} - h^{\varpi_2, \eta}) = -\varpi_1 h^{\varpi_1, \eta} + \varpi_2 h^{\varpi_2, \eta}, & \text{in } \Omega \times \mathbb{R}^3, \\ (h^{\varpi_1, \eta} - h^{\varpi_2, \eta}) = 0, & \text{on } \partial\Omega_1^-, \\ (h^{\varpi_1, \eta} - h^{\varpi_2, \eta}) = \eta P_w (h^{\varpi_1, \eta} - h^{\varpi_2, \eta}), & \text{on } \partial\Omega_2^-. \end{cases}$$

Regard  $-\varpi_1 h^{\varpi_1, \eta} + \varpi_2 h^{\varpi_2, \eta}$  as  $\phi_1$ . By repeating the same process as above, we obtain

$$\begin{aligned} & \|h^{\varpi_1, \eta} - h^{\varpi_2, \eta}\|_{L^\infty(\Omega \times \mathbb{R}^3)} \\ & \leq C_\zeta \cdot \kappa \cdot \max\{\varpi_1, \varpi_2\} (\|\langle v \rangle^{-1} h^{\varpi_1, \eta}\|_{L^\infty(\Omega \times \mathbb{R}^3)} + \|\langle v \rangle^{-1} h^{\varpi_2, \eta}\|_{L^\infty(\Omega \times \mathbb{R}^3)}) \\ & \quad + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\varpi_1, \eta} - f^{\varpi_2, \eta}\|_{L^2_{0, \gamma/2}(\Omega \times \mathbb{R}^3)} \\ & \leq C_\zeta \cdot \kappa \cdot \max\{\varpi_1, \varpi_2\} \left( \|f_{in}\|_{L^\infty_{\alpha, 0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha, 0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha, -1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\varpi_1, \eta} - f^{\varpi_2, \eta}\|_{L^2_{0, \gamma/2}(\Omega \times \mathbb{R}^3)}. \end{aligned} \tag{125}$$

This implies that  $\{h^{\varpi, \eta}\}$  is a Cauchy sequence in  $L^\infty(\Omega \times \mathbb{R}^3)$ . After we denote  $h^\eta = \lim_{\alpha \rightarrow 0} h^{\varpi, \eta}$ , note that  $h^{\eta_1} - h^{\eta_2}$  satisfies

$$\begin{cases} v \cdot \nabla_x (h^{\eta_1} - h^{\eta_2}) + \frac{1}{\kappa} L_w (h^{\eta_1} - h^{\eta_2}) = 0, & \text{in } \Omega \times \mathbb{R}^3, \\ (h^{\eta_1} - h^{\eta_2}) = 0, & \text{on } \partial\Omega_1^-, \\ (h^{\eta_1} - h^{\eta_2}) = \eta_1 P_w (h^{\eta_1} - h^{\eta_2}) + (\eta_1 - \eta_2) P_w (h^{\eta_2}), & \text{on } \partial\Omega_2^-. \end{cases}$$

We can regard  $(\eta_1 - \eta_2) P_w (h^{\eta_2})$  as  $\phi_2$  and apply the same argument again. We obtain

$$\begin{aligned} \|h^{\eta_1} - h^{\eta_2}\|_{L^\infty(\Omega \times \mathbb{R}^3)} & \leq C_\zeta |\eta_1 - \eta_2| \cdot \|P_w (h^{\eta_2})\|_{L^\infty(\partial\Omega_2^-)} + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\eta_1} - f^{\eta_2}\|_{L^2_{0, \gamma/2}(\Omega \times \mathbb{R}^3)} \\ & \leq C_\zeta |\eta_1 - \eta_2| \cdot \|h^{\eta_2}\|_{L^\infty(\partial\Omega^+)} + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\eta_1} - f^{\eta_2}\|_{L^2_{0, \gamma/2}(\Omega \times \mathbb{R}^3)} \\ & \leq C_\zeta |\eta_1 - \eta_2| \cdot \left( \|f_{in}\|_{L^\infty_{\alpha, 0}(\partial\Omega_1^-)} + \|\phi_2\|_{L^\infty_{\alpha, 0}(\partial\Omega_2^-)} + \kappa \|\phi_1\|_{L^\infty_{\alpha, -1}(\Omega \times \mathbb{R}^3)} \right) \\ & \quad + C_\zeta \min\{1, \kappa^{-1}\} \|f^{\eta_1} - f^{\eta_2}\|_{L^2_{0, \gamma/2}(\Omega \times \mathbb{R}^3)}. \end{aligned} \tag{126}$$

The last inequality follows from a similar argument showing that  $\|h^{\varpi, \eta}\|_{L^\infty}$  is bounded with respect to  $\varpi, \eta$ . Thus  $\{h^\eta\}_\eta$  is a Cauchy sequence in  $L^\infty$  and  $h := \lim_{\eta \rightarrow 1} h^\eta$  is a mild solution of (80). The proof of the estimate (86) is similar to the proof of (124).  $\square$

**Remark 4.2.** *the proof for the unseal set is similar to the case of the complete unsealed set. We just take  $\lfloor \frac{n}{m} \rfloor$  sufficiently large in the proof of **Proposition 4.1** to make  $\zeta^{\lfloor \frac{n}{m} \rfloor}$  is small enough.*

## 4.2. the proof of Theorem 1.1

Now, we conclude this section by proving **Theorem 1.1**.

*Proof. the proof of theorem 1.1* We consider the following iteration with  $f_1 = 0$ .

$$\begin{cases} v \cdot \nabla_x f_{k+1} + \frac{1}{\kappa} L(f_{k+1}) = \frac{1}{\kappa} \Gamma(f_k, f_k), & \text{in } \Omega \times \mathbb{R}^3, \\ f_{k+1} = f_{in}, & \text{on } \partial\Omega_1^-, \\ f_{k+1} = P(f_{k+1}) + m^{-1} (m_{T(x)} - m) P(f_k) \\ \quad + \frac{1}{\sqrt{2\pi}} m^{-1/2} (m_{T(x)} - m), & \text{on } \partial\Omega_2^-. \end{cases} \tag{127}$$

By **Property 4.1**, we obtain

$$\begin{aligned} \|f_{k+1}\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_{k+1}\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} &\leq C_\zeta \left( \begin{array}{l} \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} \\ + \|m^{-1}(m_{T(x)} - m)P(f_k)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} \\ + \|\frac{1}{\sqrt{2\pi}}m^{-1/2}(m_{T(x)} - m)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} \\ + \max\{1, \kappa^{-1}\}\|\Gamma(f_k, f_k)\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \end{array} \right) \\ &\leq C_\zeta \left( \begin{array}{l} \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)} \\ + C\|m^{-1/2}(m_{T(x)} - m)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)}\|f_k\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \\ + \|\frac{1}{\sqrt{2\pi}}m^{-1/2}(m_{T(x)} - m)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} \\ + \max\{1, \kappa^{-1}\}\|\Gamma(f_k, f_k)\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \end{array} \right). \end{aligned} \quad (128)$$

Let  $\mathcal{A} = \|f_{in}\|_{L_{\alpha,0}^\infty(\partial\Omega_1^-)}$  and  $\mathcal{B} = \|\frac{1}{\sqrt{2\pi}}m^{-1/2}(m_{T(x)} - m)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)}$ . It is well-known that

$$\|\Gamma(f, f)\|_{L_{\alpha,-1}^\infty(\Omega \times \mathbb{R}^3)} \leq C\|f\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)}^2. \quad (129)$$

and if  $0 < \alpha < 1/4$  and  $T(x) < 2$ , then

$$\|\frac{1}{\sqrt{2\pi}}m^{-1/2}(m_{T(x)} - m)\|_{L_{\alpha,0}^\infty(\partial\Omega_2^-)} \leq C_\alpha\|T(x) - 1\|_{L^\infty(\partial\Omega_2^-)}. \quad (130)$$

We show that there exists  $\delta$  such that, if  $\mathcal{A} + \mathcal{B} \leq \delta$ , then

$$\|f_{k+1}\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_{k+1}\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \leq 2C_\zeta(\mathcal{A} + \mathcal{B}) \quad (131)$$

for all  $k \in \mathbb{N}$ . The constant  $C_\zeta$  is defined in **Theorem 4.1**.

For  $k = 1$ ,

$$\|f_{k+1}\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_{k+1}\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \leq C_\zeta(\mathcal{A} + \mathcal{B})$$

by (128). Suppose the statement holds for  $k = n$ . For  $k = n + 1$ ,

$$\begin{aligned} \|f_{k+1}\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_{k+1}\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} &\leq C_\zeta \left[ (\mathcal{A} + \mathcal{B}) + C\mathcal{B}(\mathcal{A} + \mathcal{B}) + 4CC_\zeta \max\{1, \kappa^{-1}\}(\mathcal{A} + \mathcal{B})^2 \right] \\ &\leq C_\zeta \left[ 1 + C\mathcal{B} + 4CC_\zeta \max\{1, \kappa^{-1}\}(\mathcal{A} + \mathcal{B}) \right] (\mathcal{A} + \mathcal{B}). \end{aligned} \quad (132)$$

We choose

$$\delta < \min \left\{ \frac{1}{4C}, \frac{1}{16CC_\zeta \max\{1, \kappa^{-1}\}} \right\}.$$

Then, the proof is complete. Now, consider  $f_{k+1} - f_k$  for  $k \in \mathbb{N}$ . Following a similar procedure, we obtain

$$\|f_{k+1} - f_k\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_{k+1} - f_k\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \leq \frac{1}{2} \left( \|f_k - f_{k-1}\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f_k - f_{k-1}\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \right).$$

This implies that  $\{f_k\}$  is a Cauchy sequence in  $L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)$ . Hence, we obtain the solution by taking the limit  $f = \lim_{k \rightarrow \infty} f_k$ . Uniqueness follows by same argument. Further the upper bound of  $f$ , by (132), we have

$$\|f\|_{L_{\alpha,0}^\infty(\Omega \times \mathbb{R}^3)} + \|f\|_{L_{\alpha,0}^\infty(\partial\Omega_2^+)} \leq C(\mathcal{A} + \mathcal{B}). \quad (133)$$

□

## Section 5. Application to Thermal transpiration and thermomolecular pressure difference

### 5.1. the model of the transpiration : the proof of Theorem 1.2 and Theorem 1.3

In this section, we prove the **Theorem 1.2**. Without loss of generality, we assume that  $a = 0$  and  $b = L$ . To simplify the calculation and to obtain a more explicit estimate, we prove **Theorem 1.2** when  $\Omega$  satisfies

$$\Omega \cap \{0 \leq x_3 \leq L\} = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1^2 + x_2^2 \leq 1, 0 \leq x_3 \leq L\}.$$

The boundary condition is imposed as follows.

$$\begin{aligned} F(x, v) &= \frac{1}{(2\pi)^{3/2}} e^{-\frac{v^2}{2}}, & \text{on } \partial\Omega_{1,a}^- &= \partial\Omega^- \cap \{x_3 < 0\}, \\ F(x, v) &= \frac{1}{(2\pi)^{3/2} T_2^{5/2}} e^{-\frac{v^2}{2T_2}}, & \text{on } \partial\Omega_{1,b}^- &= \partial\Omega^- \cap \{x_3 > L\}, \\ F(x, v) &= m_{T(x)} \int_{v \cdot n(x) > 0} F(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^- &= \partial\Omega^- \setminus (\partial\Omega_{1,a}^- \cup \partial\Omega_{1,b}^-). \end{aligned} \quad (134)$$

Without loss of generality, we assume that  $T_2 > 1$ . We rewrite

$$F(x, v) = m_{T_q}(x, v) + f(x, v) = \frac{1}{(2\pi)^{3/2} T^2(q(x, v))} e^{-\frac{|v|^2}{2T(q(x, v))}} + f(x, v). \quad (135)$$

Recall that  $m_{T_q}(x, v)$  is a solution of

$$\left\{ \begin{aligned} v \cdot \nabla_x F &= 0, & \text{in } \Omega \times \mathbb{R}^3, \\ F(x, v) &= \frac{1}{(2\pi)^{3/2}} e^{-\frac{|v|^2}{2}}, & \text{on } \partial\Omega_{1,a}^- &= \partial\Omega_1^- \cap \{x_3 < a\}, \\ F(x, v) &= \frac{1}{(2\pi)^{3/2} T_2^2} e^{-\frac{|v|^2}{2T_2}}, & \text{on } \partial\Omega_{1,b}^- &= \partial\Omega_1^- \cap \{x_3 > b\}, \\ F(x, v) &= m_T(x, v) \int_{v \cdot n(x) > 0} F(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^- &= \partial\Omega^- \setminus (\partial\Omega_{1,a}^- \cup \partial\Omega_{1,b}^-). \end{aligned} \right. \quad (136)$$

The proof appears in the [38]. Now, the equation (7) becomes

$$\left\{ \begin{aligned} v \cdot \nabla_x f &= \frac{2}{\kappa} Q(m_{T_q}, f) + \frac{1}{\kappa} Q(f, f) + \frac{1}{\kappa} Q(m_{T_q}, m_{T_q}), & \text{in } \Omega \times \mathbb{R}^3, \\ f &= 0, & \text{on } \partial\Omega_{1,a}^-, \\ f &= \frac{1}{\sqrt{2\pi}} \left( \frac{1}{\sqrt{T_2}} - 1 \right) \frac{1}{2\pi T_2^2} e^{-\frac{|v|^2}{2T_2}} := -C_{T_2} m_{T_2}, & \text{on } \partial\Omega_{1,b}^-, \\ f &= \frac{1}{(2\pi) T^2(x)} e^{-\frac{|v|^2}{2T(x)}} \int_{v \cdot n(x) > 0} f(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^-. \end{aligned} \right. \quad (137)$$

Note that  $C_{T_2} > 0$ . Recall that the local flux  $u(x)$  and the total flux  $U(z)$  in the  $x_3$ -direction are defined as

$$u(x) := \int_{\mathbb{R}^3} v_3 F(x, v) dv, \quad U(z) := \int_{\{x_1^2 + x_2^2 \leq 1, x_3 = z\}} u(x) dS. \quad (138)$$

Now, we establish a lemma concerning  $\Omega$  and  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$ .

**Lemma 5.1.**  $\Omega$  is completely unsealed with respect to  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$ .

*Proof.* For any fixed  $x \in \overline{\Omega \cap \{0 \leq x_3 \leq L\}}$ , we classify  $v \in \mathbb{R}^3$  as follows.

$$\begin{aligned} A_1 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid (q(x, v), v) \in \partial\Omega_{1,a}^- \right\}, \\ A_2 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid (q(x, v), v) \in \partial\Omega_{1,b}^- \right\}, \\ A_3 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid (q(x, v), v) \in \partial\Omega_2^- \right\}, \end{aligned} \quad (139)$$

Let  $v = (v_1, v_2, v_3) = (\rho \cos \phi \sin \theta, \rho \sin \phi \sin \theta, \rho \cos \theta)$  and  $\hat{v} = (v_1, v_2)$ ,  $\hat{x} = (x_1, x_2)$ . Then the sets  $A_i$  can be written more explicitly as follow.

$$\begin{aligned} A_1 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid \frac{x_3}{\sqrt{x_3^2 + r^2}} < \cos \theta \leq 1 \right\}, \\ A_2 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid -1 \leq \cos \theta < \frac{x_3 - L}{\sqrt{(x_3 - L)^2 + r^2}} \right\}, \\ A_3 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid \frac{x_3 - L}{\sqrt{(x_3 - L)^2 + r^2}} \leq \cos \theta \leq \frac{x_3}{\sqrt{x_3^2 + r^2}} \right\}, \end{aligned} \quad (140)$$

where

$$r := h + \sqrt{h^2 + k^2}, \quad h := \hat{x} \cdot \frac{\hat{v}}{|\hat{v}|}, \quad k := \sqrt{1 - |\hat{x}|^2}.$$

Recalling **Remark 4.1**, we now consider the case  $x \in \partial\Omega_2^-$ . Without loss of generality, we assume that  $(x_1, x_2) = (1, 0)$ . Then,

$$\begin{aligned} \int_{v \cdot n(x) > 0} \mathbf{1}_{\partial\Omega_2} \circ q(x, v) m_T(x, v) v \cdot n(x) dv &= 2 \int_{-\pi/2}^{\pi/2} \int_{\cos^{-1}\left(\frac{x_3}{\sqrt{x_3^2 + 4 \sin^2 \phi}}\right)}^{\cos^{-1}\left(\frac{x_3 - L}{\sqrt{(x_3 - L)^2 + 4 \sin^2 \phi}}\right)} \cos \phi \sin^2 \theta d\theta d\phi \\ &:= \Theta(x_3). \end{aligned}$$

Observe that  $0 \leq \Theta(x_3) \leq 1$  and  $\Theta(x_3)$  is a continuous on  $[0, L]$ . Therefore, there exists  $z_0 \in [0, L]$  such that  $\max_{[0, L]} \Theta(x_3) = \Theta(z_0)$ . However,  $\Theta(z_0)$  is equal to 1 if and only if

$$\cos^{-1}\left(\frac{z_0 - L}{\sqrt{(z_0 - L)^2 + 4 \sin^2 \phi}}\right) = \pi \quad \text{and} \quad \cos^{-1}\left(\frac{z_0}{\sqrt{z_0^2 + 4 \sin^2 \phi}}\right) = 0 \quad (141)$$

for all  $\phi$ . This is impossible. Hence,

$$\int_{v \cdot n(x) > 0} \mathbf{1}_{\partial\Omega_2} \circ q(x, v) m_T(x, v) v \cdot n(x) dv \leq \Theta(z_0) < 1 \quad \text{for all } x \in \partial\Omega_2.$$

□

Note that if  $T_2$  is sufficiently close to 1 and  $\Omega$  has  $C^2$  boundary, then **Theorem 1.1** guarantees the existence of  $F$  in  $L_{0,\beta}^\infty(\Omega \times \mathbb{R}^3)$ .

*Proof. The proof of theorem 1.2 for the pipe case.*

Note that the  $f$  in 137 has the form

$$\begin{aligned} f(x, v) &= -\mathbf{1}_{\partial\Omega_{1,b}^-}(q(x, v)) C_{T_2} m_{T_2} \\ &\quad + \mathbf{1}_{\partial\Omega_2^-}(q(x, v)) m_T(q(x, v), v) \int_{v \cdot n(x) > 0} f(q(x, v), v) v \cdot n(x) dv \\ &\quad + \frac{1}{\kappa} \left( \int_0^{\tau_-(x,v)} [2Q(m_{T_q}, f) + Q(f, f) + Q(m_{T_q}, m_{T_q})] (x - sv, v) ds \right) \\ &:= \mathbf{1}_{\partial\Omega_{1,b}^-} g(x, v) + J_{\partial\Omega_2^-}^T \bar{P}(f)(x, v) + \frac{1}{\kappa} h[f](x, v). \end{aligned} \quad (142)$$

Observe that

$$\int_{\mathbb{R}^3} v_3 m_{T_q} dv = \int_0^\infty \int_0^{2\pi} \int_0^\pi \frac{e^{-\frac{\rho^2}{2T(\theta, \phi)}}}{2\pi T^2(\theta, \phi)} \rho^3 \cos \theta \sin \theta d\theta d\phi d\rho = 0.$$

We now use a procedure similar to the one in **section 4**. When tracing back  $n + 1$  time for  $n \geq 1$ , we obtain

$$\begin{aligned} \int_{\mathbb{R}^3} v_3 F(x, v) dv &= \int_{\mathbb{R}^3} v_3 f(x, v) dv & (143) \\ &= \int_{\mathbb{R}^3} v_3 \mathbf{1}_{\partial\Omega_{1,b}^-} g dv + \frac{1}{\kappa} \int_{\mathbb{R}^3} v_3 h[f] dv + \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \bar{P}(f) dv \\ &= \int_{\mathbb{R}^3} v_3 \mathbf{1}_{\partial\Omega_{1,b}^-} g dv + \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \bar{P}(\mathbf{1}_{\partial\Omega_{1,b}^-} g) dv \\ &\quad + \sum_{j=1}^n \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^j \circ \bar{P}(\mathbf{1}_{\partial\Omega_{1,b}^-} g) + \frac{1}{\kappa} \int_{\mathbb{R}^3} v_3 h[f] dv \\ &\quad + \frac{1}{\kappa} \int_{\mathbb{R}^3} J_{\partial\Omega_2^-}^T \bar{P}(h[f]) dv + \frac{1}{\kappa} \sum_{j=1}^n \int_{\mathbb{R}^3} J_{\partial\Omega_2^-}^T \hat{P}^j \circ \bar{P}(h[f]) dv \\ &\quad + \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^{n+1} \circ \bar{P}(f) dv \\ &:= A_1 + A_2 + \sum_{j=1}^n B_j + \frac{1}{\kappa} (C_1 + C_2) + D. \end{aligned}$$

The definitions of  $\bar{P}(f)$ ,  $\hat{P}^j(f)$  and  $J_{\partial A}^T$  are as follows.

$$\begin{aligned} J_A^T(f)(x, v) &:= \mathbf{1}_A(q(x, v)) m_T(q(x, v), v) f(q(x, v)), \\ \bar{P}(f)(x, v) &:= \left( \int_{v_0 \cdot n(x) > 0} f(x, v_0) v_0 \cdot n(x) dv_0 \right)(x, v), \\ \hat{P}^j(f)(x, v) &:= \left( \int \cdots \int f d\sigma_{j-1,j}^T d\sigma_{j-2,j-1}^T \cdots d\sigma_{0,1}^T \right)(x, v), \\ \left( \int f d\sigma_{j-1,j}^T \right)(x_{j-1}, v_{j-1}) &:= \int_{v_j \cdot n(x_{j-1}) > 0} J_{\partial\Omega_2^-}^T(f)(x_{j-1}, v_j) v_j \cdot n(x_{j-1}) dv_j. \end{aligned}$$

Here  $x_0 = q(x, v)$ ,  $v_0 = v$ , and  $x_i = q(x_{i-1}, v_i)$  for all  $i \in \mathbb{N}$ .

Now, we estimate each term in (143). The definition of  $A_1$  to  $D$  appearing in the last line of (143) are:

$$A_1 = \int_{\mathbb{R}^3} v_3 \mathbf{1}_{\partial\Omega_{1,b}^-} g dv, \quad (144)$$

$$A_2 = \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \bar{P}(\mathbf{1}_{\partial\Omega_{1,b}^-} g) dv, \quad (145)$$

$$B_j = \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^j \circ \bar{P}(\mathbf{1}_{\partial\Omega_{1,b}^-} g), \quad (146)$$

$$C_1 = \int_{\mathbb{R}^3} v_3 h[f] dv, \quad (147)$$

$$C_2 = \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \bar{P}(h[f]) dv + \sum_{j=1}^n \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^j \circ \bar{P}(h[f]) dv, \quad (148)$$

$$D = \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^{n+1} \circ \bar{P}(f) dv. \quad (149)$$

For  $A_1$ , using (140), we obtain

$$\begin{aligned} A_1 &= - \int_0^\infty \int_0^{2\pi} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+r^2}}\right)}^{\pi} C_{T_2} \frac{e^{-\frac{\rho^2}{2T_2}}}{2\pi T_2} \rho^3 \cos \theta \sin \theta d\theta d\phi d\rho \\ &= C \left( \int_0^{2\pi} \frac{r^2}{(x_3-L)^2+r^2} d\phi \right) \left( 1 - \frac{1}{\sqrt{T_2}} \right) \\ &> 0. \end{aligned}$$

For  $A_2$ , we observe that

$$\begin{aligned} \bar{P}(\mathbf{1}_{\partial\Omega_{1,b}^-} g)(x) &= \int_{v_0 \cdot n(x) > 0} \mathbf{1}_{\partial\Omega_{1,b}^-} g v_0 \cdot n(x) dv_0, \\ &= 2C \int_{-\pi/2}^{\pi/2} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+4\sin^2\phi}}\right)}^{\pi} \cos \phi \sin^2 \theta d\theta d\phi \\ &:= -\Theta(x_3) \left( 1 - \frac{1}{\sqrt{T_2}} \right). \end{aligned}$$

Let  $v = (\rho \cos \phi \sin \theta, \rho \sin \phi \sin \theta, \rho \cos \theta)$ . Then, we have

$$\begin{aligned} A_2 &= \frac{1}{T_2} \left( 1 - \frac{1}{\sqrt{T_2}} \right) \int_0^\infty \int_0^{2\pi} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+r^2}}\right)}^{\cos^{-1}\left(\frac{x_3}{\sqrt{x_3^2+r^2}}\right)} m_T \Theta(q(x, v)) \rho^3 \cos \theta \sin \theta d\theta d\phi d\rho \\ &:= \bar{\Theta}_1(x) \left( 1 - \frac{1}{\sqrt{T_2}} \right). \end{aligned}$$

Similarly, there exists  $\bar{\Theta}_{2,n}(x)$  such that

$$\sum_{j=1}^n B_j = \bar{\Theta}_{2,n}(x) \left( 1 - \frac{1}{\sqrt{T_2}} \right).$$

The relation between the explicit forms of  $\Theta(x)$ ,  $\bar{\Theta}_1(x)$  and  $\bar{\Theta}_2(x)$  is given by

$$\Theta(x) = 2 \int_{-\pi/2}^{\pi/2} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+4\sin^2\phi}}\right)}^{\pi} \cos \phi \sin^2 \theta d\theta d\phi, \quad (150)$$

$$\begin{aligned} \bar{\Theta}_1(x) &= \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \Theta dv \\ &= \int_0^\infty \int_0^{2\pi} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+r^2}}\right)}^{\cos^{-1}\left(\frac{x_3}{\sqrt{x_3^2+r^2}}\right)} m_T \Theta(q(x, v)) \rho^3 \cos \theta \sin \theta d\theta d\phi d\rho, \end{aligned} \quad (151)$$

$$\begin{aligned} \bar{\Theta}_2(x) &= \sum_{j=1}^n \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^j(\Theta(x)). \\ &= \sum_{j=1}^n \int_0^\infty \int_0^{2\pi} \int_{\cos^{-1}\left(\frac{x_3-L}{\sqrt{(x_3-L)^2+r^2}}\right)}^{\cos^{-1}\left(\frac{x_3}{\sqrt{x_3^2+r^2}}\right)} m_T \hat{P}^j(\Theta)(q(x, v)) \rho^3 \cos \theta \sin \theta d\theta d\phi d\rho, \end{aligned} \quad (152)$$

where  $v = (\rho \cos \phi \sin \theta, \rho \sin \phi \sin \theta, \rho \cos \theta)$ . Since  $\Omega$  is completely unsealed with respect to  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$  and applying **Proposition 2.9**, we obtain

$$\frac{1}{\kappa} (|C_1| + |C_2|) \leq \frac{C_\zeta}{\kappa} \|f\|_{L_{0,\beta}^\infty(\Omega \times \mathbb{R}^3)}^2, \quad |D| \leq C\zeta^{n+1} \|f\|_{L_{0,\beta}^\infty(\Omega \times \mathbb{R}^3)}.$$

Observe that  $\bar{\Theta}_1(x)$  and  $\bar{\Theta}_{2,n}(x)$  are positive for any  $x \in \{x_3 = 0\}$ . After combining all estimates and taking  $n$  and  $\kappa$  sufficiently large, there exists a constant  $C$  such that

$$U(0) = \int_{\{x_1^2+x_2^2 \leq 1, x_3=0\}} u(x)dS \geq C \left(1 - \frac{1}{\sqrt{T_2}}\right). \quad (153)$$

Then, we finish the proof by invoking the conservation law of mass.  $\square$

For the general case, the terms  $A_1, A_2$  and  $B_j$  are very difficult to write down in explicit form, since (140) depends on  $\partial\Omega \cap \{x_3 = 0\}$  and  $\partial\Omega \cap \{x_3 = L\}$ . However, we can still prove that  $\Omega$  is a completely unsealed set with respect to  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$ , and the proof of the general case proceed in the same way as before.

**Lemma 5.2.**  *$\Omega$  is completely unsealed set with respect to  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$*

*Proof.* Since  $\Omega$  is an open set, there exist  $\rho > 0$  and  $a_1 \in \{x_3 = 0\}, a_2 \in \{x_3 = L\}$  such that

$$B(a_1, \rho) \cap \{x_3 = 0\} \subset \Omega \cap \{x_3 = 0\} \quad \text{and} \quad B(a_2, \rho) \cap \{x_3 = L\} \subset \Omega \cap \{x_3 = L\}. \quad (154)$$

Then we define

$$\Pi := B(a_1, \rho) \cap \{x_3 = 0\} \cup B(a_2, \rho) \cap \{x_3 = L\}. \quad (155)$$

We show that  $\Omega \cap \{0 \leq x_3 \leq L\}$  is completely unsealed with respect to  $\Pi$ . We can then deduce the lemma from this conclusion. For any fixed  $x \in \Xi := \partial(\Omega \cap \{0 < x_3 < L\})$ , without loss of generality, we may assume that  $a_1 = (0, 0, 0)$ ,  $x = (x_1, x_2, x_3)$ , and  $a_2 = (a_{2,1}, a_{2,2}, L)$ . Then we classify  $v \in \mathbb{R}^3$  as follows.

$$\begin{aligned} A_1 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid q(x, v) \in B(0, \rho) \cap \{x_3 = 0\} \right\}, \\ A_2 &= \left\{ (v_1, v_2, v_3) \in \mathbb{R}^3 \mid q(x, v) \in B(a_2, \rho) \cap \{x_3 = L\} \right\}, \\ A_3 &= \mathbb{R}^3 \setminus (A_1 \cup A_2). \end{aligned}$$

Let  $v = (\hat{\rho} \sin \theta \cos \phi, \hat{\rho} \sin \theta \sin \phi, \hat{\rho} \cos \theta)$ . Then the sets  $A_1$  and  $A_2$  can be written more explicitly as follow.

$$A_1 = \left\{ (\hat{\rho}, \phi, \theta) \in \mathbb{R}_+ \times [0, 2\pi] \times [0, \frac{\pi}{2}] \mid 0 \leq \alpha_1^2 + \gamma_1, \quad \frac{\alpha_1 - \sqrt{\alpha_1^2 + \gamma_1}}{x_3} < \tan \theta < \frac{\alpha_1 + \sqrt{\alpha_1^2 + \gamma_1}}{x_3} \right\} \quad (156)$$

$$A_2 = \left\{ (\hat{\rho}, \phi, \theta) \in \mathbb{R}_+ \times [0, 2\pi] \times [\frac{\pi}{2}, \pi] \mid 0 \leq \alpha_2^2 + \gamma, \quad \frac{-\alpha_2 - \sqrt{\alpha_2^2 + \gamma_2}}{L - x_3} < \tan \theta < \frac{-\alpha_2 + \sqrt{\alpha_2^2 + \gamma_2}}{L - x_3} \right\} \quad (157)$$

where

$$\begin{aligned} \alpha_1 &= x_1 \cos \phi + x_2 \sin \phi, & \gamma_1 &= \rho^2 - (x_1^2 + x_2^2)^2 \\ \alpha_2 &= [x_1 - a_{2,1}] \cos \phi + [x_2 - a_{2,2}] \sin \phi, & \gamma_2 &= \rho^2 - [(x_1 - a_{2,1})^2 + (x_2 - a_{2,2})^2]. \end{aligned}$$

Let

$$W_1(x) := 1 - \int_{u \cdot n(x) > 0} 1_{A_1}(u) m_{T(q(x,u))} u \cdot n(x) du$$

and

$$W_2(x) := 1 - \int_{u \cdot n(x) > 0} 1_{A_2}(u) m_{T(q(x,u))} u \cdot n(x) du.$$

Since  $\Omega \cap \{0 \leq x_3 \leq L\}$  is a convex set and  $\partial\Omega \cap \{0 \leq x_3 \leq L\}$  is a  $C^1$  boundary,  $W_1$  is continuous on  $\partial\Omega \cap \{\frac{L}{2} \leq x_3 \leq L\}$  and  $W_2$  is continuous on  $\partial\Omega \cap \{0 \leq x_3 \leq \frac{L}{2}\}$ . Moreover,  $W_1$  and  $W_2$  are continuous on  $\overline{\Omega \cap \{x_3 = L\}} \setminus \overline{\Pi}$  and  $\overline{\Omega \cap \{x_3 = 0\}} \setminus \overline{\Pi}$  respectively. Hence, we have

$$\begin{aligned} & \sup_{x \in \Xi \setminus \Pi} \int_{u \cdot n(x) > 0} 1_{\Xi \setminus \Pi}(q(x, u)) m_{T(q(x, u))} u \cdot n(x) du \\ & \leq \max \left\{ \max_{x \in \partial\Omega \cap \{\frac{L}{2} \leq x_3 \leq L\}} W_1(x), \max_{x \in \Omega \cap \{x_3 = L\} \setminus \Pi} W_1(x), \max_{x \in \partial\Omega \cap \{\frac{L}{2} \leq x_3 \leq L\}} W_2(x), \max_{x \in \Omega \cap \{x_3 = 0\} \setminus \Pi} W_2(x) \right\} \\ & := \zeta. \end{aligned}$$

Similar to the argument of lemma 5.1, we deduce that  $\zeta$  is less than one. Thus, we conclude that  $\Omega$  is completely unsealed set with respect to  $\partial\Omega_{1,a} \cup \partial\Omega_{1,b}$ .  $\square$

The proof of **Theorem 1.3** is similar to the proof of **Theorem 1.2** because, after rewriting (24) by equation (135), (24) becomes

$$\begin{cases} v \cdot \nabla_x f = \frac{2}{\kappa} Q(m_{T_q}, f) + \frac{1}{\kappa} Q(f, f) + \frac{1}{\kappa} Q(m_{T_q}, m_{T_q}), & \text{in } \Omega \times \mathbb{R}^3, \\ f = 0, & \text{on } \partial\Omega_{1,a}^-, \\ f = 0, & \text{on } \partial\Omega_{1,b}^-, \\ f = \frac{1}{2\pi T^2(x)} e^{-\frac{v^2}{2T(x)}} \int_{v \cdot n(x) > 0} f(x, v) v \cdot n(x) dv, & \text{on } \partial\Omega_2^-. \end{cases} \quad (158)$$

Then, equation (143) becomes

$$\begin{aligned} \int_{\mathbb{R}^3} v_3 F(x, v) dv &= \int_{\mathbb{R}^3} v_3 f(x, v) dv \\ &= \frac{1}{\kappa} \int_{\mathbb{R}^3} v_3 h dv + \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \overline{P}(f) dv \\ &= \frac{1}{\kappa} \int_{\mathbb{R}^3} v_3 h[f] dv + \frac{1}{\kappa} \int_{\mathbb{R}^3} J_{\partial\Omega_2^-}^T \overline{P}(h[f]) dv + \frac{1}{\kappa} \sum_{j=1}^n \int_{\mathbb{R}^3} J_{\partial\Omega_2^-}^T \hat{P}^j \circ \overline{P}(h[f]) dv \\ &\quad + \int_{\mathbb{R}^3} v_3 J_{\partial\Omega_2^-}^T \hat{P}^{n+1} \circ \overline{P}(f) dv \\ &:= \frac{1}{\kappa} (C_1 + C_2) + D. \end{aligned} \quad (159)$$

Since the upper bound of  $|C_1| + |C_2|$  can be estimated independently of  $n$  and  $D$  can be made arbitrary small when  $n$  is sufficiently large, the proof is complete.

## Acknowledgments

The authors would like to thank professor Kazuo Aoki and professor Tai-Ping Liu for their interest and insightful suggestions. Also, the authors would like to thank professor Chun-Hsiung Hsia and researcher Jhe-Kuan Su for giving fruitful comments. The research of the first author is supported in part by NSTC Grant 112-2115-M-002-009-MY3.

## Section 6. Appendix A: The $H^2$ estimate of Laplace equation with Robin boundary condition

In this section, we prove the inequality (39). That is, we discuss about the following elliptic equation

$$\begin{cases} -\Delta U = F, & \text{in } \Omega, \\ \chi(x)U + \partial_\nu U = 0, & \text{on } \partial\Omega. \end{cases} \quad (160)$$

Here  $\chi(x)$  is a non-negative  $C^2$  function and  $\chi(x) = 0$  on  $\partial\Omega_2$ . We require that  $\chi > 0$  on a subset of  $\partial\Omega$  of positive measure. Before we start our argument, we recall several classical trace theorems and elliptic estimate. Here  $\gamma$  is the trace operator.

**Theorem 6.1.** Let  $u, v \in W^1(\Omega)$  and  $\Delta u, \Delta v \in L^2(\Omega)$ . Then:

(i)

$$\int_{\Omega} \Delta u \, dx = \int_{\partial\Omega} \gamma \partial_{\nu} u \, dS.$$

(ii)

$$\int_{\Omega} \nabla v \cdot \nabla u \, dx = - \int_{\Omega} u \Delta v \, dx + \int_{\partial\Omega} u \gamma \partial_{\nu} v \, dS.$$

(iii)

$$\int_{\Omega} (u \Delta v - v \Delta u) \, dx = \int_{\partial\Omega} (u \gamma \partial_{\nu} v - v \gamma \partial_{\nu} u) \, dS.$$

**Theorem 6.2** ([19] Theorem 1.5.1.3). Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with a Lipschitz boundary  $\Gamma$ . Then the mapping

$$u \mapsto \gamma u$$

which is defined for  $u \in C^{0,1}(\overline{\Omega})$ , has a unique continuous extension as an operator from  $W_p^1(\Omega)$  onto  $W_p^{1-\frac{1}{p}}(\Gamma)$ . This operator has a right continuous inverse independent of  $p$ .

The following theorem is a general version of the above theorem

**Theorem 6.3** ([19] Theorem 1.5.1.2). Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with boundary  $\Gamma = \partial\Omega$  of class  $C^{k,1}$ . Assume that

$$s - \frac{1}{p} \notin \mathbb{N}, \quad s \leq k + 1, \quad s - \frac{1}{p} = l + \sigma,$$

where  $l \in \mathbb{N}_0$  and  $0 < \sigma < 1$ . Then the mapping

$$u \mapsto (\gamma u, \gamma \partial_{\nu} u, \dots, \gamma \partial_{\nu}^l u),$$

initially defined for  $u \in C^{k,1}(\overline{\Omega})$ , admits a unique continuous extension as an operator

$$W_p^s(\Omega) \longrightarrow \prod_{j=0}^l W_p^{s-j-\frac{1}{p}}(\Gamma),$$

which is onto. Moreover, this operator possesses a continuous right inverse which does not depend on  $p$ .

**Theorem 6.4** ([19] Boundary  $H^2$ -regularity). Assume

$$f \in L^2(\Omega).$$

Suppose that  $u \in H^1(U)$  is a weak solution of the elliptic boundary-value problem

$$\begin{cases} \Delta u = f & \text{in } \Omega, \\ \gamma \partial_{\nu} u = 0 & \text{on } \partial\Omega. \end{cases}$$

Assume finally

$$\partial\Omega \text{ is } C^2 \text{ and } \int_{\Omega} f = 0.$$

Then

$$u \in H^2(\Omega),$$

and we have the estimate

$$\|u\|_{H^2(\Omega)} \leq C \|f\|_{L^2(\Omega)},$$

the constant  $C$  depending only on  $\Omega$ .

Now, we prove the following estimate

**Lemma 6.1.** *Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with  $C^2$  boundary. Assume that  $F \in L^2(\Omega)$ . Suppose that  $U \in H^1(\Omega)$  is a weak solution of (160). Then*

$$U \in H^2(\Omega),$$

and we have the estimate

$$\|u\|_{H^2(\Omega)} \leq C (\|F\|_{L^2(\Omega)} + \|u\|_{H^1(\Omega)}),$$

the constant  $C$  depending only on  $\Omega$ .

*Proof.* By Green's identity, we have

$$\int_{\Omega} F dx = \int_{\partial\Omega} \chi(x) U dS. \quad (161)$$

We consider the following elliptic problem

$$\begin{cases} -\Delta U = F, & \text{in } \Omega, \\ \partial_n U = g := -\chi(x)U, & \text{on } \partial\Omega. \end{cases} \quad (162)$$

After applying theorem 6.3, there exists  $G \in W^2(\Omega)$  such that  $\partial_\nu G = -\chi(x)U$ . Then,  $U - G$  satisfies

$$\begin{cases} -\Delta(U - G) = F + \Delta G, & \text{in } \Omega, \\ \partial_n(U - G) = 0, & \text{on } \partial\Omega. \end{cases} \quad (163)$$

By (161) and Green's identity, we have

$$\int_{\Omega} F + \Delta G dx = \int_{\Omega} F dx + \int_{\Omega} \Delta G dx = \int_{\Omega} F dx + \int_{\partial\Omega} \partial_\nu G dS = \int_{\Omega} F dx - \int_{\partial\Omega} \chi(x) U dS = 0.$$

Applying theorem 6.4, we yield

$$\|U - G\|_{H^2(\Omega)} \leq C \|F + \Delta G\|_{L^2(\Omega)}.$$

Using theorem 6.3 and theorem 6.2, we obtain

$$\begin{aligned} \|U\|_{H^2(\Omega)} &\leq C (\|F + \Delta G\|_{L^2(\Omega)} + \|G\|_{H^2(\Omega)}) \\ &\leq C (\|F\|_{L^2(\Omega)} + \|G\|_{H^2(\Omega)}) \\ &\leq C (\|F\|_{L^2(\Omega)} + \|\chi(x)U\|_{H^{1/2}(\partial\Omega)}) \\ &\leq C (\|F\|_{L^2(\Omega)} + \|U\|_{H^{1/2}(\partial\Omega)}) \\ &\leq C (\|F\|_{L^2(\Omega)} + \|U\|_{H^1(\Omega)}) \end{aligned}$$

The proof is complete. □

**Lemma 6.2.** *Let  $\Omega$  be a bounded connected open subset of  $\mathbb{R}^n$  with  $C^2$  boundary. Assume that  $F \in L^2(\Omega)$ . Then, (160) admits a unique weak solution  $U$ . Furthermore,*

$$\|U\|_{H^1(\Omega)} \leq C \|F\|_{L^2(\Omega)}, \quad (164)$$

the constant  $C$  depending only on  $\Omega$ .

*Proof.* Note that the weak solution of (160) means

$$\int_{\Omega} \nabla U \cdot \nabla v \, dx + \int_{\partial\Omega} \chi U v \, dS = \int_{\Omega} F v \, dx. \quad (165)$$

Let

$$a(u, v) := \int_{\Omega} \nabla u \cdot \nabla v \, dx + \int_{\partial\Omega} \chi u v \, dS.$$

Since  $\chi \in L^{\infty}(\partial\Omega)$  and  $u, v \in H^1(\Omega)$ , we have

$$|a(u, v)| \leq C \|u\|_{H^1(\Omega)} \|v\|_{H^1(\Omega)}.$$

Assume that  $a$  is not coercive, There exists a sequence in  $H^1(\Omega)$  such that

$$\|u_n\|_{H^1(\Omega)}^2 > n \left( \int_{\Omega} |\nabla u_n|^2 \, dx + \int_{\partial\Omega} \chi u_n^2 \, dS \right). \quad (166)$$

Without loss of generality, we assume that  $\|u_n\|_{L^2(\Omega)} = 1$  for any  $n \in \mathbb{N}$ . By (166) and  $\|u_n\|_{L^2(\Omega)} = 1$ , we discover that  $\{u_n\}$  is a bounded sequence in  $H^1(\Omega)$ . Hence  $\{u_n\}$  is weakly compact in  $H^1(\Omega)$  and is compact in  $L^2(\Omega)$ . Let  $\{u_{n_j}\}$  be a convergent subsequence of  $\{u_n\}$  and  $u$  be its limit. That is,

$$\begin{aligned} u_{n_j} &\rightharpoonup u && \text{in } H^1(\Omega), \\ u_{n_j} &\rightarrow u && \text{in } L^2(\Omega). \end{aligned} \quad (167)$$

We observe that

$$0 = \lim_{n_j \rightarrow \infty} \frac{1}{n_j} \|\nabla u\|_{L^2(\Omega)} \geq \lim_{n_j \rightarrow \infty} \|\nabla u_{n_j}\|_{L^2(\Omega)} \|\nabla u\|_{L^2(\Omega)} \geq \lim_{n_j \rightarrow \infty} \int_{\Omega} \nabla u_{n_j} \cdot \nabla u \, dx = \int_{\Omega} |\nabla u|^2 \, dx.$$

It implies that  $u_{n_j}$  converges strongly to  $u$  in  $H^1(\Omega)$ ; hence,

$$\lim_{n_j \rightarrow \infty} \int_{\partial\Omega} \chi u_{n_j}^2 \, dS = \int_{\partial\Omega} \chi u^2 \, dS = 0.$$

Thus, we deduce that  $u = 0$ , which yields a contradiction since  $\|u_n\|_{L^2(\Omega)} = 1$ . By Lax-Milgram theorem, the proof of the existence is complete. Assume that (164) does not hold, there exists a sequence  $\{U_n\}$  such that

$$\|U_n\|_{H^1(\Omega)} \geq n \|F_n\|_{L^2(\Omega)}. \quad (168)$$

Note that

$$\int_{\Omega} |\nabla U_n|^2 \, dx + \int_{\partial\Omega} \chi U_n^2 \, dS = \int_{\Omega} F_n U_n \, dx \leq \|F_n\|_{L^2(\Omega)} \|U_n\|_{L^2(\Omega)}. \quad (169)$$

Without loss of generality, we assume that  $\|U_n\|_{L^2(\Omega)} = 1$  for any  $n \in \mathbb{N}$ . By (168) and (169), we can immediately conclude that  $\{U_n\}$  is bounded in  $H^1(\Omega)$  and there exists a subsequence of  $U_n$  which strongly converge to 0 in  $H^1(\Omega)$  by similar argument. It is a contradiction since  $\|U_n\|_{L^2(\Omega)} = 1$  for any  $n \in \mathbb{N}$ . The proof is complete.  $\square$

## Section 7. Appendix B: The sufficient conditions for unsealed sets

Let  $\Omega$  be a subset in  $\mathbb{R}^3$  and  $\partial B$  be a subset of  $\partial\Omega$ . Determining whether  $\Omega$  is unsealed with respect to  $\partial B$  is a difficult problem. In this section, we provide some sufficient conditions concerning this problem.

**Lemma 7.1.** *Let  $\Omega \in \mathbb{R}^3$  be a bounded convex domain with  $C^1$  boundary and  $\partial\Omega_{1,a}, \partial\Omega_{1,b}$  be subsets of  $\partial\Omega$ . If  $\partial\Omega_{1,a} \subseteq \partial\Omega_{1,b}$  and  $\Omega$  is unsealed with respect to  $\partial\Omega_{1,a}$ , then  $\Omega$  is unsealed with respect to  $\partial\Omega_{1,b}$ .*

*Proof.* Since  $\partial\Omega_{1,a} \subseteq \partial\Omega_{1,b}$ ,  $(\partial\Omega \setminus \partial\Omega_{1,b}) \subseteq (\partial\Omega \setminus \partial\Omega_{1,a})$ . Then, by **definition 4.1**, the proof is complete.  $\square$

**Proposition 7.1.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with  $C^1$  boundary, and let  $\partial\Omega_1$  be an open subset of  $\partial\Omega$ . If there exist  $\partial C_1, \partial C_2 \subset \partial\Omega \setminus \partial\Omega_1$  and  $0 < \zeta < 1$  such that*

1.  $\partial C_1 \cup \partial C_2 = \partial\Omega \setminus \partial\Omega_1$ .
2.  $\int_{v \cdot n(x) > 0} \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(q(x, v)) m(v) v \cdot n(x) dv \leq \zeta \quad \forall x \in \partial C_1$ .
3.  $\int_{v \cdot n(x) > 0} \mathbf{1}_{\partial C_2}(q(x, v)) m(v) v \cdot n(x) dv = 0 \quad \forall x \in \partial C_2$ .

*Then,  $\Omega$  is unsealed with respect to  $\partial\Omega_1$ .*

*Proof.* For convenience, we use same symbol in **section 4**

$$\begin{aligned} & \int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_1) d\sigma_{0,1} \\ &= \int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_1}(x_1) d\sigma_{0,1} + \int \mathbf{1}_{\partial C_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_2}(x_1) d\sigma_{0,1} + \int \mathbf{1}_{\partial C_2}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_2}(x_1) d\sigma_{0,1} \\ &\leq \int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_1}(x_1) d\sigma_{0,1} + \int \mathbf{1}_{\partial C_2}(x_1) d\sigma_{0,1}. \end{aligned}$$

If  $x \in \partial C_2$ , then

$$\int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_1}(x_1) d\sigma_{0,1} \leq \zeta.$$

Otherwise,

$$\begin{aligned} \int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_1) d\sigma_{0,1} &\leq \int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial C_1}(x_1) d\sigma_{0,1} + \int \mathbf{1}_{\partial C_2}(x_1) d\sigma_{0,1} \\ &\leq \int \mathbf{1}_{\partial C_1}(x_1) d\sigma_{0,1} + \int \mathbf{1}_{\partial C_2}(x_1) d\sigma_{0,1} \\ &= \int_{v \cdot n(x,v) > 0} \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(q(x, v)) m(v) v \cdot n(x) dv \\ &\leq \zeta. \end{aligned}$$

So

$$\int \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_2) d\sigma_{2,1} \mathbf{1}_{\partial\Omega \setminus \partial\Omega_1}(x_1) d\sigma_{0,1} \leq \zeta \quad \forall x \in \partial\Omega \setminus \partial\Omega_1. \quad (170)$$

$\square$

## Section 8. Appendix C: The equivalent definition of solution

In this section, we discuss the definition of a solution of (7), (11) and valid that (142) exists when **Theorem 1.1** holds. Also, we discuss the relation between the mild solution and  $v \cdot \nabla f$ .

**Definition 8.1.**  *$F$  is an mild solution of (7) if*

$$\begin{aligned} F &= \mathbf{1}_{\partial\Omega_1^-} F_{in}(q(x, v), v) + \mathbf{1}_{\partial\Omega_2^-} m_{T(q(x, v))} \int_{u \cdot n(q(x, v)) > 0} F(q(x, v), v) u \cdot n(q(x, v)) du \\ &\quad + \frac{1}{\kappa} \int_0^{\tau_-(x, v)} Q(F, F)(x - tv) dt \end{aligned}$$

*a.e.  $(x, v) \in \Omega \times \mathbb{R}^3$ .*

Let

$$\nu_F(x, v) = \int_{\mathbb{R}^3} \int_{S^2} \mathbf{B}(|v - u|, \sigma) F(x, v) d\sigma du.$$

We have another type of definition to solution of (7). That is,

**Definition 8.2.**  $F$  is an exponential solution of (7) if

$$\begin{aligned} F &= \mathbf{1}_{\partial\Omega_1^-} e^{-\int_0^{\tau_-(x,v)} \frac{1}{\kappa} \nu_F(x-sv, v) ds} F_{in}(q(x, v), v) \\ &+ \mathbf{1}_{\partial\Omega_2^-} e^{-\int_0^{\tau_-(x,v)} \frac{1}{\kappa} \nu_F(x-sv, v) ds} m_{T(q(x,v))} \int_{u \cdot n(q(x,v)) > 0} F(q(x, v), v) u \cdot n(q(x, v)) du \\ &+ \frac{1}{\kappa} \int_0^{\tau_-(x,v)} e^{-\int_0^t \frac{1}{\kappa} \nu_F(x-sv, v) ds} Q_+(F, F)(x - tv) dt. \end{aligned}$$

a.e.  $(x, v) \in \Omega \times \mathbb{R}^3$ . Here,

$$Q_+(F, G)(x, v) = \int_{\mathbb{R}^3} \int_{S^2} [F(v')G(u') + F(u')G(v')] B(|v - u|, \omega) d\omega du,$$

After linearizing (7), we obtain (11). Similar (7), we can define the mild solution as follow.

**Definition 8.3.**  $f$  is an mild solution of (11) if

$$\begin{aligned} f &= \mathbf{1}_{\partial\Omega_1^-} f_{in}(q(x, v), v) \\ &+ \mathbf{1}_{\partial\Omega_2^-} \left[ m^{-1} m_{T(q(x,v))} P(f)(q(x, v), v) + \frac{1}{\sqrt{2\pi}} m^{-1/2} (m_{T(q(x,v))} - m) \right] \\ &+ \frac{1}{\kappa} \int_0^{\tau_-(x,v)} Q(M + m^{1/2} f, M + m^{1/2} f)(x - tv) dt. \end{aligned}$$

a.e.  $(x, v) \in \Omega \times \mathbb{R}^3$ .

Also, we can define the exponential solution

**Definition 8.4.**  $f$  is an exponential solution of (11) if

$$f = J^\kappa \left[ f_{in} \mathbf{1}_{\partial\Omega_1^-} \right] + J^\kappa \left[ m^{-1} m_{T(x)} P(f) + \frac{1}{\sqrt{2\pi}} m^{-1/2} (m_{T(x)} - m) \mathbf{1}_{\partial\Omega_2^-} \right] + \frac{1}{\kappa} S_\Omega^\kappa K f + \frac{1}{\kappa} S_\Omega^\kappa \Gamma(f, f).$$

a.e.  $(x, v) \in \Omega \times \mathbb{R}^3$ .

It is obvious that the existence of a mild solution to (7) is equivalent to the existence of a mild solution to (11). Since the proof of the equivalence between **Definition 8.1** and **Definition 8.2** is similar to that between **Definition 8.3** and **Definition 8.4**, we only present the proof for the first case. The statement is as follows.

**Theorem 8.1.** Suppose that  $\nu_F(x - sv, v)$  and  $Q_+(F, F)(x - sv, v)$  belong to  $L_{loc}^1(-\tau_-(x, -v), \tau_-(x, v))$  for almost all  $(x, v) \in \Omega \times \mathbb{R}^3$  and  $F \in L^1(\Omega \times \mathbb{R}^3)$ . Then, the definition of 8.1 is equivalent to the definition of 8.2.

*Proof.* Let

$$g(x, v) = \mathbf{1}_{\partial\Omega_1^-} F_{in}(q(x, v), v) + \mathbf{1}_{\partial\Omega_2^-} m_{T(q(x,v))} \int_{u \cdot n(q(x,v)) > 0} F(q(x, v), v) u \cdot n(q(x, v)) du$$

and

$$G(s) = g(x, v) + \frac{1}{\kappa} \int_s^{\tau_-(x,v)} Q(F, F)(x - tv) dt,$$

for any  $s \in [-\tau_-(x, -v), \tau_-(x, v)]$ . By the definition,  $G(0) = F(x, v)$ . Moreover, we have

$$\int_{\Omega \times \mathbb{R}^3} F(x, v) dx dv = \int_{\partial\Omega^-} \int_0^{\tau_-(x, -v)} F(x + sv, v) |v \cdot n(x)| ds dS dv.$$

and

$$\int_{\Omega \times \mathbb{R}^3} G(0) dx dv = \int_{\partial\Omega^-} \int_0^{\tau_-(x, -v)} G(-s) |v \cdot n(x)| ds dS dv.$$

Since  $G(0) = F(x, v)$ ,  $G(s) = F(x - sv)$  for almost all  $s \in [-\tau_-(x, -v), \tau_-(x, v)]$ .

$$\begin{cases} \frac{d}{ds} G(s) = -\frac{1}{\kappa} Q(F, F)(x - sv) \\ \quad = -\frac{1}{\kappa} Q_+(F, F)(x - sv) + \frac{1}{\kappa} \nu_F(x - sv) F(x - sv) \\ \quad = -\frac{1}{\kappa} Q_+(F, F)(x - sv) + \frac{1}{\kappa} \nu_F(x - sv) G(s), & \text{in } s \in [-\tau_-(x, -v), \tau_-(x, v)]. \\ G(0) = F(x, v). \end{cases} \quad (171)$$

Note that the solution of (171) is unique and  $G$  has two representations. One of them is

$$G(s) = g(x, v) + \frac{1}{\kappa} \int_s^{\tau_-(x, v)} Q(F, F)(x - tv) dt.$$

The other is

$$G(s) = e^{-\int_0^s \frac{1}{\kappa} \nu_F(x - sv, v) ds} g(x, v) + \frac{1}{\kappa} \int_s^{\tau_-(x, v)} e^{-\int_0^t \frac{1}{\kappa} \nu_F(x - sv, v) ds} Q_+(F, F)(x - tv) dt.$$

By uniqueness, the proof is complete.  $\square$

Now, we consider the following boundary value problem.

$$\begin{cases} v \cdot \nabla_x F = G, & \text{in } \Omega \times \mathbb{R}^3, \\ F = F_{in}, & \text{on } \partial\Omega_1^-, \end{cases} \quad (172)$$

We pay attention on the following question

”Does the existence of a mild solution implies  $v \cdot \nabla_x F$  exists in distribution sense.”

The mild solution of (172) is

$$F(x, v) = F_{in}(g(x, v), v) + \int_0^{\tau_-(x, v)} G(x - sv, v) ds.$$

Let  $\varphi$  be a smooth function with compact support. We show the following statement

**Theorem 8.2.** *If  $F$  is a mild solution of (172) and  $F, G \in L^p(\Omega \times \mathbb{R}^3)$  for some  $p \geq 1$ , then*

$$\int_{\Omega \times \mathbb{R}^3} F(x, v) v \cdot \nabla_x \varphi dx dv = - \int_{\Omega \times \mathbb{R}^3} G(x, v) \varphi dx dv$$

for any  $\varphi \in C_c^\infty(\Omega \times \mathbb{R}^3)$ .

*Proof.* Notice that

$$\begin{aligned}
F(x + hv, v) - F(x, v) &= \int_0^{\tau_-(x+hv, v)} G(x + hv - sv, v) ds - \int_0^{\tau_-(x, v)} G(x - sv, v) ds \\
&= \int_0^{\tau_-(x, v) + h} G(x - (s - h)v, v) ds - \int_0^{\tau_-(x, v)} G(x - sv, v) ds \\
&= \int_{-h}^{\tau_-(x, v)} G(x - sv, v) ds - \int_0^{\tau_-(x, v)} G(x - sv, v) ds \\
&= \int_0^h G(x + sv, v) ds
\end{aligned}$$

for any  $h > 0$ . Let  $y = x + hv$ . Since  $\Omega$  is open and  $\varphi$  has a compact support, we yield

$$\begin{aligned}
\lim_{h \rightarrow 0} \int_{\Omega \times \mathbb{R}^3} \frac{1}{h} \int_0^h G(x + sv, v) ds \varphi dx dv &= \lim_{h \rightarrow 0} \int_{\Omega \times \mathbb{R}^3} \frac{F(x + hv, v) - F(x, v)}{h} \varphi dx dv \\
&= \lim_{h \rightarrow 0} \int_{\Omega \times \mathbb{R}^3} \frac{\varphi(x - hv, v) - \varphi(x, v)}{h} F(x, v) dx dv \\
&+ \lim_{h \rightarrow 0} \int_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{\mathbf{1}_\Omega(x - hv) - \mathbf{1}_\Omega(x)}{h} \varphi(x - hv, v) F(x, v) dx dv \\
&= - \int_{\Omega \times \mathbb{R}^3} F(x, v) v \cdot \nabla_x \varphi dx dv.
\end{aligned}$$

Moreover, we have

$$\lim_{h \rightarrow 0} \int_{\Omega \times \mathbb{R}^3} \frac{1}{h} \int_0^h G(x + sv, v) ds \varphi dx dv = \int_{\Omega \times \mathbb{R}^3} G(x, v) \varphi dx dv.$$

Hence the proof is complete.  $\square$

## References

- [1] Kazuo Aoki, Yoshio Sone, and Yorifumi Waniguchi. "A rarefied gas flow induced by a temperature field: Numerical analysis of the flow between two coaxial elliptic cylinders with different uniform temperatures." *Computers & Mathematics with Applications* 35.1-2 (1998): 15-28.
- [2] M, J Bennett,, F. C Tompkins. (1957). Thermal transpiration: application of Liang's equation. *Transactions of the Faraday Society*, 53, 185-192.
- [3] Graeme Austin Bird. (1994). *Molecular gas dynamics and the direct simulation of gas flows*. Oxford university press.
- [4] Russel Caflisch. The Boltzmann equation with a soft potential. I. Linear, spatiallyhomogeneous. *Comm. Math. Phys.* 74 (1980), no. 1, 71–95.
- [5] Rojas Cardenas, Marcos, et al. "Thermal transpiration flow: A circular cross-section microtube submitted to a temperature gradient." *Physics of Fluids* 23.3 (2011).
- [6] Cercignani, Carlo, Reinhard Illner, and Mario Pulvirenti. *The mathematical theory of dilute gases*. Vol. 106. Springer Science Business Media, 2013.
- [7] Chiun-Chuan Chen, I-Kun Chen, Tai-Ping Liu, Yoshio Sone. (2007). Thermal transpiration for the linearized Boltzmann equation. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 60(2), 147-163.
- [8] Mourad Choulli, Plamen Stefanov. An inverse boundary value problem for the stationary transport equation. *Osaka J. Math.* 36 (1999), no.1, 87-104.

- [9] I-Kun Chen, Chun-Hsiung Hsia, Kawagoe, Daisuke, Jhe-Kuan Su. Geometric effects on  $W^{1,p}$  regularity of the stationary linearized Boltzmann equation, *Indiana University mathematical Journal* 74(6) 1749-1793.
- [10] I-Kun Chen, Tai-Ping Liu, Shigeru Takata. (2014). Boundary singularity for thermal transpiration problem of the linearized Boltzmann equation. *Archive for Rational Mechanics and Analysis*, 212(2), 575-595.
- [11] I-Kun Chen, Chun-Hsiung Hsia, Kawagoe, Daisuke. (2025). On the existence and regularity of weakly nonlinear stationary Boltzmann equations: a Fredholm alternative approach. preprint arXiv:2501.02419.
- [12] William Crookes. (1874). XV. On attraction and repulsion resulting from radiation. *Philosophical transactions of the Royal society of London*, (164), 501-527.
- [13] Renjun Duan, Feimin Huang, Yong Wang, Tong Yang. Global Well-Posedness of the Boltzmann Equation with Large Amplitude Initial Data. *Arch Rational Mech Anal* 225, 375–424 (2017).
- [14] Renjun Duan, Feimin Huang, Yong Wang, Zhu Zhang. Effects of Soft Interaction and Non-isothermal Boundary Upon Long-Time Dynamics of Rarefied Gas. *Arch Rational Mech Anal.* **234** (2019), 925–1006.
- [15] Raffaele Esposito, Yan Guo, Chanwoo Kim, Rossana Marra. Non-Isothermal Boundary in the Boltzmann Theory and Fourier Law. *Commun. Math. Phys.* 323, 177–239 (2013).
- [16] Lawrence Craig Evans. *Partial differential equations*. Vol. 19. American mathematical society, 2022.
- [17] Gazzola, Filippo, Hans-Christoph Grunau, and Guido Sweers. *Polyharmonic boundary value problems: positivity preserving and nonlinear higher order elliptic equations in bounded domains*. Springer, 2010.
- [18] Robert Theodore Glassey. *The Cauchy problem in kinetic theory*. Society for Industrial and Applied Mathematics, 1996.
- [19] Pierre Grisvard. *Elliptic problems in nonsmooth domains*. Society for Industrial and Applied Mathematics, 2011
- [20] Xiaohui Guo, Dhruv Singh, Jayathi Murthy, Alina A. Alexeenko1. (2009). Numerical simulation of gas-phonon coupling in thermal transpiration flows. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 80(4), 046310.
- [21] Yan Guo. Classical Solutions to the Boltzmann Equation for Molecules with an Angular Cutoff. *Arch. Rational Mech. Anal.* 169, 305–353 (2003).
- [22] Yen-Lin Han, Eric Phillip Muntz, Alina A. Alexeenko, Marcus Young. (2007). Experimental and computational studies of temperature gradient-driven molecular transport in gas flows through nano/microscale channels. *Nanoscale and Microscale Thermophysical Engineering*, 11(1-2), 151-175.
- [23] Clément Mouhot and Robert Michael Strain. "Spectral gap and coercivity estimates for linearized Boltzmann collision operators without angular cutoff." *Journal de mathématiques pures et appliquées* 87.5 (2007): 515-535.
- [24] Dongqing Li, Michael R. King, Satish G. Kandlikar, Stéphane Colin, Srinivas Garimella. (2006). *Heat transfer and fluid flow in minichannels and microchannels*. elsevier.
- [25] Kawagoe, Daisuke.  $H_x^s$  regularity of solutions to the stationary Boltzmann equation with the incoming boundary condition, preprint, arXiv:2507.18211.

- [26] Martin Knudsen. (1909). Eine revision der gleichgewichtsbedingung der gase. Thermische molekularströmung. *Annalen der Physik*, 336(1), 205-229.
- [27] Martin Knudsen. (1910). Thermischer molekulardruck der gase in röhren. *Annalen der Physik*, 338(16), 1435-1448.
- [28] Shih Ching Liang. (1953). On the calculation of thermal transpiration. *The Journal of Physical Chemistry*, 57(9), 910-911.
- [29] Shankar K. Loyalka. (1971). Kinetic theory of thermal transpiration and mechanocaloric effect. I. *The Journal of Chemical Physics*, 55(9), 4497-4503.
- [30] James Clerk Maxwell. (1879). VII. On stresses in rarified gases arising from inequalities of temperature. *Philosophical Transactions of the royal society of London*, (170), 231-256.
- [31] Hideyuki Niimi. (1971). Thermal creep flow of rarefied gas between two parallel plates. *Journal of the Physical Society of Japan*, 30(2), 572-574.
- [32] Taku Ohwada, Yoshio Sone, Kazuo Aoki. (1989). Numerical analysis of the Poiseuille and thermal transpiration flows between two parallel plates on the basis of the Boltzmann equation for hard-sphere molecules. *Physics of Fluids A: Fluid Dynamics*, 1(12), 2042-2049.
- [33] Sarantis Pantazisa, Stergios Narisa, Christos Tantosa, Dimitris Valougeorgisa, Julien Andréb, Francois Milletb, Jean Paul Perin (2013). Nonlinear vacuum gas flow through a short tube due to pressure and temperature gradients. *Fusion Engineering and Design*, 88(9-10), 2384-2387.
- [34] Ali Passian, Robert J. Warmack, Thomas L. Ferrell, Thomas Thundat. (2003). Thermal transpiration at the microscale: a Crookes cantilever. *Physical review letters*, 90(12), 124503.
- [35] Gregg A. Radtke, Nicolas G. Hadjiconstantinou, Wolfgang Wagner. (2011). Low-noise Monte Carlo simulation of the variable hard sphere gas. *Physics of fluids*, 23(3).
- [36] Michael Renardy, Robert C. Rogers. *An introduction to partial differential equations*. New York, NY: Springer New York, 2004.
- [37] Osborne Reynolds. (1879). Xviii. on certain dimensional properties of matter in the gaseous state.-Part I. Experimental researches on thermal transpiration of gases through porous plates and on the laws of transpiration and impulsion, including an experimental proof that gas is not a continuous plenum.-part ii. on an extension of the dynamical theory of gas, which includes the stresses, tangential and normal, caused by a varying condition of gas, and affords an explanation of the phenomena of transpiration and impulsion. *Philosophical Transactions of the Royal Society of London*, (170), 727-845.
- [38] Yoshio Sone, (Ed.). (2007). *Molecular gas dynamics: theory, techniques, and applications*. Boston, MA: Birkhäuser Boston.
- [39] Andrew Strongrich, Alexander Pikus, Inderjit B. Sebastiao, Alina Alexeenko. (2017). Microscale in-plane knudsen radiometric actuator: Design, characterization, and performance modeling. *Journal of Microelectromechanical Systems*, 26(3), 528-538.
- [40] Wei Su, Zhu, Liang Zhu, Peng Wang, Yonghao Zhang, Lei Wu. (2020). Can we find steady-state solutions to multiscale rarefied gas flows within dozens of iterations?. *Journal of Computational Physics*, 407, 109245.
- [41] Gui Hua Tang, Yonghao Zhang, Xiao Jun Gu, Robert W. Barber, David R. (2009). Lattice Boltzmann model for thermal transpiration. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 79(2), 027701.
- [42] Tetsuo Takaishi, Yoshihiko Sensui. (1963). Thermal transpiration effect of hydrogen, rare gases and methane. *Transactions of the Faraday Society*, 59, 2503-2514.

- [43] Shigeru Takata, Hitoshi Funagane. (2011). Poiseuille and thermal transpiration flows of a highly rarefied gas: over-concentration in the velocity distribution function. *Journal of fluid mechanics*, 669, 242-259.
- [44] Felix G. Tcheremissine. (2005, May). Direct numerical solution of the Boltzmann equation. In *AIP Conference Proceedings* (Vol. 762, No. 1, pp. 677-685). American Institute of Physics.
- [45] Seiji Ukai. "Solutions of the Boltzmann equation." *Studies in Mathematics and its Applications*. Vol. 18. Elsevier, 1986. 37-96.
- [46] Stephen E. Vargo, E. Phillip Muntz, Gerald R. Shiflett, William C. Tang. (1999). Knudsen compressor as a micro-and macroscale vacuum pump without moving parts or fluids. *Journal of Vacuum Science Technology A: Vacuum, Surfaces, and Films*, 17(4), 2308-2313.
- [47] Kung-Chien Wu, Kuan-Hsiang Wang. Hölder regularity of solutions of the steady Boltzmann equation with soft potentials, preprint arXiv:2409.12513.
- [48] Lei Wu, Reese, Jason M. Reese, Zhang, Yonghao Zhang. (2014). Solving the Boltzmann equation deterministically by the fast spectral method: application to gas microflows. *Journal of Fluid Mechanics*, 746, 53-84.