

BRUNN-MINKOWSKI INEQUALITY FOR p -HARMONIC MEASURES

ARIEL A. AGUAS-BARRENO, MURAT AKMAN, AND SHIRSHO MUKHERJEE

ABSTRACT. We prove a local Brunn-Minkowski inequality for a functional corresponding to p -harmonic measures for $2 < p < n + 1$.

1. INTRODUCTION

The Brunn-Minkowski inequalities and Minkowski problems of prescribing measures on the unit sphere goes back to the works of Minkowski [31, 32], Alexandrov [5, 6], etc. and has been one of the most prominent case to study in geometric analysis. They are also intimately connected to other inequalities including the isoperimetric inequality, Sobolev inequalities, etc. The classical Brunn-Minkowski inequality states that for any compact convex sets $E_1, E_2 \subset \mathbb{R}^n$ with nonempty interiors, we have that,

$$|\lambda E_1 + (1 - \lambda)E_2|^{\frac{1}{n}} \geq \lambda |E_1|^{\frac{1}{n}} + (1 - \lambda) |E_2|^{\frac{1}{n}}, \quad (1.1)$$

for any $\lambda \in [0, 1]$, where $|\cdot|$ is the Lebesgue measure and $+$ on the left denotes the Minkowski addition. Thus, the inequality (1.1) implies that $|\cdot|^{\frac{1}{n}}$ is a concave function with respect to Minkowski addition. Moreover, equality in (1.1) holds if and only if E_1 is a translation and dilation of E_2 . For numerous applications of (1.1) to problems in geometry and analysis, we refer to the classical book by Schneider [33] and the survey paper by Gardner [18].

Over the years, inequalities of Brunn-Minkowski type have also been proved for many other measures and homogeneous functionals. For example, one can replace the Lebesgue measure in (1.1) by capacity; in this case, it was shown by Borell [7] that

$$[\text{Cap}_2(\lambda E_1 + (1 - \lambda)E_2)]^{\frac{1}{n-2}} \geq \lambda [\text{Cap}_2(E_1)]^{\frac{1}{n-2}} + (1 - \lambda) [\text{Cap}_2(E_2)]^{\frac{1}{n-2}}, \quad (1.2)$$

whenever E_1, E_2 are compact convex sets with nonempty interiors in \mathbb{R}^n , $n \geq 3$. Here, Cap_2 denotes the *Newtonian capacity*. The exponents in this inequality and (1.1) differ because $|\cdot|$ is homogeneous of degree n , whereas $\text{Cap}_2(\cdot)$ is homogeneous of degree $n - 2$. Borell [8] proved a Brunn-Minkowski-type inequality for *logarithmic capacity*. The equality case in (1.2) was studied by Caffarelli, Jerison, and Lieb in [9], where it was shown that equality in (1.2) holds if and only if E_2 is a translate and dilate of E_1 for $n \geq 3$. Jerison in [22] used that result to prove uniqueness of the Minkowski problem. In [14], Colesanti and Salani proved the p -capacitary version of (1.2) for $1 < p < n$, i.e.

$$[\text{Cap}_p(\lambda E_1 + (1 - \lambda)E_2)]^{\frac{1}{n-p}} \geq \lambda [\text{Cap}_p(E_1)]^{\frac{1}{n-p}} + (1 - \lambda) [\text{Cap}_p(E_2)]^{\frac{1}{n-p}} \quad (1.3)$$

whenever E_1, E_2 are compact convex sets with nonempty interiors in \mathbb{R}^n , where $\text{Cap}_p(\cdot)$ denotes the p -capacity of a set defined as

$$\text{Cap}_p(E) = \inf \left\{ \int_{\mathbb{R}^n} |\nabla v|^p dx : v \in C_0^\infty(\mathbb{R}^n), v(x) \geq 1 \text{ for } x \in E \right\}. \quad (1.4)$$

Date: March 24, 2026.

2010 Mathematics Subject Classification. 31B05, 35J08, 35J25, 42B37, 42B25, 42B99.

Key words and phrases. Convex Geometry, p -Laplacian, p -harmonic measure, Minkowski problem.

M. Akman and S. Mukherjee have been supported by the EPSRC New Investigator award [grant number EP/W001586/1]. A. Aguas-Barreno has been partially supported by the School of Mathematics, Statistics and Actuarial Science at the University of Essex.

It was also shown in the same paper that equality in (1.3) holds if and only if E_2 is a translate and dilate of E_1 . In [13], Colesanti and Cuoghi defined a logarithmic capacity for $p = n, n \geq 3$, and proved a Brunn-Minkowski-type inequality for this capacity. In [11], a Minkowski problem was studied for p -capacity, $1 < p < 2$, using (1.3). For $1 < p < n$, the second author of the present article, along with others in [1], showed that (1.3) holds for any convex set with positive p -capacity and for capacities associated with more general elliptic PDEs (known as \mathcal{A} -harmonic PDEs), with the same conclusion if equality holds. In [2], the existence and uniqueness of the so-called \mathcal{A} -harmonic Green's function for the complement of a convex compact set in \mathbb{R}^n were established; using this, it was shown that a quantity related with the behavior of this function near infinity satisfies a Brunn-Minkowski inequality for $n \leq p < \infty$. It was also shown in the same article that if equality holds E_1 and E_2 , then under certain regularity and structural assumptions on \mathcal{A} (no additional assumption in the case of p -Laplace equation), then these two sets are homothetic. The torsional rigidity and first eigenvalue of the Laplacian versions of (1.1) were studied in [12]. We refer to [2, 1, 3] etc. for more results.

Harmonic measures has been studied by Dahlberg [15] and Jerison [21] and as a generalization of it, the p -harmonic measures have been studied by Lewis and Nyström, see [24, 29, 26], etc. (see Section 2 for details). Recently, the existence of domains solving the Minkowski problem for p -harmonic measures has been established by the second and the third author in [4]. In this paper, we prove a Brunn-Minkowski-type inequality curated towards some form of uniqueness of domains that solve the Minkowski problem for p -harmonic measures.

The main result of this paper is a local Brunn-Minkowski inequality for p -harmonic measures.

Theorem 1.5. *Let $K_0 \subset \mathbb{R}^n$ be a compact convex set of non-empty interior and ω_0 be any given p -harmonic measure on ∂K_0 for $2 < p < n + 1$. There exists a neighborhood \mathcal{N} of convex sets of non-empty interior containing K_0 such that for any $K \in \mathcal{N}$, there exists a p -harmonic measure ω_K on ∂K with $\omega_{K_0} = \omega_0$ and for the functional $\mathcal{F} : \mathcal{N} \rightarrow \mathbb{R}$ defined by*

$$\mathcal{F}(K) = \int_{\partial K} (h_K \circ \mathbf{g}_K) d\omega_K, \quad (1.6)$$

where $h_K : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ is the support function of K and $\mathbf{g}_K : \partial K \rightarrow \mathbb{S}^{n-1}$ is the Gauss map, we have the inequality

$$\mathcal{F}((1 - \lambda)K_1 + \lambda K_2)^{\frac{1}{n-p+1}} \geq (1 - \lambda) \mathcal{F}(K_1)^{\frac{1}{n-p+1}} + \lambda \mathcal{F}(K_2)^{\frac{1}{n-p+1}}, \quad (1.7)$$

for every $K_1, K_2 \in \mathcal{N}$ and $\lambda \in [0, 1]$.

Here we provide a brief overview on the technical novelty and key ingredients in the proof of the above theorem. The first step towards the proof is to express the p -Laplacian on convex rings in terms of the quasi-convex function formed out of the support function on sublevel sets. This is done in Proposition 3.14 following the techniques of [14]; our proof is invariant of choice of coordinates. This is done so that linearity of support functions with respect to Minkowski addition can be used to generate a subsolution on the convex combination set in the next subsection. The next and most important step is a limiting characterization of the functional \mathcal{F} that is devoid of the gradient. In the Capacitary case of [14], the identity

$$\text{Cap}_p(\{u > 0\}) = c(n, p) \left(\lim_{|x| \rightarrow \infty} \frac{u(x)}{G(x)} \right)^{p-1}$$

with $G(x) = 1/|x|^{\frac{n-p}{p-1}}$ and u as the p -capacitary function of $\{u > 0\}$, follows relatively easily, as the capacitary function asymptotically decays like G at infinity. However, it is not so in our case for twofold reasons; firstly our functional has a degeneracy in the weight of the full gradient and secondly, the boundary on which the p -harmonic function vanishes is finite. It requires deeper analysis on the boundary behavior of p -harmonic functions, which, thankfully, has been

previously carried out by Lewis and Nyström in [25]. Resting on a boundary Harnack inequality for p -harmonic functions shown by Lewis-Nyström [25], we prove the limiting characterization

$$\mathcal{F}(\{u > 0\}) = c(n, p, \gamma, \text{diam}(\{u > 0\})) \lim_{s \rightarrow 0^+} \int_{\{u=s\}} \left(\frac{u(x)}{\text{dist}(x, \partial\Omega)} \right)^{p-1} d\mathcal{H}^{n-1}(x)$$

in Proposition 3.27, where γ is a modulus of convexity. This characterization complicates the proof of the Brunn-Minkowski inequality significantly. In the capacity case, the denominator G remains fixed in the limit; we encounter an opposite situation as the numerator within the surface integral in our case remains fixed (equals s). The proof of the theorem is achieved from a completely new geometric argument to prove Theorem 3.78 using projection identities related to the distance function. These tools can also be used in other similar problems with p -harmonic functions vanishing on outer boundaries of finite convex rings.

Lastly, we remark that the case for equality seems to be difficult also due to the structure of the above characterization. It adds to the difficulty for the case of local uniqueness for the corresponding Minkowski problem which is already involved due to the non-uniqueness of the p -harmonic functions prescribing the measures. These are a part of our ongoing work.

2. PRELIMINARIES

Here, we review well-known properties of convex domains, p -harmonic measures and Minkowski problems. For $n \geq 1$, points in \mathbb{R}^n are denoted as $x = (x_1, \dots, x_n) = \sum_{i=1}^n x_i e_i$ where $\{e_1, \dots, e_n\}$ is the standard basis. The standard inner product on \mathbb{R}^n are denoted as $\langle \cdot, \cdot \rangle$ and the Euclidean norm as $|\cdot|$. For functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $F = (f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$, the gradient defined by $\nabla f = \sum_{i=1}^n (\partial_i f) e_i$, the Jacobian defined by $DF = \sum_{i=1}^m \sum_{j=1}^n (\partial_i f_j) e_i \otimes e_j$ and the Hessian is defined by $D^2 f = D(\nabla f) = \sum_{i,j=1}^n (\partial_{i,j}^2 f) e_i \otimes e_j$. For any $A \subset \mathbb{R}^n$, its topological closure and interior shall be denoted as \bar{A} and \mathring{A} and $\partial A = \bar{A} \setminus \mathring{A}$.

Given a measure space (X, μ) and a map $f : X \rightarrow Y$, the pushforward measure $f_*\mu$ on Y is defined on any measurable subset $E \subseteq Y$ as

$$(f_*\mu)(E) = \mu(f^{-1}(E)),$$

which is absolutely continuous with respect to μ and in the infinitesimal form, for any function $g : Y \rightarrow \mathbb{R}$ it is written as $g d(f_*\mu) = (g \circ f) d\mu$. We shall denote \mathcal{H}^k as the k -dimensional Hausdorff measure, $B_r(x)$ as the standard metric ball in \mathbb{R}^n and $\mathbb{S}^{n-1} = \partial B_1(0)$. We shall denote $\omega_n = \mathcal{H}^{n-1}(\mathbb{S}^{n-1})$ as the surface area of the unit sphere.

2.1. Convex domains and support functions. We recall some notions and properties of convex functions and domains. For further details we refer to Schneider [33], Gardner [18], etc.

For any $A \subset \mathbb{R}^n$, the support function $h_A : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as

$$h_A(y) = \sup \{ \langle x, y \rangle : x \in A \}. \quad (2.1)$$

Thus, we have $h_K(y) = \max_{x \in K} \langle x, y \rangle$ for a compact convex subset $K \subset \mathbb{R}^n$. Note that h_K is sub-linear in general and linear (i.e. $h_K(y) = \langle x, y \rangle$ for every $y \in \mathbb{R}^n$) iff $K = \{x\}$. The supporting hyperplane of K with outer normal $y \in \mathbb{R}^n \setminus \{0\}$, is given by

$$H_K(y) := \{ x \in \mathbb{R}^n : \langle x, y \rangle = h_K(y) \}. \quad (2.2)$$

Thus, h_K is differentiable at $y \in \mathbb{R}^n \setminus \{0\}$ with $\nabla h_K(y) = x$ if and only if $K \cap H_K(y) = \{x\}$ and in this case K is called strictly convex. Note that h_K is homogeneous of degree 1 and hence $y \mapsto \nabla h_K(y)$ is homogeneous of degree 0, i.e.

$$h_K(\lambda y) = \lambda h_K(y) \quad \text{and} \quad \nabla h_K(\lambda y) = \nabla h_K(y) \quad \forall y \in \mathbb{R}^n \setminus \{0\}, \lambda \geq 0. \quad (2.3)$$

The support functions are sub-linear (sub-additive and 1-homogeneous) and also conversely, all sub-linear functions are support functions for a corresponding domain, see [33, Theorem 1.7.1]. This leads to characterizing convex set K as

$$K = \bigcap_{\xi \in \mathbb{S}^{n-1}} \{x \in \mathbb{R}^n : \langle x, \xi \rangle \leq h_K(\xi)\}; \quad (2.4)$$

thus convex bodies correspond to Wulff shapes of their support functions. Also note that (2.4) implies $K_1 \subseteq K_2$ iff $h_{K_1} \leq h_{K_2}$. Recall that the Minkowski sum of $E, F \subseteq \mathbb{R}^n$ is defined by

$$E + F := \{x + y : x \in E, y \in F\}; \quad (2.5)$$

and the scalar multiplication $cE := \{cy : y \in E\}$ for any $c \in \mathbb{R}^n$. It is not hard to check that if E, F are convex then so is $E + F$ and αE for all $\alpha \geq 0$. Furthermore, for compact convex sets, the decomposition to Minkowski sums is unique, i.e. if $E + K_1 = E + K_2$ then $K_1 = K_2$, which allows the definition of $E - F = \{x - y : x \in E, y \in F\}$ as well. In particular, we note that $B_r(x) = \{x\} + r B_1(0)$ is the standard Euclidean metric ball. Most importantly, the support functions are Minkowski additive, i.e. for compact convex sets $E, F \subseteq \mathbb{R}^n$, we have

$$h_{\alpha E + \beta F} = \alpha h_E + \beta h_F \quad \forall \alpha, \beta \geq 0, \quad (2.6)$$

and hence, for supporting hyperplanes as in (2.2), $H_{\alpha E + \beta F}(\cdot) = \alpha H_E(\cdot) + \beta H_F(\cdot)$ holds as well. The classical Brunn-Minkowski inequality is given by

$$|E + F|^{1/n} \geq |E|^{1/n} + |F|^{1/n}, \quad (2.7)$$

for any measurable $E, F \subseteq \mathbb{R}^n$, with equality if E and F are homothetic.

The notion of smoothness of boundaries is standard, a domain $\Omega \subset \mathbb{R}^n$ is said to be of class C^k (resp. $C^{k,\alpha}$ for $k \in \mathbb{N}$ and $0 < \alpha \leq 1$) if the domain can be expressed locally as a sub-graph of a function $\phi \in C^k(\mathbb{R}^{n-1})$ (resp. $C^{k,\alpha}(\mathbb{R}^{n-1})$) and the boundary as a graph of it, after a possible rotation. If ∂K is of class C^1 for a convex domain K , then for every $x \in \partial K$ there exists a normal vector at x unique up to scaling. This gives rise to the Gauss map, $\mathbf{g}_K : \partial K \rightarrow \mathbb{S}^{n-1}$, where $\mathbf{g}_K(x)$ is the outer unit normal at $x \in \partial K$ and in terms of support function, we have

$$h_K(\mathbf{g}_K(x)) = \langle x, \mathbf{g}_K(x) \rangle, \quad \forall x \in \partial K. \quad (2.8)$$

Recall that the tangent space at $x \in \partial K$ is given by $T_x(\partial K) = \{y \in \mathbb{R}^n : \langle y, \mathbf{g}_K(x) \rangle = 0\}$ and from (2.8), we have $\mathbf{g}_K \in C^1(\partial K, \mathbb{S}^{n-1})$ and the Weingarten map is defined by

$$\mathcal{W}_K(x) := d\mathbf{g}_K(x) : T_x(\partial K) \rightarrow T_x(\partial K).$$

The principal curvatures are the eigenvalues $\kappa_1(x), \dots, \kappa_{n-1}(x)$ of $\mathcal{W}_K(x)$ and the Gaussian curvature is given by

$$\mathcal{K}_{\partial K}(x) = \det(\mathcal{W}_K(x)) = \kappa_1(x) \dots \kappa_{n-1}(x).$$

If K is strictly convex then $K \cap H_K(\mathbf{g}_K(x)) = \{x\}$ for any $x \in \partial K$ and $\mathbf{g}_K^{-1} : \mathbb{S}^{n-1} \rightarrow \partial K$ is well defined; moreover, h_K being differentiable everywhere in $\mathbb{R}^n \setminus \{0\}$, from (2.8) we have

$$h_K(\xi) = \langle \xi, \mathbf{g}_K^{-1}(\xi) \rangle \quad \text{and} \quad \nabla h_K(\xi) = \mathbf{g}_K^{-1}(\xi), \quad \forall \xi \in \mathbb{S}^{n-1}. \quad (2.9)$$

A convex set K is said to be of class C_+^k for any $k \in \mathbb{N}$, if ∂K is of class C^k and the Gauss map $\mathbf{g}_K : \partial K \rightarrow \mathbb{S}^{n-1}$ is a diffeomorphism. Given any convex set K , there exists a nested sequence K_i of C_+^2 convex sets with $K_{i+1} \subset K_i$ such that

$$K = \bigcap_{i=1}^{\infty} K_i, \quad (2.10)$$

(see [33]), so restricting to the C_+^2 class is sufficient for the arguments of most parts. If K is of class C_+^2 , note that $\mathbf{g}_K^{-1} \in C^1(\mathbb{S}^{n-1}, \partial K)$ and $h_K \in C^2(\mathbb{R}^n \setminus \{0\})$. Hence, from homogeneity of h_K , (2.9) and (2.3), we have

$$h_K(y) = \langle y, \nabla h_K(y) \rangle \quad \text{and} \quad D^2 h_K(y)y = 0, \quad \forall y \in \mathbb{R}^n \setminus \{0\}, \quad (2.11)$$

where the second equality of the above follows from the first. The inverse Weingarten map is denoted as

$$\mathcal{W}_K^{-1}(\xi) := (\mathcal{W}_K(\mathbf{g}_K^{-1}(\xi)))^{-1} = d\mathbf{g}_K^{-1}(\xi) : T_\xi(\mathbb{S}^{n-1}) \rightarrow T_\xi(\mathbb{S}^{n-1})$$

defined for all $\xi \in \mathbb{S}^{n-1}$, which is the non-singular part of $D^2h_K(\xi)$. For any $\xi \in \mathbb{S}^{n-1}$ being fixed, there exists an orthonormal basis $\{e^1, \dots, e^{n-1}, \xi\}$ of \mathbb{R}^n and hence, $\{e^i\}$ span the tangent space $T_\xi(\mathbb{S}^{n-1})$. Any $x \in \mathbb{R}^n$ in this basis is given by

$$x = \sum_{k=1}^{n-1} x^k e^k + \langle x, \xi \rangle \xi \quad \text{with} \quad x^k = \langle x, e^k \rangle. \quad (2.12)$$

Since $D^2h_K(\xi)\xi = 0$ from (2.11), the only non-zero entries of $D^2h_K(\xi)$ are $\langle D^2h_K(\xi)e^j, e^i \rangle$ for $i, j \in \{1, \dots, n-1\}$ which, from (2.9), are the entries of $d\nabla h_K(\xi) = d\mathbf{g}_K^{-1}(\xi) = \mathcal{W}_K^{-1}(\xi)$, with respect to the above basis. In other words, we have

$$D^2h_K(\xi) = \sum_{i,j=1}^{n-1} \langle \mathcal{W}_K^{-1}(\xi)e^j, e^i \rangle e^i \otimes e^j. \quad (2.13)$$

Thus, $D^2h_K(\xi)$ has eigenvalues $\{1/\kappa_1(\mathbf{g}_K^{-1}(\xi)), \dots, 1/\kappa_{n-1}(\mathbf{g}_K^{-1}(\xi)), 0\}$, where κ_i 's are the principal curvatures of ∂K . With $\xi \in U \subset \mathbb{S}^{n-1}$ and a coordinate chart $\varphi : U \rightarrow V \subset \mathbb{R}^{n-1}$, the covariant derivatives of $f : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ of first and second orders are locally defined by

$$(i) \quad \nabla f := \sum_{i=1}^{n-1} (\nabla_i f) e^i, \quad \text{where} \quad \nabla_i f(x) := \partial_i(f \circ \varphi^{-1})(\varphi(x)),$$

$$(ii) \quad \nabla^2 f := \sum_{i,j=1}^{n-1} (\nabla_{i,j} f) e^i \otimes e^j, \quad \text{where} \quad \nabla_{i,j} f(x) := \partial_{i,j}(f \circ \varphi^{-1})(\varphi(x)),$$
(2.14)

In particular, for the choice of $U = \mathbb{S}_+^{n-1} = \{x \in \mathbb{R}^n : \langle x, \xi \rangle = \sqrt{1 - \sum_{i=1}^{n-1} |x^i|^2}\}$ (or the other hemisphere $U = \mathbb{S}_-^{n-1}$ corresponding to the negative square root) and $\varphi = (x^1, \dots, x^{n-1})$, taking h_K restricted to \mathbb{S}^{n-1} , (2.11) can be written in terms of local coordinates which, together with (2.12) and (2.13), leads to

$$\nabla h_K(\xi) = h_K(\xi)\xi + \nabla h_K(\xi), \quad \text{and} \quad \mathcal{W}_K^{-1}(\xi) = h_K(\xi)\mathbb{I} + \nabla^2 h_K(\xi), \quad (2.15)$$

see [4, 11], etc. Therefore, for C_+^2 domains, we have

$$\det(\nabla^2 h_K(\xi) + h_K(\xi)\mathbb{I}) = \det(\mathcal{W}_K^{-1}(\xi)) = 1/\mathcal{K}(\mathbf{g}_K^{-1}(\xi)), \quad (2.16)$$

where \mathcal{K} is the Gaussian curvature. Hence, from transformation rule of the Jacobian, we have

$$\int_{\partial K} f(x) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{S}^{n-1}} f(\mathbf{g}_K^{-1}(\xi)) \det(\nabla^2 h_K(\xi) + h_K(\xi)\mathbb{I}) d\xi. \quad (2.17)$$

As in [4], we denote $\mathcal{C}[A] = \text{cofactor matrix of } A$ so that, from (2.16),

$$\mathcal{C}\nabla^2 h_K + h_K\mathbb{I} = \det(\nabla^2 h_K + h_K\mathbb{I})\mathbb{I} = 1/(\mathcal{K} \circ \mathbf{g}_K^{-1})\mathbb{I}. \quad (2.18)$$

Since $\mathcal{W}_K^{-1} = \nabla^2 h_K + h_K\mathbb{I}$ from (2.15), hence (2.18) can be rewritten as

$$\mathcal{W}_K(x) = \mathcal{K}(x)\mathcal{C}[\nabla^2 h_K + h_K\mathbb{I}](\mathbf{g}_K(x)), \quad \forall x \in \partial K. \quad (2.19)$$

We state following lemma from [4, Lemma 3.44].

Lemma 2.20. *Let $u \in C^2(\partial\Omega)$ and $\{e^1, \dots, e^{n-1}\}$ be an orthonormal frame field of \mathbb{S}^{n-1} such that for any $\xi \in \mathbb{S}^{n-1}$ the unit vectors $e^i = e^i(\xi) \in \mathbb{S}^{n-1}$ span the tangent space $T_\xi(\mathbb{S}^{n-1})$. The covariant derivatives being defined as in (2.14) with respect to the suitable local coordinate charts related to the frame, we have the following:*

$$(1) \quad \langle D^2u(F(\xi))e^i, e^j \rangle = -\mathcal{K}(F(\xi))|\nabla u(F(\xi))|\mathcal{C}_{i,j}[\nabla^2 h + h\mathbb{I}];$$

$$(2) \quad \langle D^2u(F(\xi))\xi, e^i \rangle = -\mathcal{K}(F(\xi)) \sum_j \mathcal{C}_{i,j} [\nabla^2 h + h\mathbb{I}] \nabla_j (|\nabla u(F(\xi))|);$$

where $\mathcal{C}_{i,j}[\cdot] = \langle \mathcal{C}[\cdot]e^j, e^i \rangle$ are entries of the cofactor matrix as in (2.18) for $i, j \in \{1, \dots, n-1\}$ with respect to this frame and $F(\xi) = \mathbf{g}_\Omega^{-1}(\xi) = \nabla h_\Omega(\xi)$.

Remark 2.21. In [4, Lemma 3.44], the function u is assumed to be p -harmonic but it is evident in its proof that the two assertions of the above works for any smooth functions. The p -harmonicity in [4, Lemma 3.44] is required only for proving a third assertion of computing the last term $\langle D^2u(F(\xi))\xi, \xi \rangle$, which is omitted here in the statement of the above lemma.

It is also easy to see that the Gaussian curvature decreases with Minkowski addition at points of respective boundaries with common unit normal i.e.

$$\mathcal{K}_{\partial K_1 + \partial K_2} \leq \mathcal{K}_{\partial K_1} + \mathcal{K}_{\partial K_2}, \quad (2.22)$$

which follows from (2.16) and superadditivity of determinant within positive matrices.

The Hausdorff distance between Borel sets $E, E' \subset \mathbb{R}^n$ is defined as

$$d_{\mathcal{H}}(E, E') = \max \left(\sup\{\text{dist}(y, E) : y \in E'\}, \sup\{\text{dist}(y, E') : y \in E\} \right); \quad (2.23)$$

equivalently, we have $d_{\mathcal{H}}(E, E') = \sup_{y \in \mathbb{R}^n} |\text{dist}(y, E) - \text{dist}(y, E')|$. The Hausdorff distance of convex sets can be characterized by support functions as

$$d_{\mathcal{H}}(E, F) = d_{\mathcal{H}}(\partial E, \partial F) = \|h_E - h_F\|_{L^\infty(\mathbb{S}^{n-1})}, \quad (2.24)$$

see [33, Lemma 1.8.1, Lemma 1.8.14].

2.2. The p -Laplacian and p -harmonic measures. Here we recall p -harmonic measures and Minkowski problems and review some previous results in these directions.

The minimizers of the p -Dirichlet energy functional $w \mapsto \int_\Omega |\nabla w|^p dx$ are weak solutions to the p -Laplacian equation $\Delta_p u = \text{div}(|\nabla u|^{p-2} \nabla u) = 0$ in Ω , satisfying

$$\int_\Omega |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = 0, \quad \forall \phi \in C_0^\infty(\Omega),$$

and are called p -harmonic functions for $1 \leq p < \infty$ which coincide with harmonic functions for $p = 2$. The existence of weak solution $u \in W^{1,p}(\Omega)$ is classical and follows from direct methods. The p -Laplacian is a monotone operator that satisfies comparison principle and thereby uniqueness, furthermore the weak solutions are locally $C^{1,\alpha}$. We collect these in the following and refer to [20, 19, 16, 28, 34, 30], etc. for more details.

Theorem 2.25. *For any bounded domain $\Omega \subset \mathbb{R}^n$ and $1 < p < \infty$, we have the following:*

- (1) (Comparison Principle) *if $\Delta_p v \leq \Delta_p u$ in the weak sense on Ω and $u \leq v$ on $\partial\Omega$ in the trace sense then $u \leq v$ on Ω ;*
- (2) (Regularity) *if $\Delta_p u = 0$ in the weak sense on Ω , then $u \in C_{\text{loc}}^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$.*

Furthermore, the continuity of the gradient of the p -Laplacian of the above implies that if $\nabla u \neq 0$ in $\Omega' \subset \Omega$, then we can conclude $u \in C^\infty(\Omega')$ from Schauder estimates, see [19].

Given a bounded and sufficiently regular domain $\Omega \subset \mathbb{R}^n$, and a neighbourhood N of $\partial\Omega$, if $u \in W^{1,p}(\Omega \cap N)$ is a positive weak solution to the p -Laplacian in $\Omega \cap N$, then upon zero extension $u \in W^{1,p}(N)$. Since p -superharmonic functions form a non-negative distribution as shown in [20], from a theorem of Schwartz using the Riesz representation theorem, there exists a non-negative Radon measure ω_p on $\partial\Omega$ such that

$$\int_\Omega |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = - \int_{\partial\Omega} \phi d\omega_p,$$

for any $\phi \in C_0^\infty(N)$; the measure ω_p is called the p -harmonic measure associated to u . Such measures can also be defined for \mathcal{A} -harmonic functions that are, in general, referred as Riesz

measures, see [20, 23]. If $\partial\Omega$ is $C^{0,1}$, then the unit normal $\nu_{\partial\Omega} = -\nabla u/|\nabla u|$ is defined almost everywhere on $\partial\Omega = \partial\{u > 0\}$ and hence

$$d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega}. \quad (2.26)$$

The function u may have a blow-up inside $\Omega \setminus N$ but the p -harmonicity in $\Omega \cap N$ implies that $u \in C_{\text{loc}}^{1,\alpha}(\Omega \cap N)$ from Theorem 2.25 above. Also, the neighborhood N of $\partial\Omega$ can be chosen up to possible reduction so that we can assume without loss of generality that $\nabla u \neq 0$ in $\Omega \cap N$ so that $u \in C^\infty(\Omega \cap N)$ as stated above, and furthermore, we can assume

$$\|u\|_{L^\infty(\partial N \cap \Omega)} + \|\nabla u\|_{L^\infty(\partial N \cap \Omega)} < \infty,$$

in other words, all possible singularities of u are strictly the interior of $\Omega \setminus \bar{N}$.

The above notion of p -harmonic measure is defined for any open connected domains Ω , including convex domains. The notion extends to a convex set K of non-empty interior in the same way taking Ω as the interior of K so that given a neighbourhood N of ∂K and a function $u \in W^{1,p}(K \cap N)$, we have the notion of a p -harmonic measure ω_p associated to u given by $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial K}$. Then, this can be further extended naturally to the case of K being a convex set of empty interior i.e. $K = \partial K$; given N is a neighborhood of K and $u \in W^{1,p}(N)$ being a positive weak solution to the p -Laplacian in $N \setminus K$ and u vanish on K , there exists a non-negative Radon measure ω_p on K such that

$$\int_N |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = - \int_K \phi d\omega_p,$$

for any $\phi \in C_0^\infty(N)$ similarly as above which is called the p -harmonic measure associated to u , and $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_K$ if K is Lipschitz. We refer to [29] for further details on quasi-linear equations on low dimensional sets.

2.3. Minkowski problem. For $n \geq 3$, we consider convex domains $\Omega \subset \mathbb{R}^n$ and the Gauss map $\mathbf{g}_\Omega : \partial\Omega \rightarrow \mathbb{S}^{n-1}$ where $\mathbf{g}_\Omega(x)$ is the outer unit normal at $x \in \partial\Omega$, defined almost everywhere on the boundary since $\partial\Omega$ is locally Lipschitz. Consider a positive finite Borel measure μ on \mathbb{S}^{n-1} which satisfies the following (necessary) conditions:

$$\begin{aligned} (i) \quad & \int_{\mathbb{S}^{n-1}} |\langle \zeta, \xi \rangle| d\mu(\xi) > 0, \quad \forall \zeta \in \mathbb{S}^{n-1}, \\ (ii) \quad & \int_{\mathbb{S}^{n-1}} \xi d\mu(\xi) = 0. \end{aligned} \quad (2.27)$$

Minkowski problems seek the existence up to a translation of a unique compact convex domain Ω with non-empty interior such that $\mu_\Omega = \mu$, where μ satisfies (i) and (ii) of the above and $\mu_\Omega = (\mathbf{g}_\Omega)_* \eta$ for some prescribed measure η on $\partial\Omega$ and \mathbf{g}_Ω is the Gauss map. The prescribed measure η is typically absolutely continuous with respect to $\mathcal{H}^{n-1} \llcorner_{\partial\Omega}$ and with respect to the Gauss map, we can express the density of the induced measure on \mathbb{S}^{n-1} as

$$d\mu_\Omega(\xi) = \mathcal{F}[h_\Omega](\xi) d\xi \quad (2.28)$$

for a functional \mathcal{F} , where h_Ω is the support function. For the original Minkowski problem, we have $\eta = \mathcal{H}^{n-1}$ and $\mu_\Omega = (\mathbf{g}_\Omega)_* \mathcal{H}^{n-1}$. The case of the given measure being discrete was considered in [32, 31], where the corresponding convex domains are polytopes. The continuous case has been shown in [5, 6, 17] and the smooth case in [10]. If Ω is of class C_+^2 and the covariant derivatives as in (2.14) defined by the charts as shown in the previous subsection, then recalling (2.16) and (2.17), we note that the density (2.28) is the reciprocal of the Gauss curvature, i.e.

$$d\mu_\Omega(\xi) = \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}) d\xi = \frac{d\xi}{\mathcal{K}(\mathbf{g}_\Omega^{-1}(\xi))},$$

and furthermore, if $\mu_\Omega = (\mathbf{g}_\Omega)_*\eta$ where $d\eta = fd\mathcal{H}^{n-1}$ for a function $f : \partial\Omega \rightarrow \mathbb{R}$, then we have

$$\mathcal{F}[h_\Omega](\xi) = \frac{(f \circ \mathbf{g}_\Omega^{-1})(\xi)}{\mathcal{K}(\mathbf{g}_\Omega^{-1}(\xi))} = f(\nabla h_\Omega) \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}). \quad (2.29)$$

The above formulae hold for S_Ω -a.e. $\xi \in \mathbb{S}^{n-1}$ for a general convex domain. As examples of the above, the prescribed measure is the harmonic measure ω at origin in [21] where we have $\mu_\Omega = (\mathbf{g}_\Omega)_*\omega$ and $f = (\partial G/\partial \nu)$ where G is the Green's function with pole at 0; in the case of capacitary measures in [11], we have $f = |\nabla U|^p$ where U is the capacitary function.

In the case of p -harmonic measures, $\mu_\Omega = (\mathbf{g}_\Omega)_*\omega_p$ where ω_p is the p -harmonic measure with respect to a function $u = u_\Omega \in W^{1,p}(\Omega \cap N)$ given by $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega}$ where u is p -harmonic in $\Omega \cap N$ and satisfies

$$\begin{cases} \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0, & \text{in } \Omega \cap N; \\ u > 0, & \text{in } \Omega; \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (2.30)$$

where N is a neighbourhood of $\partial\Omega$; thus, $u \in W^{1,p}(N)$ upon zero extension. It corresponds to the density $f = |\nabla u|^{p-1}$ in (2.29) for $1 < p < \infty$. Henceforth, we shall fix

$$\mu_\Omega = (\mathbf{g}_\Omega)_*\omega_p, \quad \text{where } \omega_p(E) = \int_{E \cap \partial\Omega} |\nabla u|^{p-1} d\mathcal{H}^{n-1}. \quad (2.31)$$

The choice of N is made so that $\nabla u \neq 0$ in $\Omega \cap N$ and $\|u\|_{L^\infty(\bar{N} \cap \Omega)} + \|\nabla u\|_{L^\infty(\bar{N} \cap \Omega)} < \infty$ up to possible reduction, and without loss of generality, we also assume that ∂N is C^∞ . Using the notations (2.31) and (2.26), we have

$$d\mu_\Omega = |\nabla u(F_\Omega(\xi))|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega} = \mathcal{F}[h_\Omega](\xi) d\xi,$$

where $F_\Omega(\xi) := \mathbf{g}_\Omega^{-1}(\xi) = \nabla h_\Omega(\xi)$ and from (2.29), we have

$$\mathcal{F}[h_\Omega](\xi) = |\nabla u(F_\Omega(\xi))|^{p-1} \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}).$$

If $K = \bar{\Omega}$ for a convex domain Ω then we define $\mu_K = \mu_\Omega$. Given a function u satisfying (2.30), the p -harmonic measures for the variations $\Omega^t = \Omega + t\Omega'$ are uniquely defined by measures corresponding to $u(\cdot, t) \in W^{1,p}(\Omega^t \cap N)$, the weak solution of the following Dirichlet problem

$$\begin{cases} \operatorname{div}(|\nabla u(\cdot, t)|^{p-2} \nabla u(\cdot, t)) = 0, & \text{in } \Omega^t \cap N; \\ u(x, t) = 0, & \forall x \in \partial\Omega^t \cap N; \\ u(x, t) = u\left(\frac{x}{1+t}\right), & \forall x \in \partial N \cap \Omega^t; \end{cases} \quad (2.32)$$

for $t \in [-\tau, \tau]$ for $\tau > 0$ small enough, so that upon zero extension, $u(\cdot, t) \in W^{1,p}(N)$. Then the corresponding measures are uniquely defined by

$$d\mu_{\Omega^t} = |\nabla u(F(\xi, t), t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega^t} = \mathcal{F}[h_{\Omega^t}](\xi) d\xi,$$

where $F(\xi, t) := F_{\Omega^t}(\xi) = \mathbf{g}_{\Omega^t}^{-1}(\xi) = \nabla h_\Omega(\xi) + t\nabla h_{\Omega'}(\xi)$ and

$$\mathcal{F}[h_\Omega + th_{\Omega'}](\xi) = |\nabla u(F(\xi, t), t)|^{p-1} \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I} + t(\nabla^2 h_{\Omega'} + h_{\Omega'} \mathbb{I})). \quad (2.33)$$

Hence, the special case $\Omega = \Omega'$ on (2.33) for $\lambda = (1+t) \in (1-\tau, 1+\tau)$ leads to

$$\mathcal{F}[\lambda h_\Omega] = \lambda^{n-p} \mathcal{F}[h_\Omega], \quad (2.34)$$

see [4, Lemma 3.12]. As in [4], letting $\mathcal{N}^\tau(\Omega) = \{\Omega + t\Omega' : 0 \leq |t| < \tau\}$, we denote the functional $\Gamma : \mathcal{N}^\tau(\Omega) \rightarrow \mathbb{R}$ as

$$\Gamma(\Omega + t\Omega') := \int_{\mathbb{S}^{n-1}} h_{\Omega + t\Omega'}(\xi) d\mu_{\Omega^t}(\xi) = \int_{\mathbb{S}^{n-1}} (h_\Omega + th_{\Omega'}) (\xi) \mathcal{F}[h_{\Omega^t}](\xi) d\xi \quad (2.35)$$

and $\Gamma(\bar{\Omega}) := \Gamma(\Omega)$. Then from the homogeneity of support function and (2.34),

$$\Gamma(\lambda\Omega) = \lambda^{n-p+1} \Gamma(\Omega), \quad \forall \lambda = (1+t) \in (1-\tau, 1+\tau). \quad (2.36)$$

Furthermore, we have the following Hadamard type formula, see [4, Proposition 3.76].

Lemma 2.37. *Given convex domains $\Omega + t\Omega'$ with $0 \leq |t| < \tau$ for $\tau > 0$ small enough, $h_{\Omega'} > 0$ and with $d\mu_{\Omega+t\Omega'} = |\nabla u(\cdot, t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial(\Omega+t\Omega')}$ where $u(\cdot, t)$ is the unique solution of the Dirichlet problem (2.32), we have*

$$\frac{d}{dt} \Big|_{t=0} \Gamma(\Omega + t\Omega') = (n - p + 1) \int_{\mathbb{S}^{n-1}} h_{\Omega'} d\mu_{\Omega}, \quad (2.38)$$

where Γ as in (2.35) and μ_{Ω} is as in (2.31).

Using these results, the existence theorem for the Minkowski problem for p -harmonic measures has been proved in [4]. We quote the following from [4, Theorem 1.2].

Theorem 2.39 (Existence). *Given a finite regular Borel measure μ on \mathbb{S}^{n-1} satisfying conditions (2.27), there exists a bounded convex domain Ω with non-empty interior such that $\mu_{\Omega} = \mu$ and $\int_{\mathbb{S}^{n-1}} h_{\Omega} d\mu_{\Omega} = 1$, where μ_{Ω} is as in (2.31) for $1 < p < \infty$.*

Remark 2.40. The constraint $\Gamma(\Omega) = \int_{\mathbb{S}^{n-1}} h_{\Omega} d\mu_{\Omega} = 1$ in the above theorem is evident in the proof of [4, Theorem 1.2].

We complete this section with some clarifying remarks in the following.

Remark 2.41. The existence of domains solving the Minkowski problem for harmonic measures was shown by Jerison [21]. It was also shown in [21] that the corresponding functional may not satisfy its corresponding Brunn-Minkowski inequality. Although our result is stated for $p > 2$, the results in [21] are not in contradiction to $p = 2$ of our case for the following reasons:

- (1) The measures in [21] have the constraint that they are probability measures unlike [4] and our case. This alters the degree of homogeneity of the functional making the corresponding inequality in [21] different from ours.
- (2) The principal difference between [21] and [4] and our case is the fact that the definition of classical harmonic measures is too restrictive. In our notation it corresponds to a particular 2-harmonic measure with $N = \mathbb{R}^n \setminus \{0\}$ strictly and $u = G$, which is the Green's function that blows up at 0, i.e. the inner boundary reduces to a point where the harmonic function blows up. This case does not satisfy the finiteness assumption $\|u\|_{L^\infty(\partial N \cap \Omega)} + \|\nabla u\|_{L^\infty(\partial N \cap \Omega)} < \infty$, which we are allowed to freely choose otherwise given our choice of the neighborhood can be taken with any possible reductions (as a consequence of this, the first variation of \mathcal{F} is self-adjoint in L^2 for [4] but not self-adjoint for [21]). A drawback for the case of p -harmonic measures is that they cannot be uniquely defined and depend on the choice of an a priori p -harmonic function.

3. THE BRUNN-MINKOWSKI INEQUALITY

Here onwards, we fix a compact convex set $K_0 \subset \mathbb{R}^n$ of non-empty interior, a neighbourhood N (open and connected) of ∂K_0 and a function $u_0 \in W^{1,p}(K_0 \cap N)$ that satisfies

$$\begin{cases} \operatorname{div}(|\nabla u_0|^{p-2} \nabla u_0) = 0, & \text{in } \overset{\circ}{K}_0 \cap N; \\ u_0 > 0, & \text{in } \overset{\circ}{K}_0; \\ u_0 = 0, & \text{on } \partial K_0. \end{cases} \quad (3.1)$$

Up to a possible reduction of the choice of N , we can assume without loss of generality that $\nabla u_0 \neq 0$ in $K_0 \cap N$ so that $u_0 \in C^\infty(K_0 \cap N)$. Furthermore, since K_0 is convex, we can assume that N is a convex ring (see [27] for definition) and

$$\|u_0\|_{L^\infty(\partial N \cap K_0)} + \|\nabla u_0\|_{L^\infty(\partial N \cap K_0)} < \infty.$$

We shall establish a local Brunn-Minkowski inequality for a functional \mathcal{F} as in (1.6). For the inequality, we shall consider convex bodies in a neighbourhood of K_0 given by

$$\mathcal{N}_\tau(K_0) = \left\{ \frac{K_0 + tK'}{(1+t)} : 0 \leq |t| < \tau \right\}, \quad (3.2)$$

with $0 \in K'$ and $\tau > 0$ small enough such that N is also a neighbourhood of ∂K i.e. $\partial K \subset N$ and hence $\partial N \cap K = \partial N \cap K_0$ for any $K \in \mathcal{N}_\tau(K_0)$. We label $u_{K_0} = u_0$ and $u_K \in W^{1,p}(K \cap N)$ for any $K \in \mathcal{N}_\tau(K_0)$, as the weak solution of the Dirichlet problem

$$\begin{cases} \operatorname{div}(|\nabla u_K|^{p-2} \nabla u_K) = 0, & \text{in } \overset{\circ}{K} \cap N; \\ u_K(x) = 0, & \forall x \in \partial K \cap N; \\ u_K(x) = u_0(x), & \forall x \in \partial N \cap \overset{\circ}{K}. \end{cases} \quad (3.3)$$

Upon zero extension $u_K \in W^{1,p}(N)$. As in Section 2, we henceforth denote the image of the p -harmonic measure μ_K on \mathbb{S}^{n-1} associated to u_K of (3.3) as

$$\mu_K(E) := \int_{\mathbf{g}_K^{-1}(E)} |\nabla u_K|^{p-1} d\mathcal{H}^{n-1}, \quad \text{for any measurable } E \subseteq \mathbb{S}^{n-1}, \quad (3.4)$$

where \mathbf{g}_K is the Gauss map. In other words, for the p -harmonic measure $d\omega_K = |\nabla u_K|^{p-1} d\mathcal{H}^{n-1}$ on ∂K , we have $\mu_K = (\mathbf{g}_K)_* \omega_K$. The functional $\mathcal{F} : \mathcal{N}_\tau(K_0) \rightarrow \mathbb{R}$ of (1.6) is defined by

$$\mathcal{F}(K) = \int_{\partial K} (h_K \circ \mathbf{g}_K) |\nabla u_K|^{p-1} d\mathcal{H}^{n-1} = \int_{\mathbb{S}^{n-1}} h_K(\xi) d\mu_K(\xi), \quad (3.5)$$

for which we prove the Brunn-Minkowski inequality (2.7) for $K_1, K_2 \in \mathcal{N}(K_0) \subseteq \mathcal{N}_\tau(K_0)$.

A key step towards the proof is to show that supremal convolution of u_{K_1} and u_{K_2} is a subsolution of the p -Laplacian in $(1-\lambda)K_1 + \lambda K_2$. Towards this, we need to obtain the p -Laplacian in terms of the support functions of sub-level sets, which is essential to figure the behavior of p -harmonicity with respect to the Minkowski addition of convex sets. This is the content of the following subsection. It follows along the direction of [14].

3.1. Support functions on sub-level sets. Here we assume Ω as a convex domain of class C_+^2 containing the origin, a neighborhood N of $\partial\Omega$ so that $\Omega \cap N$ is a convex ring, and $u \in C^2(\Omega \cap N)$ such that $u > 0$ and $|\nabla u| > 0$ in $\Omega \cap N$, $u = 0$ on $\partial\Omega$ and $\|u\|_{L^\infty(\partial N \cap \Omega)} + \|\nabla u\|_{L^\infty(\partial N \cap \Omega)} < \infty$. Moreover, we assume

$$\Omega_t := \{x \in \Omega : u(x) > t\}, \quad \forall 0 \leq t \leq \|u\|_{L^\infty(N \cap \Omega)}, \quad (3.6)$$

are convex bodies of class C_+^2 . Note that $\Omega_0 = \Omega$ and for any $0 \leq t \leq \|u\|_{L^\infty(N \cap \Omega)}$, we have $\partial\Omega_t = \{x \in \Omega : u(x) = t\}$ and $\mathbf{g}_{\Omega_t}(x) = -\nabla u(x)/|\nabla u(x)|$ for all $x \in \partial\Omega_t$. Since Ω_t 's are C_+^2 hence $\mathbf{g}_{\Omega_t}^{-1}$ exists and we have the following decomposition of the convex ring,

$$\Omega \cap N = \bigcup_{0 < t < \|u\|_{L^\infty(N \cap \Omega)}} \partial\Omega_t, \quad (3.7)$$

with each $\partial\Omega_t$ being diffeomorphic to \mathbb{S}^{n-1} via $\Theta_u : \Omega \cap N \rightarrow \mathbb{S}^{n-1} \times [0, \|u\|_{L^\infty(N \cap \Omega)}]$, given by

$$\Theta_u(x) = (-\nabla u(x)/|\nabla u(x)|, u(x)) \quad \text{and} \quad \Theta_u^{-1}(\xi, t) = \mathbf{g}_{\Omega_t}^{-1}(\xi). \quad (3.8)$$

We denote the support function corresponding to $\{u > t\}$ as $h_u : \mathbb{R}^n \times [0, \|u\|_{L^\infty}] \rightarrow \mathbb{R}$ as

$$h_u(y, t) := h_{\Omega_t}(y), \quad (3.9)$$

with h_{Ω_t} as in (2.1) and for derivatives of h_u , the notations ∇ and D^2 shall be used as usual for the first coordinates and ∂_t, ∂_t^2 , etc. for the last coordinate.

Now, the goal is to express $\Delta_p u$ entirely in terms of h_u and its derivatives. Towards this, we require the following identities.

Lemma 3.10. *Given Ω and $u \in C^2(\Omega \cap N)$ as above, we have $h_u \in C^2(\mathbb{R}^n \setminus \{0\} \times [0, \|u\|_{L^\infty}))$ and the following holds at any $x \in \Omega \cap N$:*

- (1) $h_u(-\nabla u/|\nabla u|, u) = \langle x, -\nabla u/|\nabla u| \rangle$;
- (2) $\nabla h_u(-\nabla u/|\nabla u|, u) = x$, and $D^2 h_u(-\nabla u/|\nabla u|, u) \nabla u/|\nabla u| = 0$;
- (3) $\partial_t h_u(-\nabla u/|\nabla u|, u) = -1/|\nabla u|$.

Proof. At $x \in \partial\Omega_t$, using $\mathbf{g}_{\Omega_t}(x) = -\nabla u(x)/|\nabla u(x)|$, (3.9) and recalling (2.8), we have

$$h_u(-\nabla u/|\nabla u|, u) = h_u(\mathbf{g}_{\Omega_t}(x), t) = h_{\Omega_t}(\mathbf{g}_{\Omega_t}(x)) = \langle x, \mathbf{g}_{\Omega_t}(x) \rangle = \langle x, -\nabla u(x)/|\nabla u(x)| \rangle,$$

for any $t \in (0, \|u\|_{L^\infty})$. This, together with (3.7), concludes (1). Similarly, (2) follows from (2.9) and (2.11) in the same way using $\nabla h_{\Omega_t}(\xi) = \mathbf{g}_{\Omega_t}^{-1}(\xi)$ and $D^2 h_{\Omega_t}(\xi)\xi = 0$ for $\xi = -\nabla u/|\nabla u|$. To prove (3), first note that the using homogeneity (2.3) on (1) and (2), we have

$$h_u(-\nabla u, u) = \langle x, -\nabla u \rangle \quad \text{and} \quad \nabla h_u(-\nabla u, u) = x. \quad (3.11)$$

Differentiating the first identity of (3.11), the full gradients of both sides yield

$$-D^2 u \nabla h_u(-\nabla u, u) + \partial_t h_u(-\nabla u, u) \nabla u = -(D^2 u)x - \nabla u,$$

and the second identity of (3.11) cancels the first terms of the above, leading to

$$(\partial_t h_u(-\nabla u, u) + 1)\nabla u = 0.$$

Since $|\nabla u| > 0$ in $\Omega \cap N$, it implies $\partial_t h_u(-\nabla u, u) = -1$ which is enough to conclude (3) as ∂_t does not change the homogeneity of the first variable of h_u . This completes the proof. \square

Similarly as (3.9), we denote the Weingarten map and its inverse corresponding to u as

$$\mathcal{W}_u(x, t) := \mathcal{W}_{\Omega_t}(x), \quad \text{and} \quad \mathcal{W}_u^{-1}(y, t) := \mathcal{W}_{\Omega_t}^{-1}(y), \quad (3.12)$$

for $0 \leq t \leq \|u\|_{L^\infty}$. As in Section 2, we choose an orthonormal basis $\{e^1, \dots, e^{n-1}, -\nabla u/|\nabla u|\}$ where $\{e^i : i \in \{1, \dots, n-1\}\}$ span the tangent space $T_{-\nabla u/|\nabla u|}(\mathbb{S}^{n-1})$. As in Section 2, we restrict to $h_u : \mathbb{S}^{n-1} \times [0, \|u\|_{L^\infty}] \rightarrow \mathbb{R}$ and find the identities on second derivatives of h_u with respect to the covariant derivatives as (2.14) on suitable coordinate charts that yield (2.15).

Lemma 3.13. *Let Ω and $u \in C^2(\Omega \cap N)$ be as above, $\{e^1, \dots, e^{n-1}\}$ be an orthonormal frame field of \mathbb{S}^{n-1} such that the unit vectors $e^i = e^i(-\nabla u/|\nabla u|) \in \mathbb{S}^{n-1}$ span the tangent space $T_{-\nabla u/|\nabla u|}(\mathbb{S}^{n-1})$, and the covariant derivatives be defined as in (2.14) with respect to suitable local coordinate charts related to the frame. Then the following holds at any $x \in \Omega \cap N$:*

- (1) $D^2 h_u(-\nabla u/|\nabla u|, u) = \sum_{i,j=1}^{n-1} \langle \mathcal{W}_u^{-1}(-\nabla u/|\nabla u|, u) e^j, e^i \rangle e^i \otimes e^j$;
- (2) $\nabla h_u(-\nabla u/|\nabla u|, u) = x + h_u(-\nabla u/|\nabla u|, u) \nabla u/|\nabla u|$;
- (3) $\nabla^2 h_u(-\nabla u/|\nabla u|, u) = \mathcal{W}_u^{-1}(-\nabla u/|\nabla u|, u) - h_u(-\nabla u/|\nabla u|, u) \mathbb{I}$.

Proof. Similarly as in the proof of the above lemma, from (2.13), (3.12) and (3.7), (1) follows directly. For the rest, note that from (2.15), (3.9) and (3.12), we have

$$\begin{aligned} \nabla h_u(-\nabla u/|\nabla u|, u) &= -h_u(-\nabla u/|\nabla u|, u) \nabla u/|\nabla u| + \nabla h_u(-\nabla u/|\nabla u|, u); \\ \mathcal{W}_u^{-1}(-\nabla u/|\nabla u|, u) &= h_u(-\nabla u/|\nabla u|, u) \mathbb{I} + \nabla^2 h_u(-\nabla u/|\nabla u|, u), \end{aligned}$$

and use Lemma 3.10 and (3.7) to conclude (2) and (3) and complete the proof. \square

Now, we evaluate the p -Laplacian in terms of the support function in the following. It shall be used to generate a sub-solution in the next subsection.

Proposition 3.14. *Given Ω, N and $u \in C^2(\Omega \cap N)$ as above, we have*

$$\Delta_p u = \frac{-1}{(-\partial_t h_u)^{p-1}} \left[\text{Tr} \left[(\nabla^2 h_u + h_u \mathbb{I})^{-1} \right] + \frac{(p-1)}{(\partial_t h_u)^2} \left\{ \langle (\nabla^2 h_u + h_u \mathbb{I})^{-1} \nabla \partial_t h_u, \nabla \partial_t h_u \rangle - \partial_t^2 h_u \right\} \right]$$

with h_u evaluated at $(-\nabla u(x)/|\nabla u(x)|, u(x)) \in \mathbb{S}^{n-1} \times (0, \|u\|_{L^\infty})$ for any $x \in \Omega \cap N$.

Proof. We evaluate u and its derivatives at every $\partial\Omega_t$ and make use of the identities of the above lemmas to obtain relations that hold identically for each $t \in (0, \|u\|_{L^\infty})$, thereby conclude them on $\Omega \cap N$ from (3.7), similarly as in the proof of the above lemmas.

First, with respect to the orthonormal basis $\{e^1, \dots, e^{n-1}, -\nabla u/|\nabla u|\}$ as in Lemma 3.13, we decompose the trace to obtain

$$\begin{aligned}\Delta u &= \text{Tr}(D^2u) = \sum_{i=1}^{n-1} \langle D^2u e^i, e^i \rangle + \langle D^2u \nabla u/|\nabla u|, \nabla u/|\nabla u| \rangle \\ &= -|\nabla u| \text{Tr}(\mathcal{W}_u) + \langle D^2u \nabla u/|\nabla u|, \nabla u/|\nabla u| \rangle,\end{aligned}$$

where the last equality follows from Lemma 2.20, (2.19) and (3.12). Using the above together with the fact that $\nabla(|\nabla u|) = D^2u \nabla u/|\nabla u|$, we obtain the p -Laplacian as

$$\begin{aligned}\Delta_p u &= |\nabla u|^{p-2} \left[\Delta u + (p-2) \langle D^2u \nabla u, \nabla u \rangle / |\nabla u|^2 \right] \\ &= |\nabla u|^{p-2} \left[-|\nabla u| \text{Tr}(\mathcal{W}_u) + (p-1) \langle D^2u \nabla u/|\nabla u|, \nabla u/|\nabla u| \rangle \right] \\ &= -|\nabla u|^{p-1} \left[\text{Tr}(\mathcal{W}_u) - \frac{(p-1)}{|\nabla u|} \langle \nabla(|\nabla u|), \nabla u/|\nabla u| \rangle \right].\end{aligned}\tag{3.15}$$

Now, recalling (3) of Lemma 3.10, we shall use $|\nabla u| = -1/\partial_t h_u(-\nabla u/|\nabla u|, u)$ along with the other identities. To evaluate the last term of (3.15) in terms of h_u , we note that

$$\begin{aligned}\nabla(|\nabla u|) &= \nabla \left(-1/\partial_t h_u(-\nabla u/|\nabla u|, u) \right) = \frac{1}{(\partial_t h_u)^2} \nabla \left(\partial_t h_u(-\nabla u/|\nabla u|, u) \right) \\ &= \frac{1}{(\partial_t h_u)^2} \left[-D(\nabla u/|\nabla u|) \nabla \partial_t h_u(-\nabla u/|\nabla u|, u) + \partial_t^2 h_u(-\nabla u/|\nabla u|, u) \nabla u \right].\end{aligned}$$

Taking the inner product with $\nabla u/|\nabla u|$ on the above, we obtain

$$\begin{aligned}\langle \nabla(|\nabla u|), \nabla u/|\nabla u| \rangle &= \frac{1}{(\partial_t h_u)^2} \left[-\langle D(\nabla u/|\nabla u|) \nabla \partial_t h_u, \nabla u/|\nabla u| \rangle + |\nabla u| \partial_t^2 h_u \right] \\ &= \frac{1}{(\partial_t h_u)^3} \left[-(\partial_t h_u) \langle D(\nabla u/|\nabla u|) \nabla \partial_t h_u, \nabla u/|\nabla u| \rangle - \partial_t^2 h_u \right],\end{aligned}$$

which is applied to (3.15) along with using $\partial_t h_u = -1/|\nabla u|$, to obtain

$$\Delta_p u = \frac{-1}{(-\partial_t h_u)^{p-1}} \left[\text{Tr}(\mathcal{W}_u) + \frac{(p-1)}{(\partial_t h_u)^2} \left[-(\partial_t h_u) \langle D(\nabla u/|\nabla u|) \nabla \partial_t h_u, \nabla u/|\nabla u| \rangle - \partial_t^2 h_u \right] \right].$$

To complete the proof, we need to look into the term in the right hand side of the above containing the Jacobian of $\nabla u/|\nabla u|$, which can be computed as

$$D(\nabla u/|\nabla u|) = \frac{1}{|\nabla u|} \left(D^2u - \frac{1}{|\nabla u|^2} D^2u \nabla u \otimes \nabla u \right).$$

Notice that $D(\nabla u/|\nabla u|) \nabla u = 0$. Furthermore, since $\partial_t h_u$ is also 1-homogeneous, we have $\partial_t h_u(y, t) = \langle y, \nabla \partial_t h_u(y, t) \rangle$ and therefore using (2.12) and (2.14) one can similarly obtain

$$\nabla \partial_t h_u(-\nabla u/|\nabla u|, u) = -\partial_t h_u(-\nabla u/|\nabla u|, u) \nabla u/|\nabla u| + \nabla \partial_t h_u(-\nabla u/|\nabla u|, u),\tag{3.16}$$

for the chosen coordinate chart just as in the case of h_u . Using these, we have that

$$\begin{aligned}\langle D(\nabla u/|\nabla u|) \nabla \partial_t h_u, \nabla u/|\nabla u| \rangle &= \langle D(\nabla u/|\nabla u|) \nabla \partial_t h_u, \nabla u/|\nabla u| \rangle \\ &= \langle D(\nabla u/|\nabla u|)^T \nabla u/|\nabla u|, \nabla \partial_t h_u \rangle.\end{aligned}\tag{3.17}$$

We eliminate the last term by computing the derivative of $\nabla h_u(-\nabla u/|\nabla u|, u) = x$, from (2) of Lemma 3.10. On its j -th component i.e. $\partial_j h_u(\nabla u/|\nabla u|, u) = x_j$, differentiating with ∂_i yields

$$\begin{aligned}\delta_{i,j} &= \partial_i \left(\partial_j h_u(-\nabla u/|\nabla u|, u) \right) \\ &= \sum_k \partial_k \partial_j h_u(-\nabla u/|\nabla u|, u) \partial_i \left(\frac{-\partial_k u}{|\nabla u|} \right) + \partial_t \partial_j h_u(-\nabla u/|\nabla u|, u) \partial_i u.\end{aligned}$$

In matrix form, the above reads as

$$D(-\nabla u/|\nabla u|) D^2 h_u(-\nabla u/|\nabla u|, u) + \nabla u \otimes \nabla \partial_t h_u(\nabla u/|\nabla u|, u) = \mathbb{I},$$

which, upon transposition, leads to

$$D^2 h_u(-\nabla u/|\nabla u|, u) D(-\nabla u/|\nabla u|)^T + \nabla \partial_t h_u(\nabla u/|\nabla u|, u) \otimes \nabla u = \mathbb{I}. \quad (3.18)$$

To generate the rightmost term of (3.17), first we multiply (3.18) with $\nabla u/|\nabla u|$ to get

$$-D^2 h_u(-\nabla u/|\nabla u|, u) D(\nabla u/|\nabla u|)^T \nabla u/|\nabla u| + |\nabla u| \nabla \partial_t h_u(-\nabla u/|\nabla u|, u) = \nabla u/|\nabla u|,$$

and then take inner product with respect to any e^j to obtain

$$\langle D^2 h_u(-\nabla u/|\nabla u|, u) D(\nabla u/|\nabla u|)^T \nabla u/|\nabla u|, e^j \rangle = |\nabla u| \langle \nabla \partial_t h_u(-\nabla u/|\nabla u|, u), e^j \rangle.$$

Now, from (1) of Lemma 3.13 and (2.12), it is easy to see that $D^2 h_u(-\nabla u/|\nabla u|, u) e^j = \mathcal{W}_u^{-1} e^j$ and recalling (2) of Lemma 3.10, we have $D^2 h_u(-\nabla u/|\nabla u|, u) \nabla u/|\nabla u| = 0$. Therefore, taking summation over $j \in \{1, \dots, n-1\}$ of the above and using from (2.12) and (3.16), we have

$$\mathcal{W}_u^{-1} D(\nabla u/|\nabla u|)^T \nabla u/|\nabla u| = |\nabla u| \nabla \partial_t h_u.$$

Thus, we have obtained

$$D(\nabla u/|\nabla u|)^T \nabla u/|\nabla u| = |\nabla u| \mathcal{W}_u \nabla \partial_t h_u. \quad (3.19)$$

The p -Laplacian formula above, together with (3.17) and (3.19), can be re-written as

$$\Delta_p u = \frac{-1}{(-\partial_t h_u)^{p-1}} \left[\text{Tr}(\mathcal{W}_u) + \frac{(p-1)}{(\partial_t h_u)^2} [\langle \mathcal{W}_u \nabla \partial_t h_u, \nabla \partial_t h_u \rangle - \partial_t^2 h_u] \right],$$

and the proof is concluded from $\mathcal{W}_u = (\nabla^2 h_u + h_u \mathbb{I})^{-1}$ on above, recalling Lemma 3.13. \square

Now, in addition to the above, we make a further assumption that the given function u_0 is constant on the inner boundary, i.e. $\partial N \cap \Omega = \{u_0 = \varepsilon_0\}$ for some $\varepsilon_0 > 0$, so that

$$\mathcal{F}(\Omega) = \int_{\partial \Omega} (h_\Omega \circ \mathbf{g}_\Omega) |\nabla u|^{p-1} d\mathcal{H}^{n-1} \quad (3.20)$$

is consistent with (3.5), if $u = u_\Omega$ solves the Dirichlet problem

$$\begin{cases} \text{div}(|\nabla u|^{p-2} \nabla u) = 0, & \text{in } \Omega \cap N; \\ u(x) = 0, & \forall x \in \partial \Omega \cap N; \\ u(x) = \varepsilon_0, & \forall x \in \partial N \cap \Omega. \end{cases} \quad (3.21)$$

The following lemma relates the p -harmonic functions on sub-level sets solving

$$\begin{cases} \text{div}(|\nabla u_{\Omega_t}|^{p-2} \nabla u_{\Omega_t}) = 0, & \text{in } \Omega_t \cap N; \\ u_{\Omega_t}(x) = 0, & \forall x \in \partial \Omega_t \cap N; \\ u_{\Omega_t}(x) = \varepsilon_0, & \forall x \in \partial N \cap \Omega_t, \end{cases} \quad (3.22)$$

where $\Omega_t = \{u > t\}$ is as in (3.6) with u solving (3.21). The neighborhood N is chosen upto possible reduction such that $\partial N \cap \Omega_t = \partial N \cap \Omega$ for all $0 \leq t \leq \varepsilon_0/2$.

Lemma 3.23. *Given u that solves (3.21) and $\Omega_t = \{u > t\}$, let u_{Ω_t} be the solution of the Dirichlet problem (3.22). Then, we have the following for $0 \leq t \leq \varepsilon_0/2$:*

$$(i) \quad u_{\Omega_t} = \frac{u-t}{1-t/\varepsilon_0}, \quad (ii) \quad \dot{u} := \frac{\partial}{\partial t} \Big|_{t=0} u_{\Omega_t} = u/\varepsilon_0 - 1 \quad (3.24)$$

Proof. Let $u_t = (u-t)/(1-t/\varepsilon_0)$, we note that $\Delta_p u_t = 0$ in $\Omega_t \cap N$. Also, since $\partial\Omega_t = \{u = t\}$, hence $u_t = 0$ on $\partial\Omega_t \cap N$ and from boundary condition of (3.21), $u_t = \varepsilon_0$ on $\partial N \cap \Omega_t = \partial N \cap \Omega$. Therefore, u_t solves (3.24) and (i) follows from uniqueness; (ii) follows from computation. \square

To proceed to prove the Brunn-Minkowski inequality of (3.20), we require a limiting characterization of the functional (3.20) in terms of the defining function instead of the gradient. This does not follow easily as in the capacity case in [14, 11] etc. since our case involves a finite boundary where the p -harmonic functions vanish. It requires deeper analysis of the boundary behavior of p -harmonic functions. We require the following boundary Harnack inequality for the p -Laplacian, which is a deep result due to Lewis-Nyström [25, Lemma 4.28].

Lemma 3.25 (Lewis-Nyström). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, $x_0 \in \partial\Omega$ and $u : \Omega \cap \bar{B}_r(x_0) \rightarrow \mathbb{R}$ be a positive, p -harmonic function in $\Omega \cap B_r(x_0)$ with $u = 0$ on $\partial\Omega \cap B_r(x_0)$, for $0 < r < r_0$ and $1 < p < \infty$. Then there exists $\xi \in \mathbb{S}^{n-1}$ and constants $c, \Lambda > 1$ depending on n, p , Lipschitz constant of Ω , such that we have,*

$$\frac{1}{\Lambda} \frac{u(x)}{\text{dist}(x, \partial\Omega)} \leq \langle \nabla u(x), \xi \rangle \leq |\nabla u(x)| \leq \Lambda \frac{u(x)}{\text{dist}(x, \partial\Omega)},$$

for all $x \in \Omega \cap B_{r/c}(x_0)$. Moreover, $\xi \in \mathbb{S}^{n-1}$ can be chosen independently of u .

The following is a technical lemma which would be required.

Lemma 3.26. *Given $a, b \in \mathbb{R}^n$, we have the following:*

- (1) if $\langle a, b \rangle \geq 0$ then we have $0 \leq \int_{\mathbb{S}^{n-1}} \langle a, \omega \rangle \langle \omega, b \rangle d\mathcal{H}^{n-1}(\omega) \leq \omega_n \langle a, b \rangle$;
- (2) if $\langle a, b \rangle \leq 0$ then we have $\omega_n \langle a, b \rangle \leq \int_{\mathbb{S}^{n-1}} \langle a, \omega \rangle \langle \omega, b \rangle d\mathcal{H}^{n-1}(\omega) \leq 0$.

Proof. It follows from easily the fact that $\langle a, \omega \rangle \langle \omega, b \rangle = \langle (a \otimes b)\omega, \omega \rangle$ forms the Rayleigh quotient for any $\omega \in \mathbb{S}^{n-1}$ and $a \otimes b$ is a rank one matrix with eigenvalues $0, \dots, 0, \langle a, b \rangle$. \square

Now, we show the limiting characterization in the following.

Proposition 3.27. *Let Ω be a convex domain of class C_+^2 containing the origin, u be the solution of (3.21) for some $\varepsilon_0 > 0$ and $p > 2$. Let $\Omega_t = \{u > t\}$ as in (3.6) be convex for every $0 \leq t < \varepsilon_0$. Then, given \mathcal{F} as in (3.20), there exists $c = c(n, p, \varepsilon_0, \gamma, \text{diam}(\Omega)) > 0$, where γ is modulus of convexity of Ω , such that we have*

$$\mathcal{F}(\Omega) = c \lim_{s \rightarrow 0^+} \int_{\partial\Omega_s} \left(\frac{u(x)}{\text{dist}(x, \partial\Omega)} \right)^{p-1} d\mathcal{H}^{n-1}(x). \quad (3.28)$$

Proof. In the following, let us use the notation \sim to denote equality up to multiplication by a constant $c = c(n, p, \gamma, \text{diam}(\Omega)) > 0$, equivalently being bounded from above and below by two similar constants. Then, note that it is enough to prove

$$\mathcal{F}(\Omega) \sim \int_{\partial\Omega} |\nabla u|^{p-1} d\mathcal{H}^{n-1} = \lim_{s \rightarrow 0^+} \int_{\partial\Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1}; \quad (3.29)$$

indeed, for any $x \in \partial\Omega_s$ close enough to $\partial\Omega$ with $s > 0$ is small enough, we invoke Lemma 3.25 to have $\Lambda = \Lambda(n, p, \gamma, \text{diam}(\Omega)) > 0$ such that

$$\frac{1}{\Lambda} \frac{u(x)}{\text{dist}(x, \partial\Omega)} \leq |\nabla u(x)| \leq \Lambda \frac{u(x)}{\text{dist}(x, \partial\Omega)},$$

so that the limit (3.28) exists and follows from (3.29). The proof of (3.29) is in the following.

Let u_{Ω_t} be the solution of (3.22) for $0 \leq t \leq \varepsilon_0/2$. Recalling (3.5) and (1) of Lemma 3.10, note that we have

$$\begin{aligned} \mathcal{T}(\{u > t\}) &= \int_{\{u=t\}} (h_{\Omega_t} \circ \mathbf{g}_{\Omega_t}) |\nabla u_{\Omega_t}|^{p-1} d\mathcal{H}^{n-1} = \int_{\{u=t\}} \left\langle x, \frac{-\nabla u}{|\nabla u|} \right\rangle |\nabla u_{\Omega_t}|^{p-1} d\mathcal{H}^{n-1} \\ &= \frac{d}{dt} \left(\int_{\{u>t\}} \langle x, \nabla u \rangle |\nabla u_{\Omega_t}|^{p-1} dx \right), \end{aligned}$$

where the last equality follows from the co-area formula. Integrating both sides of the above from 0 to s , we get

$$\int_0^s \mathcal{T}(\{u > t\}) dt = \int_{\{u>s\}} \langle x, \nabla u \rangle |\nabla u_{\Omega_s}|^{p-1} dx - \int_{\{u>0\}} \langle x, \nabla u \rangle |\nabla u|^{p-1} dx. \quad (3.30)$$

We wish to replace $|\nabla u_{\Omega_s}|^{p-1}$ with $|\nabla u|^{p-1}$ in the first term of (3.30) modulo small error terms. From Lemma 3.23, since $\frac{\partial}{\partial t}|_{t=0} \nabla u_{\Omega_t} = \nabla \dot{u} = \nabla u / \varepsilon_0$, we have the following;

$$\begin{aligned} & \left| \int_{\{u>s\}} \langle x, \nabla u \rangle \left(|\nabla u_{\Omega_s}|^{p-1} - |\nabla u|^{p-1} \right) dx \right| \\ & \leq C \|\nabla u\|_{L^\infty} \|\nabla u_{\Omega_s} - \nabla u\|_{L^\infty}^{p-1} \leq C s^{p-1} \|\nabla u\|_{L^\infty} \|\nabla \dot{u}\|_{L^\infty}^{p-1} = \frac{C}{\varepsilon_0^{p-1}} s^{p-1} \|\nabla u\|_{L^\infty}^p, \end{aligned} \quad (3.31)$$

for some $C = C(n, \text{diam}(\Omega)) > 0$. Using (3.31) we can rewrite (3.30) as follows

$$\begin{aligned} \int_0^s \mathcal{T}(\{u > t\}) dt &= \int_{\{u>s\}} \langle x, \nabla u \rangle |\nabla u|^{p-1} dx - \int_{\{u>0\}} \langle x, \nabla u \rangle |\nabla u|^{p-1} dx + O(s^{p-1}) \\ &= - \int_{\{0 < u < s\}} \langle x, \nabla u \rangle |\nabla u|^{p-1} dx + O(s^{p-1}). \end{aligned}$$

In other words, we have shown that

$$\int_0^s \mathcal{T}(\Omega_t) dt = - \int_{\Omega \setminus \Omega_s} \langle x, \nabla u \rangle |\nabla u|^{p-1} dx + O(s^{p-1}). \quad (3.32)$$

The right hand side of (3.32) is to be estimated using p -harmonicity in $\Omega \setminus \Omega_s$. For small $s > 0$, since $\Omega \setminus \Omega_s \subset \Omega \cap N$, and the outer normal of $\partial(\Omega \setminus \Omega_s)$ being $-\nabla u / |\nabla u|$ for $\partial\Omega$ and $\nabla u / |\nabla u|$ for $\partial\Omega_s$, the p -harmonicity of u implies that for any $\phi \in C^1(\Omega \cap N)$, we have

$$\int_{\Omega \setminus \Omega_s} |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = - \int_{\partial\Omega} \phi |\nabla u|^{p-1} d\mathcal{H}^{n-1} + \int_{\partial\Omega_s} \phi |\nabla u|^{p-1} d\mathcal{H}^{n-1}. \quad (3.33)$$

The right hand side of (3.32) can be generated from the left hand side of the above for a choice of ϕ whose gradient equals $x|\nabla u|$. But it may not exist in general as $x|\nabla u|$ may not be a conservative field. A choice of ϕ close to this can be obtained using the Helmholtz-decomposition $x|\nabla u| = \nabla \phi + R$ where $\text{div}(R) = 0$. Hence, $\Delta \phi = \text{div}(x|\nabla u|)$ and by prescribing boundary values, $\phi = u$ in $\partial(\Omega \setminus \Omega_s)$ the choice is unique. In other words, we choose ϕ_s as the solution of

$$\begin{cases} \Delta \phi_s = \text{div}(x|\nabla u|), & \text{in } \Omega \setminus \Omega_s; \\ \phi_s = 0, & \text{on } \partial\Omega; \\ \phi_s = u, & \text{on } \partial\Omega_s, \end{cases} \quad (3.34)$$

and $R_s = x|\nabla u| - \nabla \phi_s$ is the divergence free component. If we have the Green's kernel $G_s(x, x')$ that solves the Dirichlet problem

$$\begin{cases} -\Delta_{x'} G_s(x, x') = \delta(x' - x), & \text{in } \Omega \setminus \Omega_s; \\ G_s(x, x') = 0, & \text{on } \partial(\Omega \setminus \Omega_s), \end{cases} \quad (3.35)$$

then, it is well-known that ϕ_s solving (3.34) has the following representation

$$\begin{aligned}\phi_s(x) &= - \int_{\Omega \setminus \Omega_s} \operatorname{div}(x' |\nabla u(x')|) G_s(x, x') dx' - \int_{\partial \Omega_s} u(x') \langle \nabla_{x'} G_s(x, x'), \mathbf{g}_{\Omega_s}(x') \rangle d\mathcal{H}^{n-1}(x') \\ &= - \int_{\Omega \setminus \Omega_s} \operatorname{div}(x' |\nabla u(x')|) G_s(x, x') dx' - s \int_{\partial \Omega_s} \langle \nabla_{x'} G_s(x, x'), \mathbf{g}_{\Omega_s}(x') \rangle d\mathcal{H}^{n-1}(x') \\ &= - \int_{\Omega \setminus \Omega_s} \operatorname{div}(x' |\nabla u(x')|) G_s(x, x') dx' - s,\end{aligned}$$

where the second equality follows since $\partial \Omega_s = \{u = s\}$ and the third equality is not hard to see from $-\Delta_{x'} G_s(x, x') = \delta(x' - x)$, as in (3.35). Although $G_s(x, x')$ has a singularity at $x = x'$, integration by parts still can be done for the above, due to the decay rate of the Green's function. Precisely, with any $\varepsilon > 0$ small enough, using standard integral by parts we have

$$\begin{aligned}- \int_{(\Omega \setminus \Omega_s) \setminus B_\varepsilon(x)} \operatorname{div}(x' |\nabla u(x')|) G_s(x, x') dx' &= \int_{(\Omega \setminus \Omega_s) \setminus B_\varepsilon(x)} |\nabla u(x')| \langle x', \nabla_{x'} G_s(x, x') \rangle dx' \\ &\quad - \int_{\partial B_\varepsilon(x)} |\nabla u(x')| G_s(x, x') \langle x', \nu(x') \rangle d\mathcal{H}^{n-1}(x'),\end{aligned}$$

since G_s vanish on $\partial(\Omega \setminus \Omega_s)$ as in (3.35). Now, it is well known that

$$G_s(x, x') = \Gamma(x, x') - \psi_s(x, x'), \quad \text{whenever } x, x' \in \Omega \setminus \Omega_s, x \neq x', \quad (3.36)$$

where $\Gamma(x, x') = \Phi(|x - x'|)$ corresponds to the singular kernel of the Newtonian potential with Φ as the fundamental solution of the Laplacian, and ψ_s is a smooth function that solves the Dirichlet problem $\Delta_{x'} \psi_s(x, x') = 0$ in $\Omega \setminus \Omega_s$ and $\psi_s(x, x') = \Gamma(x, x')$ on $\partial(\Omega \setminus \Omega_s)$. Hence,

$$\begin{aligned}\lim_{\varepsilon \rightarrow 0^+} \left| \int_{\partial B_\varepsilon(x)} |\nabla u(x')| G_s(x, x') \langle x', \nu(x') \rangle d\mathcal{H}^{n-1}(x') \right| \\ \leq \lim_{\varepsilon \rightarrow 0^+} c(n, s, \Omega) \|\nabla u\|_{L^\infty} |\Phi(\varepsilon)| \mathcal{H}^{n-1}(\partial B_\varepsilon(x)) \\ = \lim_{\varepsilon \rightarrow 0^+} c(n, s, \Omega) \|\nabla u\|_{L^\infty} \begin{cases} \varepsilon & \text{for } n \neq 2 \\ \varepsilon \log(\varepsilon) & \text{for } n = 2 \end{cases} = 0.\end{aligned}$$

Therefore, from all of the above, we have obtained

$$\begin{aligned}\phi_s(x) &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\Omega \setminus \Omega_s) \setminus B_\varepsilon(x)} |\nabla u(x')| \langle x', \nabla_{x'} G_s(x, x') \rangle dx' - s \\ &= \text{p.v.} \int_{(\Omega \setminus \Omega_s)} |\nabla u(x')| \langle x', \nabla_{x'} G_s(x, x') \rangle dx' - s.\end{aligned} \quad (3.37)$$

The derivative of ϕ_s can be obtained first by differentiating inside the integral of its Green's representation above. We refer to [19, Lemma 4.1] for the differentiation of the Newtonian potential and conclude

$$\partial_{x_i} \phi_s(x) = - \int_{\Omega \setminus \Omega_s} \operatorname{div}(x' |\nabla u(x')|) \partial_{x_i} G_s(x, x') dx'. \quad (3.38)$$

But now, performing integral by parts on (3.38) similarly as above, would leave a residue since the decay rate has altered with differentiation. Since $\partial_{x_i} \Gamma(x, x') = -\frac{1}{\omega_n} (x_i - x'_i) / |x - x'|^n$ for

all $n \geq 1$, hence we compute the residue as

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \int_{\partial B_\varepsilon(x)} |\nabla u(x')| \partial_{x_i} G_s(x, x') \langle x', \nu(x') \rangle d\mathcal{H}^{n-1}(x') \\
&= \lim_{\varepsilon \rightarrow 0^+} \int_{\partial B_\varepsilon(x)} |\nabla u(x')| \partial_{x_i} \Gamma(x, x') \langle x', \nu(x') \rangle d\mathcal{H}^{n-1}(x') \\
&= \lim_{\varepsilon \rightarrow 0^+} \int_{\partial B_\varepsilon(x)} -\frac{1}{\omega_n} \frac{x_i - x'_i}{|x - x'|^n} |\nabla u(x')| \langle x', \nu(x') \rangle d\mathcal{H}^{n-1}(x') \\
&= -\frac{1}{\omega_n} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{S}^{n-1}} \frac{\omega_i}{\varepsilon^{n-1}} |\nabla u(x - \varepsilon\omega)| \langle x - \varepsilon\omega, -\omega \rangle \varepsilon^{n-1} d\mathcal{H}^{n-1}(\omega) \\
&= \frac{1}{\omega_n} \int_{\mathbb{S}^{n-1}} \omega_i |\nabla u(x)| \langle x, \omega \rangle d\mathcal{H}^{n-1}(\omega),
\end{aligned}$$

from dominated convergence theorem. Therefore, carrying out integral by parts on (3.38) similarly as the case of ϕ_s before and incorporating the above residue leads to

$$\begin{aligned}
\partial_{x_i} \phi_s(x) &= \text{p.v.} \int_{(\Omega \setminus \Omega_s)} |\nabla u(x')| \langle x', \nabla_{x'} \partial_{x_i} G_s(x, x') \rangle dx' - \frac{1}{\omega_n} |\nabla u(x)| \int_{\mathbb{S}^{n-1}} \omega_i \langle x, \omega \rangle d\mathcal{H}^{n-1}(\omega) \\
&= -\text{p.v.} \int_{(\Omega \setminus \Omega_s)} |\nabla u(x')| \langle x', \nabla_{x'} \partial_{x'_i} G_s(x, x') \rangle dx' - \frac{1}{\omega_n} |\nabla u(x)| \int_{\mathbb{S}^{n-1}} \omega_i \langle x, \omega \rangle d\mathcal{H}^{n-1}(\omega) \\
&\quad - \int_{(\Omega \setminus \Omega_s)} |\nabla u(x')| \langle x', \nabla_{x'} (\partial_{x_i} \psi_s(x, x') + \partial_{x'_i} \psi_s(x, x')) \rangle dx'
\end{aligned}$$

where the latter equality holds due to $\partial_{x_i} \Gamma(x, x') = -\partial_{x'_i} \Gamma(x, x')$. Now, let us recall (3.33) with the choice of $\phi = \phi_s$, so that we have

$$\int_{\Omega \setminus \Omega_s} |\nabla u|^{p-2} \langle \nabla u, \nabla \phi_s \rangle dx = \int_{\partial \Omega_s} u |\nabla u|^{p-1} d\mathcal{H}^{n-1}. \quad (3.39)$$

Using the expression of $\partial_{x_i} \phi_s$ from above on (3.39), we obtain the following,

$$\begin{aligned}
& - \int_{(\Omega \setminus \Omega_s)} \text{p.v.} \int_{(\Omega \setminus \Omega_s)} |\nabla u(x)|^{p-2} |\nabla u(x')| \langle D_{x'}^2 G_s(x, x') x', \nabla u(x) \rangle dx' dx - J \\
& - \frac{1}{\omega_n} \int_{\Omega \setminus \Omega_s} \int_{\mathbb{S}^{n-1}} |\nabla u|^{p-1} \langle \nabla u, \omega \rangle \langle \omega, x \rangle d\mathcal{H}^{n-1}(\omega) dx = s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1},
\end{aligned} \quad (3.40)$$

where J is a double integral term containing ψ_s . Now, note that $D_{x'}^2 G_s(x, \cdot)$ is in general a compactly supported distribution satisfying the following distributional identity,

$$\frac{\partial^2 G_s}{\partial x'_i \partial x'_j} = \text{p.v.} \left(\frac{\partial^2 G_s}{\partial x'_i \partial x'_j} \right) - \frac{\delta_{i,j}}{n} \delta(x' - x),$$

which we use to rewrite (3.40) as the following,

$$\begin{aligned}
& -\frac{1}{n} \int_{\Omega \setminus \Omega_s} |\nabla u|^{p-1} \langle x, \nabla u \rangle dx - \frac{1}{\omega_n} \int_{\Omega \setminus \Omega_s} \int_{\mathbb{S}^{n-1}} |\nabla u|^{p-1} \langle \nabla u, \omega \rangle \langle \omega, x \rangle d\mathcal{H}^{n-1}(\omega) dx \\
&= s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + J + \int_{\Omega \setminus \Omega_s} |\nabla u|^{p-2} \sum_{i,j} \partial_{x_i} u \int_{\Omega \setminus \Omega_s} G_s(x, x') \partial_{x'_i x'_j}^2 (x'_j |\nabla u(x')|) dx' dx \\
&= s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + J + \int_{\Omega \setminus \Omega_s} |\nabla u|^{p-2} \langle \nabla u, \int_{\Omega \setminus \Omega_s} G_s(x, x') \nabla_{x'} \text{div}(x' |\nabla u(x')|) dx' \rangle dx
\end{aligned}$$

Since $|G_s(x, x')| \leq C(1 + \Phi(|x - x'|))$, the last term is bounded despite the singularity of G_s and it is of order $O(s^2)$, which can be seen from the following decomposition using coarea formula,

$$\begin{aligned} & \int_{\Omega \setminus \Omega_s} |\nabla u|^{p-2} \langle \nabla u, \int_{\Omega \setminus \Omega_s} G_s(x, x') \nabla_{x'} \operatorname{div}(x' |\nabla u(x')|) dx' \rangle dx \\ &= \int_0^s \int_{\{u=t\}} |\nabla u|^{p-2} \left\langle \frac{\nabla u}{|\nabla u|}, \int_0^s \int_{\{u=t'\}} \frac{G_s(x, x')}{|\nabla u(x')|} \nabla_{x'} \operatorname{div}(x' |\nabla u(x')|) d\mathcal{H}^{n-1}(x') \right\rangle d\mathcal{H}^{n-1}(x) dt' dt. \end{aligned}$$

By a similar decomposition as above, it can be seen the double integral term J is also of $O(s^2)$. Therefore, using this on the above, we can conclude

$$\begin{aligned} & -\frac{1}{n} \int_{\Omega \setminus \Omega_s} |\nabla u|^{p-1} \langle x, \nabla u \rangle dx - \frac{1}{\omega_n} \int_{\Omega \setminus \Omega_s} \int_{\mathbb{S}^{n-1}} |\nabla u|^{p-1} \langle \nabla u, \omega \rangle \langle \omega, x \rangle d\mathcal{H}^{n-1}(\omega) dx \\ &= s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + O(s^2). \end{aligned} \quad (3.41)$$

Furthermore, since $\langle x, -\nabla u \rangle \geq 0$, using Lemma 3.26 to estimate the second term of (3.41), we can finally conclude

$$-\int_{\Omega \setminus \Omega_s} |\nabla u|^{p-1} \langle x, \nabla u \rangle dx \sim s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + O(s^2),$$

as desired. Using the above in (3.32), we get

$$\int_0^s \mathcal{F}(\Omega_t) dt \sim s \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + O(s^2) + O(s^{p-1}).$$

Therefore, we have obtained

$$\int_0^s \mathcal{F}(\Omega_t) dt \sim \int_{\partial \Omega_s} |\nabla u|^{p-1} d\mathcal{H}^{n-1} + O(s) + O(s^{p-2}).$$

Since $p > 2$, letting $s \rightarrow 0^+$ on the above and using Lebesgue differentiation theorem, is enough to establish (3.29), thereby concluding the proof. \square

Remark 3.42. The proof of the above proposition is the only place where $p > 2$ is required. For every other results in this paper $p > 1$ is enough.

3.2. Sub-solutions from supremal convolution. Given u_0 satisfying (3.1) for the convex body $K_0 = \{u_0 \geq 0\}$ and N being an open neighborhood of $\partial K_0 = \{u_0 = 0\}$, we can select ε_0 small enough (depending on modulus of convexity of K_0), such that $\{u_0 \geq t\}$ is convex for all $0 \leq t \leq \varepsilon_0$ and $\{u_0 = \varepsilon_0\} \subseteq N$. Therefore, we can take a reduction of N so that

$$\partial N \cap K_0 = \{u_0 = \varepsilon_0\},$$

which we assume henceforth without loss of generality. Note that N is still a convex ring and for any $K \in \mathcal{N}_\tau(K_0)$, since $\partial N \cap K = \partial N \cap K_0$, (3.3) becomes

$$\begin{cases} \operatorname{div}(|\nabla u_K|^{p-2} \nabla u_K) = 0, & \text{in } \overset{\circ}{K} \cap N; \\ u_K(x) = 0, & \forall x \in \partial K \cap N; \\ u_K(x) = \varepsilon_0, & \forall x \in \partial N \cap \overset{\circ}{K}. \end{cases} \quad (3.43)$$

The maximum principle for the p -Laplacian for (3.43) leads to $\|u_K\|_{L^\infty(K \cap N)} = \varepsilon_0$. The convexity of the sub-level sets of u_K 's for any $K \in \mathcal{N}_\tau(K_0)$, is established in the following. The proof is similar to that of [27, Theorem 1(i)] and is omitted.

Theorem 3.44. *Let $u_K \in C^\infty(\overset{\circ}{K} \cap N)$ be the solution of (3.43) for any $K \in \mathcal{N}_\tau(K_0)$. Then, the set $\{x \in K : u_K(x) > t\}$ is convex for all $0 \leq t \leq \varepsilon_0$.*

Now we are in a position to use Proposition 3.14. Given u_{K_i} as the solutions of (3.43) for $K = K_i$ for $i \in \{1, 2\}$ and any $\lambda \in [0, 1]$, the supremal convolution of u_{K_1} and u_{K_2} is given by

$$u_\lambda^*(x) := \sup_{\substack{y, z \in \mathbb{R}^n \\ x = (1-\lambda)y + \lambda z}} [\min\{u_{K_1}(y), u_{K_2}(z)\}]. \quad (3.45)$$

This is defined so that for any $0 \leq t \leq \varepsilon_0$ and $\lambda \in [0, 1]$, we have

$$\{u_\lambda^* \geq t\} = (1-\lambda)\{u_{K_1} \geq t\} + \lambda\{u_{K_2} \geq t\}. \quad (3.46)$$

Using the p -Laplacian expression in Proposition 3.14, we shall show that u_λ^* is a sub-solution. We need the following technical lemma, we refer to [14, p. 472-473]) for the proof.

Lemma 3.47. *Given any symmetric and positive definite matrices $M_1, M_2 \in \mathbb{R}^n \otimes \mathbb{R}^n$, vectors $z_1, z_2 \in \mathbb{R}^n$ and $t_1, t_2 \in \mathbb{R}$, let $M_\lambda = (1-\lambda)M_1 + \lambda M_2$, $z_\lambda = (1-\lambda)z_1 + \lambda z_2$ and $t_\lambda = (1-\lambda)t_1 + \lambda t_2$ for any $\lambda \in [0, 1]$. Then we have the following:*

$$(i) \quad \langle M_\lambda^{-1} z_\lambda, z_\lambda \rangle \leq (1-\lambda) \langle M_1^{-1} z_1, z_1 \rangle + \lambda \langle M_2^{-1} z_2, z_2 \rangle; \quad (3.48)$$

$$(ii) \quad t_\lambda^2 \operatorname{Tr}(M_\lambda^{-1}) \leq (1-\lambda) t_1^2 \operatorname{Tr}(M_1^{-1}) + \lambda t_2^2 \operatorname{Tr}(M_2^{-1}). \quad (3.49)$$

Using the above Lemma 3.47 and Proposition 3.14, we have the following.

Proposition 3.50. *For any $K_1, K_2 \in \mathcal{N}_\tau(K_0) \cap C_+^2$ and $\lambda \in [0, 1]$, let $K_\lambda = (1-\lambda)K_1 + \lambda K_2$ and u_λ^* be as in (3.45). Then we have $-\Delta_p u_\lambda^* \leq 0$ in $K_\lambda \cap N$.*

Proof. For $i \in \{1, 2\}$, let $h_i(\xi, t) = h_{u_i}(\xi, t)$ as in (3.9) with $u_i = u_{K_i}$ and let us denote

$$M_i = \nabla^2 h_i(\xi, t) + h_i(\xi, t)\mathbb{I}, \quad t_i = \partial_t h_i(\xi, t), \quad z_i = \nabla \partial_t h_i(\xi, t).$$

Let $h_\lambda^*(\cdot, t) = h_{\{u_\lambda^* \geq t\}}$ with u_λ^* as in (3.45) so that from (3.46) and (2.6), we have

$$h_\lambda^* = (1-\lambda)h_1 + \lambda h_2.$$

Hence, the convex combinations as in Lemma 3.47 are given by

$$M_\lambda = \nabla^2 h_\lambda^*(\xi, t) + h_\lambda^*(\xi, t)\mathbb{I}, \quad t_\lambda = \partial_t h_\lambda^*(\xi, t), \quad z_\lambda = \nabla \partial_t h_\lambda^*(\xi, t),$$

for any $\lambda \in [0, 1]$. Also, recalling (3.8), let $\Theta_i = \Theta_{u_i}$ and $\Theta_\lambda^* = \Theta_{u_\lambda^*}$. Then from (3.8) and (3.9),

$$\Theta_\lambda^{*-1} = \nabla h_\lambda^* = (1-\lambda)\nabla h_1 + \lambda\nabla h_2 = (1-\lambda)\Theta_1^{-1} + \lambda\Theta_2^{-1}. \quad (3.51)$$

Note that $K_\lambda \cap N = \{u_\lambda^* \geq 0\}$ from (3.46). Hence, for any $x \in K_\lambda \cap N$, if $(\xi, t) = \Theta_\lambda^*(x)$, then from (3.51), we have $x = (1-\lambda)\Theta_1^{-1}(\xi, t) + \lambda\Theta_2^{-1}(\xi, t)$ which implies

$$\Theta_\lambda^*(x) = (\xi, t) = \Theta_1(x_1) = \Theta_2(x_2), \quad \text{for some } x_1 \in K_1 \cap N, x_2 \in K_2 \cap N.$$

Thus, we can use Proposition 3.14 uniformly at any $(\xi, t) \in \mathbb{S}^{n-1} \times [0, \varepsilon_0]$. We have, $\Delta_p u_1 = 0$ in $K_1 \cap N$ and $\Delta_p u_2 = 0$ in $K_2 \cap N$ from (3.43). Hence, using Proposition 3.14, at any $(\xi, t) \in \Theta_\lambda^*(K_\lambda \cap N)$, we have

$$\begin{aligned} t_1^2 \operatorname{Tr}(M_1^{-1}) + (p-1)[\langle M_1 z_1, z_1 \rangle - \partial_t^2 h_1] &= 0, \quad \text{in } K_1 \cap N, \\ t_2^2 \operatorname{Tr}(M_2^{-1}) + (p-1)[\langle M_2 z_2, z_2 \rangle - \partial_t^2 h_2] &= 0, \quad \text{in } K_2 \cap N. \end{aligned} \quad (3.52)$$

Using Proposition 3.14, Lemma 3.47 and (3.52) respectively, we obtain

$$\begin{aligned} [(-\partial_t h_\lambda^*)^{p-1}](-\Delta_p u_\lambda^*) &= t_\lambda^2 \operatorname{Tr}(M_\lambda^{-1}) + (p-1) [\langle M_\lambda^{-1} z_\lambda, z_\lambda \rangle - \partial_t^2 h_\lambda^*] \\ &\leq (1-\lambda) t_1^2 \operatorname{Tr}(M_1^{-1}) + \lambda t_2^2 \operatorname{Tr}(M_2^{-1}) \\ &\quad + (p-1) [(1-\lambda) \langle M_1^{-1} z_1, z_1 \rangle + \lambda \langle M_2^{-1} z_2, z_2 \rangle] \\ &\quad - (p-1) [(1-\lambda) \partial_t^2 h_1 + \lambda \partial_t^2 h_2] \\ &= (1-\lambda) \{t_1^2 \operatorname{Tr}(M_1^{-1}) + (p-1) [\langle M_1^{-1} z_1, z_1 \rangle - \partial_t^2 h_1]\} \\ &\quad + \lambda \{t_2^2 \operatorname{Tr}(M_2^{-1}) + (p-1) [\langle M_2^{-1} z_2, z_2 \rangle - \partial_t^2 h_2]\} = 0. \end{aligned}$$

Recalling Lemma 3.10, $-\partial_t h_\lambda^* \geq 0$ and the proof is finished. \square

Given any function u defined on a convex ring, let us denote the following,

$$\tilde{u}(x) := \sup_{\substack{y, z \in \mathbb{R}^n \\ x \in [y, z]}} [\min\{u(y), u(z)\}]. \quad (3.53)$$

as in [27]. Then, $u \leq \tilde{u}$ follows directly. Moreover, it is not hard to show that the sub-level sets $\{u \geq t\}$ are convex for $0 \leq t \leq \|u\|_{L^\infty}$ iff $\tilde{u} \leq u$ and hence, $\tilde{u} = u$. This leads to the following.

Corollary 3.54. *For any $K_1, K_2 \in \mathcal{N}_\tau(K_0) \cap C_+^2$ and $\lambda \in [0, 1]$, let $K_\lambda = (1 - \lambda)K_1 + \lambda K_2$ and u_λ^* be as in (3.45). Then we have $u_\lambda^* \leq u_{K_\lambda}$ in $K_\lambda \cap N$.*

Proof. From (3.3) and Proposition 3.50, $-\Delta_p u_\lambda^* \leq 0 = -\Delta_p u_{K_\lambda}$ in \mathring{K}_λ . From the definition (3.45) and boundary conditions of (3.3), since

$$\min_{y \in \partial K_1, z \in \partial K_2} \{u_{K_1}(y), u_{K_2}(z)\} = 0,$$

we have $u_\lambda^* = 0 = u_{K_\lambda}$ in ∂K_λ . Also since $\partial N \cap K = \partial N \cap K_0$ for any $K \in \mathcal{N}_\tau(K_0)$, hence from (3.3), (3.45), and (3.53), note that $u_\lambda^* = \tilde{u}_0 = u_{K_\lambda}$ in $\partial N \cap K_\lambda$. Hence, $u_\lambda^* \leq u_{K_\lambda}$ in $K_\lambda \cap N$ follows from comparison principle (Theorem 2.25) and the proof is finished. \square

Thus, from Corollary 3.54, for any $K_1, K_2 \in \mathcal{N}_\tau(K_0) \cap C_+^2$, we have

$$u_{K_\lambda} \geq \min\{u_{K_1}, u_{K_2}\}, \quad \text{for any } \lambda \in [0, 1]. \quad (3.55)$$

3.3. Homogeneity and Hadamard's formula. We fix any convex $K \in \mathcal{N}_\tau(K_0)$ and recall some notions from subsection 2.3 of Section 2, fitted to this setting. It includes results previously shown in [4] provided in Section 2 and their consequences in our present set up.

Letting $\mathcal{N}^\tau(K) = \{K + tK' : 0 \leq |t| < \tau\}$, $K^t = K + tK'$ for the convex K' as in (3.2) and $u_{K, K'}(\cdot, t) \in W^{1,p}(\mathring{K}^t \cap N)$ as the weak solution of the Dirichlet problem

$$\begin{cases} \operatorname{div}(|\nabla u_{K, K'}(\cdot, t)|^{p-2} \nabla u_{K, K'}(\cdot, t)) = 0, & \text{in } \mathring{K}^t \cap N; \\ u_{K, K'}(x, t) = 0, & \forall x \in \partial K^t \cap N; \\ u_{K, K'}(x, t) = u_0\left(\frac{x}{1+t}\right), & \forall x \in \partial N \cap \mathring{K}^t; \end{cases} \quad (3.56)$$

for any $t \in (-\tau, \tau)$. We may also assume $K, K' \in C_+^2$ without loss of generality, whenever needed. Recall the functional $\Gamma : \mathcal{N}^\tau(K) \rightarrow \mathbb{R}$ as in (2.35), in this case given by

$$\Gamma(K^t) = \int_{\mathbb{S}^{n-1}} h_{K^t}(\xi) \tilde{\mu}_{K^t}(\xi), \quad \text{where } d\tilde{\mu}_{K^t}(\xi) = |\nabla u_{K, K'}(\nabla h_{K^t}(\xi), t)|^{p-1} \det(\nabla^2 h_{K^t} + h_{K^t} \mathbb{I}) d\xi,$$

for which, as noted in subsection 2.3 of Section 2 (Lemma 2.37), it is shown previously that

$$(i) \Gamma(\lambda K) = \lambda^{n-p+1} \Gamma(K), \quad (ii) \left. \frac{d}{dt} \right|_{t=0} \Gamma(K + tK') = (n - p + 1) \int_{\mathbb{S}^{n-1}} h_{K'} d\tilde{\mu}_K, \quad (3.57)$$

for $|t| < \tau$ and $\lambda \in (1 - \tau, 1 + \tau)$. Comparing (3.56) at $t = 0$ with (3.1), note that from uniqueness $u_{K, K'}(x, 0) = u_0(x)$, and from (3.4) and (3.5), we have

$$\tilde{\mu}_K = \mu_K, \quad \text{and} \quad \Gamma(K) = \mathcal{I}(K). \quad (3.58)$$

It is not so in general for $t \neq 0$ and we require a different local variation of convex sets satisfying (3.3) which yields the density of the measure (3.4) corresponding to them. Denoting $\mathcal{N}_\tau(K)$ similarly as (3.2), for small enough $\tau > 0$, let us denote

$$K_t = \frac{K + tK'}{(1+t)} \in \mathcal{N}_\tau(K) \quad \text{for } t \in (-\tau, \tau). \quad (3.59)$$

We establish a relation between $\mathcal{I}(K_t)$ and $\Gamma(K^t)$ which, from (3.57), we show that $\mathcal{I}(K_t)$ have the same local homogeneity and satisfy a similar Hadamard-type variational formula.

Lemma 3.60. *Let $u_{K,K'}(\cdot, t) \in C^1(K^t \cap N)$ be the solution of the Dirichlet problem (3.56) for all $K^t = K + tK' \in \mathcal{N}^\tau(K)$ and $u_{K_t} \in C^1(K_t \cap N)$ be the solution of the Dirichlet problem (3.3) for any $K_t \in \mathcal{N}_\tau(K)$ as in (3.59). Then, for any $y \in K_t \cap N$, we have*

$$u_{K_t}(y) = u_{K,K'}((1+t)y, t), \quad \forall |t| < \tau. \quad (3.61)$$

Proof. Let us denote $v_t(y) = u_{K,K'}((1+t)y, t) = u_{K,K'}(x, t)$ with $x = (1+t)y \in K^t \cap N$ for any $y \in K_t \cap N$. We show that

$$\begin{cases} \operatorname{div}(|\nabla v_t|^{p-2} \nabla v_t) = 0, & \text{in } \mathring{K}_t \cap N; \\ v_t(y) = 0, & \forall y \in \partial K_t \cap N; \\ v_t(y) = u_0(y), & \forall x \in \partial N \cap \mathring{K}_t. \end{cases} \quad (3.62)$$

Indeed, note that from (3.56), $-\Delta_p v_t = 0$ in $\mathring{K}_t \cap N$, $v_t(y) = 0$ for $y \in \partial K_t \cap N$, and when $\tau > 0$ is small enough, $\partial N \cap \mathring{K}^t = \partial N \cap \mathring{K} = \partial N \cap \mathring{K}_t$ for all $|t| < \tau$. Therefore, for any $y \in \partial N \cap \mathring{K}_t$ we have $x = (1+t)y \in \partial N \cap \mathring{K}^t$ and from (3.56),

$$v_t(y) = u_{K,K'}(x, t) = u_0(x/(1+t)) = u_0(y).$$

Comparing (3.62) with (3.3) and using uniqueness, we conclude $v_t = u_{K_t}$ in $K_t \cap N$. \square

Here onwards, we denote a smaller neighborhood $\mathcal{N}(K) \subseteq \mathcal{N}_\tau(K) \cap \mathcal{N}^\tau(K)$.

Corollary 3.63. *Given \mathcal{S} as in (3.5), Γ and convex sets K^t and K_t as above,*

$$\mathcal{S}(K_t) = \frac{\Gamma(K^t)}{(1+t)^{n-p+1}} \quad (3.64)$$

holds whenever $K_t, K^t \in \mathcal{N}(K)$.

Proof. From Lemma 3.60, note that $\nabla u_{K_t}(y) = (1+t)\nabla u_{K,K'}((1+t)y, t)$ for any $y \in K_t \cap N$ and from homogeneity of support function, $h_{K_t}(\cdot) = h_{K^t}(\cdot)/(1+t)$. Using these and recalling (3.4), we obtain

$$\begin{aligned} d\mu_{K_t}(\xi) &= |\nabla u_{K_t}(\nabla h_{K_t}(\xi))|^{p-1} \det(\nabla^2 h_{K_t}(\xi) + h_{K_t}(\xi)\mathbb{I}) d\xi \\ &= \frac{1}{(1+t)^{n-p}} |\nabla u(\nabla h_{K^t}(\xi), t)|^{p-1} \det(\nabla^2 h_{K^t}(\xi) + h_{K^t}(\xi)\mathbb{I}) d\xi = \frac{1}{(1+t)^{n-p}} d\tilde{\mu}_{K^t}(\xi) \end{aligned}$$

which, together with (3.5) and the above, completes the proof. \square

This leads to local homogeneity and a similar Hadamard-type variational formula for \mathcal{S} .

Proposition 3.65. *Given $\mathcal{N}(K) \subseteq \mathcal{N}_\tau(K) \cap \mathcal{N}^\tau(K)$ for $\tau > 0$ small enough, there exists $0 < \tau' \leq \tau$ such that we have the following:*

$$(1) \mathcal{S}(\lambda K) = \lambda^{n-p+1} \mathcal{S}(K) \text{ for all } \lambda \in (1 - \tau', 1 + \tau');$$

$$(2) \text{ for any } K_t = (K + tK')/(1+t) \in \mathcal{N}(K) \text{ with } |t| < \tau', \text{ we have}$$

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{S}(K_t) = (n-p+1) \int_{\mathbb{S}^{n-1}} (h_{K'} - h_K) d\mu_K. \quad (3.66)$$

Proof. Let $\lambda = (1 + \delta t)/(1 + t)$ so that taking $K' = \delta K$ on (3.64) and using (3.57) and (3.58), we obtain

$$\mathcal{S}(\lambda K) = \frac{\Gamma((1 + \delta t)K)}{(1+t)^{n-p+1}} = \frac{(1 + \delta t)^{n-p+1}}{(1+t)^{n-p+1}} \Gamma(K) = \lambda^{n-p+1} \mathcal{S}(K),$$

which proves the first part. To prove the next, by differentiating (3.64), we obtain

$$\frac{d}{dt} \mathcal{S}(K_t) = \frac{1}{(1+t)^{n-p+1}} \frac{d}{dt} \Gamma(K^t) - \frac{(n-p+1)}{(1+t)^{n-p+2}} \Gamma(K^t),$$

which at $t = 0$ together with (3.57) and (3.58), completes the proof. \square

3.4. Proof of the Theorem. The proof of Brunn-Minkowski inequality shall follow as a consequence of the limiting characterization in Proposition 3.27 and Corollary 3.54. To use (3.28) of Proposition 3.27, we require comparison between the infima of distance to the boundaries with respect to convex combinations. Such comparison can be established by looking into the projected points on the boundaries where the infima are achieved.

To this end, we require some preparation. Given any $K \in \mathcal{N}(K_0)$, let us denote

$$\Omega_s(K) = \{x \in K : u_K(x) > s\} \quad (3.67)$$

for any $s \geq 0$, where u_K solves the Dirichlet problem (3.3) or (3.43). Thus $\bar{\Omega}_0(K) = K$ and $\partial\Omega_s(K) = \{u_K = s\}$. For any $x \in \partial\Omega_s(K)$, $\mathbf{g}_{\Omega_s(K)}(x) = -\nabla u_K(x)/|\nabla u_K(x)|$ for all $s \geq 0$. Convexity of $\Omega_s(K)$ for all $0 \leq s < \varepsilon_0$ from Theorem 3.44, implies that for any $x \in \mathbb{R}^n$ there exists a unique projection $p_{\partial K}(x) \in \partial K$ such that

$$|x - p_{\partial K}(x)| = \text{dist}(x, \partial K) \quad \text{and} \quad \mathbf{g}_{\Omega_s}(x) = \frac{x - p_{\partial K}(x)}{|x - p_{\partial K}(x)|}. \quad (3.68)$$

Therefore, from (2.8) and (3.68), note that,

$$h_{\Omega_s(K)}(\mathbf{g}_{\Omega_s(K)}(x)) = \langle x, \mathbf{g}_{\Omega_s(K)}(x) \rangle = \frac{\langle x, x - p_{\partial K}(x) \rangle}{\text{dist}(x, \partial K)}. \quad (3.69)$$

Thus we obtain the following identity,

$$\frac{1}{\text{dist}(x, \partial K)} = \frac{h_{\Omega_s(K)}(\mathbf{g}_{\Omega_s(K)}(x))}{\langle x, x - p_{\partial K}(x) \rangle}, \quad (3.70)$$

which shall be used together with Proposition 3.27, in the proof of Brunn-Minkowski inequality. To estimate the denominator of the right hand side of (3.70), we require the following lemma, which is a variant of standard projection inequalities.

Lemma 3.71. *Given any $x \in \mathbb{R}^n$ and a tangent vector $v \in T_{p_{\partial K}(x)}(\partial K)$, for a convex domain K of class C^1 , we have the following inequality,*

$$\langle x - p_{\partial K}(x), x \rangle \leq \langle x - p_{\partial K}(x), x - v \rangle. \quad (3.72)$$

Proof. Given a tangent vector $v \in T_{p_{\partial K}(x)}(\partial K)$, let $\gamma : [0, \delta] \rightarrow \partial K$, for some small $\delta > 0$, be any curve on ∂K , such that $\gamma(0) = p_{\partial K}(x)$ and $\gamma'(0) = v$, (e.g. $\gamma(t) = \exp_{p_{\partial K}(x)}(tv)$, the flow exponential of ∂K). Let $f : [0, \delta] \rightarrow \mathbb{R}$ be given by

$$f(t) = |x - \gamma(t)|^2,$$

so that $f'(t) = -2\langle x - \gamma(t), \gamma'(t) \rangle$. Since f is a non-negative function that satisfies

$$f(0) = |x - p_{\partial K}(x)|^2 = \text{dist}(x, \partial K)^2 = \min_{t \in [0, \delta]} f(t),$$

from (3.68), therefore we must have $f'(0) \geq 0$. This implies

$$-\langle x - p_{\partial K}(x), v \rangle \geq 0,$$

and by adding $\langle x - p_{\partial K}(x), x \rangle$ to both sides the proof is finished. \square

In the following, we prove another technical lemma to establish the decay rate of (3.70) at points in the boundary of sub-level sets (3.67).

Lemma 3.73. *Given any $K \in \mathcal{N}(K_0)$ and $x \in \partial\Omega_s(K)$ with $\Omega_s(K)$ as in (3.67), we have*

$$\frac{1}{\text{dist}(x, \partial K)} \geq \frac{1}{s\|\nabla u_K\|_{L^\infty(\partial K)}}, \quad (3.74)$$

where $s > 0$ and u_K is the solution of the Dirichlet problem (3.3).

Proof. For any $x \in \partial\Omega_s(K)$, we have $\text{dist}(x, \partial\Omega_s(K)) = 0$. Therefore, recalling the definition of Hausdorff distance (2.23) and it's relation to support functions (2.24), note that

$$\begin{aligned} \text{dist}(x, \partial K) &\leq d_{\mathcal{H}}(\partial\Omega_s(K), \partial K) = d_{\mathcal{H}}(\Omega_s(K), K) = \|h_{\Omega_s(K)} - h_K\|_{L^\infty(\mathbb{S}^{n-1})} \\ &= \|h_{u_K}(\cdot, s) - h_{u_K}(\cdot, 0)\|_{L^\infty(\mathbb{S}^{n-1})} \leq s \|\partial_t h_{u_K}(\cdot, 0)\|_{L^\infty(\mathbb{S}^{n-1})}, \end{aligned} \quad (3.75)$$

where h_{u_K} is as in (3.9). Then (3.74) follows from (3) of Lemma 3.10 to finish the proof. \square

Now we are ready to prove the Brunn-Minkowski inequality. We choose any $K_1, K_2 \in \mathcal{N}(K_0)$. Recalling the definition (3.2) of $\mathcal{N}_\tau(K_0)$, there exists $t_1, t_2 \in (-\tau, \tau)$ such that

$$K_1 = K_{t_1} = \frac{K_0 + t_1 K'}{(1 + t_1)}, \quad \text{and} \quad K_2 = K_{t_2} = \frac{K_0 + t_2 K'}{(1 + t_2)}. \quad (3.76)$$

For any $\lambda \in [0, 1]$, let us denote the linear combination $K_\lambda = (1 - \lambda)K_1 + \lambda K_2$. It is easy to see that we have $K_\lambda \in \mathcal{N}(K_0)$ and from (3.76), the following holds,

$$K_\lambda = K_{t_\lambda} = \frac{K_0 + t_\lambda K'}{(1 + t_\lambda)}, \quad \text{where,} \quad \frac{1}{1 + t_\lambda} = \frac{1 - \lambda}{1 + t_1} + \frac{\lambda}{1 + t_2}. \quad (3.77)$$

The following is an equivalent form of the Brunn-Minkowski inequality.

Theorem 3.78. *For any convex $K_1, K_2 \in \mathcal{N}(K_0)$ and any $\lambda \in [0, 1]$ we have*

$$\mathcal{T}((1 - \lambda)K_1 + \lambda K_2) \geq \min\{\mathcal{T}(K_1), \mathcal{T}(K_2)\}, \quad (3.79)$$

where \mathcal{T} is as in (3.5).

Proof. We can assume that the domains are C_+^2 without loss of generality since (2.7) for general domains can be shown using approximation (2.10) by C_+^2 domains.

Now, recalling (3.28) in Proposition 3.27, we have the following,

$$\mathcal{T}(K) = c \lim_{s \rightarrow 0^+} \int_{\partial\Omega_s(K)} \left(\frac{u_K(x)}{\text{dist}(x, \partial K)} \right)^{p-1} d\mathcal{H}^{n-1}(x), \quad (3.80)$$

where we can regard $c = c(n, p, \gamma_0, \text{diam}(K_0)) > 0$, where γ_0 is modulus of convexity of K_0 , whenever $K \in \mathcal{N}(K_0)$. Therefore, to prove the theorem, it is enough to show

$$\int_{\partial\Omega_s(K_\lambda)} \left(\frac{u_{K_\lambda}(x)}{\text{dist}(x, \partial K_\lambda)} \right)^{p-1} d\mathcal{H}^{n-1}(x) \geq \min_{i \in \{1, 2\}} \int_{\partial\Omega_s(K_i)} \left(\frac{u_{K_i}(x)}{\text{dist}(x, \partial K_i)} \right)^{p-1} d\mathcal{H}^{n-1}(x) + o(1).$$

But from (3.67), since $u_K = s$ on $\partial\Omega_s(K)$ for any K , therefore the above is equivalent to

$$\int_{\partial\Omega_s(K_\lambda)} \frac{d\mathcal{H}^{n-1}(x)}{\text{dist}(x, \partial K_\lambda)^{p-1}} \geq \min_{i \in \{1, 2\}} \int_{\partial\Omega_s(K_i)} \frac{d\mathcal{H}^{n-1}(x)}{\text{dist}(x, \partial K_i)^{p-1}} + \frac{o(1)}{s^{p-1}} \quad (3.81)$$

Towards the proof of (3.81), first we recall that we have shown in Corollary 3.54 that $u_{K_\lambda} \geq u_\lambda^*$. Recalling (3.45) and (3.46), it is equivalent to

$$\{u_{K_\lambda} \geq s\} \supseteq \{u_\lambda^* \geq s\} = (1 - \lambda)\{u_{K_1} \geq s\} + \lambda\{u_{K_2} \geq s\}, \quad (3.82)$$

for every $s \geq 0$. In other words, in terms of the sub-level sets (3.67), we have

$$\overline{\Omega_s(K_\lambda)} \supseteq (1 - \lambda)\overline{\Omega_s(K_1)} + \lambda\overline{\Omega_s(K_2)}. \quad (3.83)$$

We shall prove (3.81) using the identities of projections and the previous lemmas. To this end, first we find the relation of points on the boundaries with respect to a common normal. Since the domains are $C^{2,+}$, the map $\Psi_{\lambda, i}^s : \partial\Omega_s(K_\lambda) \rightarrow \partial\Omega_s(K_i)$ given by

$$\Psi_{\lambda, i}^s := \mathbf{g}_{\Omega_s(K_i)}^{-1} \circ \mathbf{g}_{\Omega_s(K_\lambda)},$$

is a diffeomorphism. Given any $x \in \partial\Omega_s(K_\lambda)$, let $x_i = \Psi_{\lambda, i}^s(x) \in \partial\Omega_s(K_i)$ so that we have




FIGURE 1

$$\mathbf{g}_{\Omega_s(K_\lambda)}(x) = \mathbf{g}_{\Omega_s(K_i)}(x_i) =: \xi,$$

for $i \in \{1, 2\}$, see Figure 1. As $p > 1$, using (3.70) followed by (3.83) and (2.6), we have

$$\frac{1}{\text{dist}(x, \partial K_\lambda)^{p-1}} = \left(\frac{h_{\Omega_s(K_\lambda)}(\mathbf{g}_{\Omega_s(K_\lambda)}(x))}{\langle x, x - p_{\partial K_\lambda}(x) \rangle} \right)^{p-1} \geq \left[\frac{(1-\lambda)h_{\Omega_s(K_1)}(\xi) + \lambda h_{\Omega_s(K_2)}(\xi)}{\langle x, x - p_{\partial K_\lambda}(x) \rangle} \right]^{p-1}.$$

To continue, we use Lemma 3.71 to estimate the denominator of the above. For each $i \in \{1, 2\}$, we pick a tangent vector $v_i \in T_{p_{\partial K_\lambda}(x)}(\partial K_\lambda) \cap H_{i,\lambda}^+$, where $H_{i,\lambda}^+$ is a half-space defined by

$$H_{i,\lambda}^+ = \left\{ v \in \mathbb{R}^n : \langle v, \xi \rangle \geq \langle x - \frac{d_i}{d_\lambda} x_i, \xi \rangle \right\},$$

where $d_i = \text{dist}(x_i, \partial K_i)$, $d_\lambda = \text{dist}(x, \partial K_\lambda)$. Hence, we have $\langle x - v_i, d_\lambda \xi \rangle \leq \langle x_i, d_i \xi \rangle$. Recalling (3.68), we know that

$$\frac{x - p_{\partial K_\lambda}(x)}{d_\lambda} = \xi = \frac{x_i - p_{\partial K_i}(x_i)}{d_i}. \quad (3.84)$$

Therefore, we have $\langle x - v_i, x - p_{\partial K_\lambda}(x) \rangle \leq \langle x_i, x_i - p_{\partial K_i}(x_i) \rangle$, which together with (3.72) of Lemma 3.71, leads to

$$\frac{1}{\langle x, x - p_{\partial K_\lambda}(x) \rangle} \geq \frac{1}{\langle x - v_i, x - p_{\partial K_\lambda}(x) \rangle} \geq \frac{1}{\langle x_i, x_i - p_{\partial K_i}(x_i) \rangle}. \quad (3.85)$$

Using (3.85) and (3.70) to continue the above estimate, we obtain

$$\begin{aligned} \frac{1}{\text{dist}(x, \partial K_\lambda)^{p-1}} &\geq \left[\frac{(1-\lambda)h_{\Omega_s(K_1)}(\mathbf{g}_{\Omega_s(K_1)}(x_1))}{\langle x_1, x_1 - p_{\partial K_1}(x_1) \rangle} + \lambda \frac{h_{\Omega_s(K_2)}(\mathbf{g}_{\Omega_s(K_2)}(x_2))}{\langle x_2, x_2 - p_{\partial K_2}(x_2) \rangle} \right]^{p-1} \\ &= \left[\frac{1-\lambda}{\text{dist}(x_1, \partial K_1)} + \frac{\lambda}{\text{dist}(x_2, \partial K_2)} \right]^{p-1} \geq \min_{i \in \{1, 2\}} \frac{1}{\text{dist}(x_i, \partial K_i)^{p-1}}. \end{aligned} \quad (3.86)$$

To complete the proof, we integrate (3.86) to obtain the following,

$$\begin{aligned} \int_{\partial \Omega_s(K_\lambda)} \frac{d\mathcal{H}^{n-1}(x)}{\text{dist}(x, \partial K_\lambda)^{p-1}} &\geq \min_{i \in \{1, 2\}} \int_{\partial \Omega_s(K_\lambda)} \frac{d\mathcal{H}^{n-1}(x)}{\text{dist}(\Psi_{\lambda,i}^s(x), \partial K_i)^{p-1}} \\ &= \min_{i \in \{1, 2\}} \int_{\partial \Omega_s(K_i)} \frac{|\det(d\Psi_{\lambda,i}^s)^{-1}|}{\text{dist}(x', \partial K_i)^{p-1}} d\mathcal{H}^{n-1}(x'), \end{aligned} \quad (3.87)$$

Now, recall that $\mathcal{W}_{\Omega_s(K)} = d\mathbf{g}_{\Omega_s(K)}$ is the Weingarten map whose determinant equals Gaussian curvature $\mathcal{K}_{\partial\Omega_s(K)}$ (see Section 2). Therefore, we have

$$\det(d\Psi_{\lambda,i}^s)^{-1} = \det(d\mathbf{g}_{\Omega_s(K_\lambda)}^{-1} \circ d\mathbf{g}_{\Omega_s(K_i)}) = \det(\mathcal{W}_{\Omega_s(K_\lambda)}^{-1}) \det(\mathcal{W}_{\Omega_s(K_i)}) = \frac{\mathcal{K}_{\partial\Omega_s(K_i)}}{\mathcal{K}_{\partial\Omega_s(K_\lambda)}}.$$

Now, recalling (2.22), we know that the Gaussian curvature decreases with Minkowski addition, hence $\mathcal{K}_{\partial K_\lambda} \leq \mathcal{K}_{\partial K_i}$. From (3.75), we have $d_{\mathcal{H}}(\Omega_s(K), K) \rightarrow 0^+$ and it is a standard result of Hausdorff convergence of C_+^2 domains that $\mathcal{K}_{\partial\Omega_s(K)} \rightarrow \mathcal{K}_{\partial K}$, $1/\mathcal{K}_{\partial\Omega_s(K)} \rightarrow 1/\mathcal{K}_{\partial K}$ pointwise \mathcal{H}^{n-1} -a.e. as $s \rightarrow 0^+$, see Weil [35, 36]. Therefore, we have

$$\det(d\Psi_{\lambda,i}^s)^{-1} = \frac{\mathcal{K}_{\partial\Omega_s(K_i)}}{\mathcal{K}_{\partial\Omega_s(K_\lambda)}} = \frac{\mathcal{K}_{\partial K_i}}{\mathcal{K}_{\partial K_\lambda}} + o(1) \geq 1 + o(1), \quad \mathcal{H}^{n-1} - a.e. \quad (3.88)$$

Using (3.88) on (3.87) together with (3.74) of Lemma 3.73, we finally obtain

$$\int_{\partial\Omega_s(K_\lambda)} \frac{d\mathcal{H}^{n-1}(x)}{\text{dist}(x, \partial K_\lambda)^{p-1}} \geq \min_{i \in \{1,2\}} \int_{\partial\Omega_s(K_i)} \frac{d\mathcal{H}^{n-1}(x')}{\text{dist}(x', \partial K_i)^{p-1}} + \frac{o(1)}{s^{p-1}}.$$

Thus we have established (3.81) as desired and the proof is complete. \square

The proof of the classical Brunn-Minkowski inequality (2.7) follows from Theorem 3.78 in a standard way as described by Gardner [18], which we provide below for completeness.

Proof of Theorem 1.5. Given $K_1, K_2 \in \mathcal{N}(K_0)$, we take the dilates $K'_1 = \delta_1 K_1$ and $K'_2 = \delta_2 K_2$ such that $\delta_1, \delta_2 \in (1 - \tau', 1 + \tau')$ for $\tau' > 0$ small enough ensuring $K'_1, K'_2 \in \mathcal{N}(K_0)$. Hence, from local homogeneity of Proposition 3.65, we have $\mathcal{F}(K'_i) = \delta_i^{n-p+1} \mathcal{F}(K_i)$ for each $i \in \{1, 2\}$. Now, let

$$m = \min \{ \mathcal{F}(K'_1), \mathcal{F}(K'_2) \} \quad \text{and} \quad \nu = \frac{t/\delta_2}{s/\delta_1 + t/\delta_2},$$

for any $s, t > 0$. Hence, using (3.79) of Theorem 3.78, we get

$$m \leq \mathcal{F}((1 - \nu)K'_1 + \nu K'_2) = \mathcal{F}\left(\frac{(s/\delta_1)K'_1 + (t/\delta_2)K'_2}{s/\delta_1 + t/\delta_2}\right) = \mathcal{F}\left(\frac{sK_1 + tK_2}{s/\delta_1 + t/\delta_2}\right) \quad (3.89)$$

Now given any $\lambda \in [0, 1]$, we choose $s = 1 - \lambda$ and $t = \lambda$ so that when $\tau' > 0$ is small enough, we have $1/[(1 - \lambda)/\delta_1 + \lambda/\delta_2] \in (1 - \tau', 1 + \tau')$. This allows us to use the local homogeneity of Proposition 3.65 on (3.89) again to conclude

$$m \left(\frac{1 - \lambda}{\delta_1} + \frac{\lambda}{\delta_2} \right)^{n-p+1} \leq \mathcal{F}((1 - \lambda)K_1 + \lambda K_2) \quad (3.90)$$

Now, for $p < n + 1$ and the neighborhood $\mathcal{N}(K_0)$ being chosen small enough, the choice of

$$\delta_i = (\mathcal{F}(K_0)/\mathcal{F}(K_i))^{1/(n-p+1)}, \quad \text{for each } i \in \{1, 2\},$$

is admissible for $K_1, K_2 \in \mathcal{N}(K_0)$. In this case, $\mathcal{F}(K'_i) = \mathcal{F}(K_0)$ for each $i \in \{1, 2\}$, hence $m = \mathcal{F}(K_0)$ and it is easy to see that the inequality (1.7) is obtained from (3.90). \square

REFERENCES

- [1] Murat Akman, Jasun Gong, Jay Hineman, John Lewis, and Andrew Vogel. The Brunn-Minkowski inequality and a Minkowski problem for nonlinear capacity. *Mem. Amer. Math. Soc.*, 275(1348):vi+115, 2022. (Cited on page 2.)
- [2] Murat Akman, John Lewis, Olli Saari, and Andrew Vogel. The Brunn-Minkowski inequality and a Minkowski problem for \mathcal{A} -harmonic Green's function. *Adv. Calc. Var.*, 14(2):247–302, 2021. (Cited on page 2.)
- [3] Murat Akman, John Lewis, and Andrew Vogel. Note on an eigenvalue problem with applications to a Minkowski type regularity problem in \mathbb{R}^n . *Calc. Var. Partial Differential Equations*, 59(2):Paper No. 47, 36, 2020. (Cited on page 2.)

- [4] Murat Akman and Shirsho Mukherjee. On the minkowski problem for p -harmonic measures. *Calculus of Variations and Partial Differential Equations*, 2024. (Cited on pages 2, 5, 6, 8, 9, and 20.)
- [5] A. D. Aleksandrov. On the theory of mixed volumes. iii. extension of two theorems of minkowski on convex polyhedra to arbitrary convex bodies. *Mat. Sb. (N.S.)*, 3:27–46, 1938. (Cited on pages 1 and 7.)
- [6] A. D. Aleksandrov. On the surface area measure of convex bodies. *Mat. Sb. (N.S.)*, 6:167–174, 1939. (Cited on pages 1 and 7.)
- [7] Christer Borell. Capacitary inequalities of the Brunn-Minkowski type. *Math. Ann.*, 263(2):179–184, 1983. (Cited on page 1.)
- [8] Christer Borell. Hitting probabilities of killed Brownian motion: a study on geometric regularity. *Ann. Sci. École Norm. Sup. (4)*, 17(3):451–467, 1984. (Cited on page 1.)
- [9] Luis A. Caffarelli, David Jerison, and Elliott H. Lieb. On the case of equality in the Brunn-Minkowski inequality for capacity. *Adv. Math.*, 117(2):193–207, 1996. (Cited on page 1.)
- [10] Shiu Yuen Cheng and Shing Tung Yau. On the regularity of the solution of the n -dimensional Minkowski problem. *Comm. Pure Appl. Math.*, 29(5):495–516, 1976. (Cited on page 7.)
- [11] A. Colesanti, K. Nyström, P. Salani, J. Xiao, D. Yang, and G. Zhang. The Hadamard variational formula and the Minkowski problem for p -capacity. *Adv. Math.*, 285:1511–1588, 2015. (Cited on pages 2, 5, 8, and 14.)
- [12] Andrea Colesanti. Brunn-Minkowski inequalities for variational functionals and related problems. *Adv. Math.*, 194(1):105–140, 2005. (Cited on page 2.)
- [13] Andrea Colesanti and Paola Cuoghi. The Brunn-Minkowski inequality for the n -dimensional logarithmic capacity of convex bodies. *Potential Anal.*, 22(3):289–304, 2005. (Cited on page 2.)
- [14] Andrea Colesanti and Paolo Salani. The Brunn-Minkowski inequality for p -capacity of convex bodies. *Math. Ann.*, 327(3):459–479, 2003. (Cited on pages 1, 2, 10, 14, and 19.)
- [15] Björn E. J. Dahlberg. Estimates of harmonic measure. *Arch. Rational Mech. Anal.*, 65(3):275–288, 1977. (Cited on page 2.)
- [16] E. DiBenedetto. $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. *Nonlinear Anal.*, 7(8):827–850, 1983. (Cited on page 6.)
- [17] W. Fenchel and B. Jessen. Mengenfunktionen und konvexe körper, danske vid. selsk. *Mat.-Fys. Medd.*, 16:1–31, 1938. (Cited on page 7.)
- [18] R. J. Gardner. The Brunn-Minkowski inequality. *Bull. Amer. Math. Soc. (N.S.)*, 39(3):355–405, 2002. (Cited on pages 1, 3, and 25.)
- [19] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition. (Cited on pages 6 and 16.)
- [20] Juha Heinonen, Tero Kilpeläinen, and Olli Martio. *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Dover Publications Inc., 2006. (Cited on pages 6 and 7.)
- [21] David Jerison. Prescribing harmonic measure on convex domains. *Invent. Math.*, 105(2):375–400, 1991. (Cited on pages 2, 8, and 9.)
- [22] David Jerison. A Minkowski problem for electrostatic capacity. *Acta Math.*, 176(1):1–47, 1996. (Cited on page 1.)
- [23] Tero Kilpeläinen and Xiao Zhong. Growth of entire \mathcal{A} -subharmonic functions. *Ann. Acad. Sci. Fenn. Math.*, 28(1):181–192, 2003. (Cited on page 7.)
- [24] John Lewis and Kaj Nyström. Boundary behaviour for p harmonic functions in Lipschitz and starlike Lipschitz ring domains. *Ann. Sci. École Norm. Sup. (4)*, 40(5):765–813, 2007. (Cited on page 2.)
- [25] John Lewis and Kaj Nyström. Boundary behavior and the Martin boundary problem for p harmonic functions in Lipschitz domains. *Ann. of Math. (2)*, 172(3):1907–1948, 2010. (Cited on pages 3 and 14.)
- [26] John Lewis, Kaj Nyström, and Andrew Vogel. On the dimension of p -harmonic measure in space. *J. Eur. Math. Soc. (JEMS)*, 15(6):2197–2256, 2013. (Cited on page 2.)
- [27] John L. Lewis. Capacitary functions in convex rings. *Arch. Rational Mech. Anal.*, 66(3):201–224, 1977. (Cited on pages 9, 18, and 20.)
- [28] John L. Lewis. Regularity of the derivatives of solutions to certain degenerate elliptic equations. *Indiana Univ. Math. J.*, 32(6):849–858, 1983. (Cited on page 6.)
- [29] John L. Lewis and Kaj Nyström. Quasi-linear PDEs and low-dimensional sets. *J. Eur. Math. Soc. (JEMS)*, 20(7):1689–1746, 2018. (Cited on pages 2 and 7.)
- [30] Gary M. Lieberman. Boundary regularity for solutions of degenerate elliptic equations. *Nonlinear Anal.*, 12(11):1203–1219, 1988. (Cited on page 6.)
- [31] Hermann Minkowski. Allgemeine lehrsätze über die convexen polyeder. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, 1897:198–220, 1897. (Cited on pages 1 and 7.)
- [32] Hermann Minkowski. Volumen und Oberfläche. *Math. Ann.*, 57(4):447–495, 1903. (Cited on pages 1 and 7.)
- [33] Rolf Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993. (Cited on pages 1, 3, 4, and 6.)

- [34] Peter Tolksdorf. Regularity for a more general class of quasilinear elliptic equations. *J. Differential Equations*, 51(1):126–150, 1984. (Cited on page 6.)
- [35] Wolfgang Weil. Über die Projektionenkörper konvexer Polytope. *Arch. Math. (Basel)*, 22:664–672, 1971. (Cited on page 25.)
- [36] Wolfgang Weil. Ein Approximationssatz für konvexe Körper. *Manuscripta Math.*, 8:335–362, 1973. (Cited on page 25.)

ARIEL A. AGUAS-BARRENO, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ESSEX, COLCHESTER CO4 3SQ, UNITED KINGDOM

Email address: `aa23688@essex.ac.uk`

MURAT AKMAN, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ESSEX, COLCHESTER CO4 3SQ, UNITED KINGDOM

Email address: `murat.akman@essex.ac.uk`

SHIRSHO MUKHERJEE, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ESSEX, COLCHESTER CO4 3SQ, UNITED KINGDOM

Email address: `shirsho.mukherjee@essex.ac.uk`, `m.shirsho@gmail.com`