

Nonlinear Lebesgue spaces: Curves and geometry

Guillaume Sériey's^{a,*}

^aUniversité Paris Cité, CNRS, MAP5, F-75006, Paris, France

Abstract

This paper is the second in a series by the author and collaborators devoted to the study of geometric and analytic properties of *nonlinear Lebesgue spaces*, that is, L^p spaces of mappings taking values in arbitrary metric spaces. The present article formalizes the pointwise description of their geometric properties—their length structure, bounds on their Alexandrov curvature as well as the definition of a speed for absolutely continuous curves despite the lack of differential structure. To obtain this pointwise description, we first prove a nonlinear analogue of the Fubini–Lebesgue theorem, which yields an identification of L^p curves in nonlinear Lebesgue spaces to mappings taking values in the space of L^p curves. This identification of L^p curves then enables a similar identification for absolutely continuous curves, from which the pointwise description of the geometric properties of nonlinear Lebesgue spaces follows.

Keywords: Nonlinear Lebesgue spaces, Absolutely continuous curves, Càdlàg curves of bounded variation, Length structure, Geodesics, Curvature bounds, Finsler manifolds

Contents

1	Introduction	2
2	Preliminaries	3
2.1	Nonlinear Lebesgue spaces	3
2.1.1	Basic assumptions	3
2.1.2	Measurable mappings	4
2.1.3	Lebesgue mappings	4
2.2	Curves in metric spaces	6
2.2.1	Basic assumptions	6
2.2.2	Sobolev curves and continuous representations	7
2.2.3	Curves of bounded variation and càdlàg representations	8
3	Main results	11
3.1	Characterization of L^p curves in nonlinear Lebesgue spaces	11
3.2	Characterization of absolutely continuous curves for $p > 1$	17
3.3	Characterization of càdlàg curves of bounded variation for $p = 1$	23
4	Applications	27
4.1	Geometry of nonlinear Lebesgue spaces for $p > 1$	27
4.1.1	Characterization of geodesics	27
4.1.2	Characterization of the length structure	28
4.1.3	Characterization of curvature bounds for $p = 2$	31
4.2	Speed of absolutely continuous curves for $p > 1$	33

*Corresponding author

Email address: guillaume.serieys@u-paris.fr (Guillaume Sériey's)


4.2.1	Measurable and Lebesgue sections	33
4.2.2	Speed of absolutely continuous curves in Finsler manifolds	33
4.2.3	Speed of absolutely continuous curves in nonlinear Lebesgue spaces	37
Appendix A	Omitted results	44
Appendix A.1	Omitted results of Section 3.1	44
Appendix B	Reminders	44
Appendix B.1	Length spaces	44
Appendix B.2	Measurable multifunctions	46
Appendix B.3	Differentiable manifolds	46

1. Introduction

Physical phenomena inherently involve noncontinuous signals with values in nonlinear spaces. For instance, in medical imaging the images cannot be assumed continuous due to the multiphase nature of anatomy and can be valued in non-Euclidean spaces, such as the space of symmetric positive definite matrices (as in diffusion tensor imaging, to measure the diffusion of water in tissues [BML94]), the probability simplex (to account for uncertainty in the identification of the nature of tissues [AF05] or to construct probabilistic atlases of organs [MTE+95]) or the space of probability measures on the sphere (as encountered in Q-ball imaging [Tuc04] or constrained spherical deconvolution [TCGC04]). When working with this type of data, be it for reconstruction or statistical analysis purposes, there is a stake on the choice of a *meaningful* metric on the space of values, which usually makes it nonlinear. In addition, while images are signals supported on flat spaces, signals supported on curved spaces such as functional shapes [Cha13, Definition 2.1.1.] are also commonly encountered. For all those reasons, studying merely measurable mappings defined on measurable spaces with values in arbitrary metric spaces is of importance from an applicative viewpoint.

From a more theoretical perspective, L^p spaces of measurable mappings taking values in arbitrary metric spaces, which we call *nonlinear Lebesgue spaces*, first emerged in the study of Sobolev spaces of measurable mappings taking values in arbitrary metric spaces and, in particular, of harmonic mappings, starting with the seminal work of N. J. Korevaar and R. M. Schoen [KS93] followed by the works of K.-T. Sturm [Stu01, Stu02] and J. Jost [Jos94, Jos97a, Jos97b]. Nonlinear Lebesgue spaces typically model the space of transport maps in optimal transport theory [AGS05, Section 5.4], but are also useful to study curves in arbitrary metric spaces such as Wasserstein spaces, as studied by S. Lisini [Lis07, Section 2.3], and recently emerged in a generalization, proposed by M. Bauer, F. Mémoli, T. Needham and M. Nishino [BMNN25, Section 2.1.3], of the Gromov–Wasserstein metric to measure spaces equipped with a kernel taking values in an arbitrary metric space instead of a standard metric. The analytical properties of these spaces derived in particular cases in the mentioned works—the characterizations of their completeness and separability, as well as the density of the spaces of simple, continuous, and smooth mappings—are gathered and generalized under a unified exposition in a recent work by the author and A. Trouvé [ST25]. The main behaviour of nonlinear Lebesgue spaces that emerge from these previous studies is the fact that their properties are essentially pointwise in nature, in the sense that they are mostly determined by the properties of the space of values and vice versa. This is particularly true for their geometric properties, as first highlighted by K.-T. Sturm [Stu01, Stu02] and J. Jost [Jos94, Jos97a, Jos97b], who derived curvature properties when the target space is nonpositively curved (in the sense of Alexandrov, see [BBI22, Chapter 4] for a definition) through the construction of geodesics using geodesics in the space of values. Yet, this pointwise behaviour is, to the best of our knowledge, not rigorously formalized, so that the study of the geometry of nonlinear Lebesgue spaces is limited by a lack of a proper characterization of their curves. In fact, one would expect a behaviour that is analogous to that of curves in Wasserstein spaces, studied by S. Lisini [Lis06, Lis07], who exhibited a superposition principle for absolutely continuous curves (i.e., any such curve can be identified to a probability measure on absolutely continuous curves in the space supporting the measures). Let us conclude with a note on terminology. As highlighted in [Bou13, TVS V.86], L^p spaces of real-valued mappings were first introduced by F. Riesz in [Rie10]. Yet, they are commonly referred to as “Lebesgue” spaces [DS88, Chapter III, Section 3] or “Bochner” spaces when mappings are valued in a Banach space [HVN VW16, Section 1.2.b]. For consistency with the literature, we use the terminology of “nonlinear Lebesgue spaces” to designate L^p spaces of mappings taking values in arbitrary metric spaces, terminology that was first used by M. Bačák in [Bac14, p. 18].

In this article, we formalize the pointwise description of the geometric properties of nonlinear Lebesgue spaces resulting from a characterization of absolutely continuous (a.c.) curves, thus providing a complementary perspective to the analytic properties of nonlinear Lebesgue spaces studied in [ST25]. To do so, we first prove a nonlinear analogue of the Fubini–Lebesgue theorem (Theorem 3.9) that allows for the characterization of L^p curves in nonlinear Lebesgue spaces through the identification of a canonical isometry (Theorem 3.13). Using Theorem 3.13, we characterize a.c. curves (Theorem 3.19) as well as càdlàg curves of bounded variation (Theorem 3.23), depending on the value of the parameter p of nonlinear Lebesgue spaces. The characterization of a.c. curves given by Theorem 3.19 notably allows for the pointwise description of the length structure of nonlinear Lebesgue spaces (Theorem 4.9), for the definition of a notion of speed for a.c. curves in nonlinear Lebesgue spaces despite the lack of a differential structure (Theorem 4.44), and for the identification of geodesics as mappings into the space of geodesics (Theorem 4.2). This particular characterization of geodesics then enables the pointwise description of the curvature, in the sense of Alexandrov, of nonlinear Lebesgue spaces (Theorem 4.15). To put it visually:



The paper is structured as follows:

- Section 2 introduces the basic setting of this article regarding minimal assumptions and introduces the core definitions and results needed to construct and study nonlinear Lebesgue spaces (Section 2.1) as well as Sobolev curves and curves of bounded variation in metric spaces (Section 2.2).
- Section 3 gathers the main results of this article, including the characterization of L^p curves in nonlinear Lebesgue spaces (Section 3.1) and its consequence for the characterization of a.c. curves (Section 3.2) and càdlàg curves of bounded variation (Section 3.3) in these spaces.
- Section 4 exposes applications of the main results established in Section 3, including the consequences of Section 3.2 for the pointwise characterization of the geometry of nonlinear Lebesgue spaces (Section 4.1) as well as the definition of a notion of speed for a.c. curves in these spaces (Section 4.2.1).

2. Preliminaries

2.1. Nonlinear Lebesgue spaces

This section introduces notations, definitions and results that were presented in the paper [ST25] by the author and A. Trouvé, only keeping the ones that will be needed in the present article. The reader already familiar with [ST25] may skip to Section 2.2.

2.1.1. Basic assumptions

Let us first introduce the basic assumptions of this article.

Assumption 2.1 (Basic assumptions). Throughout this article, we consider mappings from a nonempty set M , called the *base space*, to a nonempty set N , called the *target space*. M and N are assumed to carry the following structures:

- (i) (M, Σ_M, μ_M) is a measure space, that is, the set M is paired with a σ -algebra Σ_M on M and a measure $\mu_M : \Sigma_M \rightarrow [0, \infty]$ on (M, Σ_M) .
- (ii) (N, d_N) is a metric space with finite metric $d_N : N^2 \rightarrow [0, \infty)$.

Throughout this article, we shall not recall these basic assumptions in the statements of the results; yet, any additional assumption relative to Assumption 2.1 will be specified.

Notation and terminology are progressively introduced throughout the article and are also gathered in a dedicated section (p. 42) at the end of this article.

2.1.2. Measurable mappings

In this section, we introduce several classes of mappings that will serve as core ingredients in the construction of nonlinear Lebesgue spaces, namely, the set of measurable mappings and its variant.

Definition 2.2 (Measurable mappings). Define:

- (i) $\mathcal{L}^0(M, N)$ the set of *measurable mappings*, that is, all mappings $f : M \rightarrow N$ such that for all set B in $\mathcal{B}(N)$, the Borel σ -algebra of N , it satisfies $f^{-1}(B) \in \Sigma_M$.
- (ii) $\mathcal{L}_s^0(M, N)$ the set of *separably valued measurable mappings*, that is, all mappings $f \in \mathcal{L}^0(M, N)$ such that its range, denoted by $f(M)$, is separable.

Remark 2.3 (“Borel measurable” mappings). When M is a topological space with $\Sigma_M = \mathcal{B}(M)$, the Borel σ -algebra of M , a mapping belonging to $\mathcal{L}^0(M, N)$ is usually called “Borel measurable”.

A first well-known result on this class of mappings is that the pointwise limit of measurable and separably valued mappings, when it exists, is itself measurable and separably valued [ST25, Proposition 2.6].

Proposition 2.4 ($\mathcal{L}_s^0(M, N)$ is closed under pointwise limit). Let $(f_n)_{n \in \mathbb{N}}$ be a sequence in $\mathcal{L}_s^0(M, N)$ and assume that $f(x) := \lim_{n \rightarrow \infty} f_n(x)$ exists in N for all $x \in M$. Then, $f \in \mathcal{L}_s^0(M, N)$.

In particular, Proposition 2.4 holds for sequences of *simple mappings*, that is, measurable mappings with a finite range.

Definition 2.5 (Simple mappings). Define $\mathcal{E}(M, N)$ the set of *simple mappings*, that is, all mappings $f \in \mathcal{L}^0(M, N)$ such that $|f(M)|$, the cardinality of the range of f , is finite.

Showing that a mapping belongs to $\mathcal{L}_s^0(M, N)$ thus usually comes down to verifying that it is the pointwise limit of a sequence of simple mappings.

Now, one can define an equivalence relation on the set of measurable mappings.

Proposition 2.6 (Equivalence relation on $\mathcal{L}^0(M, N)$). The relation $f \sim f'$ if and only if $f(x) = f'(x)$ holds for μ_M -a.e. $x \in M$ is an equivalence relation on $\mathcal{L}^0(M, N)$.

Using this equivalence relation, one can define the sets of equivalence classes of measurable and simple mappings.

Definition 2.7 (Equivalence classes of measurable mappings). Define:

- (i) $L^0(M, N) := \mathcal{L}_s^0(M, N) / \sim$ the set of equivalence classes of separably valued measurable mappings, that is, all $[f] := \{f' \in \mathcal{L}^0(M, N) : f \sim f'\}$ with $f \in \mathcal{L}_s^0(M, N)$.
- (ii) $E(M, N) := \mathcal{E}(M, N) / \sim$ the set of equivalence classes of simple mappings.

From these first building blocks, we move on to the construction of nonlinear Lebesgue spaces.

2.1.3. Lebesgue mappings

Before defining nonlinear Lebesgue spaces, let us first define the \mathcal{L}^p semi-metrics (see [BBI22, Definition 1.1.4] for a definition of the notion of semi-metric).

Definition 2.8 (\mathcal{L}^p semi-metrics). Let $p \in [1, \infty]$. Then, define the mapping $D_p : \mathcal{L}_s^0(M, N)^2 \rightarrow [0, \infty]$ as $D_p(f, f') := \|d_N(f, f')\|_{p, \mu_M}$, where $d_N(f, f')(x) := d_N(f(x), f'(x))$. Precisely, for $p \in [1, \infty)$,

$$D_p(f, f') = \left(\int_M d_N(f(x), f'(x))^p d\mu_M(x) \right)^{1/p}$$

and, for $p = \infty$,

$$D_\infty(f, f') = \mu_M\text{-ess sup}_{x \in M} d_N(f(x), f'(x))$$

with the μ_M -essential supremum defined as

$$\mu_M\text{-ess sup}_{x \in M} d_N(f(x), f'(x)) := \inf \{C \in \mathbb{R} : d_N(f(x), f'(x)) \leq C \text{ for } \mu_M\text{-a.e. } x \in M\}.$$

At this point, D_p satisfies the symmetry and triangle inequality axioms of a metric on $\mathcal{L}_s^0(M, N)$ thanks to the symmetry of d_N and the fact that both d_N and $\|\cdot\|_{p, \mu_M}$ satisfy the triangle inequality for $p \in [1, \infty]$. However, D_p is not necessarily finite and cannot distinguish two distinct measurable mappings that agree μ_M -almost everywhere. In fact, D_p separates equivalence classes in $L^0(M, N)$.

Proposition 2.9 (\mathcal{L}^p semi-metrics separate equivalence classes). *Let $p \in [1, \infty]$ and $(f, f') \in \mathcal{L}_s^0(M, N)$. Then, $D_p(f, f') = 0$ if and only if $f \sim f'$.*

Remark 2.10 (\mathcal{L}^p semi-metrics metrize $L^0(M, N)$). Note that $L^0(M, N)$ equipped with D_p , for any choice of $p \in [1, \infty]$, satisfies all the axioms of a metric space, but with a metric D_p that might be infinite.

First, to make D_p finite, consider its restriction to \mathcal{L}^p mappings, which are defined as mappings at a finite D_p -distance from a base mapping h in $L^0(M, N)$.

Definition 2.11 (\mathcal{L}^p spaces). Let $h \in L^0(M, N)$ (recall that h has a representative with separable range by Definition 2.2) and $p \in [1, \infty]$. Then, define the set of measurable mappings at a finite D_p -distance from h as

$$\mathcal{L}_h^p(M, N) := \{f \in \mathcal{L}_s^0(M, N) : D_p(f, h) < \infty\}.$$

Remark 2.12 (Choice of the base mapping in the linear case). When N is a normed vector space, the base mapping h is usually taken as $h \equiv 0_N$, with 0_N the identity element of N .

Then, $(\mathcal{L}_h^p(M, N), D_p)$ satisfies all axioms of a metric space, except separation: it is a *semi-metric* space (see [BB122, Definition 1.1.4] for a definition). We will often write $\mathcal{L}_h^p(M, N)$ instead of $(\mathcal{L}_h^p(M, N), D_p)$.

Now, to make it a metric space, μ_M -a.e. identical mappings should be identified through the equivalence relation defined in Proposition 2.6.

Definition 2.13 (Nonlinear Lebesgue spaces). Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Then, the nonlinear p -Lebesgue space $L_h^p(M, N)$ is defined as the quotient space

$$L_h^p(M, N) := \mathcal{L}_h^p(M, N) / \sim,$$

where we recall that $f \sim f'$ if and only if $D_p(f, f') = 0$ (Proposition 2.9). Also, we will usually omit p when referring to nonlinear p -Lebesgue spaces and call ‘‘Lebesgue mappings’’ the elements of nonlinear Lebesgue spaces.

Remark 2.14 (Extension to μ_M -measurable mappings). Nonlinear Lebesgue spaces can equivalently be defined using μ_M -measurable mappings, that is, mappings f satisfying for all B in $\mathcal{B}(N)$ that $f^{-1}(B)$ belongs to $\overline{\Sigma}_M$, the completion of the σ -algebra Σ_M under μ_M [ST25, Proposition 2.22].

The construction of nonlinear Lebesgue spaces developed above eventually yields that they are metric spaces [ST25, Proposition 2.25].

Proposition 2.15 (Nonlinear Lebesgue spaces are metric spaces). *Let $p \in [1, \infty]$. When equipped with D_p , $L_h^p(M, N)$ becomes a metric space with finite metric. Also, we will usually only write $L_h^p(M, N)$ when referring to the metric space $(L_h^p(M, N), D_p)$.*

We highlight the fact that we will frequently not distinguish representatives from their associated equivalence class, so that we can write $f \in L_h^p(M, N)$ while treating f as a \mathcal{L}^p mapping when it is clear from the context. Also, to ensure that our notation remains consistent with the literature on linear Lebesgue spaces, we denote $\mathcal{L}^p(M, N) := \mathcal{L}_h^p(M, N)$ and $L^p(M, N) := L_h^p(M, N)$ when N is a normed vector space and the base mapping is set to $h \equiv 0_N$ with 0_N the identity element of N .

An important matter to know about nonlinear Lebesgue spaces is whether they are reduced to the equivalence class of their base mapping, that is, $L_h^p(M, N) = \{h\}$ for $h \in L^0(M, N)$. In such cases, nonlinear Lebesgue spaces are called *trivial*. Nonlinear Lebesgue spaces are then called *nontrivial* when they are strictly bigger, in the sense of inclusion, than the equivalence class of the base mapping. This is the case if and only if μ_M is not purely infinite, that is, the range of μ_M is not reduced to $\{0, \infty\}$, and $|N| > 1$ [ST25, Proposition 3.3].

Proposition 2.16 (Nontrivial nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Then, the following assertions are equivalent:*

- (i) μ_M is not purely infinite and $|N| > 1$.
- (ii) $L_h^p(M, N)$ is nontrivial.

Also, nonlinear Lebesgue spaces inherit completeness from their target space (see [ST25, Proposition 4.1] or [Stu01, Proposition 3.3]).

Proposition 2.17 (Completeness of nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Suppose that N is complete. Then, $L_h^p(M, N)$ is complete.*

A subsidiary result of the proof of Proposition 2.17 is that the set of equivalence classes of measurable mappings also inherits completeness from the target space.

Corollary 2.18 (Completeness of the space of measurable mappings). *Let $p \in [1, \infty]$. Suppose that N is complete. Then, $(L^0(M, N), D_p)$ is complete.*

In addition, when μ_M is not purely infinite, the target space is isometric to a closed subset of nonlinear Lebesgue spaces [ST25, Corollary 3.8].

Proposition 2.19 (Closed isometric image of the target space into nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Suppose that μ_M is not purely infinite. Then, there exists an isometric (up to a scaling factor) mapping $\iota : N \rightarrow L_h^p(M, N)$ such that $\iota(N)$ is closed in $L_h^p(M, N)$ and there exists a measurable set A with $\mu_M(A) \in (0, \infty)$ for which $\iota(y)|_A \equiv y$ and $\iota(y)|_{M \setminus A} = h|_{M \setminus A}$ for all $y \in N$.*

Remark 2.20 (On a stronger conclusion when μ_M is finite and h is bounded). When it is additionally assumed that μ_M is finite and that h is bounded, we can actually choose $A = M$, so that $\iota(y) \equiv y$ for all $y \in N$ [ST25, Proposition 3.7].

Finally, a class of mappings that will play an important role in propagating properties to nonlinear Lebesgue spaces through density arguments is the set of *almost simple* mappings, that is, measurable mappings that are simple on a measurable set of finite μ_M -measure and (μ_M -a.e.) equal to the base mapping otherwise.

Definition 2.21 (Almost simple mappings). Let $h \in L^0(M, N)$. Then, define:

- (i) $\mathcal{E}_h(M, N)$ the set of *almost simple mappings*, that is, all mappings $f \in \mathcal{L}_s^0(M, N)$ such that there exists $B \in \mathcal{F}_{\mu_M}$, that is, a measurable set of finite μ_M -measure, for which f satisfies $f|_B \in \mathcal{E}(B, N)$ and $f|_{M \setminus B} \sim h|_{M \setminus B}$.
- (ii) $E_h(M, N) := \mathcal{E}_h(M, N) / \sim$ the set of equivalence classes of almost simple mappings.

Indeed, almost simple mappings are dense in nonlinear Lebesgue spaces for $p \geq 1$ [ST25, Proposition 3.10].

Proposition 2.22 (Density of almost simple mappings for $p \geq 1$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Then, $E_h(M, N) \cap L_h^p(M, N)$ is a dense subspace of $L_h^p(M, N)$.*

We now move to a reminder on curves in metric spaces.

2.2. Curves in metric spaces

2.2.1. Basic assumptions

Before delving into the study of curves in metric spaces, we clarify the assumptions on the set representing time.

Assumption 2.23 (Time interval). In the rest of the article, the nonempty set I , representing time, is such that $I = [a, b]$ with $(a, b) \in \mathbb{R}^2$ satisfying $a < b$.

We also clarify the assumptions on the generic metric space used in this section.

Assumption 2.24 (Generic metric space). In this section only, the generic nonempty set X is assumed to carry the structure of a metric space with finite metric $d_X : X^2 \rightarrow [0, \infty)$.

Throughout this article and this section, we shall not recall these assumptions in the statements of the results; yet, any additional assumption relative to Assumption 2.23 or Assumption 2.24 will be specified.

2.2.2. Sobolev curves and continuous representations

This section follows [Lis07, Section 2.4], but with slight relaxations on the assumptions of the results that are used in the present article. A first important class of curves in metric spaces is the set of *Sobolev curves*.

Definition 2.25 (Sobolev curves). Let $p \in (1, \infty)$. Then, the set of *Sobolev curves* is defined as

$$W^{1,p}(I, X) := \left\{ c \in L^p(I, X) : \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{X,\tau} c(t))^p dt < \infty \right\},$$

where $\Delta_{X,\tau} c(t) := d_X(c(t), c(t + \tau))/\tau$. Also, define the finite metric $d_p : L^p(I, X)^2 \rightarrow [0, \infty)$ as

$$d_p(c, c') := \left(\int_I d_X(c(t), c'(t))^p dt \right)^{1/p}.$$

Remark 2.26 (On the definition of $L^p(I, X)$). Note that unlike in [Lis07, Section 2.4], the definition of $L^p(I, X)$ used in Definition 2.25 is the one of Definition 2.13, so that we can always pick a separably valued representative instead of assuming that X is separable in most proofs. Also, the base mapping is a mapping constant equal to some $z_0 \in N$ and thus any $y \in N$, since I has finite \mathcal{L}^1 -measure.

Remark 2.27 ($W^{1,p}(I, X)$ is a Borel subset of $L^p(I, X)$). Let $p \in (1, \infty)$. As a direct consequence of Definition 2.25 and the fact that the functional $c \mapsto \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{X,\tau} c(t))^p dt$ is lower semi-continuous from $L^p(I, X)$ to $[0, \infty]$, $W^{1,p}(I, X)$ is a Borel subset of $L^p(I, X)$.

We will see that Sobolev curves are related to another important class of curves in metric spaces, namely the set of *absolutely continuous curves*, whose definition is recalled below and borrowed from [AGS05, Definition 1.1.1].

Definition 2.28 (Absolutely continuous curves). A curve $c : I \rightarrow X$ belongs to $\mathcal{AC}^p(I, X)$, for $p \in [1, \infty]$, if and only if there exists $m \in L^p(I)$ such that for all $(s, t) \in I^2$ such that $s \leq t$ we have

$$d_X(c(s), c(t)) \leq \int_s^t m(r) dr. \quad (1)$$

Absolutely continuous curves have the nice property of being almost everywhere differentiable in the metric sense [AGS05, Theorem 1.1.2].

Theorem 2.29 (Metric derivative). Let $p \in [1, \infty]$. Then, if $c \in \mathcal{AC}^p(I, X)$, the limit

$$|c'|_X(t) := \lim_{s \rightarrow t} \frac{d_X(c(s), c(t))}{|s - t|}$$

exists for \mathcal{L}^1 -a.e. $t \in I$. In addition, the mapping $t \rightarrow |c'|_X(t)$ belongs to $L^p(I)$, is an admissible integrand for the right-hand side of Equation (1) and is minimal in the sense that for all $m \in L^p(I)$ verifying Equation (1) it satisfies

$$|c'|_X(t) \leq m(t) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I.$$

Now, when $p \in (1, \infty)$ and X is complete, the set of absolutely continuous curves is isometric to the set of Sobolev curves for the metric d_p .

Proposition 2.30 (Absolutely continuous representations of Sobolev curves). Let $p \in (1, \infty)$. Suppose that X is complete. Then, the restriction of the quotient mapping

$$q : \begin{cases} \mathcal{L}_s^0(I, X) & \longrightarrow L^0(I, X) \\ c & \longmapsto [c] \end{cases}$$

to $\mathcal{AC}^p(I, X)$, denoted by q_p , is an isometry between $(\mathcal{AC}^p(I, X), d_p)$ and $(W^{1,p}(I, X), d_p)$.

Proof. Let $p \in [1, \infty)$.

Step 1: Definition of q_p . Note that q_p is well-defined since any curve in $\mathcal{AC}^p(I, X)$ has separable range as the continuous image of a separable set and since Borel measurability follows from continuity. Also, the fact that $q_p(\mathcal{AC}^p(I, X)) \subset W^{1,p}(I, X)$ follows from the first part of the proof of [Lis07, Lemma 1].

Step 2: Bijectivity of q_p . The bijectivity of q_p from $\mathcal{AC}^p(I, X)$ to $W^{1,p}(I, X)$ follows from the second part of the proof of [Lis07, Lemma 1] which shows that to each element $c \in W^{1,p}(I, X)$ we can associate a unique element $\tilde{c} \in \mathcal{AC}^p(I, X)$ such that $q_p(\tilde{c}) = c$. The only variation in our case is that, instead of assuming that X is separable, the proof can be carried out the same way by using the fact that elements of $W^{1,p}(I, X)$ have separably valued representatives (Remark 2.26).

Step 3: Isometric property of q_p . The isometric property of q_p for d_p follows from the invariance of d_p to variations on \mathcal{L}^1 -null sets. \square

Proposition 2.30 and Definition 2.25 yield an equivalent definition of the set of absolutely continuous curves, when $p \in (1, \infty)$ and X is complete.

Corollary 2.31 (Equivalent definition of absolutely continuous curves for $p > 1$). *Let $p \in (1, \infty)$. Suppose that X is complete. Then, the set of absolutely continuous is equivalently defined as*

$$\mathcal{AC}^p(I, X) = \left\{ c \in C(I, X) : \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{X,\tau} c(t))^p dt < \infty \right\}.$$

Remark 2.32 ($\mathcal{AC}^p(I, X)$ is a Borel subset of $C(I, X)$). Let $p \in (1, \infty)$. As a direct consequence of Corollary 2.31 and the fact that the functional $c \mapsto \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{X,\tau} c(t))^p dt$ is lower semi-continuous from $C(I, X)$ to $[0, \infty]$, $\mathcal{AC}^p(I, X)$ is a Borel subset of $C(I, X)$.

Finally, note that the evaluation mapping is measurable on the set of absolutely continuous curves equipped with the metric d_p as soon as $p \in (1, \infty)$ and X is complete and separable.

Proposition 2.33 (Measurability of the evaluation mapping for d_p). *Let $p \in (1, \infty)$. Suppose that X is complete and separable. Then, for all $t \in I$, the restriction of the evaluation mapping*

$$e_t : \begin{cases} C(I, X) & \longrightarrow X \\ c & \longmapsto c(t) \end{cases}$$

to $\mathcal{AC}^p(I, X)$, denoted by $e_{t,p}$, is Borel measurable from $(\mathcal{AC}^p(I, X), d_p)$ to X .

Proof. Let $p \in [1, \infty]$.

Step 1: Borel measurability of $e_{t,p}$ for d_∞ . Since e_t is continuous, its restriction to $\mathcal{AC}^p(I, X)$, denoted by $e_{t,p}$, is Borel measurable from $(\mathcal{AC}^p(I, X), d_\infty)$ to X for all $t \in I$.

Step 2: Coincidence of the Borel sets of $(\mathcal{AC}^p(I, X), d_p)$ and $(\mathcal{AC}^p(I, X), d_\infty)$. Recall that $W^{1,p}(I, X)$ and $\mathcal{AC}^p(I, X)$ are Borel sets of $L^p(I, X)$ and $C(I, X)$, respectively (see Remarks 2.27 and 2.32). As such, they are Lusin sets, hence Suslin sets as soon as X is Polish (i.e., complete and separable), since, under such assumptions on X , $L^p(I, X)$ and $C(I, X)$ become Polish (see [Sch73, Theorem 2, p. 95] for the proof that a Borel subset of a Lusin space is Lusin). Furthermore, $(\mathcal{AC}^p(I, X), d_p)$ being isometric to $(W^{1,p}(I, X), d_p)$ by q_p (Proposition 2.30), it is also Suslin. Hence, the conclusion follows from the fact that d_p induces on $\mathcal{AC}^p(I, X)$ a Suslin topology that is weaker than the one induced by d_∞ [Sch73, Corollary 2, p. 101]. \square

We now continue with another important class of curves in metric spaces beyond continuous curves, namely the curves of bounded variation.

2.2.3. Curves of bounded variation and càdlàg representations

This section follows [ALS24, Section 2.2], but with slight relaxations on the assumptions of the results that are used in the present article. First, let us recall the definition of the *variation* of curves.

Definition 2.34 (Variation of curves). The *variation* of a curve $c : I \rightarrow X$ on any subset $J \subset I$ is defined as

$$\text{Var}_X(c; J) := \sup \left\{ \sum_{i=0}^n d_X(c(t_i), c(t_{i+1})) : \{t_i\}_{0 \leq i \leq n+1} \subset J \text{ and } t_0 < \dots < t_{n+1} \right\}$$

and its *essential variation* is defined as

$$\text{ess Var}_X(c; J) := \inf \{ \text{Var}_X(\tilde{c}) : \tilde{c} = c \text{ } \mathcal{L}^1\text{-a.e.} \}.$$

Using [Definition 2.34](#), the set of *curves of bounded variation* can be defined.

Definition 2.35 (Curves of bounded variation). The set of *curves of bounded variation*, which we call BV-curves for short, is defined as

$$BV(I, X) := \{ c \in L^1(I, X) : \text{ess Var}(c) < \infty \}.$$

Also, define the finite metric $d_1 : L^1(I, X)^2 \rightarrow [0, \infty)$ as

$$d_1(c, c') := \int_I d_X(c(t), c'(t)) dt.$$

Remark 2.36 (On the definition of $L^1(I, X)$). Note that unlike in [\[ALS24, Section 2.2\]](#), the definition of $L^1(I, X)$ used in [Definition 2.35](#) is the one of [Definition 2.13](#), so that we can always pick a separably valued representative instead of assuming that X is separable in most proofs. Also, the base mapping is a mapping constant equal to some $z_0 \in N$ and thus any $y \in N$, since I has finite \mathcal{L}^1 -measure.

Remark 2.37 ($BV(I, X)$ is a Borel subset of $L^1(I, X)$). As a direct consequence of [Definition 2.35](#) and the fact that the essential variation is lower semi-continuous from $L^1(I, X)$ to $[0, \infty]$ [\[ALS24, Lemma 2.7\]](#), $BV(I, X)$ is a Borel subset of $L^1(I, X)$.

Using the definition of the essential variation of a curve, one can define the *variation measure* associated to a BV-curve.

Definition 2.38 (Variation measure). The *variation measure* of a BV-curve $c \in BV(I, X)$ is defined as the Lebesgue–Stieltjes measure $|Dc|_X$ on I induced by the non-decreasing function $V : I \rightarrow \mathbb{R}$ defined as $V(t) := \text{ess Var}_X(c; (0, t))$ (see [\[SS09, Chapter 6, Section 3.3\]](#) for the construction of $|Dc|_X$), that is, $|Dc|_X$ is the unique measure on I satisfying for all $(s, t) \in I^2$ with $s < t$ that

$$|Dc|_X((s, t)) = \lim_{u \rightarrow t^-} V(u) - \lim_{u \rightarrow s^+} V(u).$$

Now, when X is complete, we have an equivalent definition of the set of BV-curves.

Proposition 2.39 (Equivalent definition of $BV(I, X)$). *Suppose that X is complete. Then, the set of curves of bounded variation is equivalently defined as*

$$BV(I, X) = \left\{ c \in L^1(I, X) : \sup_{0 < \tau < b-a} \int_a^{b-\tau} \Delta_{X,\tau} c(t) dt < \infty \right\}$$

and for all $c \in BV(I, X)$ we have for all $(s, t) \in I^2$ with $s < t$ that

$$|Dc|_X((s, t)) = \text{ess Var}_X(c; (s, t)) = \sup_{0 < \tau < t-s} \int_s^{t-\tau} \Delta_{X,\tau} c(t) dt.$$

Proof. The only variation with respect to the proof of [\[ALS24, Theorem 2.17\]](#) is the fact that, instead of assuming that X is separable, we can use the fact that elements of $BV(I, X)$ have separably valued representatives ([Remark 2.36](#)), so that, rather than taking a dense set in X in the proof that (4) \implies (1), it is sufficient to take a dense set in the range of a separably valued representative. \square

We will see that BV-curves are related to another important class of curves in metric spaces, namely the set of *càdlàg curves of bounded variation*. Let us first recall the definition of *càdlàg curves*.

Definition 2.40 (Càdlàg curves). A curve $c : I \rightarrow X$ belongs to $\mathcal{D}(I, X)$ if it is right continuous with left limits, that is, for all $t \in I$ it satisfies $c(t) = \lim_{s \rightarrow t^+} c(s)$ and $\lim_{s \rightarrow t^-} c(s)$ exists in X . This set becomes a metric space when equipped with the Skorokhod metric $d_{Sk} : \mathcal{D}(I, X)^2 \rightarrow [0, \infty)$ defined as

$$d_{Sk}(c, c') := \inf_{\lambda} \max\{\|\lambda\|_B, d_{\infty}(c, c' \circ \lambda)\},$$

where the infimum is taken over all increasing homeomorphisms $\lambda : I \rightarrow I$ and

$$\|\lambda\|_B := \sup_{(s,t) \in I^2, s < t} \left| \log \left(\frac{\lambda(t) - \lambda(s)}{t - s} \right) \right|.$$

Then, one can define the set of *càdlàg curves of bounded variation*.

Definition 2.41 (Càdlàg curves of bounded variation). A curve $c : I \rightarrow X$ belongs to $\mathcal{BV}(I, X)$ if it is càdlàg, continuous on the left at $t = b$ and has finite variation, that is, $c \in \mathcal{D}(I, X)$, $c(b) = \lim_{t \rightarrow b^-} c(t)$ and $\text{Var}_X(c) < \infty$.

Remark 2.42 (On the condition of left-continuity at $t = b$). Note that unlike [ALS24, p. 16] we define $\mathcal{BV}(I, X)$ by adding the condition of left-continuity at $t = b$. Indeed, for notational consistency, we argue that it makes more sense for $\mathcal{BV}(I, X)$ to denote the set that is in bijection with $BV(I, X)$ via the quotient mapping on $\mathcal{L}_s^0(I, X)$ (Proposition 2.44), rather than a strictly larger set. In addition, using the definition of $\mathcal{BV}(I, X)$ from Definition 2.41 instead of the one from [ALS24, p. 16] does not affect the exposition of this section since the definition of $\mathcal{BV}(I, X)$ from Definition 2.41 is closed under limits for the Skorokhod metric, so that the results on the set of all càdlàg curves of finite variation mentioned in [ALS24] all hold for this definition.

Remark 2.43 ($\mathcal{BV}(I, X)$ is a Borel subset of $\mathcal{D}(I, X)$). As a direct consequence of Definition 2.41 and the fact that the variation is lower semi-continuous from $\mathcal{D}(I, X)$ to $[0, \infty]$ [ALS24, Lemma 2.13], $\mathcal{BV}(I, X)$ is a Borel subset of $\mathcal{D}(I, X)$.

When X is complete, the set of BV-curves is isometric to the set of càdlàg curves of bounded variation for the metric d_1 [ALS24, Lemma 2.5].

Proposition 2.44 (Càdlàg representations of BV-curves). *Suppose that X is complete. Then, the following assertions hold:*

(i) *for all $c \in \mathcal{D}(I, X)$ and all $(s, t) \in I^2$ we have*

$$|Dc|_X((s, t)) = \text{ess Var}_X(c; (s, t)) = \text{Var}_X(c; (s, t)).$$

(ii) *the restriction of the quotient mapping*

$$q : \begin{cases} \mathcal{L}_s^0(I, X) & \longrightarrow L^0(I, X) \\ c & \longmapsto [c] \end{cases}$$

to $\mathcal{BV}(I, X)$, denoted by q_1 , is an isometry between $(\mathcal{BV}(I, X), d_1)$ and $(BV(I, X), d_1)$.

Finally, note that the evaluation mapping is measurable on the set of càdlàg curves of bounded variation equipped with the metric d_1 as soon as X is complete and separable.

Proposition 2.45 (Measurability of the evaluation mapping for d_1). *Suppose that X is complete and separable. For all $t \in I$, the restriction of the evaluation mapping*

$$e_t : \begin{cases} \mathcal{D}(I, X) & \longrightarrow X \\ c & \longmapsto c(t) \end{cases}$$

to $\mathcal{BV}(I, X)$, denoted by $e_{t,1}$, is Borel measurable from $(\mathcal{BV}(I, X), d_1)$ to X .

Proof. Step 1: Borel measurability of $e_{t,1}$ for d_{S_k} . The mapping e_t being Borel measurable from $(\mathcal{D}(I, X), d_{S_k})$ to X for all $t \in I$ [ALS24, Proposition 2.15], its restriction to $\mathcal{BV}(I, X)$, denoted by $e_{t,1}$, is Borel measurable from $(\mathcal{BV}(I, X), d_{S_k})$ to X for all $t \in I$.

Step 2: Coincidence of the Borel sets of $(\mathcal{BV}(I, X), d_1)$ and $(\mathcal{BV}(I, X), d_{S_k})$. Recall that $BV(I, X)$ and $\mathcal{BV}(I, X)$ are Borel sets of $L^1(I, X)$ and $\mathcal{D}(I, X)$, respectively (see Remarks 2.37 and 2.43). As such, they are Lusin sets, hence Suslin sets as soon as X is Polish (i.e., complete and separable), since, under such assumptions on X , $L^1(I, X)$ and $\mathcal{D}(I, X)$ become Polish (see [Sch73, Theorem 2, p. 95] for the proof that a Borel subset of a Lusin space is Lusin and [ALS24, Theorem 2.10] for the proof that $\mathcal{D}(I, X)$ is Polish when X is). Furthermore, $(\mathcal{BV}(I, X), d_1)$ being isometric to $(BV(I, X), d_1)$ by q_1 (Proposition 2.44), it is also Suslin. Hence, the conclusion follows from the fact that d_1 induces on $\mathcal{BV}(I, X)$ a Suslin topology that is weaker than the one induced by d_∞ [Sch73, Corollary 2, p. 101]. \square

To characterize the behaviours of all the classes of curves presented in Section 2.2 for the particular case of nonlinear Lebesgue spaces, we will need to characterize L^p curves in nonlinear Lebesgue spaces since Sobolev and BV curves are defined using L^p curves (Definitions 2.25 and 2.35). The next section thus present such a characterization of L^p curves and its consequence for the characterization of absolutely continuous curves and càdlàg curves of bounded variation in nonlinear Lebesgue spaces.

3. Main results

3.1. Characterization of L^p curves in nonlinear Lebesgue spaces

We first clarify some recurrent notation used in this section and thereafter.

Notation 3.1. Let $h \in L^0(M, N)$, $p \in [1, \infty)$ and $\mathcal{L}^1 \otimes \mu_M$ be the product measure obtained using Carathéodory's process [DiB02, Chapter II]. From thereon:

- (i) \bar{h} denotes the mapping from I to $L_h^p(M, N)$ defined as $\bar{h} \equiv h$.
- (ii) \underline{h} denotes the mapping from M to $L^p(I, N)$ defined as $\underline{h} := \iota \circ h$, which is measurable from M to $L^p(I, N)$ as the composition of the measurable mapping h and the isometric mapping ι defined in Remark 2.20. The mapping \underline{h} is also measurable from M to $(E(I, N), d_p)$ since the range of ι is closed in $L^p(I, N)$ and included in $E(I, N)$.
- (iii) \hat{h} denotes the mapping from $I \times M$ to N defined as $\hat{h}(t, x) = h(x)$ for $\mathcal{L}^1 \otimes \mu_M$ -a.e. $(t, x) \in I \times M$.
- (iv) $L^p(I, L_h^p(M, N))$ denotes the nonlinear Lebesgue space with base mapping set as \bar{h} .
- (v) $d_{p,p}$ denotes the finite metric on $L^p(I, L_h^p(M, N))$ defined as

$$d_{p,p}(c, c') := \left(\int_I D_p(c(t), c'(t))^p dt \right)^{1/p}.$$

- (vi) $D_{p,p}$ denotes the finite metric on $L_{\underline{h}}^p(M, L^p(I, N))$ defined as

$$D_{p,p}(f, f') := \left(\int_M d_p(f(x), f'(x))^p d\mu_M(x) \right)^{1/p}.$$

- (vii) \hat{D}_p denotes the finite metric on $L_{\hat{h}}^p(I \times M, N)$ defined as

$$\hat{D}_p(\hat{c}, \hat{c}') := \left(\int_{I \times M} d_N(\hat{c}(t, x), \hat{c}'(t, x))^p d\mathcal{L}^1 \otimes \mu_M(t, x) \right)^{1/p}.$$

Establishing a characterization of L^p curves in nonlinear Lebesgue spaces essentially consists in the identification of a canonical isometry of spaces of simple mappings that extends to nonlinear Lebesgue spaces through a density argument. To identify such a canonical isometry, a core ingredient is the set of *rectangular almost simple mappings* defined below.

Definition 3.2 (Rectangular almost simple mappings). Let $h \in L^0(M, N)$. Then, define:

(i) $\mathcal{E}_{\hat{h},\sigma}(I \times M, N)$ the set of *rectangular almost simple mappings*, that is, all mappings $\hat{c} \in \mathcal{L}^0(I \times M, N)$ such that there exist a finite collection $(y_j)_{j \in J}$ of elements of N and a finite collection $(I_j \times M_j)_{j \in J}$ of disjoint measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure such that $\hat{c}|_{I_j \times M_j} \equiv y_j$ for all $j \in J$ and $\hat{c}|_{(I \times M) \setminus R} \sim \hat{h}|_{(I \times M) \setminus R}$ with $R := \cup_{j \in J} I_j \times M_j$.

(ii) $E_{\hat{h},\sigma}(I \times M, N) := \mathcal{E}_{\hat{h},\sigma}(I \times M, N) \setminus \sim$ the set of equivalence classes of rectangular almost simple mappings. Note that, by definition, $E_{\hat{h},\sigma}(I \times M, N) \subset E_{\hat{h}}(I \times M, N)$.

On the set of rectangular almost simple mappings, the section mappings on I and M define isometries.

Proposition 3.3 (Isometries induced by the section mappings on I and M). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Then, the following assertions hold:*

(i) *the section mapping on I*

$$\text{sec}_{I,p} : \begin{cases} E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N) & \longrightarrow E(I, E_h(M, N) \cap L_h^p(M, N)) \\ [\hat{c}] & \longmapsto [t \mapsto [\hat{c}(t, \cdot)]] \end{cases}$$

is an isometry between those sets equipped with the metrics \hat{D}_p and $d_{p,p}$, respectively.

(ii) *the section mapping on M*

$$\text{sec}_{M,p} : \begin{cases} E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N) & \longrightarrow E_{\hat{h}}(M, E(I, N)) \cap L_{\hat{h}}^p(M, L^p(I, N)) \\ [\hat{c}] & \longmapsto [x \mapsto [\hat{c}(\cdot, x)]] \end{cases}$$

is an isometry between those sets equipped with the metrics \hat{D}_p and $D_{p,p}$, respectively.

Remark 3.4 (On the choice of product measure $\mathcal{L}^1 \otimes \mu_M$). The choice of product measure $\mathcal{L}^1 \otimes \mu_M$ does not matter, since we can always reduce to integration over measurable rectangles on which all product measures coincide.

Proof. Let $h \in L^0(M, N)$ and $p \in [1, \infty)$.

(i) *Step 1: Definition of $\text{sec}_{I,p}$.* Let $\hat{c} \in E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$. Then, there is $\tilde{c} \in \mathcal{E}_{\hat{h},\sigma}(I \times M, N) \cap \mathcal{L}_{\hat{h}}^p(I \times M, N)$ such that $\hat{D}_p(\tilde{c}, \hat{c}) = 0$. By definition of \tilde{c} , there exists $n \in \mathbb{N}$, a collection $(y_i)_{0 \leq i \leq n}$ of elements of N and a collection $(I_i \times M_i)_{0 \leq i \leq n}$ of disjoint measurable rectangles of finite μ_M -measure such that $\tilde{c}|_{I_i \times M_i} \equiv y_i$ for all $0 \leq i \leq n$ and $\tilde{c}|_{(I \times M) \setminus R} = h|_{(I \times M) \setminus R}$ with $R := \cup_{i=0}^n (I_i \times M_i)$. Then, $t \mapsto [\tilde{c}(t, \cdot)]$ is simple from I to $(E_h(M, N), D_p)$ and, since $\tilde{c} \in \mathcal{L}_{\hat{h}}^p(I \times M, N)$ and R is the finite union of measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure, we have, by the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1], that $D_p(\tilde{c}(t, \cdot), h) < \infty$ for \mathcal{L}^1 -a.e. $t \in I$, hence $[\tilde{c}(t, \cdot)] \in E_h(M, N) \cap L_h^p(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$. This shows that $\text{sec}_{I,p}(\hat{c}) = [t \mapsto [\tilde{c}(t, \cdot)]]$ belongs to $E(I, E_h(M, N) \cap L_h^p(M, N))$.

Step 2: Isometric property of $\text{sec}_{I,p}$. Let $(\hat{c}, \hat{c}') \in (E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N))^2$. Since \hat{c} and \hat{c}' differ on the finite union of measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure, we have, by the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1], that

$$d_{p,p}(\text{sec}_{I,p}(\hat{c}), \text{sec}_{I,p}(\hat{c}')) = \hat{D}_p(\hat{c}, \hat{c}').$$

Step 3: Surjectivity of $\text{sec}_{I,p}$. Let $c \in E(I, E_h(M, N) \cap L_h^p(M, N))$. Then, there is $\tilde{c} \in \mathcal{E}(I, E_h(M, N) \cap \mathcal{L}_h^p(M, N))$ such that $d_{p,p}(\tilde{c}, c) = 0$. By definition of \tilde{c} , there exists a finite indexing set I_c such that $\tilde{c}(I) = \{\tilde{s}_i \in E_h(M, N) \cap \mathcal{L}_h^p(M, N) : i \in I_c\}$. Also, define $I_i := \tilde{c}^{-1}(\{\tilde{s}_i\})$ for all $i \in I_c$. In addition, there exist $B_i \in \mathcal{F}_{\mu_M}$ on which \tilde{s}_i differs from h and a finite indexing set J_i such that $\tilde{s}_i(B_i) := \{y_{i,j} \in N : j \in J_i\}$ and $\tilde{s}_i|_{M \setminus B_i} = h|_{M \setminus B_i}$. Then, define $B_{i,j} := \tilde{s}_i^{-1}(\{y_{i,j}\}) \subset B_i$ for all $j \in J_i$. Thus, assuming without loss of generality that the J_i 's are disjoint and defining the finite set $J := \cup_{i \in I_c} J_i$, we observe that $(I_i \times B_{i,j})_{(i,j) \in I_c \times J}$ is a finite collection of disjoint measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure and we can define the mapping $\hat{c} : I \times M \rightarrow N$ as $\hat{c}|_{I_i \times B_{i,j}} \equiv y_{i,j}$ for all $(i, j) \in I_c \times J$ and $\hat{c}(t, x) = h(x)$ for all $(t, x) \in (I \times M) \setminus P$ with $P := \cup_{(i,j) \in I_c \times J} I_i \times B_{i,j}$. By construction, $D_p(\hat{c}(t, \cdot), c(t)) = 0$ for \mathcal{L}^1 -a.e. $t \in I$ and \hat{c} differs from \hat{h} on the finite union of measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure, hence, by the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1], $\hat{D}_p(\hat{c}, \hat{h}) = d_{p,p}(c, \bar{h}) < \infty$. Thus, up to the identification with its equivalence class, \hat{c} belongs to $E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$ and $c = \text{sec}_{I,p}(\hat{c})$, that is, c belongs to the range of $\text{sec}_{I,p}$.

(ii) The proof follows from the same arguments as the proof of (i), but using almost simple mappings. \square

Remark 3.5 (On relaxing the assumptions on I). Note that the proof of [Proposition 3.3](#) does not rely on any specific property of I other than the finiteness of the measure on it to ensure that $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^p(I, N))$ is nonempty. Hence, [Proposition 3.3](#) holds for a measure space (I, Σ_I, μ_I) such that μ_I is finite. Alternatively, to ensure that $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^p(I, N))$ is nonempty, we can replace the finiteness assumption on μ_I by the assumption that $h \in E(M, N)$. In that case, $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^p(I, N))$ should be replaced by $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^0(I, N))$ in assertion (ii) of [Proposition 3.3](#), with $L_{\underline{h}}^p(M, L^0(I, N))$ denoting the set of equivalence classes of separably valued measurable mappings from M to $(L^0(M, N), D_p)$ at a finite $D_{p,p}$ -distance from \underline{h} . If h is not only simple, but constant, the assertion (ii) of [Proposition 3.3](#) holds as is. The assertion (i) of [Proposition 3.3](#) even holds when (I, Σ_I, μ_I) is just a measure space, with no further assumptions, by reasoning with almost simple mappings instead of simple mappings.

As a direct consequence of [Proposition 3.3](#), we can define an isometry between the target spaces of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$.

Corollary 3.6 (Isometry between the target spaces of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Then, the isometries $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$ defined in [Proposition 3.3](#) induce the isometry*

$$i_p := \text{sec}_{M,p} \circ \text{sec}_{I,p}^{-1} : E(I, E_h(M, N) \cap L_h^p(M, N)) \rightarrow E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^p(I, N)).$$

Let us now identify the closures of the target and base spaces of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$.

Proposition 3.7 (Closures of the base and target spaces of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Then, the following assertions hold:*

(i) *the closure of the base space of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$ is*

$$\overline{E_{\hat{h}, \sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)}^{\hat{D}_p} = L_{\hat{h}}^p(I \times M, N).$$

(ii) *the closure of the target space of $\text{sec}_{I,p}$ is*

$$\overline{E(I, E_h(M, N) \cap L_h^p(M, N))}^{d_{p,p}} = L^p(I, L_h^p(M, N)).$$

(iii) *the closure of the target space of $\text{sec}_{M,p}$ is*

$$\overline{E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^p(I, N))}^{D_{p,p}} = L_{\underline{h}}^p(M, L^p(I, N)).$$

Proof. Let $h \in \mathcal{L}_s^0(M, N)$, $p \in [1, \infty)$ and $\varepsilon > 0$.

(i) Let $\hat{c} \in \mathcal{L}_{\hat{h}}^p(I \times M, N)$. Then, \hat{c} differs from \hat{h} on a measurable set \hat{P} of σ -finite $\mathcal{L}^1 \otimes \mu_M$ -measure, that is, there exists a sequence $(P_n)_{n \in \mathbb{N}}$ of measurable sets, which can be assumed increasing, such that $\hat{P} = \bigcup_{n \in \mathbb{N}} P_n$.

Step 1: Approximation by a Lebesgue mapping such that the set where it differs from \hat{h} is contained in a measurable set where \hat{h} is bounded. Let $z_0 \in N$ and $H_n := \{x \in M : d_N(z_0, h(x)) \leq n\}$ for all $n \in \mathbb{N}$, so that, d_N being assumed finite, $M = \bigcup_{n \in \mathbb{N}} H_n$. Also, define the mapping $\hat{c}_n : I \times M \rightarrow N$ as

$$\hat{c}_n(t, x) := \begin{cases} \hat{c}(t, x), & \text{if } x \in H_n \\ \hat{h}(t, x), & \text{otherwise} \end{cases}$$

Then, since $(H_n)_{n \in \mathbb{N}}$ is an increasing sequence of sets in M , we have for all $(t, x) \in I \times M$ that $\hat{c}_n(t, x) \rightarrow \hat{c}(t, x)$ as $n \rightarrow \infty$ and $d_N(\hat{c}_n(t, x), \hat{c}(t, x)) \leq d_N(\hat{h}(t, x), \hat{c}(t, x))$ for all $n \in \mathbb{N}$. Since $\hat{c} \in \mathcal{L}_{\hat{h}}^p(I \times M, N)$, we have, by Lebesgue's dominated convergence theorem [[Coh13](#), Theorem 2.4.5], that $\hat{D}_p(\hat{c}_n, \hat{c}) \rightarrow 0$ as $n \rightarrow \infty$. Hence, there exists $n_0 \in \mathbb{N}$ such that $\hat{D}_p(\hat{c}_{n_0}, \hat{c}) < \varepsilon/4$. In addition, \hat{c}_{n_0} differs from \hat{h} on a measurable set contained in the measurable rectangle $I \times H_{n_0}$.

Step 2: Approximation by a Lebesgue mapping such that the set where it differs from \hat{h} is contained in a measurable set of finite $\mathcal{L}^1 \otimes \mu_M$ -measure. For all $n \in \mathbb{N}$, define the mapping $\hat{c}'_n : I \times M \rightarrow N$ as

$$\hat{c}'_n(t, x) := \begin{cases} \hat{c}_{n_0}(t, x), & \text{if } (t, x) \in P_n \\ \hat{h}(t, x), & \text{otherwise} \end{cases}$$

Then, since $(P_n)_{n \in \mathbb{N}}$ is an increasing sequence of sets in \hat{P} , we have for all $(t, x) \in I \times M$ that $\hat{c}'_n(t, x) \rightarrow \hat{c}_{n_0}(t, x)$ as $n \rightarrow \infty$ and $d_N(\hat{c}'_n(t, x), \hat{c}_{n_0}(t, x)) \leq d_N(\hat{h}(t, x), \hat{c}_{n_0}(t, x))$ for all $n \in \mathbb{N}$. Since $\hat{c}_{n_0} \in \mathcal{L}^p_{\hat{h}}(I \times M, N)$, we have, by Lebesgue's dominated convergence theorem [Coh13, Theorem 2.4.5], that $\hat{D}_p(\hat{c}'_n, \hat{c}_{n_0}) \rightarrow 0$ as $n \rightarrow \infty$. Hence, there exists $n_1 \in \mathbb{N}$ such that $\hat{D}_p(\hat{c}'_{n_1}, \hat{c}_{n_0}) < \varepsilon/4$. In addition, \hat{c}'_{n_1} differs from \hat{h} on a measurable set contained in $P'_{n_1} := P_{n_1} \cap (I \times H_{n_0})$, which has finite $\mathcal{L}^1 \otimes \mu_M$ -measure.

Step 3: Approximation by an almost simple mapping such that the set where it differs from \hat{h} is contained in a measurable set of finite $\mathcal{L}^1 \otimes \mu_M$ -measure. Using Proposition 2.22, we can construct $\hat{s} \in \mathcal{E}_{\hat{h}}(I \times M, N)$ such that it differs from \hat{h} on a measurable set \hat{S} contained in P'_{n_1} and $\hat{D}_p(\hat{c}'_{n_1}, \hat{s}) < \varepsilon/4$. Hence, we can assume that there exists a finite indexing set J such that $\hat{s}(\hat{S}) = \{y_j \in N : j \in J\}$ and let $\hat{S}_j := \hat{s}|_{\hat{S}_j}^{-1}(\{y_j\})$ for all $j \in J$. Also, define

$$C := \max \left\{ \max_{i \neq j} d_N(y_i, y_j), \max_{j \in J} \{d_N(y_j, z_0) + n_0\} \right\}.$$

Step 4: Approximation by a rectangular almost simple mapping. For all $j \in J$, \hat{S}_j has finite $\mathcal{L}^1 \otimes \mu_M$ -measure, hence, by construction of $\mathcal{L}^1 \otimes \mu_M$, there exists a countable collection of measurable rectangles $(\hat{I}_{j,n} \times \hat{M}_{j,n})_{n \in \mathbb{N}}$ such that, defining $\hat{R}_j := \cup_{n \in \mathbb{N}} (\hat{I}_{j,n} \times \hat{M}_{j,n})$, it satisfies $\hat{S}_j \subset \hat{R}_j$ and $\mathcal{L}^1 \otimes \mu_M(\hat{R}_j \setminus \hat{S}_j) < \varepsilon^p / (2|J|C^p 4^p)$ [DiB02, Chapter II, Proposition 10.2]. In addition, since $\hat{S} \subset I \times H_{n_0}$, it can be assumed that $\hat{M}_{j,n} \subset H_{n_0}$ for all $(j, n) \in J \times \mathbb{N}$ since the intersection of measurable rectangles remains a measurable rectangle. Then, define $\hat{R}_{j,n} := \cup_{k=0}^n (\hat{I}_{j,k} \times \hat{M}_{j,k})$, so that $(\cup_{j \in J} \hat{R}_{j,n})_{n \in \mathbb{N}}$ is an increasing sequence of sets in $\cup_{j \in J} \hat{R}_j$. Hence, J being finite, there exists $n_2 \in \mathbb{N}$ such that $\sum_{j \in J} \mathcal{L}^1 \otimes \mu_M(\hat{R}_j \setminus \hat{R}_{j,n_2}) < \varepsilon^p / (2C^p 4^p)$. Since $(\hat{R}_{j,n_2})_{j \in J}$ is a finite collection of finite unions of measurable rectangles and since the set of all measurable rectangles is a semi-algebra [DiB02, Chapter III, Section 12], the \hat{R}_{j,n_2} 's can be assumed disjoint, up to replacing them with the collection $(\hat{R}'_{j,n_2})_{j \in J}$ such that, assuming that $J = \{0, 1, \dots, m\}$ for some $m \in \mathbb{N}$, $\hat{R}'_{0,n_2} := \hat{R}_{0,n_2}$ and $\hat{R}'_{j,n_2} := \hat{R}_{j,n_2} \setminus (\cup_{k=0}^{j-1} \hat{R}_{k,n_2})$, which can be decomposed in the finite union of measurable rectangles, for all $0 < j \leq m$. Now, define the mapping $\hat{s}_{n_2} : I \times M \rightarrow N$ as $\hat{s}_{n_2}|_{\hat{R}'_{j,n_2}} \equiv y_j$ for all $j \in J$ and $\hat{s}_{n_2}|_{(I \times M) \setminus \hat{R}} = \hat{h}|_{(I \times M) \setminus \hat{R}}$ with $\hat{R} := \cup_{j \in J} \hat{R}_{j,n_2}$. Since \hat{R} is the finite union of measurable rectangles, it can be decomposed into the finite union of disjoint measurable rectangles. Hence, by construction, \hat{s}_{n_2} belongs to $\mathcal{E}_{\hat{h}, \sigma}(I \times M, N)$ and satisfies

$$\begin{aligned} \hat{D}_p(\hat{s}, \hat{s}_{n_2})^p &\leq C^p \sum_{j \in J} \mathcal{L}^1 \otimes \mu_M(\hat{S}_j \Delta \hat{R}'_{j,n_2}) \\ &\leq C^p \left(\sum_{j \in J} \mathcal{L}^1 \otimes \mu_M(\hat{R}_j \setminus \hat{R}'_{j,n_2}) + \sum_{j \in J} \mathcal{L}^1 \otimes \mu_M(\hat{R}_j \setminus \hat{S}_j) \right) < (\varepsilon/4)^p. \end{aligned}$$

Finally, we have, by the triangle inequality, that

$$\hat{D}_p(\hat{c}, \hat{s}') \leq \hat{D}_p(\hat{c}, \hat{c}_{n_0}) + \hat{D}_p(\hat{c}_{n_0}, \hat{c}'_{n_1}) + \hat{D}_p(\hat{c}'_{n_1}, \hat{s}) + \hat{D}_p(\hat{s}, \hat{s}_{n_2}) < \varepsilon.$$

(ii) Let $c \in L^p(I, L^p_h(M, N))$.

Step 1: Approximation by a simple mapping. By Proposition 2.22, there exists $s \in E(I, L^p_h(M, N))$ such that $d_{p,p}(c, s) < \varepsilon/2$. Up to its identification with an appropriate representative, we can assume that s is a simple mapping, hence there exists a finite indexing set I_s such that $s(I) := \{s_i \in L^p_h(M, N) : i \in I_s\}$ and define $I_i := g^{-1}(\{s_i\})$.

Step 2: Approximation of the values of the simple mapping by almost simple mappings. For each $i \in I_s$, still using Proposition 2.22, there exists $\tilde{s}_i \in E_h(M, N) \cap L^p_h(M, N)$ such that $D_p(s_i, \tilde{s}_i) < \varepsilon / (2^p |I_s| \mathcal{L}^1(I_i))^{1/p}$. Then, define the mapping $\tilde{s} : I \rightarrow E_h(M, N) \cap L^p_h(M, N)$ as $\tilde{s}|_{I_i} \equiv s_i$ for all $i \in I_s$, so that we get

$$d_{p,p}(s, \tilde{s})^p = \sum_{i \in I_s} D_p(s_i, \tilde{s}_i)^p \mathcal{L}^1(I_i) < (\varepsilon/2)^p.$$

Finally, we have, by the triangle inequality, that

$$d_{p,p}(c, \tilde{s}) \leq d_{p,p}(c, s) + d_{p,p}(s, \tilde{s}) < \varepsilon.$$

(iii) The proof follows from the same arguments as the proof of (ii), but using almost simple mappings. \square

Remark 3.8 (On relaxing the assumptions on I). Note that, similarly to what was highlighted in [Remark 3.5](#) for [Proposition 3.3](#), the proof of [Proposition 3.7](#) does not rely on any specific property of I other than the finiteness of the measure on it to ensure that $E_{\hat{h}}(M, E(I, N)) \cap L_{\hat{h}}^p(M, L^p(I, N))$ is nonempty. Hence, the result holds for a measure space (I, Σ_I, μ_I) such that μ_I is finite. The assumption of finiteness of μ_I can be replaced by the assumption that $h \in E(M, N)$. In that case, the assertion (iii) of [Proposition 3.7](#) should be replaced by [Proposition A.1](#). If h is not only simple, but constant, the assertion (iii) of [Proposition 3.7](#) holds as is. The assertions (i) and (ii) of [Proposition 3.7](#) even hold when (I, Σ_I, μ_I) is just a measure space, with no further assumptions, by reasoning with almost simple mappings instead of simple mappings.

Using [Proposition 3.7](#), the isometries defined in [Proposition 3.3](#) can be uniquely extended to nonlinear Lebesgue spaces as soon as N is complete. The latter statement actually yields a nonlinear analogue of the Fubini–Lebesgue theorem (see [Remark 3.12](#) for a further discussion on that matter).

Theorem 3.9 (Extension of $\text{sec}_{I,p}$ and $\text{sec}_{M,p}$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Suppose that N is complete. Then, the following assertions hold:*

(i) *the isometry $\text{sec}_{I,p}$ defined in (i) of [Proposition 3.3](#) extends uniquely to the isometry*

$$\overline{\text{sec}}_{I,p} : \begin{cases} L_{\hat{h}}^p(I \times M, N) & \longrightarrow & L^p(I, L_{\hat{h}}^p(M, N)) \\ [\hat{c}] & \longmapsto & [t \mapsto [\hat{c}(t, \cdot)]] \end{cases}.$$

(ii) *the isometry $\text{sec}_{M,p}$ defined in (ii) of [Proposition 3.3](#) extends uniquely to the isometry*

$$\overline{\text{sec}}_{M,p} : \begin{cases} L_{\hat{h}}^p(I \times M, N) & \longrightarrow & L_{\hat{h}}^p(M, L^p(I, N)) \\ [\hat{c}] & \longmapsto & [x \mapsto [\hat{c}(\cdot, x)]] \end{cases}.$$

Remark 3.10 (On relaxing the finiteness assumption on d_N). [Theorem 3.9](#) extends to the case where d_N might take infinite values (see [Assumption 2.1](#) for the assumption on d_N) by assuming that h is constant equal to some $z_0 \in N$. In that case, we can define the open set, hence Borel set, $N_0 := \{y \in N : d_N(z_0, y) < \infty\}$ in N on which d_N is finite. Then, any mapping in $L_{\hat{h}}^p(I \times M, N)$ takes its values $\mathcal{L}^1 \otimes \mu_M$ -a.e. in N_0 and $L_{\hat{h}}^p(I \times M, N)$ can be identified with $L_{\hat{h}}^p(I \times M, N_0)$. Hence, when h is constant, the extension to the case where d_N might take infinite values reduces to [Theorem 3.9](#) by replacing N with N_0 .

Proof. Let $h \in L^0(M, N)$ and $p \in [1, \infty)$.

(i) *Step 1: Extension of $\text{sec}_{I,p}$.* Recall that $\text{sec}_{I,p}$ is an isometry from $(E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N), \hat{D}_p)$ to $(E(I, E_h(M, N)) \cap L_{\hat{h}}^p(M, N), d_{p,p})$. Now, $L_{\hat{h}}^p(I \times M, N)$ being the completion of $E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$, by (i) of [Proposition 3.7](#) and [Proposition 2.17](#), and $L^p(I, L_{\hat{h}}^p(M, N))$ being the completion of $E(I, E_h(M, N)) \cap L_{\hat{h}}^p(M, N)$, by (ii) of [Proposition 3.7](#) and [Proposition 2.17](#), we get that $\text{sec}_{I,p}$ extends uniquely to an isometry $\overline{\text{sec}}_{I,p}$ from $L_{\hat{h}}^p(I \times M, N)$ to $L^p(I, L_{\hat{h}}^p(M, N))$ by setting $\overline{\text{sec}}_{I,p}(\hat{c}) := \text{sec}_{I,p}(\hat{c})$ for all $\hat{c} \in E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$ and $\overline{\text{sec}}_{I,p}(\hat{c}) := \lim_{n \rightarrow \infty} \text{sec}_{I,p}(\hat{c}_n)$ for all $\hat{c} \in L_{\hat{h}}^p(I \times M, N) \setminus (E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N))$ with $(\hat{c}_n)_{n \in \mathbb{N}}$ being any sequence in $E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$ converging to \hat{c} , since, $\text{sec}_{I,p}$ being an isometry, the choice of sequence does not matter.

Step 2: Explicit expression of the extension of $\text{sec}_{I,p}$. Let $\hat{c} \in L_{\hat{h}}^p(I \times M, N)$. Then, there is $\hat{c}' \in L_{\hat{h}}^p(I \times M, N)$ such that $\hat{D}_p(\hat{c}, \hat{c}') = 0$. By (i) of [Proposition 3.7](#), there is a sequence $(\hat{c}'_n)_{n \in \mathbb{N}}$ in $E_{\hat{h},\sigma}(I \times M, N) \cap L_{\hat{h}}^p(I \times M, N)$ such that $\hat{D}_p(\hat{c}'_n, \hat{c}') \rightarrow 0$ as $n \rightarrow \infty$ and define $\hat{c}_n := [\hat{c}'_n]$ as well as $c_n := \text{sec}_{I,p}(\hat{c}'_n)$. By definition of the \hat{c}'_n 's ([Definition 3.2](#)), the union of the sets on which the \hat{c}'_n 's differ from \hat{h} is contained in the countable union of a collection $(I_m \times M_m)_{m \in \mathbb{N}}$ of measurable rectangles of finite $\mathcal{L}^1 \otimes \mu_M$ -measure. Thus, the set $\hat{M} := \cup_{m \in \mathbb{N}} M_m$ has σ -finite μ_M -measure. In addition, since $\hat{D}_p(\hat{c}'_n, \hat{c}') \rightarrow 0$ as $n \rightarrow \infty$, we get that

$$\int_{I \times (M \setminus \hat{M})} d_N(\hat{h}(t, x), \hat{c}'(t, x))^p \, d\mathcal{L}^1 \otimes \mu_M(t, x) = 0,$$

so that, up to the modification on a $\mathcal{L}^1 \otimes \mu_M$ -null set, we can assume that $\hat{c}'(t, x) = \hat{h}(t, x)$ for all $(t, x) \in I \times (M \setminus \hat{M})$. Now, define $c : t \mapsto \hat{c}'(t, \cdot)$ for which we know that $c(t) \in \mathcal{L}_s^0(M, N)$ for all $t \in I$. Then, since I has finite \mathcal{L}^1 -measure and \hat{M} has σ -finite μ_M -measure, we have, by the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1], that $d_{p,p}(c_n, c) = \hat{D}_p(\hat{c}'_n, \hat{c}') \rightarrow 0$ as $n \rightarrow \infty$, so that, up to the extraction of a subsequence, we can assume for \mathcal{L}^1 -a.e. $t \in I$ that $D_p(c_n(t), c(t)) \rightarrow 0$ as $n \rightarrow \infty$. Finally, using the triangle inequality and the definition of $\text{sec}_{I,p}$, we get that for \mathcal{L}^1 -a.e. $t \in I$

$$D_p(\overline{\text{sec}}_{I,p}(\hat{c})(t), \hat{c}'(t, \cdot)) \leq D_p(\overline{\text{sec}}_{I,p}(\hat{c})(t), \text{sec}_{I,p}(\hat{c}_n)(t)) + D_p(c_n(t), c(t))$$

holds for all $n \in \mathbb{N}$. Thus, using the definition of $\overline{\text{sec}}_{I,p}$ and up to the further extraction of subsequence to obtain for \mathcal{L}^1 -a.e. $t \in I$ that $D_p(\overline{\text{sec}}_{I,p}(\hat{c})(t), \text{sec}_{I,p}(\hat{c}_n)(t)) \rightarrow 0$ as $n \rightarrow \infty$, we get that

$$D_p(\overline{\text{sec}}_{I,p}(\hat{c})(t), [\hat{c}'(t, \cdot)]) = 0 \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I.$$

- (ii) The proof follows from the same arguments as the proof of (i), but using (iii) of [Proposition 3.7](#) instead of (ii) of [Proposition 3.7](#). \square

Remark 3.11 (On relaxing the assumptions on I). Since the proof of [Theorem 3.9](#) only relies on the additional use of [Proposition 3.7](#) relatively to [Proposition 3.3](#) and we can define, if necessary, $\hat{I} := \cup_{m \in \mathbb{N}} I_m$ to use the Fubini–Tonelli theorem in step 2 of the proof of assertion (i) of [Theorem 3.9](#), the relaxations of [Propositions 3.3](#) and [3.7](#) given by [Remarks 3.5](#) and [3.8](#) ensure that [Proposition 3.7](#) holds when (I, Σ_I, μ_I) is just a measure space and either μ_I is finite or h is simple while replacing $L_{\underline{h}}^p(M, L^p(I, N))$ with $L_{\underline{h}}^p(M, L^0(I, N))$ (see [Remark 3.5](#) for a definition of this space) in assertion (ii) of [Theorem 3.9](#) for that last case and recalling that $(L^0(I, N), d_p)$ is complete when N is ([Corollary 2.18](#)). If h is not only simple, but constant, the assertion (ii) of [Theorem 3.9](#) holds as is. The assertion (i) of [Theorem 3.9](#) even holds when (I, Σ_I, μ_I) is just a measure space, with no further assumptions.

Remark 3.12 (Related results in the literature). [Theorem 3.9](#) is a well-known result for linear Lebesgue spaces in the case where N is a Banach space, h is constant, and both μ_I (see [Remark 3.11](#) for the definition of this measure) and μ_M are assumed σ -finite [[HVN VW16](#), Proposition 1.2.24]. When (I, Σ_I, μ_I) is just a measure space, with no further assumptions, d_N is non-finite and h is constant ([Remarks 3.10](#) and [3.11](#)), [Theorem 3.9](#) is actually a nonlinear version of the Fubini–Lebesgue theorem that requires neither that μ_I and μ_M are both σ -finite [[Coh13](#), Theorem 5.2.2], nor that they are both complete [[DiB02](#), Chapter II, Theorem 14.1].

As a direct consequence of [Theorem 3.9](#), the isometry defined in [Corollary 3.6](#) extends uniquely to an isometry yielding a characterization of L^p curves in nonlinear Lebesgue spaces.

Theorem 3.13 (Characterization of L^p curves in nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty)$. Suppose that N is complete. Then, the isometry i_p defined in [Corollary 3.6](#) extends uniquely to the isometry*

$$\bar{i}_p := \overline{\text{sec}}_{M,p} \circ \overline{\text{sec}}_{I,p}^{-1} : L^p(I, L_{\underline{h}}^p(M, N)) \rightarrow L_{\underline{h}}^p(M, L^p(I, N)).$$

Remark 3.14 (On relaxing the finiteness assumption on d_N). Thanks to [Remark 3.10](#), the conclusion of [Theorem 3.13](#) holds when d_N is not assumed finite (see [Assumption 2.1](#) for the assumption d_N), but requires h being constant in that case.

Remark 3.15 (On relaxing the assumptions on I). Thanks to [Remark 3.11](#), the conclusion of [Theorem 3.13](#) holds when (I, Σ_I, μ_I) is just a measure space such that either μ_I is finite or h is simple while replacing $L_{\underline{h}}^p(M, L^p(I, N))$ with $L_{\underline{h}}^p(M, L^0(I, N))$ (see [Remark 3.5](#) for a definition of this space) in that last case. If h is not only simple, but constant, the conclusion of [Theorem 3.13](#) holds as is.

Remark 3.16 (Related results in the literature). This result is well-known in the context of linear Lebesgue spaces where N is a Banach space and h is constant [[HVN VW16](#), Corollary 1.2.23].

Let us end this section with a discussion on a edge case to better understand the scope of application of [Theorems 3.9](#) and [3.13](#).

Example 3.17 (Edge case for Theorems 3.9 and 3.13). One sometimes encounters the claim that the Fubini–Lebesgue theorem, like the Fubini–Tonelli theorem, requires the measures to be σ -finite. In this context, the indicator function of the diagonal with the Lebesgue and counting measures is frequently mentioned as an edge case [HVN VW16, Remark 1.2.8]. In fact, this example contradicts neither Theorem 3.9 nor Theorem 3.13, but remains an interesting edge case to better understand the scope of application of Theorems 3.9 and 3.13. Indeed, suppose that $I = M = [0, 1]$ with its standard topology and its Borel σ -algebra, that $\mu_I = \mathcal{L}^1|_I$, that $\mu_M = \#$, that $N = \mathbb{R}$ with its standard topology and that $h \equiv 0$. Also, let $D := \{(x, x) : x \in [0, 1]\}$ be the diagonal, which is a Borel set, and let $f := \mathbb{1}_D$. Then, we have

$$\int_I \left(\int_M \mathbb{1}_D(t, x) d\mu_M(x) \right) d\mu_I(t) = \int_I \left(\int_M \mathbb{1}_{\{t\}}(x) d\mu_M(x) \right) d\mu_I(t) = \int_{[0,1]} \#(\{t\}) d\mathcal{L}^1_I(t) = 1$$

and

$$\int_M \left(\int_I \mathbb{1}_D(t, x) d\mu_I(t) \right) d\mu_M(x) = \int_M \left(\int_I \mathbb{1}_{\{x\}}(t) d\mu_I(t) \right) d\mu_M(x) = \int_{[0,1]} \mathcal{L}^1|_I(\{x\}) d\#(x) = 0.$$

Hence, we get

$$\int_I \left(\int_M \mathbb{1}_D(t, x) d\mu_M(x) \right) d\mu_I(t) \neq \int_M \left(\int_I \mathbb{1}_D(t, x) d\mu_I(t) \right) d\mu_M(x).$$

However, the above conclusion contradicts neither Theorem 3.9 nor Theorem 3.13. Indeed, the diagonal D , which is uncountable, has infinite $\mu_I \otimes \mu_M$ -measure, as, otherwise, D would be contained in the countable union of measurable rectangles of finite $\mu_I \otimes \mu_M$ -measure [DiB02, Chapter II, Proposition 10.2]; hence, it would be countable since μ_M is the counting measure, which yields a contradiction. Thus, the equivalence class of the indicator function of D does not belong to $L^1(I \times M, \mathbb{R})$ and neither $\overline{\text{sec}}_{I,p}$ nor $\overline{\text{sec}}_{M,p}$ are defined on f (Theorem 3.9). Furthermore, $f_I : t \mapsto f(t, \cdot) = \mathbb{1}_{\{t\}}$ does not have a separable range in $L^1(M, \mathbb{R})$ since the family $\{\mathbb{1}_{\{t\}} : t \in [0, 1]\}$ is uncountable and $\int_M |\mathbb{1}_{\{t\}}(x) - \mathbb{1}_{\{s\}}(x)| d\mu_M(x) = 2 > 0$ as soon as $s \neq t$, so that $\mathbb{1}_{\{t\}}$ and $\mathbb{1}_{\{s\}}$ cannot belong to the same equivalence class. Also, f_I does not admit an equivalent mapping with separable range in $L^1(M, \mathbb{R})$ since the counting measure imposes a pointwise equality on M for the values of f_I , so it cannot be approximated by simple mappings and $\overline{\text{sec}}_{I,p}^{-1}$ is actually not defined on the equivalence class of f_I (Theorem 3.9). On the contrary, $f_M : x \mapsto f(\cdot, x) = \mathbb{1}_{\{x\}}$ does satisfy the conditions of Theorems 3.9 and 3.13. Indeed, $D_{p,p}(f_M, \underline{h}) = 0$, so that f_M is assimilated to the equivalence class of \underline{h} whose image by $\overline{\text{sec}}_{M,p}^{-1}$ is the equivalence class of the mapping constant equal to 0 on $I \times M$ and by $\overline{\text{sec}}_{I,p}^{-1}$ is the equivalence class of the mapping constant equal to h on I . The mappings f and f_I are thus in fact not related to f_M through the sections mappings on I and M defined in Theorem 3.9.

We now continue with the characterization of curves in nonlinear Lebesgue spaces using the isometry defined in Theorem 3.13.

3.2. Characterization of absolutely continuous curves for $p > 1$

We first clarify some recurrent notation used in this section and thereafter.

Notation 3.18. Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. From thereon:

- (i) $L_h^p(M, \mathcal{A}C^p(I, N))$ denotes the set of equivalence classes of separably valued measurable mapping from M to $(\mathcal{A}C^p(I, N), d_p)$ at a finite $D_{p,p}$ -distance from \underline{h} (see Notation 3.1 for the definition of \underline{h}).
- (ii) $|c'|_p := |c'|_{L_h^p(M, N)}$ denotes the metric derivative of the curve $c \in \mathcal{A}C^p(I, L_h^p(M, N))$ (Theorem 2.29).
- (iii) Q_p denotes the restriction to $\mathcal{A}C^p(I, L_h^p(M, N))$ of the quotient mapping

$$Q : \begin{cases} \mathcal{L}_s^0(I, L_h^p(M, N)) & \longrightarrow L^0(I, L_h^p(M, N)) \\ c & \longmapsto [c] \end{cases}.$$

Then, the isometry given by Theorem 3.13 yields a characterization of absolutely continuous curves in nonlinear Lebesgue spaces for $p > 1$.

Theorem 3.19 (Characterization of absolutely continuous curves for $p > 1$). Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that N is complete. Then, the isometry \tilde{i}_p defined in [Theorem 3.13](#) induces the isometric mapping

$$\tilde{i}_p := (q_p^{-1})_* \circ \bar{i}_p \circ Q_p : (\mathcal{AC}^p(I, L_h^p(M, N)), d_{p,p}) \rightarrow (L_h^p(M, \mathcal{AC}^p(I, N)), D_{p,p})$$

whose inverse mapping on its range is $\tilde{i}_p^{-1} := Q_p^{-1} \circ \bar{i}_p^{-1} \circ (q_p)_*$ and satisfies

$$\tilde{i}_p^{-1}(f)(t) = (e_{t,p})_*(f) \quad \text{for all } t \in I \text{ and } f \in \tilde{i}_p(\mathcal{AC}^p(I, L_h^p(M, N))). \quad (2)$$

In addition, \tilde{i}_p satisfies for all $c \in \mathcal{AC}^p(I, L_h^p(M, N))$ that

$$|c'|_p^p(t) = \int_M \tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I \quad (3)$$

and its range is the set of all $f \in L_h^p(M, \mathcal{AC}^p(I, N))$ such that

$$\int_M \int_I |f(x)'|_N^p(t) dt d\mu_M(x) < \infty.$$

Remark 3.20 (On the definition of pushforwards for equivalence classes). The pushforward of an equivalence class by a mapping is meant as the equivalence class of the composition of the given mapping with a compatible representative of the input equivalence class.

Remark 3.21 (On relaxing the finiteness assumption on d_N). The conclusion of [Theorem 3.19](#) holds when d_N is not assumed finite (see [Assumption 2.1](#) for the assumption on d_N) by assuming that h is constant equal to some $z_0 \in N$. In that case, we can define the open set, hence Borel set, $N_0 := \{y \in N : d_N(z_0, y) < \infty\}$ in N on which d_N is finite. Then, any mapping in $L_h^p(M, N)$ takes its values μ_M -a.e. in N_0 and $L_h^p(M, N)$ can be identified with $L_h^p(M, N_0)$. Hence, when h is constant, the extension to the case where d_N might take infinite values reduces to [Theorem 3.19](#) by replacing N with N_0 .

Proof. The proof follows the same approach used in [[Lis07](#), Theorems 4 and 5] for Wasserstein spaces, which consists in using piecewise constant approximations of absolutely continuous curves to prove inequalities and propagate the results to absolutely continuous curves by passing to the limit.

Let $h \in \mathcal{L}_s^0(M, N)$, $p \in (1, \infty)$.

Step 1: Reduction to the case where N is separable. Let us first reduce the proof to the case where N is separable. Indeed, when $c \in \mathcal{AC}^p(I, L_h^p(M, N))$, the set $N_0 := \overline{\cup_{t \in \mathbb{Q} \cap I} c(t)(M)}$ is separable as the closure of the countable union of separable sets. Then, since c is continuous, we have for all $t \in I$ that $c(t) = \lim_{s \rightarrow t, s \in \mathbb{Q} \cap I} c(s)$ in $L_h^p(M, N)$, hence, up to the extraction of a subsequence, $c(t)(x) = \lim_{s \rightarrow t, s \in \mathbb{Q} \cap I} c(s)(x)$ in N for μ_M -a.e. $x \in M$, that is, $c(t)(x) \in N_0$ for μ_M -a.e. $x \in M$. Thus, we can replace N with N_0 in the rest of the proof, so that $\mathcal{L}^0(M, N) = \mathcal{L}_s^0(M, N)$.

Step 2: Definition of \tilde{i}_p . Let $c \in \mathcal{AC}^p(I, L_h^p(M, N))$. Then, $Q_p(c)$ belongs to $W^{1,p}(I, L_h^p(M, N))$, by [Proposition 2.30](#), hence $(\bar{i}_p \circ Q_p)(c)$ belongs to $L_h^p(M, L^p(I, N))$, by [Theorem 3.13](#). Define for all $n \in \mathbb{N}^*$ the set $I_n := \{0, 1, \dots, 2^n\}$ and the sequence $(t_{n,i})_{i \in I_n}$ such that $t_{n,i} := a + \mathcal{L}^1(I) i 2^{-n}$. Also, define the simple mapping $c_n : I \rightarrow L_h^p(M, N)$ as $c_n|_{[t_{n,i}, t_{n,i+1})} \equiv c(t_{n,i})$ for all $i \in I_{n-1}$ and $c_n(b) := c(b)$. Since the set of dyadic rationals (scaled by $\mathcal{L}^1(I)$ and translated by a) is dense in I and c is continuous on $[a, b)$ and $\mathcal{L}^1(\{b\}) = 0$, we have for \mathcal{L}^1 -a.e. $t \in I$ that $c_n(t) \rightarrow c(t)$ in $L_h^p(M, N)$ as $n \rightarrow \infty$. In addition, if $t \in [t_{n,i}, t_{n,i+1})$,

$$D_p(c_n(t), c(t)) = D_p(c(t_{n,i}), c(t)) \leq \int_{t_{n,i}}^t |c'|_p(u) du \leq \int_I |c'|_p(u) du.$$

Hence, we get, using Hölder's inequality, that

$$D_p(c_n(t), c(t))^p \leq \mathcal{L}^1(I)^{p-1} \int_I |c'|_p^p(u) du \quad \text{for all } (t, n) \in I \times \mathbb{N}^*.$$

Since I has finite \mathcal{L}^1 -measure, the mapping constant equal to $\mathcal{L}^1(I)^{p-1} \int_I |c'|_p^p(u) du < \infty$ on I is integrable, so that, by Lebesgue's dominated convergence theorem [Coh13, Theorem 2.4.5], we have $d_{p,p}(c_n, c) \rightarrow 0$ as $n \rightarrow \infty$. In addition, for all $n \in \mathbb{N}^*$ and all $i \in I_n$ there is $c_{n,i} \in \mathcal{L}_h^p(M, N)$ such that $D_p(c_{n,i}, c(t_{n,i})) = 0$, so we can define $c'_n : I \rightarrow \mathcal{L}_h^p(M, N)$ as $c'_n|_{[t_{n,i}, t_{n,i+1})} \equiv c_{n,i}$ for all $i \in I_{n-1}$ and $c'_n(b) := c_{n,2^n}$, which satisfies $D_p(c_n(t), c'_n(t)) = 0$ for all $t \in I$. Thus, we can define $\hat{c}'_n : I \times M \rightarrow N$ as $\hat{c}'_n(t, x) := c'_n(t)(x)$, which belongs to $\mathcal{L}_h^p(I \times M, N)$ (simple on I and by definition of the $c_{n,i}$'s) and satisfies $D_p(\hat{c}'_n(t, \cdot), c_n(t)) = 0$ for all $t \in I$. Hence, $\overline{\text{sec}}_{I,p}$ being an isometry (Theorem 3.9), we get that $\hat{D}_p(\hat{c}'_n, \overline{\text{sec}}_{I,p}^{-1}(c_n)) = 0$. Now, defining the mapping $f'_n : M \rightarrow \mathcal{L}^p(I, N)$ as $f'_n(x) := \hat{c}'_n(\cdot, x)$ and recalling that $\overline{\text{sec}}_{M,p}$ is an isometry (Theorem 3.9), we get that $D_{p,p}(f'_n, \overline{\text{sec}}_{M,p}^{-1}(c_n)) = 0$, hence that $d_p(f'_n(x), \overline{\text{sec}}_{M,p}^{-1}(c_n)(x)) = 0$ for μ_M -a.e. $x \in M$. Then, define for all $n \in \mathbb{N}^*$ the functional $\psi_n : L^p(I, N) \rightarrow [0, \infty]$ as

$$\psi_n(\eta) := \sup_{\mathcal{L}^1(I)2^{-n} < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau} \eta(t))^p dt,$$

so that $\psi_n(\eta) \leq \psi_{n+1}(\eta)$ for all $(n, \eta) \in \mathbb{N}^* \times L^p(I, N)$ and let $\psi_\infty(\eta) := \lim_{n \rightarrow \infty} \psi_n(\eta)$, which is lower-semicontinuous. Now, observe that for all $(m, n) \in (\mathbb{N}^*)^2$ such that $m \geq n$ we have for μ_M -a.e. $x \in M$ that

$$\begin{aligned} \psi_m(\overline{\text{sec}}_{I,p}(c_n)(x)) &= \sup_{\mathcal{L}^1(I)2^{-m} < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau} \overline{\text{sec}}_{I,p}(c_n)(x)(t))^p dt \\ &\leq \sup_{\mathcal{L}^1(I)2^{-n} < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau} \overline{\text{sec}}_{I,p}(c_n)(x)(t))^p dt \\ &= \sup_{\mathcal{L}^1(I)2^{-n} < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau} f'_n(x)(t))^p dt. \end{aligned}$$

Then, take the integer $k \geq 1$ such that $k \mathcal{L}^1(I)2^{-n} \leq \tau \leq (k+1) \mathcal{L}^1(I)2^{-n}$ and define $u_{n,i}(t) := t + \mathcal{L}^1(I)i2^{-n}$, so that, using the triangle inequality, we get

$$\begin{aligned} d_N(f'_n(x)(t), f'_n(x)(t+\tau)) &\leq \sum_{i=0}^k d_N(f'_n(x)(u_{n,i}(t)), f'_n(x)(u_{n,i+1}(t))) \\ &= \sum_{i=0}^k d_N(c'_n(u_{n,i}(t))(x), c'_n(u_{n,i+1}(t))(x)). \end{aligned}$$

Hence, Hödler's discrete inequality yields that

$$d_N(f'_n(x)(t), f'_n(x)(t+\tau))^p \leq (k+1)^{p-1} \sum_{i=0}^k d_N(c'_n(u_{n,i}(t))(x), c'_n(u_{n,i+1}(t))(x))^p.$$

Thus, we get

$$\begin{aligned} \int_a^{b-\tau} (\Delta_{N,\tau} f'_n(x)(t))^p dt &\leq \tau^{-p} \int_a^{b-k \mathcal{L}^1(I)2^{-n}} d_N(f'_n(x)(t), f'_n(x)(t+\tau))^p dt \\ &\leq \tau^{-p} \int_a^{a+(2^n-k) \mathcal{L}^1(I)2^{-n}} (k+1)^{p-1} \sum_{i=0}^k d_N(c'_n(u_{n,i}(t))(x), c'_n(u_{n,i+1}(t))(x))^p dt \\ &= \tau^{-p} (k+1)^{p-1} \sum_{i=0}^k \mathcal{L}^1(I)2^{-n} \sum_{j=0}^{2^n-k-1} d_N(c_{n,i+j}(x), c_{n,i+j+1}(x))^p. \end{aligned}$$

Since $d_N(c_{n,i+j}(x), c_{n,i+j+1}(x))^p$ is counted at most $k+1$ times, we get

$$\int_a^{b-\tau} (\Delta_{N,\tau} f'_n(x)(t))^p dt \leq \left(\frac{k+1}{\tau}\right)^p \mathcal{L}^1(I)2^{-n} \sum_{i=0}^{2^n-1} d_N(c_{n,i}(x), c_{n,i+1}(x))^p.$$

The latter inequality together with the fact that $\mathcal{L}^1(I)2^{-n} \leq (\mathcal{L}^1(I)2^{-n})^{1-p} \frac{2^n}{k^p}$ yields that

$$\begin{aligned} \psi_m(\bar{i}_p(c_n)(x)) &\leq \left(\frac{k+1}{k}\right)^p (\mathcal{L}^1(I)2^{-n})^{1-p} \sum_{i=0}^{2^n-1} d_N(c_{n,i}(x), c_{n,i+1}(x))^p \\ &\leq 2^p (\mathcal{L}^1(I)2^{-n})^{1-p} \sum_{i=0}^{2^n-1} d_N(c_{n,i}(x), c_{n,i+1}(x))^p \\ &= 2^p (\mathcal{L}^1(I)2^{-n})^{1-p} \sum_{i=0}^{2^n-1} d_N(c_{n,i}(x), c_{n,i+1}(x))^p. \end{aligned}$$

Then, passing to the limit when $m \rightarrow \infty$ and integrating over M , we get for all $n \in \mathbb{N}^*$ that

$$\begin{aligned} \int_M \psi_\infty(\bar{i}_p(c_n)(x)) \, d\mu_M(x) &\leq 2^p (\mathcal{L}^1(I)2^{-n})^{1-p} \int_M \sum_{i=0}^{2^n-1} d_N(c_{n,i}(x), c_{n,i+1}(x))^p \, d\mu_M(x) \\ &= 2^p (\mathcal{L}^1(I)2^{-n})^{1-p} \sum_{i=0}^{2^n-1} D_p(c(t_{n,i}), c(t_{n,i+1}))^p \\ &\leq 2^p (\mathcal{L}^1(I)2^{-n})^{1-p} \sum_{i=0}^{2^n-1} \left(\int_{t_{n,i}}^{t_{n,i+1}} |c'|_p(t) \, dt \right)^p \\ &\leq 2^p \sum_{i=0}^{2^n-1} \int_{t_{n,i}}^{t_{n,i+1}} |c'|_p^p(t) \, dt \\ &= 2^p \int_I |c'|_p^p(t) \, dt, \end{aligned}$$

where Hölder's inequality was used in the second to last inequality. Now, \bar{i}_p being isometric and using the definition of Q_p (Notation 3.18), we also have $D_{p,p}(\bar{i}_p(c_n), (\bar{i}_p \circ Q_p)(c)) \rightarrow 0$ as $n \rightarrow \infty$. Then, up to the extraction of a subsequence, we can assume for μ_M -a.e. $x \in M$ that $\bar{i}_p(c_n)(x) \rightarrow (\bar{i}_p \circ Q_p)(c)(x)$ in $L^p(I, N)$ as $n \rightarrow \infty$, so that, by lower-semicontinuity of ψ_∞ and using Fatou's lemma [Coh13, Theorem 2.4.4], we get

$$\begin{aligned} \int_M \psi_\infty((\bar{i}_p \circ Q_p)(c)(x)) \, d\mu_M(x) &\leq \int_M \liminf_{n \rightarrow \infty} \psi_\infty(\bar{i}_p(c_n)(x)) \, d\mu_M(x) \\ &\leq \liminf_{n \rightarrow \infty} \int_M \psi_\infty(\bar{i}_p(c_n)(x)) \, d\mu_M(x) \\ &\leq 2^p \int_I |c'|_p^p(t) \, dt < \infty. \end{aligned}$$

This yields that for μ_M -a.e. $x \in M$

$$\psi_\infty((\bar{i}_p \circ Q_p)(c)(x)) = \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau}(\bar{i}_p \circ Q_p)(c)(x)(t))^p \, dt < \infty,$$

which implies that $(\bar{i}_p \circ Q_p)(c)(x)$ belongs to $W^{1,p}(I, N)$ (Definition 2.25). Hence, $W^{1,p}(I, N)$ being a Borel subset of $L^p(I, N)$ (Remark 2.27), $(\bar{i}_p \circ Q_p)(c)$ belongs to $L^p_{\underline{h}}(M, W^{1,p}(I, N))$. Therefore, the isometric mapping

$$\tilde{i}_p := (q_p^{-1})_* \circ \bar{i}_p \circ Q_p : \mathcal{AC}^p(I, L^p_{\underline{h}}(M, N)) \rightarrow L^p_{\underline{h}}(M, \mathcal{AC}^p(I, N))$$

is well-defined and its inverse mapping on its range is $\tilde{i}_p^{-1} := Q_p^{-1} \circ \bar{i}_p^{-1} \circ (q_p)_*$.

Step 3: Proof of Equation (2). Let f be in the range of \tilde{i}_p . Then, there is a unique $c \in \mathcal{AC}^p(I, L^p_{\underline{h}}(M, N))$ such that $f = \tilde{i}_p(c)$ and define

$$\hat{f} := (\overline{\text{sec}}_{I,p}^{-1} \circ \bar{i}_p^{-1} \circ (q_p)_*)(f) = (\overline{\text{sec}}_{M,p}^{-1} \circ (q_p)_*)(f).$$

Also, note that $\hat{f} = (\overline{\text{sec}}_{I,p}^{-1} \circ Q_p)(c)$ since $\tilde{i}_p = (q_p^{-1})_* \circ \bar{i}_p \circ Q_p$. Now, there is $f' \in \mathcal{L}_h^p(M, \mathcal{AC}^p(I, N))$ such that $D_{p,p}(f', f) = 0$, which implies that $f'(x) = f(x)$ in $C(I, N)$ for μ_M -a.e. $x \in M$ (Proposition 2.30). Thus, we can define the mapping $\tilde{f} : I \times M \rightarrow N$ as $\tilde{f}(t, x) := f'(x)(t) = (e_{t,p} \circ f')(x)$. To show that \tilde{f} is measurable, consider for all $n \in \mathbb{N}^*$ the mapping $\tilde{f}_n : I \times M \rightarrow N$ defined as $\tilde{f}_n(t, x) := (e_{t_n,p} \circ f')(x)$ if $t \in [t_{n,i}, t_{n,i+1})$ with $i \in I_{n-1}$ and $\tilde{f}_n(b, x) := (e_{b,p} \circ f')(x)$, so that \tilde{f}_n belongs to $\mathcal{L}^0(I \times M, N)$ since $e_{t,p} \circ f' \in \mathcal{L}^0(M, N)$ for all $t \in I$ (Proposition 2.33). Then, the set of dyadic rationals (scaled by $\mathcal{L}^1(I)$ and translated by a) being dense in I and $f'(x)$ being continuous on $[a, b]$ for all $x \in M$, we have $\tilde{f}(t, x) = \lim_{n \rightarrow \infty} \tilde{f}_n(t, x)$ for all $(t, x) \in I \times M$. Therefore, \tilde{f} is measurable as the pointwise limit of measurable mappings (Proposition 2.4). In addition, $D_p(\tilde{f}(t, \cdot), (e_{t,p})_*(f)) = 0$ for all $t \in I$ since $\tilde{f}(\cdot, x) = f(x)$ in $C(I, N)$ for μ_M -a.e. $x \in M$. Then, using the definitions of \tilde{f} , $\overline{\text{sec}}_{M,p}$ (Theorem 3.9) and q_p (Proposition 2.30), we have $d_p(\tilde{f}(\cdot, x), \hat{f}(\cdot, x)) = 0$ for μ_M -a.e. $x \in M$. Hence, $\overline{\text{sec}}_{M,p}$ being an isometry, we get $\hat{D}_p(\tilde{f}, \hat{f}) = 0$. Thus, using the definitions of \tilde{f} , $\overline{\text{sec}}_{I,p}$ (Theorem 3.9) and Q_p (Notation 3.18), we get $D_p(\tilde{f}(t, \cdot), c(t)) = 0$ for \mathcal{L}^1 -a.e. $t \in I$, which yields $\tilde{i}_p^{-1}(f)(t) = (e_{t,p})_*(f)$ in $L_h^p(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$. It remains to prove the equality for all $t \in I$. To do so, define the set

$$\begin{aligned} A &:= \{I \times M : |\tilde{i}_p(c)(x)'|_N(t) \text{ does not exist}\} \\ &= \{I \times M : \limsup_{\tau, \tau' \rightarrow 0} |g_\tau(t, \tilde{i}_p(c)(x)) - g_{\tau'}(t, \tilde{i}_p(c)(x))| > 0\}, \end{aligned}$$

which is measurable since $\tilde{i}_p(c)$ is measurable from M to $(\mathcal{AC}^p(I, N), d_\infty)$ and the mappings

$$g_\tau : \begin{cases} I \times \mathcal{AC}^p(I, N) & \longrightarrow [0, \infty) \\ (t, u) & \longmapsto \Delta_{N, \tau} u(t) \end{cases}$$

are continuous, hence Borel measurable, for all $\tau > 0$, so that $(t, x) \mapsto |\tilde{i}_p(c)(x)'|_N(t)$ is measurable from $I \times M$ to N , as the pointwise limit of compositions of measurable mappings. In addition, $\mathcal{L}^1(A^x) = 0$ since $\tilde{i}_p(c)(x) \in \mathcal{AC}^p(I, N)$ for μ_M -a.e. $x \in M$. Then, recalling that d_p separates elements of $\mathcal{AC}^p(I, N)$ (Proposition 2.30) and defining the measurable set

$$\overline{M}_c := \{x \in M : d_p(\tilde{i}_p(c)(x), \underline{h}(x)) > 0\} = \{x \in M : d_\infty(\tilde{i}_p(c)(x), \underline{h}(x)) > 0\}$$

of σ -finite μ_M -measure, we have $|\tilde{i}_p(c)(x)'|_N(t) = 0$ for all $(x, t) \in (M \setminus \overline{M}_c) \times I$ since $\underline{h}(x)$ is constant over I . Thus, A is a subset of the measurable rectangle $I \times \overline{M}_c$ and, \overline{M}_c having σ -finite μ_M -measure and I having finite \mathcal{L}^1 -measure, we get, by the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1], that $\mathcal{L}^1 \otimes \mu_M(A) = 0$ and $\mu_M(A_t) = 0$ for \mathcal{L}^1 -a.e. $t \in I$. Hence, for all $(s, t) \in I^2$ such that $s < t$, we have

$$\begin{aligned} D_p((e_{s,p})_*(\tilde{i}_p(c)), (e_{t,p})_*(\tilde{i}_p(c)))^p &= \int_M d_N(\tilde{i}_p(c)(x)(s), \tilde{i}_p(c)(x)(t))^p d\mu_M(x) \\ &\leq \int_M \left(\int_s^t |\tilde{i}_p(c)(x)'|_N(t) dt \right)^p d\mu_M(x) \\ &\leq |t - s|^{p-1} \int_M \int_s^t |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) \\ &= |t - s|^{p-1} \int_{\overline{M}_c} \int_s^t |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) \\ &= |t - s|^{p-1} \int_s^t \int_{\overline{M}_c} |\tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) dt \\ &= |t - s|^{p-1} \int_s^t \int_M |\tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) dt, \end{aligned}$$

where the second to last equality follows from the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1]. The latter inequality yields that $t \mapsto (e_{t,p})_*(\tilde{i}_p(c))$ belongs to $\mathcal{AC}^p(I, L_h^p(M, N))$, hence, $d_{p,p}$ separating the elements of $\mathcal{AC}^p(I, L_h^p(M, N))$ (Proposition 2.30), we get $\tilde{i}_p^{-1}(f)(t) = (e_{t,p})_*(f)$ in $L_h^p(M, N)$ for all $t \in I$, which ends the proof of Equation (2).

Step 4: Proof of Equation (3). Let $c \in \mathcal{A}C^p(I, L_h^p(M, N))$. Using Lebesgue's differentiation theorem [DiB02, Theorem 11.1] in the previous inequality, we obtain that

$$|c'|_p^p(t) \leq \int_M \tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I. \quad (4)$$

In addition, using Fatou's lemma [Coh13, Theorem 2.4.4] and recalling that $c(t) = \tilde{i}_p^{-1}(\tilde{i}_p(c))(t) = (e_{t,p})_*(\tilde{i}_p(c))$ in $L_h^p(M, N)$ for all $t \in I$, we have for all $(s, t) \in I$ such that $s \leq t$

$$\begin{aligned} \int_s^t \int_M \tilde{i}_p(c)(x)'|_N^p(u) d\mu_M(x) du &= \int_s^t \int_M \liminf_{\tau \rightarrow 0} (\Delta_{N,\tau} \tilde{i}_p(c)(x)(u))^p d\mu_M(x) du \\ &\leq \int_s^t \liminf_{\tau \rightarrow 0} \int_M (\Delta_{N,\tau} \tilde{i}_p(c)(x)(u))^p d\mu_M(x) du \\ &= \int_s^t \liminf_{\tau \rightarrow 0} \int_M (\Delta_{N,\tau} c(u)(x))^p d\mu_M(x) du \\ &= \int_s^t |c'|_p^p(u) du. \end{aligned}$$

Therefore, using Lebesgue's differentiation theorem [DiB02, Theorem 11.1], we get that

$$\int_M \tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) \leq |c'|_p^p(t) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I.$$

Together with Equation (4), this yields the equality

$$|c'|_p^p(t) = \int_M \tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I,$$

so that Equation (3) is proved.

Step 5: Characterization of the range of \tilde{i}_p . By the previous step and the Fubini–Tonelli theorem [Coh13, Theorem 2.4.4], we have for all $c \in \mathcal{A}C^p(I, L_h^p(M, N))$ that

$$\int_M \int_I \tilde{i}_p(c)(x)'|_N^p(u) du d\mu_M(x) \leq \int_I |c'|_p^p(u) du < \infty.$$

Hence, the range of \tilde{i}_p is contained in the set of all $f \in L_{\underline{h}}^p(M, \mathcal{A}C^p(I, N))$ such that

$$\int_M \int_I |f(x)'|_N^p(t) dt d\mu_M(x) < \infty.$$

Let us now prove the reverse inclusion. Thus, let $f \in L_{\underline{h}}^p(M, \mathcal{A}C^p(I, N))$ such that

$$\int_M \int_I |f(x)'|_N^p(t) dt d\mu_M(x) < \infty.$$

We already know that $\tilde{i}_p^{-1} \circ (q_p)_*(f) \in L^p(I, L_h^p(M, N))$, by Proposition 2.30 and Theorem 3.13, and, by Theorem 3.9, that

$$\hat{f} := (\overline{\text{sec}}_{I,p}^{-1} \circ \tilde{i}_p^{-1} \circ (q_p)_*)(f) = (\overline{\text{sec}}_{M,p}^{-1} \circ (q_p)_*)(f)$$

belongs to $L_{\underline{h}}^p(I \times M, N)$. Also, by the definitions of $\overline{\text{sec}}_{M,p}$, $\overline{\text{sec}}_{I,p}$ and q_p (Theorem 3.9 and Proposition 2.30), we have $\hat{f}(\cdot, x) = f(x)$ in $L^p(I, N)$ for μ_M -a.e. $x \in M$ and $\hat{f}(t, \cdot) = (\tilde{i}_p^{-1} \circ (q_p)_*)(f)(t)$ in $L_h^p(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$. However, \hat{f} differs from \underline{h} on a measurable set M_f of σ -finite μ_M -measure, so that \hat{f} differs from \hat{h} on a measurable set contained

in the measurable rectangle $I \times M_f$. Hence, we get, using the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1] and Hölder’s inequality, that

$$\begin{aligned}
\sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{p,\tau}(\tilde{i}_p^{-1} \circ (q_p)_*)(f)(u))^p du &= \sup_{0 < \tau < b-a} \int_a^{b-\tau} \int_{M_f} (\Delta_{N,\tau} \hat{f}(u, x))^p d\mu_M(x) du \\
&= \sup_{0 < \tau < b-a} \int_{M_f} \int_a^{b-\tau} (\Delta_{N,\tau} \hat{f}(u, x))^p du d\mu_M(x) \\
&= \sup_{0 < \tau < b-a} \int_M \int_a^{b-\tau} (\Delta_{N,\tau} f(x)(u))^p du d\mu_M(x) \\
&\leq \int_M \sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{N,\tau} f(x)(u))^p du d\mu_M(x) \\
&\leq \int_M \sup_{0 < \tau < b-a} \int_a^{b-\tau} \tau^{-p} \left(\int_u^{u+\tau} |f(x)'|_N(\theta) d\theta \right)^p du d\mu_M(x) \\
&\leq \int_M \sup_{0 < \tau < b-a} \int_a^{b-\tau} \tau^{-1} \int_u^{u+\tau} |f(x)'|_N^p(\theta) d\theta du d\mu_M(x) \\
&= \int_M \sup_{0 < \tau < b-a} \int_I |f(x)'|_N^p(\theta) \tau^{-1} \left(\int_a^{b-\tau} \mathbb{1}_{\{u \leq \theta \leq u+\tau\}}(u) du \right) d\theta d\mu_M(x) \\
&= \int_M \sup_{0 < \tau < b-a} \int_I |f(x)'|_N^p(\theta) \tau^{-1} \left(\int_a^{b-\tau} \mathbb{1}_{\{\theta-\tau \leq u \leq \theta\}}(u) du \right) d\theta d\mu_M(x),
\end{aligned}$$

where the Fubini–Tonelli theorem [Coh13, Proposition 5.2.1] was used to interchange the order of integration over u and θ . Now, we have the following

$$\int_a^{b-\tau} \mathbb{1}_{\{\theta-\tau \leq u \leq \theta\}}(u) du = \begin{cases} \theta - a, & \text{if } a \leq \theta < a + \tau \\ \tau, & \text{if } a + \tau \leq \theta \leq b - \tau \\ b - \theta, & \text{if } b - \tau < \theta \leq b \end{cases}$$

In any case, we get that $\tau^{-1} \int_a^{b-\tau} \mathbb{1}_{\{\theta-\tau \leq u \leq \theta\}}(u) du \leq 1$ for all $\tau \in (0, b-a)$ and $\theta \in I$, so that we eventually get the following inequality

$$\sup_{0 < \tau < b-a} \int_a^{b-\tau} (\Delta_{p,\tau}(\tilde{i}_p^{-1} \circ (q_p)_*)(f)(u))^p du \leq \int_M \int_I |f(x)'|_N^p(\theta) d\theta d\mu_M(x) < \infty.$$

Thus, $(\tilde{i}_p^{-1} \circ (q_p)_*)(f)$ belongs to $W^{1,p}(I, L_h^p(M, N))$ (Definition 2.25) and we can define $c := (Q_p^{-1} \circ \tilde{i}_p^{-1} \circ (q_p)_*)(f)$, which belongs to $\mathcal{AC}^p(I, L_h^p(M, N))$ and satisfies $f = \tilde{i}_p(c)$. Hence, f belongs to the range of \tilde{i}_p . We thus proved that the range of \tilde{i}_p is the set of all $f \in L_h^p(M, \mathcal{AC}^p(I, N))$ such that

$$\int_M \int_I |f(x)'|_N^p(t) dt d\mu_M(x) < \infty. \quad \square$$

In the next section, we show that, when $p = 1$, the appropriate class of curves is the set of càdlàg curves of bounded variation.

3.3. Characterization of càdlàg curves of bounded variation for $p = 1$

We first clarify some recurrent notation used in this section and thereafter.

Notation 3.22. Let $h \in L^0(M, N)$. From thereon, it is assumed that:

- (i) $L_h^1(M, \mathcal{BV}(I, N))$ denotes the set of equivalence classes of separably valued measurable mapping from M to $(\mathcal{BV}(I, N), d_1)$ at a finite $D_{1,1}$ -distance from \underline{h} (see Notation 3.1 for the definition of \underline{h}).

- (ii) $\text{Var}_1 := \text{Var}_{L_h^1(M,N)}$ denotes the variation of curves in $L_h^1(M,N)$ (Definition 2.34).
- (iii) $|Dc|_1 := |Dc|_{L_h^1(M,N)}$ denotes the variation measure of the curve $c \in \mathcal{BV}(I, L_h^1(M,N))$ (Definition 2.38)
- (iv) Q_1 denotes the restriction to $\mathcal{BV}(I, L_h^1(M,N))$ of the quotient mapping Q (see Notation 3.18 for the definition of Q).

Then, the isometry given by Theorem 3.13 yields a characterization of càdlàg curves of bounded variation in nonlinear Lebesgue spaces for $p = 1$.

Theorem 3.23 (Characterization of càdlàg curves of bounded variation for $p = 1$). *Let $h \in L^0(M,N)$. Suppose that N is complete. Then, the isometry \tilde{i}_1 defined in Theorem 3.13 induces the isometric mapping*

$$\tilde{i}_1 := (q_1^{-1})_* \circ \bar{i}_1 \circ Q_1 : (\mathcal{BV}(I, L_h^1(M,N)), d_{1,1}) \rightarrow (L_h^1(M, \mathcal{BV}(I,N)), D_{1,1})$$

whose inverse mapping on its range is $\tilde{i}_1^{-1} := Q_1^{-1} \circ \bar{i}_1^{-1} \circ (q_1)_*$ and satisfies

$$\tilde{i}_1^{-1}(f)(t) = (e_{t,1})_*(f) \quad \text{for all } t \in I \text{ and } f \in \tilde{i}_1(\mathcal{BV}(I, L_h^1(M,N))). \quad (5)$$

In addition, \tilde{i}_1 satisfies for all $c \in \mathcal{BV}(I, L_h^1(M,N))$ that

$$|Dc|_1 = \int_M |D\tilde{i}_1(c)(x)|_N d\mu_M(x) \quad (6)$$

and its range is the set of all $f \in L_h^1(M, \mathcal{BV}(I,N))$ such that

$$\int_M |D\tilde{i}_1(c)(x)|_N(I) d\mu_M(x) < \infty.$$

Remark 3.24 (On the definition of pushforwards for equivalence classes). The pushforward of an equivalence class by a mapping is meant as the equivalence class of the compositions of the given mapping with a compatible representative of the input equivalence class.

Remark 3.25 (On relaxing the finiteness assumption on d_N). Using the same arguments as Remark 3.21, the conclusion of Theorem 3.23 holds when d_N is not assumed finite (see Assumption 2.1 for the assumption d_N), but requires h being constant in that case.

Proof. The proof follows the same approach used in [ALS24, Theorems 3.1 and 3.3] for Wasserstein spaces, which itself follows the approach of [Lis07, Theorems 4 and 5]. It consists in using piecewise constant approximations of càdlàg curves of bounded variation to prove inequalities and propagate the results to càdlàg curves of bounded variation by passing to the limit.

Let $h \in \mathcal{L}^0(M,N)$ and let $c \in \mathcal{BV}(I, L_h^1(M,N))$.

Step 1: Reduction to the case where N is separable. Let us first reduce the proof to the case where N is separable. Indeed, when $c \in \mathcal{BV}(I, L_h^1(M,N))$, the set $N_0 := \overline{\cup_{t \in \mathbb{Q} \cap I} c(t)(M)}$ is separable as the closure of the countable union of separable sets. Then, since c is right-continuous on $[a,b]$ and left-continuous at $t = b$, we have for all $t \in I$ that $c(t) = \lim_{s \rightarrow t^+, s \in \mathbb{Q} \cap I} c(s)$ and $c(b) = \lim_{s \rightarrow b^-, s \in \mathbb{Q} \cap I} c(s)$ in $L_h^p(M,N)$, hence, up to the extraction of a subsequence $c(t)(x) = \lim_{s \rightarrow t^+, s \in \mathbb{Q} \cap I} c(s)(x)$ and $c(b)(x) = \lim_{s \rightarrow b^-, s \in \mathbb{Q} \cap I} c(s)(x)$ in N for μ_M -a.e. $x \in M$, that is, $c(t)(x) \in N_0$ and $c(b)(x) \in N_0$ for μ_M -a.e. $x \in M$. Thus, we can replace N with N_0 in the rest of the proof, so that $\mathcal{L}^0(M,N) = \mathcal{L}_s^0(M,N)$.

Step 2: Definition of \tilde{i}_1 . Let $c \in \mathcal{BV}(I, L_h^1(M,N))$. Then, $Q_1(c)$ belongs to $BV(I, L_h^1(M,N))$, by Proposition 2.44, hence $(\bar{i}_1 \circ Q_1)(c)$ belongs to $L_h^p(M, L^p(I,N))$, by Theorem 3.13. Define for all $n \in \mathbb{N}^*$ the set $I_n := \{0, 1, \dots, 2^n\}$ and the sequence $(t_{n,i})_{i \in I_n}$ such that $t_{n,i} := a + \mathcal{L}^1(I) i 2^{-n}$. Also, define the simple mapping $c_n : I \rightarrow L_h^1(M,N)$ as $c_n|_{[t_{n,i}, t_{n,i+1})} \equiv c(t_{n,i})$ for all $i \in I_{n-1}$ and $c_n(b) := c(b)$. Since the set of dyadic rationals (scaled by $\mathcal{L}^1(I)$ and translated by a) is dense in I and c is right-continuous on $[a,b]$ and $\mathcal{L}^1(\{b\}) = 0$, we have for all \mathcal{L}^1 -a.e. $t \in I$ that $c_n(t) \rightarrow c(t)$ in $L_h^1(M,N)$ as $n \rightarrow \infty$. In addition, if $t \in [t_{n,i}, t_{n,i+1})$,

$$D_1(c_n(t), c(t)) = D_1(c(t_{n,i}), c(t)) \leq \text{Var}_1(c; I).$$

Hence, we get that $D_p(c_n(t), c(t))^p \leq \text{Var}_1(c; I)$ for all $(t, n) \in I \times \mathbb{N}^*$. Since I has finite \mathcal{L}^1 -measure, the mapping constant equal to $\text{Var}_1(c; I) < \infty$ on I is integrable, so that, by Lebesgue's dominated convergence theorem [Coh13, Theorem 2.4.5], we have $d_{1,1}(c_n, c) \rightarrow 0$ as $n \rightarrow \infty$. In addition, for all $n \in \mathbb{N}^*$ and all $i \in I_n$ there is $c_{n,i} \in \mathcal{L}_h^1(M, N)$ such that $D_1(c_{n,i}, c(t_{n,i})) = 0$, so we can define $c'_n : I \rightarrow \mathcal{L}_h^1(M, N)$ as $c'_n|_{[t_{n,i}, t_{n,i+1})} \equiv c_{n,i}$ for all $i \in I_{n-1}$ and $c'_n(b) := c_{n,2^n}$, which satisfies $D_1(c_n(t), c'_n(t)) = 0$ for all $t \in I$. Thus, we can define $\hat{c}'_n : I \times M \rightarrow N$ as $\hat{c}'_n(t, x) := c'_n(t)(x)$, which belongs to $\mathcal{L}_h^1(I \times M, N)$ (simple on I and by definition of the $c_{n,i}$'s) and satisfies $D_1(\hat{c}'_n(t, \cdot), c_n(t)) = 0$ for all $t \in I$. Hence, $\overline{\text{sec}}_{I,1}$ being an isometry (Theorem 3.9), we get that $\hat{D}_1(\hat{c}'_n, \overline{\text{sec}}_{I,1}^{-1}(c_n)) = 0$. Now, defining the mapping $f'_n : M \rightarrow \mathcal{L}^1(I, N)$ as $f'_n(x) := \hat{c}'_n(\cdot, x)$ and recalling that $\overline{\text{sec}}_{M,1}$ is an isometry (Theorem 3.9), we get that $D_{1,1}(f'_n, \bar{i}_1(c_n)) = 0$, hence that $d_1(f'_n(x), \bar{i}_1(c_n)(x)) = 0$ for μ_M -a.e. $x \in M$. Then, let $(s, t) \in I^2$ such that $\mathcal{L}^1(I)2^{-n} < t-s$ and define $k_s := \max\{k \in \mathbb{N} : a + \mathcal{L}^1(I)k \leq s\}$, $k_t := \min\{k \in \mathbb{N} : a + \mathcal{L}^1(I)k \geq t\}$ as well as the mapping

$$\psi_{s,t} : \gamma \mapsto \sup_{0 < \tau < t-s} \int_s^{t-\tau} \Delta_{N,\tau} \gamma(u) \, du.$$

Then, we have

$$\begin{aligned} \int_M \psi_{s,t}(\bar{i}_1(c_n)(x)) \, d\mu_M(x) &= \int_M \psi_{s,t}(f'_n(x)) \, d\mu_M(x) \\ &= \int_M \sup_{0 < \tau < \mathcal{L}^1(I)2^{-n}} \int_s^{t-\tau} \Delta_{N,\tau} f'_n(x)(u) \, du \, d\mu_M(x) \\ &= \int_M \sum_{i=k_s}^{k_t-1} d_N(f'_n(x)(t_{n,i}), f'_n(x)(t_{n,i+1})) \, d\mu_M(x) \\ &= \sum_{i=k_s}^{k_t-1} \int_M d_N(c'_n(t_{n,i})(x), c'_n(t_{n,i+1})(x)) \, d\mu_M(x) \\ &= \sum_{i=k_s}^{k_t-1} D_1(c_n(t_{n,i}), c_n(t_{n,i+1})) \\ &= \sum_{i=k_s}^{k_t-1} D_1(c(t_{n,i}), c(t_{n,i+1})) \\ &\leq \text{Var}_1(c; (s, t)) = |Dc|_1((s, t)) < \infty, \end{aligned}$$

where the second equality follows from [ALS24, Lemma 2.18] and the last equality follows from assertion (i) of Proposition 2.44. When $\mathcal{L}^1(I)2^{-n} \geq t-s$, there is nothing to prove since

$$\int_M \psi_{s,t}(\bar{i}_1(c_n)(x)) \, d\mu_M(x) = \int_M \psi_{s,t}(f'_n(x)) \, d\mu_M(x) = 0 \leq |Dc|_1((s, t)).$$

The mapping \bar{i}_1 being isometric, we also have $D_{1,1}(\bar{i}_1(c_n), (\bar{i}_1 \circ Q_1)(c)) \rightarrow 0$ as $n \rightarrow \infty$. Hence, up to the extraction of a subsequence, we can assume for μ_M -a.e. $x \in M$ that $\bar{i}_1(c_n)(x) \rightarrow (\bar{i}_1 \circ Q_1)(c)(x)$ in $L^1(I, N)$ as $n \rightarrow \infty$. Therefore, recalling that $\psi_{s,t}$ is lower-semicontinuous on $L^1(I, N)$ and using Fatou's lemma [Coh13, Theorem 2.4.4], the latter inequality yields that

$$\begin{aligned} \int_M \psi_{s,t}((\bar{i}_1 \circ Q_1)(c)(x)) \, d\mu_M(x) &\leq \int_M \liminf_{n \rightarrow \infty} \psi_{s,t}(\bar{i}_1(c_n)(x)) \, d\mu_M(x) \\ &\leq \liminf_{n \rightarrow \infty} \int_M \psi_{s,t}(\bar{i}_1(c_n)(x)) \, d\mu_M(x) \\ &\leq |Dc|_1((s, t)) < \infty. \end{aligned}$$

This yields that for μ_M -a.e. $x \in M$

$$\psi_{a,b}((\bar{i}_1 \circ Q_1)(c)(x)) = \sup_{0 < \tau < b-a} \int_a^{b-\tau} \Delta_{N,\tau}(\bar{i}_1 \circ Q_1)(c)(x)(t) \, dt < \infty,$$

which implies that $(\tilde{i}_1 \circ Q_1)(c)(x)$ belongs to $BV(I, N)$ ([Proposition 2.39](#)). Hence, $BV(I, X)$ being a Borel subset of $L^1(I, X)$ ([Remark 2.37](#)), $(\tilde{i}_1 \circ Q_1)(c)$ belongs to $L^1_{\tilde{h}}(M, BV(I, N))$. Therefore, the isometric mapping

$$\tilde{i}_1 := (q_1^{-1})_* \circ \tilde{i}_1 \circ Q_1 : \mathcal{BV}(I, L^1_{\tilde{h}}(M, N)) \rightarrow L^1_{\tilde{h}}(M, \mathcal{BV}(I, N))$$

is well-defined and its inverse mapping on its range is $\tilde{i}_1^{-1} := Q_1^{-1} \circ \tilde{i}_1^{-1} \circ (q_1)_*$. In addition, the last inequality yields that

$$\int_M |D\tilde{i}_1(c)|_N d\mu_M(x) \leq |Dc|_1. \quad (7)$$

Step 3: Characterization of the range of \tilde{i}_1 . By [Equation \(7\)](#), we know that the range of \tilde{i}_1 is contained in the set of all $f \in L^1_{\tilde{h}}(M, \mathcal{BV}(I, N))$ such that

$$\int_M |Df(x)|_N(I) d\mu_M(x) < \infty.$$

Let us now prove that the reverse inclusion. Thus, let $f \in L^1_{\tilde{h}}(M, \mathcal{BV}(I, N))$ such that

$$\int_M |Df(x)|_N(I) d\mu_M(x) < \infty.$$

We already know that $\tilde{i}_1^{-1} \circ (q_1)_*(f) \in L^1(I, L^1_{\tilde{h}}(M, N))$ and, by [Theorem 3.9](#), we know that

$$\hat{f} := (\overline{\text{sec}}_{I,1}^{-1} \circ \tilde{i}_1^{-1} \circ (q_1)_*)(f) = (\overline{\text{sec}}_{M,1}^{-1} \circ (q_1)_*)(f)$$

belongs $L^p_{\tilde{h}}(I \times M, N)$. Also, by the definitions of $\overline{\text{sec}}_{M,1}$, $\overline{\text{sec}}_{I,1}$ and q_1 ([Theorem 3.9](#) and [Proposition 2.44](#)), it satisfies $\hat{f}(\cdot, x) = f(x)$ in $L^1(I, N)$ for μ_M -a.e. $x \in M$ and $\hat{f}(t, \cdot) = (\tilde{i}_1^{-1} \circ (q_1)_*)(f)(t)$ in $L^1_{\tilde{h}}(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$. However, \hat{f} differs from \tilde{h} on a measurable set M_f of σ -finite μ_M -measure, so that \hat{f} differs from \tilde{h} on a measurable set contained in the measurable rectangle $I \times M_f$. Hence, we get, using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1], that for all $(s, t) \in I^2$ such that $s < t$

$$\begin{aligned} \sup_{0 < \tau < t-s} \int_s^{t-\tau} \Delta_{1,\tau}(\tilde{i}_1^{-1} \circ (q_1)_*)(f)(u) du &= \sup_{0 < \tau < t-s} \int_s^{t-\tau} \int_{M_f} \Delta_{N,\tau} \hat{f}(u, x) d\mu_M(x) du \\ &= \sup_{0 < \tau < t-s} \int_{M_f} \int_s^{t-\tau} \Delta_{N,\tau} \hat{f}(u, x) du d\mu_M(x) \\ &\leq \int_M \sup_{0 < \tau < t-s} \int_s^{t-\tau} \Delta_{N,\tau} f(x)(u) du d\mu_M(x) \\ &= \int_M |Df(x)|_N((s, t)) d\mu_M(x) < \infty. \end{aligned}$$

This yields that $(\tilde{i}_1^{-1} \circ (q_1)_*)(f) \in BV(I, L^1_{\tilde{h}}(M, N))$ ([Proposition 2.39](#)) and we can define $c := (Q_1^{-1} \circ \tilde{i}_1^{-1} \circ (q_1)_*)(f)$, which belongs to $\mathcal{BV}(I, L^1_{\tilde{h}}(M, N))$ and satisfies $f = \tilde{i}_1(c)$. Hence, f belongs to the range of \tilde{i}_1 . We thus proved that the range of \tilde{i}_1 is the set of all $f \in L^1_{\tilde{h}}(M, \mathcal{BV}(I, N))$ such that

$$\int_M |D\tilde{i}_1(c)(x)|_N(I) d\mu_M(x) < \infty.$$

In addition, we get, from the previous inequality, that for all $c \in \mathcal{BV}(I, L^1_{\tilde{h}}(M, N))$

$$|Dc|_1 \leq \int_M |D\tilde{i}_1(c)(x)|_N d\mu_M(x). \quad (8)$$

Step 4: Proof of [Equation \(6\)](#). As a consequence of [Equations \(7\)](#) and [\(8\)](#), we get that for all $c \in \mathcal{BV}(I, L^1_{\tilde{h}}(M, N))$

$$|Dc|_1 = \int_M |D\tilde{i}_1(c)(x)|_N d\mu_M(x).$$

Step 5: Proof of Equation (5). Let f be in the range of \tilde{i}_1 . Then, let $c \in \mathcal{BV}(I, L_h^1(M, N))$ be the unique curve such that $f = \tilde{i}_1(c)$ and define

$$\hat{f} := (\overline{\text{sec}}_{I,1}^{-1} \circ \tilde{i}_1^{-1} \circ (q_1)_*)(f) = (\overline{\text{sec}}_{M,1}^{-1} \circ (q_1)_*)(f).$$

Also, note that $\hat{f} = \overline{\text{sec}}_{I,1}^{-1} \circ Q_1(c)$ since $\tilde{i}_1 = (q_1^{-1})_* \circ \tilde{i}_1 \circ Q_1$. Now, there is $f' \in \mathcal{L}_h^1(M, \mathcal{BV}(I, N))$ such that $D_{1,1}(f', f) = 0$, which implies that $f'(x) = f(x)$ in $\mathcal{D}(I, N)$ for μ_M -a.e. $x \in M$ (Proposition 2.44). Thus, we can define the mapping $\tilde{f} : I \times M \rightarrow N$ as $\tilde{f}(t, x) := f'(x)(t) = (e_{t,1} \circ f')(x)$. To show that \tilde{f} is measurable, consider for all $n \in \mathbb{N}^*$ the mapping $\tilde{f}_n : I \times M \rightarrow N$ defined as $\tilde{f}_n(t, x) := (e_{t_{n,i},1} \circ f')(x)$ if $t \in [t_{n,i}, t_{n,i+1})$ with $i \in I_{n-1}$ and $\tilde{f}_n(b, x) := (e_{b,1} \circ f')(x)$, so that \tilde{f}_n belongs to $\mathcal{L}^0(I \times M, N)$ since $e_{t,1} \circ f' \in \mathcal{L}^0(M, N)$ for all $t \in I$ (Proposition 2.45). Then, the set of dyadic rationals (scaled by $\mathcal{L}^1(I)$ and translated by a) being dense in I and $f'(x)$ being right-continuous on $[a, b)$ for all $x \in M$, we have $\tilde{f}(t, x) = \lim_{n \rightarrow \infty} \tilde{f}_n(t, x)$ for all $(t, x) \in I \times M$. Therefore, \tilde{f} is measurable as the pointwise limit of measurable mappings (Proposition 2.4). In addition, $D_1(\tilde{f}(t, \cdot), (e_{t,1})_*(f)) = 0$ for all $t \in I$ since $\tilde{f}(t, x) = f(x)$ in $\mathcal{D}(I, N)$ for μ_M -a.e. $x \in M$. Then, using the definitions of \hat{f} , $\overline{\text{sec}}_{M,1}$ (Theorem 3.9) and q_1 (Proposition 2.44), we have $d_1(\tilde{f}(\cdot, x), \hat{f}(\cdot, x)) = 0$ for μ_M -a.e. $x \in M$. Hence, $\overline{\text{sec}}_{M,1}$ being an isometry, we get $\hat{D}_1(\tilde{f}, \hat{f}) = 0$. Thus, using the definitions of \hat{f} , $\overline{\text{sec}}_{I,1}$ (Theorem 3.9) and Q_1 (Notation 3.22), we get $D_1(\tilde{f}(t, \cdot), c(t)) = 0$ for \mathcal{L}^1 -a.e. $t \in I$, which yields $\tilde{i}_1^{-1}(f)(t) = (e_{t,1})_*(f)$ in $L_h^1(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$. Together with the equality of Equation (6), this implies that the curve $t \mapsto (e_{t,1})_*(f)$ belongs to $\mathcal{BV}(I, L_h^1(M, N))$ and since $d_{1,1}$ separates $\mathcal{BV}(I, L_h^1(M, N))$ (Proposition 2.44), $\tilde{i}_1^{-1}(f)(t) = (e_{t,1})_*(f)$ in $L_h^1(M, N)$ for all $t \in I$, which ends the proof of Equation (5). \square

To better understand why Theorem 3.19 fails when $p = 1$, we provide a simple counter-example.

Example 3.26 (Counter-example to Theorem 3.19 when $p = 1$). Let $I = M = (0, 1)$ with $\Sigma_I = \Sigma_M = \mathcal{B}((0, 1))$ and $\mu_I = \mu_M = \mathcal{L}^1|_{(0,1)}$, let $N = \mathbb{R}$ with the standard topology, and let $h \equiv 0$. Then, define $f \equiv 0$ and $f' \equiv 1$, which both belong to $L^1((0, 1), \mathbb{R})$. Also, $c : (0, 1) \rightarrow L^1((0, 1), \mathbb{R})$ as $c(t) := \mathbb{1}_{(0,t)}$. Then, for any $(s, t) \in (0, 1)^2$, $D_1(c(s), c(t)) = |\int_s^t dx| = |s - t|D_1(f, f')$, since $D_1(f, f') = 1$, so that $c \in \mathcal{AC}^1((0, 1), L^1((0, 1), \mathbb{R}))$. However, it holds for \mathcal{L}^1 -a.e. $x \in (0, 1)$ that $\tilde{i}_1(c)(x) = \mathbb{1}_{(x,1)}$, which is clearly not a continuous curve.

In the next section, we show that Theorem 3.19 allows for the pointwise characterization of the geometry of nonlinear Lebesgue spaces for $p > 1$ and that, in these spaces, a notion of speed can be defined for absolutely continuous curves despite the lack of differential structure outside the case where the target space N is linear.

4. Applications

4.1. Geometry of nonlinear Lebesgue spaces for $p > 1$

4.1.1. Characterization of geodesics

Let us first recall the definition of *constant speed geodesics*.

Definition 4.1 (Geodesics). Let X be a metric space with metric $d_X : X^2 \rightarrow [0, \infty]$. A curve $c : I \rightarrow X$ is called a *constant speed geodesic* if and only if it satisfies $d_X(c(a), c(b)) < \infty$ and for all $(s, t) \in I^2$

$$d_X(c(s), c(t)) = \left| \frac{t-s}{b-a} \right| d_X(c(a), c(b)).$$

The set of all such curves is denoted by $\mathcal{G}(I, X)$ and is a subset of $\mathcal{AC}^p(I, N)$ for all $p \in [1, \infty]$.

Then, Theorem 3.19 yields a characterization of constant speed geodesics in nonlinear Lebesgue spaces for $p > 1$.

Theorem 4.2 (Characterization of geodesics for $p > 1$). Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that N is complete. Then, the isometric mapping \tilde{i}_p defined in Theorem 3.19 satisfies

$$\tilde{i}_p(\mathcal{G}(I, L_h^p(M, N))) \subset L_h^p(M, \mathcal{G}(I, N)).$$

Remark 4.3 (On relaxing the finiteness assumption on d_N). As highlighted in Remark 3.21, Theorem 3.19 holds when d_N is not assumed finite, but h is assumed constant. In that case, Theorem 4.2 also holds.

Proof. Let $h \in \mathcal{L}_s^0(M, N)$ and $p \in (1, \infty)$. Let $c \in \mathcal{G}(I, L_h^p(M, N))$. By [Theorem 3.19](#), $\tilde{i}_p(c)$ belongs to $L_h^p(M, \mathcal{A}C^p(I, N))$ since $\mathcal{G}(I, L_h^p(M, N))$ is a subset of $\mathcal{A}C^p(I, L_h^p(M, N))$. Then, defining the measurable set

$$M_c := \{x \in M : d_p(\tilde{i}_p(c)(x), \underline{h}(x)) > 0\} = \{x \in M : d_\infty(\tilde{i}_p(c)(x), \underline{h}(x)) > 0\}$$

of σ -finite μ_M -measure, we have that $|\tilde{i}_p(c)(x)'|_N(t) = 0$ for all $(x, t) \in (M \setminus M_c) \times I$ since $\underline{h}(x)$ is constant over I . Therefore, using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1], we get that for all $(s, t) \in I^2$ such that $s < t$

$$\begin{aligned} \int_M d_N(\tilde{i}_p(c)(x)(s), \tilde{i}_p(c)(x)(t))^p d\mu_M(x) &= D_p(c(s), c(t))^p \\ &= |t - s|^{p-1} \int_s^t |c'|_p^p(u) du \\ &= |t - s|^{p-1} \int_s^t \int_M |\tilde{i}_p(c)(x)'|_N^p(u) d\mu_M(x) du \\ &= |t - s|^{p-1} \int_s^t \int_{M_c} |\tilde{i}_p(c)(x)'|_N^p(u) d\mu_M(x) du \\ &= |t - s|^{p-1} \int_{M_c} \int_s^t |\tilde{i}_p(c)(x)'|_N^p(u) du d\mu_M(x) \\ &= |t - s|^{p-1} \int_M \int_s^t |\tilde{i}_p(c)(x)'|_N^p(u) du d\mu_M(x). \end{aligned}$$

Therefore, together with the triangle inequality and Hölder's inequality, this yields that for μ_M -a.e. $x \in M$

$$d_N(\tilde{i}_p(c)(x)(a), \tilde{i}_p(c)(x)(b)) = \int_a^b |\tilde{i}_p(c)(x)'|_N(t) dt = |b - a|^{p-1} \int_a^b |\tilde{i}_p(c)(x)'|_N^p(t) dt < \infty.$$

Thus, for \mathcal{L}^1 -a.e. $t \in I$

$$|\tilde{i}_p(c)(x)'|(t) = \frac{d_N(\tilde{i}_p(c)(x)(a), \tilde{i}_p(c)(x)(b))}{|b - a|}.$$

Hence, for μ_M -a.e. $x \in M$ we have that for all $(s, t) \in (\mathbb{Q} \cap I)^2$ such that $s < t$

$$d_N(\tilde{i}_p(c)(x)(s), \tilde{i}_p(c)(x)(t)) = \left| \frac{t - s}{b - a} \right| d_N(\tilde{i}_p(c)(x)(a), \tilde{i}_p(c)(x)(b)).$$

By uniform continuity of $\tilde{i}_p(c)(x)$ and density of $\mathbb{Q} \cap I$ in I , this holds for all $(s, t) \in I^2$ such that $s < t$, hence $\tilde{i}_p(c)(x) \in \mathcal{G}(I, N)$ for μ_M -a.e. $x \in M$ and $\tilde{i}_p(c) \in L_h^p(M, \mathcal{G}(I, N))$. \square

In the next two sections, we show that [Theorem 4.2](#) allows for the pointwise characterization of the geometry of nonlinear Lebesgue spaces for $p > 1$.

4.1.2. Characterization of the length structure

Before delving into the length structure of nonlinear Lebesgue spaces, let us point out that a reminder on length spaces is provided in [Appendix Appendix B.1](#). Now, for $p \in [1, \infty]$, denote by L_p the length induced by the metric D_p on $L_h^p(M, N)$ ([Definition B.1](#)). Thus, the intrinsic metric induced by D_p ([Definition B.2](#)) is defined, for all $(f, f') \in L_h^p(M, N)^2$, as

$$\widehat{D}_p(f, f') := \inf \{L_p(c) : c \in \mathcal{A}C(I, L_h^p(M, N)), c(a) = f, c(b) = f'\}.$$

A first notable result on the length structure (see [Definition B.3](#) for a definition of length spaces) of nonlinear Lebesgue spaces is that it is induced by the length structure of the target space.

Proposition 4.4 (Length structure of nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Suppose that $L_h^p(M, N)$ is trivial ([Proposition 2.16](#)) or that N is a complete and separable length (resp. geodesic) space. Then, $L_h^p(M, N)$ is a length (resp. geodesic) space.*

Proof. The proof follows the same approach used in [Lis07, Proposition 1] for Wasserstein spaces, which essentially consists in constructing a curve whose length is controlled by the distance between its endpoints from the measurable selection of pointwise curves satisfying such a constraint.

Let $p \in [1, \infty]$, $h \in \mathcal{L}_s^0(M, N)$, $(f, f') \in \mathcal{L}_h^p(M, N)^2$ and $\kappa > 1$. In addition, let $h^2(x) := (h(x), h(x))$ and define the mapping $\pi : M \rightarrow N^2$ as $\pi(x) := (f(x), f'(x))$ which belongs to $\mathcal{L}_{h^2}^p(M, N^2)$ since $(f, f') \in \mathcal{L}_h^p(M, N)^2$. When $L_h^p(M, N)$ is trivial, that is, $L_h^p(M, N) = \{[h]\}$, it is also a geodesic space as it contains a single element. Therefore, assume that $L_h^p(M, N)$ is nontrivial and that N is a complete and separable length space.

Step 1: Reduction to the case where μ_M is σ -finite. The mappings f and f' both differ from h on a measurable set of σ -finite μ_M -measure [ST25, Proposition 3.4]. Hence, f and f' differ on a set of σ -finite μ_M -measure, so that we can assume that μ_M is σ -finite in the rest of the proof. Now, since μ_M is assumed σ -finite, there exists a finite measure which has the same null sets, for instance, $\nu_M : A \mapsto \sum_{n \in \mathbb{N}} 2^{-n} \frac{\mu_M(A)}{\mu_M(B_n)+1}$ where $(B_n)_{n \in \mathbb{N}}$ is a sequence of measurable sets of finite μ_M -measure such that $M = \cup_{n \in \mathbb{N}} B_n$.

Step 2: Construction of a Borel measurable selection of curves connecting pairs of points in N . Then, define the multi-valued application $\Gamma_\kappa : N^2 \rightarrow 2^{C(I, N)}$ as follows

$$\Gamma_\kappa(y, y') := \{\gamma \in C(I, N) : \gamma(a) = y, \gamma(b) = y', \text{Lip}(\gamma)|b - a| \leq \kappa d_M(y, y')\}.$$

Since N is a length space (Definition B.3), $\Gamma_\kappa(y, y')$ is non-empty for each $(y, y') \in N^2$. Now, using the lower semi-continuity of the Lipschitz constant, we can show that the graph of Γ_κ defined as

$$G(\Gamma_\kappa) := \{(y, y', \gamma) \in N^2 \times C(I, N) : \gamma \in \Gamma_\kappa(y, y')\}$$

is closed. Indeed, taking a sequence $((y_n, y'_n, \gamma_n))_{n \in \mathbb{N}}$ in $G(\Gamma_\kappa)$ that converges to $(y, y', \gamma) \in G(\Gamma_\kappa)$, we have that $d_N(y_n, y) \rightarrow 0$, $d_N(y'_n, y') \rightarrow 0$ and $d_\infty(\gamma_n, \gamma)$ as $n \rightarrow \infty$. Hence, by uniqueness of the limit, $\gamma(a) = y$ and $\gamma(b) = y'$ and, by lower semi-continuity of the Lipschitz constant [BBI22, Proposition 2.3.4. (iv)], we can pass to the limit in $\text{Lip}(\gamma_n)|b - a| \leq d_N(y_n, y'_n)$, so that $\text{Lip}(\gamma)|b - a| \leq d_N(y, y')$, which yields that $\gamma \in \Gamma_\kappa(y, y')$ and thus that $(y, y', \gamma) \in G(\Gamma_\kappa)$. Since $(C(I, N), d_\infty)$ is complete and separable and $G(\Gamma_\kappa)$ is closed, we have, by Aumann's measurable selection theorem (Theorem B.9), that there exists a $\pi_* \nu_M$ -measurable mapping (see Remark 2.14 for a definition) $S_\kappa : N \times N \rightarrow C(I, N)$ such that $S_\kappa(y, y') \in \Gamma_\kappa(y, y')$ for all $(y, y') \in N^2$. Since, $C(I, N)$ is separable and $\pi_* \mu_M$ has the same null sets as $\pi_* \nu_M$, there exists a Borel measurable mapping \tilde{S}_κ such that $\tilde{S}_\kappa(y, y') = S_\kappa(y, y')$ for $\pi_* \mu_M$ -a.e. $(y, y') \in N^2$ [ST25, Proposition 2.12].

Step 3: Construction of a curve in $L_h^p(M, N)$ from the Borel measurable selection. Define $c(t) := e_t \circ \tilde{S}_\kappa \circ \pi$ for all $t \in I$. Since we have $(\pi, \tilde{S}_\kappa, e_t) \in \mathcal{L}^0(M, N^2) \times \mathcal{L}^0(N^2, C(I, N)) \times \mathcal{L}^0(C(I, N), N)$ with the Borel σ -algebras on N^2 and $C(I, N)$, we get, by composition, that $c(t) \in \mathcal{L}^0(M, N)$ for all $t \in I$. Now, recalling that $\tilde{S}_\kappa(f(x), f'(x)) \in \Gamma_\kappa(f(x), f'(x))$ for μ_M -a.e. $x \in M$, we have, for all $(s, t) \in I^2$, that

$$\begin{aligned} D_p(c(s), c(t))^p &= \int_M d_N(\tilde{S}_\kappa(f(x), f'(x))(s), \tilde{S}_\kappa(f(x), f'(x))(t))^p d\mu_M(x) \\ &\leq \int_M \text{Lip}(\tilde{S}_\kappa(f(x), f'(x)))^p |s - t|^p d\mu_M(x) \\ &\leq \int_M \kappa^p d_N(f(x), f'(x))^p \left| \frac{s - t}{b - a} \right|^p d\mu_M(x) \\ &= \kappa^p D_p(f, f')^p \left| \frac{s - t}{b - a} \right|^p. \end{aligned}$$

Since $c(a) = f \in \mathcal{L}_h^p(M, N)$, this yields that $c(t) \in \mathcal{L}_h^p(M, N)$ for all $t \in I$ and that, up to the identification with an equivalence class for each $t \in I$, c determines a curve in $\mathcal{A}C(I, L_h^p(M, N))$ such that $L_p(c) \leq \kappa D_p(f, f')$. Since we can construct such a curve for an arbitrary $\kappa > 1$, this yields that $\widehat{D}_p(f, f') = D_p(f, f')$, that is, $L_h^p(M, N)$ is a length space. When N is a geodesic space, the proof works with $\kappa = 1$. \square

Remark 4.5 (On the separability assumption on N). Note that the proof of Proposition 4.4 only requires the completeness and separability assumptions on N to ensure that the pointwise selection of curves is measurable. When N is uniquely geodesic and d_N is jointly convex, for instance, when N is a global NPC space (Definition 4.10), N need

not be separable for the conclusion of [Proposition 4.4](#) to hold since there is no measurable selection of geodesics to construct as any pair of points in N is joined by a unique geodesic and geodesics in N are continuous with respect to their endpoints thanks to the joint convexity of d_N . The question of whether the assumptions of completeness and separability on N remains open.

Remark 4.6 (Related results in the literature). We emphasize that while the conclusion of [Proposition 4.4](#) in the geodesic case already appeared in a work (under the assumption that h is constant) by J. Jöst [[Jos97b](#), Lemma 4.1.2], the proof seemed incomplete as it selected pointwise geodesics without checking the measurability with respect to endpoints, which is precisely what is done in the proof of [Proposition 4.4](#), hence missing the key assumptions of completeness and separability on N . As highlighted in [Remark 4.5](#), the question of whether the completeness and separability assumptions on N could be relaxed remains open.

The reverse implication holds when restricting to $p \in (1, \infty)$ and a measure μ_M that is not purely infinite, but does not require N being separable.

Proposition 4.7 (Length structure of the target space). *Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that μ_M is not purely infinite and that N is complete. Then, if $L_h^p(M, N)$ is a length (resp. geodesic) space, so is N .*

Remark 4.8 (Why it fails when μ_M is purely infinite). When $|N| = 1$, $L_h^p(M, N)$ is trivial ([Proposition 2.16](#)), hence a geodesic space, but so is N (since it contains a single element). Thus, the conclusion of [Proposition 4.7](#) holds in that case. However, when μ_M is purely infinite and $|N| > 1$, we know that $L_h^p(M, N)$ remains trivial ([Proposition 2.16](#)), hence a geodesic space, regardless of the nature of the target space N .

Proof. Let $(y, y') \in N^2$ and suppose there exists $\kappa > 1$ such that for all $\gamma \in \mathcal{AC}^p(I, N)$ such that $\gamma(a) = y$ and $\gamma(b) = y'$ we have

$$\mathcal{L}^1(I)^{p-1} \int_I |\gamma'|_N^p(t) dt > \kappa d_N(y, y')^p.$$

Since the measure μ_M is not purely infinite, we have, by [Proposition 2.19](#), that there exists an isometric mapping $\iota : N \rightarrow L_h^p(M, N)$, so that $\iota(y)$ and $\iota(y')$ belong to $L_h^p(M, N)$. In addition, there exists a measurable set A with $\mu_M(A) \in (0, \infty)$ such that $\iota(z)|_A \equiv z$ and $\iota|_{M \setminus A} = h|_{M \setminus A}$ for all $z \in N$. Now, since $L_h^p(M, N)$ is a length space, there exists $c \in \mathcal{AC}^p(I, L_h^p(M, N))$ such that $c(a) = \iota(y)$, $c(b) = \iota(y')$ and $\mathcal{L}^1(I)^{p-1} \int_I |c'|_p^p(t) dt \leq \kappa D_p(\iota(y), \iota(y'))^p$ ([Proposition B.7](#)). Then, by [Theorem 3.19](#), $\tilde{i}_p(c) \in L_h^p(M, \mathcal{AC}^p(I, N))$ with $\tilde{i}_p(c)(a) = \iota(y)$ and $\tilde{i}_p(c)(b) = \iota(y')$, so that for μ_M -a.e. $x \in A$ the curve $\tilde{i}_p(c)(x) \in \mathcal{AC}^p(I, N)$ is such that $\tilde{i}_p(c)(x)(a) = y$ and $\tilde{i}_p(c)(x)(b) = y'$. Hence, by assumption, we have that for μ_M -a.e. $x \in A$

$$\mathcal{L}^1(I)^{p-1} \int_I |\tilde{i}_p(c)(x)'|_N^p(t) dt > \kappa d_N(\iota(y)(x), \iota(y')(x))^p.$$

Integrating over A and using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1] since A has finite μ_M -measure, we get that

$$\begin{aligned} \mathcal{L}^1(I)^{p-1} \int_I \int_A |\tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x) dt &= \mathcal{L}^1(I)^{p-1} \int_A \int_I |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) \\ &> \kappa \int_A d_N(\iota(y)(x), \iota(y')(x))^p d\mu_M(x). \end{aligned}$$

Thus, recalling that $|c'|_p^p(t) = \int_M |\tilde{i}_p(c)(x)'|_N^p(t) d\mu_M(x)$ and that $\iota(y)|_{M \setminus A} = \iota(y')|_{M \setminus A}$, we have that

$$\mathcal{L}^1(I)^{p-1} \int_I |c'|_p^p(t) dt > \kappa D_p(\iota(y), \iota(y'))^p,$$

hence a contradiction. The conclusion that d_N coincides with the induced intrinsic metric \widehat{d}_N ([Definition B.2](#)), that is, that N is a length space, then follows from the fact that the intrinsic metric can be equivalently defined as the infimum over the p -energy of curves ([Proposition B.7](#)). When N is a geodesic space, the above proof works with $\kappa = 1$. Alternatively, the geodesic space case can be proved using [Theorem 4.2](#) since, if $L_h^p(M, N)$ is a geodesic space, $\iota(y)$ and $\iota(y')$ can be connected by a geodesic c in $L_h^p(M, N)$, so that, by [Theorem 4.2](#), the curve $\tilde{i}_p(c)(x)$ is a geodesic connecting y and y' for μ_M -a.e. $x \in A$. Also, note that the proof does not require the separability assumption on N . \square

As a direct consequence of [Propositions 4.4](#) and [4.7](#), the length structure of nonlinear Lebesgue spaces is characterized by the length structure of the target space.

Theorem 4.9 (Characterization of the length structure). *Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that μ_M is not purely infinite and that N is complete and separable. Then, the following assertions are equivalent:*

- (i) $L_h^p(M, N)$ is a length (resp. geodesic) space.
- (ii) N length (resp. geodesic) space.

In the next section, we prove that the geodesic case of [Theorem 4.9](#) has some implications on the characterization of curvature bounds in nonlinear Lebesgue spaces using curvature bounds in the target space.

4.1.3. Characterization of curvature bounds for $p = 2$

A notable consequence of [Theorems 4.2](#) and [4.9](#) is that the curvature of nonlinear Lebesgue spaces, in the sense of Alexandrov (see [[BBI22](#), Chapter 4] for a definition), depends solely on the Alexandrov curvature of the target space for $p = 2$. Before stating the result, recall that in a complete metric space, the local and global Alexandrov curvatures are equivalent, so that we can evaluate it through triangle comparisons.

Definition 4.10 (Alexandrov curvature). Let X be a complete geodesic space ([Definition B.5](#)). X is said to be of global *nonnegative curvature* (NNC), respectively of global *nonpositive curvature* (NPC), in the sense of Alexandrov if and only if, for any point $z \in X$ and constant speed geodesic $\gamma : I \rightarrow X$ ([Definition 4.1](#)), the inequality (see [[LP10](#)] or [[Stu23](#), Theorem 4.4] for global NNC)

$$d_X(z, \gamma(t))^2 \geq (1-t)d_X(z, \gamma(a))^2 + t d_X(z, \gamma(b))^2 - (1-t)t d_X(\gamma(a), \gamma(b))^2$$

holds for all $t \in I$ and the converse inequality in the NPC case (see [[Stu99](#), Corollary 1.5] for global NPC). A complete geodesic space of global NNC (resp. NPC) in the sense of Alexandrov is thus called a *global NNC (resp. NPC) space*.

Then, the Alexandrov curvature of nonlinear Lebesgue spaces for $p = 2$ is determined by the Alexandrov curvature of the target space.

Proposition 4.11 (Alexandrov curvature of nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$. Then, either $L_h^2(M, N)$ is trivial ([Proposition 2.16](#)), hence is trivially both a global NNC space and a global NPC space, or the following assertions hold:*

- (i) if N is a separable and global NNC space, $L_h^2(M, N)$ is also a global NNC space.
- (ii) if N is a global NPC space, $L_h^2(M, N)$ is also a global NPC space.

Proof. Let $h \in L^0(M, N)$. When $L_h^2(M, N)$ is trivial, that is, $L_h^2(M, N) = \{[h]\}$, it is both a global NNC space and a global NPC space as it contains a single element. Therefore, assume that $L_h^2(M, N)$ is nontrivial.

- (i) *Step 1: $L_h^2(M, N)$ is complete and geodesic.* First note that, N being complete, separable and geodesic, we have, by [Proposition 4.4](#), that $L_h^2(M, N)$ is complete and, by [Proposition 4.4](#), that it is geodesic. The proof then comes down to verifying that every geodesic triangle satisfies the condition of NNC ([Definition 4.10](#)).

Step 2: NNC triangle comparison. Let $c \in \mathcal{G}(I, L_h^2(M, N))$ and $f \in L_h^2(M, N)$. Then, by [Theorem 4.2](#), $\tilde{i}_p(c)(x)$ belongs to $\mathcal{G}(I, N)$ for μ_M -a.e. $x \in M$ and, since N is a global NNC space, we get for μ_M -a.e. $x \in M$ that the inequality

$$d_N(f(x), \tilde{i}_p(c)(x)(t))^2 \geq (1-t)d_N(f(x), \tilde{i}_p(c)(x)(a))^2 + t d_N(f(x), \tilde{i}_p(c)(x)(b))^2 - (1-t)t d_N(\tilde{i}_p(c)(x)(a), \tilde{i}_p(c)(x)(b))^2$$

holds for all $t \in I$. Hence, integrating over M and recalling that $c(t) = (e_{t,2})_*(\tilde{i}_p(c))$ for all $t \in I$ ([Theorem 3.19](#)), we get that the inequality

$$D_2(f, c(t))^2 \geq (1-t)D_2(f, c(a))^2 + t D_2(f, c(b))^2 - (1-t)t D_2(c(a), c(b))^2$$

holds for all $t \in I$. We thus proved that $L_h^2(M, N)$ is a global NNC space.

- (ii) The NPC case follows from the same arguments as (i), but with reverse inequalities. In that case, the proof does not require the use of [Proposition 4.4](#), so that the conclusion holds without assuming that N is separable ([Remark 4.5](#)). \square

Remark 4.12 (Related results in the literature). We emphasize that while assertion (ii) of [Proposition 4.11](#) already appeared in works (under the assumption that h is constant for some) by J. Jöst (see [[Jos94](#), Section 2] and [[Jos97b](#), Corollary 4.1.1]), K.-T. Sturm (see [[Stu01](#), Proposition 3.3.]) and M. Bačák (see [[Bac14](#), Proposition 1.2.18]), their proofs seemed incomplete as they lacked a result stating that *all* geodesics are, up to an identification, pointwise geodesics, that is, mappings from M to the space of geodesics in N , which is precisely the statement of [Theorem 4.2](#). Indeed, all mentioned works construct geodesics in $L_h^2(M, N)$ from pointwise geodesics to prove the NPC inequality in $L_h^2(M, N)$, but this inequality should be satisfied for *every* geodesic, so that it remains to show that every geodesic in $L_h^2(M, N)$ can be identified to a pointwise geodesic, as done in [Theorem 4.2](#).

The reverse implication holds when restricting to a measure μ_M that is not purely infinite, but does not require N being separable.

Proposition 4.13 (Alexandrov curvature of the target space). *Let $h \in L^0(M, N)$. Suppose that μ_M is not purely infinite. Then, if $L_h^2(M, N)$ is a global NNC (resp. NPC) space, so is N .*

Remark 4.14 (Why it fails when μ_M is purely infinite). When $|N| = 1$, $L_h^2(M, N)$ is trivial ([Proposition 2.16](#)), hence both a global NNC and NPC space, but so is N (since it contains a single element). Thus, the conclusion of [Proposition 4.13](#) holds in that case. However, when μ_M is purely infinite and $|N| > 1$, we know that $L_h^2(M, N)$ remains trivial ([Proposition 2.16](#)), hence both a global NNC and NPC space, regardless of the nature of the target space N .

Proof. We only prove the NNC case since the NPC case follows from the same arguments, but with reversed inequalities.

Step 1: N is complete and geodesic. First note that, $L_h^2(M, N)$ being geodesic and μ_M not being purely infinite, we have, by [Proposition 4.7](#), that N is geodesic. In addition, since $L_h^2(M, N)$ is complete and μ_M is not purely infinite, N is complete, by [[ST25](#), Proposition 4.2]. The proof then comes down to verifying that every geodesic triangle satisfies the condition of NNC ([Definition 4.10](#)).

Step 2: NNC triangle comparison. Let $\gamma \in \mathcal{G}(I, N)$ and $z \in N$. By [Proposition 2.19](#), there exists an isometric mapping $\iota : N \rightarrow L_h^2(M, N)$, so that the curve $c : t \rightarrow \iota(\gamma(t))$ belongs to $\mathcal{G}(I, L_h^2(M, N))$ and $f := \iota(z) \in L_h^2(M, N)$. Since, $L_h^2(M, N)$ is NNC, we have for all $t \in I$ that

$$D_2(f, c(t))^2 \geq (1-t)D_2(f, c(a))^2 + tD_2(f, c(b))^2 - (1-t)tD_2(c(a), c(b))^2.$$

Hence, ι being isometric,

$$d_N(z, \gamma(t))^2 \geq (1-t)d_N(z, \gamma(a))^2 + td_N(z, \gamma(b))^2 - (1-t)t d_N(\gamma(a), \gamma(b))^2.$$

We thus proved that N is a global NNC space. \square

As a direct consequence of [Propositions 4.11](#) and [4.13](#), the Alexandrov curvature of nonlinear Lebesgue spaces for $p = 2$ is characterized by the Alexandrov curvature of the target space.

Theorem 4.15 (Characterization of the Alexandrov curvature). *Let $h \in L^0(M, N)$. Suppose that μ_M is not purely infinite and that N is separable. Then, the following assertions are equivalent:*

- (i) $L_h^2(M, N)$ is a global NNC (resp. NPC) space.
- (ii) N is a global NNC (resp. NPC) space.

Remark 4.16 (On the separability assumption on N). Note that, in the NPC case, the conclusion of [Theorem 4.15](#) holds without the separability assumption on N ([Proposition 4.11](#)).

In the next section, we prove that, when the target space has a differential structure satisfying appropriate conditions, a notion of speed can be defined for absolutely continuous curves in nonlinear Lebesgue spaces despite the lack of differential structure.

4.2. Speed of absolutely continuous curves for $p > 1$

4.2.1. Measurable and Lebesgue sections

First, recall the definition of a *section* of a surjective mapping.

Definition 4.17 (Sections). Let E and B be sets and $\pi_E : E \rightarrow B$ be a surjective mapping. Then, a mapping $s : B \rightarrow E$ is called a *section of π_E* if $\pi_E \circ s = \text{id}_B$. The set of all sections of π_E is denoted by $\mathcal{S}(\pi_E)$.

Asking for a metric on E might be a bit restrictive, so we only require it to be a topological space in the following sections and try to exploit its fibered structure. One can then define the set of *measurable sections*.

Definition 4.18 (Measurable sections). Suppose that E is a topological space, that (B, Σ_B) is a measurable space and that $\pi_E : E \rightarrow B$ is a surjective mapping. Then, a mapping $s : B \rightarrow E$ is called a *measurable section of π_E* if it belongs to $\mathcal{S}(\pi_E) \cap \mathcal{L}^0(B, E)$. Then, define:

- (i) $\mathcal{S}_s^0(\pi_E)$ the set of all measurable sections of π_E with separable range.
- (ii) $S^0(\pi_E)$ the set of all equivalence classes of measurable sections of π_E with separable range.

Remark 4.19 (On the definition of measurability). Note that the definition of measurable mappings given in [Definition 2.2](#) holds when the target space is just a topological space, so that $\mathcal{L}^0(B, E)$ makes sense in [Definition 4.18](#) despite E being only a topological space.

Similarly, we can define *Lebesgue sections* given that we only have a *fibered metric* on the topological space E , that is, a Borel measurable mapping $m_E : E^2 \rightarrow [0, \infty)$ such that its restriction $m_x := m_E|_{\pi_E^{-1}(\{x\})^2}$, for some $x \in B$, is a metric on the fiber $\pi_E^{-1}(\{x\})$ which induces the topology of the fiber.

Definition 4.20 (Lebesgue sections). Let $h \in S^0(\pi_E)$ and $p \in [1, \infty]$. Suppose that E is a topological space, that (B, Σ_B, μ_B) is a measure space, that $\pi_E : E \rightarrow B$ is a surjective mapping and that m_{π_E} is a fibered metric on E . Then, a mapping $s : B \rightarrow E$ is called *p -Lebesgue section* if it belongs to $\mathcal{S}_s^0(\pi_E)$ and $D_{p, \pi_E}(s, h) < \infty$ with

$$D_{p, \pi_E}(s, h) := \begin{cases} \left(\int_B m_x(s(x), h(x))^p d\mu_B(x) \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mu_B\text{-ess sup}_{x \in B} m_x(s(x), h(x)), & \text{if } p = \infty \end{cases}.$$

Then, define:

- (i) $\mathcal{S}_h^p(\pi_E)$ the set of all p -Lebesgue sections of π_E with h as base mapping.
- (ii) $S_h^p(\pi_E)$ the set of all equivalence classes of p -Lebesgue sections of π_E with h as base mapping.

Hence, D_{p, π_E} induces a finite metric on $\mathcal{S}_h^p(\pi_E)$. When m_E is a metric on E which induces its topology, $S_h^p(\pi_E)$ is a subset of $L_h^p(B, E)$ ([Definition 2.13](#)) and $D_{p, \pi_E} = D_p$ ([Definition 2.8](#)).

Remark 4.21 (On the choice of h for vector bundles). When $\pi_E : E \rightarrow B$ is a vector bundle, that is, the fibers $(\pi_E^{-1}(\{x\}))_{x \in B}$ are vector spaces, h is usually taken as the zero section (if it is measurable) of π_E , that is, h maps to each $x \in B$ the identity element of $\pi_E^{-1}(\{x\})$. In that case, we denote $\mathcal{S}^p(\pi_E) := \mathcal{S}_h^p(\pi_E)$ and $S^p(\pi_E) := S_h^p(\pi_E)$.

In the next section, we show that the speed of absolutely continuous curves in differentiable manifolds satisfying suitable conditions can be defined as Lebesgue sections.

4.2.2. Speed of absolutely continuous curves in Finsler manifolds

Let us first recall the definition of *Finsler manifolds* introduced by R. S. Palais [[Pal66](#), Definition 2.1] (see [[JSSG11](#), Definition 2.1] for a more recent exposition and a comparison with other existing definitions of Finsler manifolds).

Definition 4.22 (Finsler manifolds). Let $k \in \mathbb{N}^* \cup \{\infty\}$. A C^k manifold X modelled on a banach space $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$ ([Definition B.12](#)) is called *Finsler* when it is equipped with a mapping $b_X : TX \rightarrow [0, \infty)$ such that:

- (i) for every $x \in X$, the restriction $b_x := b_X|_{T_x X} : T_x X \rightarrow [0, \infty)$ is a norm on the tangent space $T_x X$ which induces the manifold topology, that is, for every chart (U, φ) such that $x \in U$, the norm $b_x(d\varphi^{-1}(\varphi(x))(\cdot))$ is equivalent to $\|\cdot\|_{\mathbb{B}}$.

(ii) for every $x_0 \in X$, $\varepsilon > 0$ and chart (U, φ) such that $x_0 \in U$, there is an open neighborhood $V \subset U$ of x_0 such that if $x \in V$ and $v \in \mathbb{B}$

$$\frac{1}{1 + \varepsilon} b_{x_0}(\mathrm{d}\varphi^{-1}(\varphi(x_0))(v)) \leq b_x(\mathrm{d}\varphi^{-1}(\varphi(x))(v)) \leq (1 + \varepsilon) b_{x_0}(\mathrm{d}\varphi^{-1}(\varphi(x_0))(v)).$$

Remark 4.23 (On the continuity of b_X). Note that, when (X, b_X) is a Finsler manifold as defined in [Definition 4.22](#), the mapping $b_X : TX \rightarrow [0, \infty)$ is continuous [[Pal66](#), Theorem 2.7].

A Finsler structure induces a metric space structure [[Pal66](#), Theorem 3.3].

Proposition 4.24 (Metric space structure of Finsler manifolds). *Suppose that (X, b_X) is a C^1 Finsler manifold. Then, X is metric space when equipped with the metric $d_X : X^2 \rightarrow [0, \infty]$ defined as*

$$d_X(x, x') := \inf \left\{ \int_I b_X(\dot{c}(t)) dt : c \in C^1(I, X), c(a) = x, c(b) = x' \right\}.$$

In addition, the induced metric is locally equivalent to the norm [[JJSSG14](#), Lemma 1.3].

Proposition 4.25 (Local equivalence of the norm and the metric). *Let (X, b_X) be a C^1 Finsler manifold. Then, for every $x_0 \in X$, $\varepsilon > 0$ and chart (U, φ) such that $x_0 \in U$, there exists an open neighborhood $V \subset U$ of x_0 such that for all $(x, x') \in V$ we have*

$$\frac{1}{1 + \varepsilon} d_X(x, x') \leq b_{x_0}(\mathrm{d}\varphi^{-1}(\varphi(x_0))(\varphi(x) - \varphi(x'))) \leq (1 + \varepsilon) d_X(x, x').$$

We will see that [Proposition 4.25](#) is useful to show that the norm of the speed and the metric derivative coincide for absolutely continuous curves. To define a notion of speed, let us first recall a definition of absolutely continuous curves in Banach spaces that is stronger than [Definition 2.28](#).

Definition 4.26 (Strongly absolutely continuous curves in Banach spaces). Let $p \in [1, \infty]$. When \mathbb{B} is a Banach space, define the set $\mathcal{AC}_s^p(I, \mathbb{B})$ of *strongly absolutely continuous curves*, that is, the set of all continuous curves $c : I \rightarrow \mathbb{B}$ such that there is $\eta \in L^p(I, \mathbb{B})$ satisfying

$$c(t) - c(s) = \int_s^t \eta(u) du \quad \text{for all } a \leq s \leq t \leq b,$$

where the latter integral is meant in the sense of Bochner (see [[HVN VW16](#), Section 1.2.a] for a complete exposition).

[Definition 4.26](#) extends to Banach manifolds through the use of local charts [[Sch16](#), Definition 1.2].

Definition 4.27 (Strongly absolutely continuous curves in Banach manifolds). Let $p \in [1, \infty]$. When X is a C^1 Banach manifold ([Definition B.12](#)), define the set $\mathcal{AC}_s^p(I, X)$ of all continuous curves $c : I \rightarrow X$ such that there is a partition $a = t_0 < t_1 < \dots < t_n = b$ and charts (U_i, φ_i) of X for $1 \leq i \leq n$ such that $c([t_{i-1}, t_i]) \subset U_i$ and $\varphi_i \circ c|_{[t_{i-1}, t_i]} \in \mathcal{AC}_s^p([t_{i-1}, t_i], \varphi_i(U_i))$.

Remark 4.28 (On the invariance to the choice of partition and charts). As pointed out in [[Glo15](#), Lemma 3.21], once a curve satisfies the conditions of [Definition 4.27](#) for a partition of I and a collection of charts, the conditions are also satisfied for every other partition $a = t_0 < t_1 < \dots < t_n = b$ of I and collection of charts $(U_i, \varphi_i)_{1 \leq i \leq n}$ of X such that $c([t_{i-1}, t_i]) \subset U_i$ for all $1 \leq i \leq n$.

One can then define a Banach bundle of L^p speeds over the set of strongly absolutely continuous curves in Finsler manifolds.

Proposition 4.29 (Vector bundle over $\mathcal{AC}_s^p(I, X)$ in Finsler manifolds). *Let $p \in [1, \infty]$. Suppose that (X, b_X) is a C^1 Finsler manifold. Then, the set $L^p(I, TX) := \cup_{c \in \mathcal{AC}_s^p(I, X)} S^p(\pi_{c^*TX})$ forms a Banach bundle*

$$\pi_{L^p(I, TX)} := (\pi_{TX})_* : L^p(I, TX) \rightarrow \mathcal{AC}_s^p(I, X),$$

where $c^*TX := \{(t, v) \in I \times TX : \pi_{TX}(v) = c(t)\}$ is the pullback by c of the tangent bundle TX and the projection $\pi_{c^*TX} : c^*TX \rightarrow I$ is given by the canonical projection on the first coordinate $\pi_{c^*TX} := \mathrm{proj}_1$.

Remark 4.30 (On the structure of $L^p(I, TX)$). We emphasize that this result says nothing about a potential differentiable structure of $L^p(I, TX)$. This question is left open for now and will be further investigated in future work. Calling $\pi_{L^p(I, TX)}$ a ‘‘Banach bundle’’ is thus a slight abuse of notation since it is only a surjective mapping having Banach fibers.

Proof. Let $p \in [1, \infty]$. For each $c \in \mathcal{AC}_s^p(I, X)$, the zero section of π_{c^*TX} is continuous, hence Borel measurable, as it is given by the composition of the zero section of π_{TX} , which is C^1 , and of c , which is continuous. Hence, $S^p(\pi_{c^*TX})$ is defined without ambiguity using the norm $b_{p, \pi_{c^*TX}} : S^p(\pi_{c^*TX}) \rightarrow [0, \infty)$ defined as

$$b_{p, \pi_{c^*TX}}(v) := \begin{cases} \left(\int_I b_{c(t)}(v(t))^p dt \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mathcal{L}^1\text{-ess sup}_{t \in I} b_{c(t)}(v(t)), & \text{if } p = \infty \end{cases}.$$

In addition, $S^p(\pi_{c^*TX})$ is a Banach space. Indeed, π_{c^*TX} being a Banach bundle, $S^p(\pi_{c^*TX})$ inherits its vector space structure from the pointwise operations of addition and scalar multiplication and is complete by performing the same proof as [Proposition 2.17](#). Then, we can define

$$L^p(I, TX) := \cup_{c \in \mathcal{AC}_s^p(I, X)} S^p(\pi_{c^*TX}).$$

Furthermore, the mapping

$$\pi_{L^p(I, TX)} := (\pi_{TX})_* : L^p(I, TX) \rightarrow \mathcal{AC}_s^p(I, X)$$

is surjective and its fibers are given by $\pi_{L^p(I, TX)}^{-1}(\{c\}) = S^p(\pi_{c^*TX})$ for all $c \in \mathcal{AC}_s^p(I, X)$. Hence, $\pi_{L^p(I, TX)}$ is a Banach bundle. \square

The mapping that associates its speed to each strongly absolutely continuous curve in a Finsler manifold is then defined as a section of the Banach bundle given by [Proposition 4.29](#).

Proposition 4.31 (Speed of strongly absolutely continuous curves in Finsler manifolds). *Let $p \in [1, \infty]$. Suppose that (X, b_X) is a C^1 Finsler manifold modelled. Then, the mapping*

$$\partial_X : \begin{cases} \mathcal{AC}_s^p(I, X) & \longrightarrow & L^p(I, TX) \\ c & \longmapsto & \dot{c} := \partial_X c \end{cases}$$

is a section of $\pi_{L^p(I, TX)}$.

Proof. Let $p \in [1, \infty]$ and $c \in \mathcal{AC}_s^p(I, X)$. Thus, there exists a partition $a = t_0 < t_1 < \dots < t_n = b$ and charts (U_i, φ_i) of X for $1 \leq i \leq n$ such that $c([t_{i-1}, t_i]) \subset U_i$ and $\varphi_i \circ c|_{[t_{i-1}, t_i]} \in \mathcal{AC}_s^p([t_{i-1}, t_i], \varphi_i(U_i))$. Hence, denoting \mathbb{B} the Banach space on which X is modelled, there exists $\eta_i \in L^p([t_{i-1}, t_i], \mathbb{B})$ such that for all $(s, t) \in [t_{i-1}, t_i]^2$ satisfying $s < t$ we have

$$(\varphi_i \circ c)(t) - (\varphi_i \circ c)(s) = \int_s^t \eta_i(u) du.$$

Then, we can define for \mathcal{L}^1 -a.e. $t \in I$

$$\dot{c}(t) := \sum_{i=1}^n \mathbb{1}_{[t_{i-1}, t_i]}(t) d\varphi_i^{-1}(\varphi_i(c(t)))(\eta_i(t)) \in T_{c(t)}X,$$

which does not depend on the choice of charts. Therefore, \dot{c} is measurable with separable range as the sum of measurable mappings with separable ranges and is a section of π_{c^*TX} . Thus, up to the identification with its class of equivalence, \dot{c} belongs to $S^p(\pi_{c^*TX})$. \square

Remark 4.32 (Related results in the literature). [Proposition 4.31](#) was proved by W. P. A. Klingenberg [[Kli95](#), Proposition 2.3.16] in the case of finite-dimensional Riemannian manifolds and by A. Schmeding [[Sch16](#), Proposition 2.8] in the case of (strong) infinite-dimensional Riemannian manifolds. Both results are corollaries of [Proposition 4.31](#) since any Hilbert manifold with a (strong) continuous Riemannian metric is a Finsler manifold [[Pal66](#), Theorem 2.12]. However, we emphasize that, unlike [[Kli95](#), Proposition 2.3.16] and [[Sch16](#), Proposition 2.8], [Proposition 4.31](#) tells nothing about the regularity of the section which maps any absolutely continuous curve to its L^p speed. This question is left open for now and will be further investigated in future work as it is related to [Remark 4.30](#).

Now, one could ask under which conditions on X can the above construction transpose to absolutely continuous curves in the sense of [Definition 2.28](#). To answer this question, let us first recall that, when X is a Banach space satisfying the Radon–Nikodým property (see [\[Lis06, Section 1.6.2\]](#) for a definition), the set of absolutely continuous curves defined in [Definition 2.28](#) coincides with the set of strongly absolutely continuous curves defined in [Definition 4.26](#) [\[Lis06, Theorem 1.16\]](#).

Proposition 4.33 (Coincidence of $\mathcal{AC}^p(I, X)$ and $\mathcal{AC}_s^p(I, X)$ in RNP Banach spaces). *Let $p \in [1, \infty]$. Suppose that X is a Banach space satisfying the Radon–Nikodým property. Then, $\mathcal{AC}^p(I, X) = \mathcal{AC}_s^p(I, X)$.*

Then, [Proposition 4.33](#) extends to Finsler manifolds modelled on a Banach space satisfying the Radon–Nikodým property through the use of local charts, so that the notion of strongly absolutely continuous curves given by [Definition 4.27](#) and the notion of absolutely continuous curves given by [Definition 2.28](#) also coincide.

Proposition 4.34 (Coincidence of $\mathcal{AC}^p(I, X)$ and $\mathcal{AC}_s^p(I, X)$ in RNP Finsler manifolds). *Let $p \in [1, \infty]$. Suppose that (X, b_X) is a C^1 Finsler manifold modelled on a Banach space \mathbb{B} satisfying the Radon–Nikodým property. Then, $\mathcal{AC}^p(I, X) = \mathcal{AC}_s^p(I, X)$.*

Proof. Let $p \in [1, \infty]$. The inclusion $\mathcal{AC}_s^p(I, X) \subset \mathcal{AC}^p(I, X)$ follows from [Proposition 4.25](#). Let us prove the reverse inclusion. Let $c \in \mathcal{AC}^p(I, X)$. Since I is compact and c is continuous, there exist a partition $a = t_0 < t_1 < \dots < t_n = b$ and charts (U_i, φ_i) of X for $1 \leq i \leq n$ such that $c([t_{i-1}, t_i]) \subset U_i$. By [Proposition 4.25](#), the fact that $c \in \mathcal{AC}^p(I, X)$ implies that $\varphi_i \circ c|_{[t_{i-1}, t_i]} \in \mathcal{AC}^p([t_{i-1}, t_i], \varphi_i(U_i))$ and, since \mathbb{B} has the Radon–Nikodým property, $\varphi_i \circ c|_{[t_{i-1}, t_i]} \in \mathcal{AC}_s^p([t_{i-1}, t_i], \varphi_i(U_i))$, by [Proposition 4.33](#). Hence, $c \in \mathcal{AC}_s^p(I, X)$. \square

As a consequence of [Propositions 4.25](#) and [4.34](#), the mapping that associates its speed to each absolutely continuous curves is defined as in [Proposition 4.31](#) and the norm of the speed coincide with the metric derivative, when X has a Finsler structure and is modelled on a Banach space satisfying the Radon–Nikodým property.

Theorem 4.35 (Speed of absolutely continuous curves in RNP Finsler manifolds). *Let $p \in [1, \infty]$. Suppose that (X, b_X) is a C^1 Finsler manifold modelled on a Banach space satisfying the Radon–Nikodým property. Then, the mapping*

$$\partial_X : \begin{cases} \mathcal{AC}^p(I, X) & \longrightarrow & L^p(I, TX) \\ c & \longmapsto & \dot{c} := \partial_X c \end{cases}$$

is a section of $\pi_{L^p(I, TX)}$ and satisfies for all $c \in \mathcal{AC}^p(I, X)$ that

$$|c'|_X(t) = b_{c(t)}(\dot{c}(t)) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I, \quad (9)$$

Proof. The proof follows by identifying absolutely continuous curves and strongly absolutely continuous curves using [Proposition 4.34](#), repeating the arguments in the proof of [Proposition 4.31](#) to construct the speed and using [Proposition 4.25](#) to show that the norm of the speed and the metric derivative coincide.

Step 1: Construction of the speed. Let $p \in [1, \infty]$ and $c \in \mathcal{AC}^p(I, X)$. By [Proposition 4.34](#), $c \in \mathcal{AC}_s^p(I, X)$. Hence, there exists a partition $a = t_0 < t_1 < \dots < t_n = b$ and charts (U_i, φ_i) of X for $1 \leq i \leq n$ such that $c([t_{i-1}, t_i]) \subset U_i$ and $\varphi_i \circ c|_{[t_{i-1}, t_i]} \in \mathcal{AC}_s^p([t_{i-1}, t_i], \varphi_i(U_i))$. Thus, denoting \mathbb{B} the Banach space on which X is modelled, there exists $\eta_i \in L^p([t_{i-1}, t_i], \mathbb{B})$ such that for all $(s, t) \in [t_{i-1}, t_i]^2$ satisfying $s < t$ we have

$$(\varphi_i \circ c)(t) - (\varphi_i \circ c)(s) = \int_s^t \eta_i(u) \, du.$$

Then, we can define for \mathcal{L}^1 -a.e. $t \in I$

$$\dot{c}(t) := \sum_{i=1}^n \mathbb{1}_{[t_{i-1}, t_i]}(t) \, d\varphi_i^{-1}(\varphi_i(c(t))) (\eta_i(t)) \in T_{c(t)}X,$$

which does not depend on the choice of charts. Therefore, \dot{c} is measurable with separable range as the sum of measurable mappings with separable ranges and is a section of π_{c^*TX} . Thus, up to the identification with its class of equivalence, \dot{c} belongs to $S^p(\pi_{c^*TX})$.

Step 2: Proof of Equation (9). Finally, thanks to [Proposition 4.25](#), we get that for \mathcal{L}^1 -a.e. $t \in I$ it holds for all $\varepsilon > 0$ that

$$\frac{1}{1 + \varepsilon} |c'|_X(t) \leq b_{c(t)}(\dot{c}(t)) \leq (1 + \varepsilon) |c'|_X(t).$$

Hence, we have

$$|c'|_X(t) = b_{c(t)}(\dot{c}(t)) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I. \quad \square$$

Remark 4.36 (Related results in the literature). [Theorem 4.35](#) was proved by A. Burtscher [[Bur15](#), Theorem 2.2., Proposition 4.10] in the case of finite-dimensional manifolds equipped with continuous Riemannian metrics. This result is a corollary of [Theorem 4.35](#) since any Hilbert manifold with a continuous Riemannian metric is a Finsler manifold [[Pal66](#), Theorem 2.12]. Also, since every Banach space is trivially a Finsler manifold, we get back from [Theorem 4.35](#) the well-known fact that the norm of the speed of an absolutely continuous curves in a Banach space satisfying the Radon–Nikodým property coincides with its metric derivative (see [[Lis07](#), Theorem 1.16] or [[AGS05](#), Remark 1.1.3]).

In the next section, we define the speed of absolutely continuous curves in nonlinear Lebesgue spaces in a similar manner as [Theorem 4.35](#), when the target space is a Finsler manifold and despite the lack of differential structure on nonlinear Lebesgue spaces outside the linear case.

4.2.3. Speed of absolutely continuous curves in nonlinear Lebesgue spaces

Let us first define a notion of tangent bundles over nonlinear Lebesgue spaces.

Proposition 4.37 (Tangent bundles on nonlinear Lebesgue spaces). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. When (N, b_N) is a C^1 Finsler manifold, define the tangent bundle of $L_h^p(M, N)$ as the disjoint union*

$$TL_h^p(M, N) := \cup_{f \in L_h^p(M, N)} S^p(\pi_{f^*TN}),$$

where $f^*TN := \{(x, v) \in M \times TN : \pi_{TN}(v) = f(x)\}$ is the pullback by $f \in L_h^p(M, N)$ of the tangent bundle TN and the projection $\pi_{f^*TN} : f^*TN \rightarrow M$ is given by the canonical projection on the first coordinate $\pi_{f^*TN} := \text{proj}_1$. In addition, the mapping

$$\pi_{TL_h^p(M, N)} := (\pi_{TN})_* : TL_h^p(M, N) \rightarrow L_h^p(M, N)$$

forms a Banach bundle over $L_h^p(M, N)$ and we can define the mapping $B_p : TL_h^p(M, N) \rightarrow [0, \infty)$ such that for all $f \in L_h^p(M, N)$ it induces a norm on the tangent space $T_f L_h^p(M, N) := S^p(\pi_{f^*TN})$ defined as

$$B_p|_{T_f L_h^p(M, N)}(z) := \begin{cases} \left(\int_M b_{f(x)}(z(x))^p d\mu_M(x) \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mu_M\text{-ess sup}_{x \in M} b_{f(x)}(z(x)), & \text{if } p = \infty \end{cases},$$

which we also denote by $B_f := B_p|_{T_f L_h^p(M, N)}$ for short and refer as \mathcal{L}^p fibered norm.

Remark 4.38 (On the structure of $TL_h^p(M, N)$). Recall that $L_h^p(M, N)$ does not have a differential structure despite N having one. Similarly, $TL_h^p(M, N)$ has a priori no differential structure, so that calling it a “tangent bundle” is a slight abuse of notation since it is only the disjoint union of Banach spaces and $\pi_{TL_h^p(M, N)}$ is just a surjective mapping with Banach fibers.

Proof. Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. For each $f \in L_h^p(M, N)$, the zero section of π_{f^*TN} is measurable as it is given by the composition of the zero section of π_{TN} , which is C^1 , and of f , which is measurable. Hence, $S^p(\pi_{f^*TN})$ is defined without ambiguity using the norm $B_{p, \pi_{f^*TN}} : S^p(\pi_{f^*TN}) \rightarrow [0, \infty)$ defined as

$$B_{p, \pi_{f^*TN}}(z) := \begin{cases} \left(\int_M b_{f(x)}(z(x))^p d\mu_M(x) \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mu_M\text{-ess sup}_{x \in M} b_{f(x)}(z(x)), & \text{if } p = \infty \end{cases}.$$

In addition, $S^P(\pi_{f^*TN})$ is a Banach space. Indeed, π_{f^*TN} being a Banach bundle, $S^P(\pi_{f^*TN})$ inherits its vector space structure from the pointwise operations of addition and scalar multiplication and is complete by performing the same proof as [Proposition 2.17](#). Then, we can define

$$TL_h^p(M, N) := \cup_{f \in L_h^p(M, N)} S^P(\pi_{f^*TN}).$$

Furthermore, the mapping

$$\pi_{TL_h^p(M, N)} := (\pi_{TN})_* : TL_h^p(M, N) \rightarrow L_h^p(M, N)$$

is surjective and its fibers are given by $\pi_{TL_h^p(M, N)}^{-1}(\{f\}) = S^P(\pi_{f^*TN})$ for all $f \in L_h^p(M, N)$. Hence, $\pi_{TL_h^p(M, N)}$ is a Banach bundle. Furthermore, the mapping $B_p : TL_h^p(M, N) \rightarrow [0, \infty)$ is defined as $B_p|_{T_f L_h^p(M, N)} := B_{p, \pi_{f^*TN}}$ for all $f \in L_h^p(M, N)$. \square

There is a priori no guarantee that the mapping B_p defined in [Proposition 4.37](#) is measurable in any sense, hence we might not be able to integrate functionals over $TL_h^p(M, N)$ involving B_p . To define a topology on $TL_h^p(M, N)$ and make B_p continuous on it, we will need a metric on TN , which will also be required to apply [Theorem 3.19](#) in order to define the speed of absolutely continuous curves in nonlinear Lebesgue spaces. Such a metric can be defined as soon as N is equipped with a spray or, equivalently, with a connection (see [\[Lan12, Chapter IV, §3\]](#) for the definitions of the notions of sprays and connections).

Proposition 4.39 (Splitting norm on TN). *Suppose that (N, b_N) is a C^2 Finsler manifold with a spray S_N (or, equivalently, a connection). Then, if K_N denotes the connector associated to S_N (see [\[Lan12, Lemma 4.4\]](#) for a definition) and $T\pi_{TN}$ the tangent map of π_{TN} (see [\[Lan12, Chapter III, §2\]](#) for a definition), the C^1 Banach manifold TN can be equipped with the splitting norm $m_{TN} : TTN \rightarrow [0, \infty)$ defined for all $v \in TN$ and $\xi \in T_v TN$ as*

$$m_{TN}|_{T_v TN}(\xi)^2 := b_{\pi_{TN}(v)}(T\pi_{TN}(\xi))^2 + b_{\pi_{TN}(v)}(K_N(\xi))^2, \quad (10)$$

so that TN becomes a metric space when equipped with the induced metric $d_{TN} : TN^2 \rightarrow [0, \infty)$ defined as

$$d_{TN}(v, v') := \inf \left\{ \int_I m_{TN}|_{T_u TN}(\dot{u}) dt : u \in C^1(I, TN), u(a) = v, u(b) = v' \right\}.$$

Proof. The result is a consequence of Dombrowski's splitting theorem [\[Lan12, Theorem 4.5\]](#), which states that the mapping

$$(\pi_{TN}, T\pi_{TN}, K_N) : TTN \rightarrow TN \oplus TN \oplus TN$$

is an isomorphism of fiber bundles over N . Indeed, the conclusion of [Proposition 4.39](#) follows from the same arguments exposed at the end of [\[Lan12, Chapter V, §4\]](#) or in [\[Sch16, Section 2.1\]](#), but defining the splitting norm as in [Equation \(10\)](#) using b_N instead of a Riemannian metric. In addition, the induced metric is compatible with the topology of TN . \square

Remark 4.40 (Borel measurability of the zero section of $\pi_{TL_h^p(M, N)}$). When TN is equipped with the splitting norm, its zero section $s_{0, N}$ is isometric from N to TN , so that the zero section of $\pi_{TL_h^p(M, N)}$ defined as $s_{0, p} := (s_{0, N})_*$ is isometric from $L_h^p(M, N)$ to $TL_h^p(M, N)$

As a direct consequence of [Proposition 4.39](#), the tangent bundles over nonlinear Lebesgue spaces defined in [Proposition 4.37](#) become metric spaces as soon as N is equipped with a spray or, equivalently, a connection.

Corollary 4.41 (Metric on $TL^p(M, N)$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Suppose that (N, b_N) is a C^2 Finsler manifold with a spray (or, equivalently, a connection). Then, $TL_h^p(M, N)$ is a metric space when equipped with the finite metric $D_{p, TN} : TL_h^p(M, N)^2 \rightarrow [0, \infty)$ defined as*

$$D_{p, TN}(z, z') := \begin{cases} \left(\int_M d_{TN}(z(x), z'(x))^p d\mu_M(x) \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mu_M\text{-ess sup}_{x \in M} d_{TN}(z(x), z'(x)), & \text{if } p = \infty \end{cases}.$$

Similarly to [Proposition 4.29](#) and using both [Proposition 4.37](#) and [Corollary 4.41](#), one can define a Banach bundle of L^p speeds over the set of absolutely continuous curves in nonlinear Lebesgue spaces.

Proposition 4.42 (Vector bundle over $\mathcal{AC}^p(I, L_h^p(M, N))$). *Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. Suppose that (N, b_N) is a C^2 Finsler manifold with a spray (or, equivalently, a connection). Then, the set*

$$L^p(I, TL_h^p(M, N)) := \cup_{c \in \mathcal{AC}^p(I, L_h^p(M, N))} S^p(\pi_{c^*TL_h^p(M, N)})$$

forms a Banach bundle

$$\pi_{L^p(I, TL_h^p(M, N))} := (\pi_{TL_h^p(M, N)})_* : L^p(I, TL_h^p(M, N)) \rightarrow \mathcal{AC}^p(I, L_h^p(M, N)),$$

where $c^*TL_h^p(M, N) := \{(t, z) \in I \times TL_h^p(M, N) : \pi_{TL_h^p(M, N)}(z) = c(t)\}$ is the pullback by c of the tangent bundle $TL_h^p(M, N)$ and the projection $\pi_{c^*TL_h^p(M, N)} : c^*TL_h^p(M, N) \rightarrow M$ is given by the canonical projection on the first coordinate $\pi_{c^*TL_h^p(M, N)} := \text{proj}_1$.

Remark 4.43 (On the structure of $L^p(I, TL_h^p(M, N))$). As in [Remark 4.30](#), we emphasize that this result says nothing about a potential topological manifold structure of $L^p(I, TL_h^p(M, N))$. This question is left open for now and will be further investigated in future work. Calling $\pi_{L^p(I, TL_h^p(M, N))}$ a ‘‘Banach bundle’’ is thus a slight abuse of notation since it is only a surjective mapping having Banach fibers.

Proof. Let $h \in L^0(M, N)$ and $p \in [1, \infty]$. For each $c \in \mathcal{AC}^p(I, L_h^p(M, N))$. The zero section of $\pi_{c^*TL_h^p(M, N)}$ is continuous, hence Borel measurable, as the composition of the zero section of $\pi_{TL_h^p(M, N)}$, which is continuous ([Remark 4.40](#)), and of c , which is also continuous. Hence, $S^p(\pi_{c^*TL_h^p(M, N)})$ is defined without ambiguity using the norm $B_{p, \pi_{c^*TL_h^p(M, N)}} : S^p(\pi_{c^*TL_h^p(M, N)}) \rightarrow [0, \infty)$ defined as

$$B_{p, \pi_{c^*TL_h^p(M, N)}}(z) := \begin{cases} \left(\int_I B_{c(t)}(z(t))^p dt \right)^{1/p}, & \text{if } p \in [1, \infty) \\ \mathcal{L}^1\text{-ess sup}_{t \in I} B_{c(t)}(z(t)), & \text{if } p = \infty \end{cases}.$$

In addition, $S^p(\pi_{c^*TL_h^p(M, N)})$ is a Banach space. Indeed, $\pi_{c^*TL_h^p(M, N)}$ being a Banach bundle, $S^p(\pi_{c^*TL_h^p(M, N)})$ inherits its vector space structure from the pointwise operations of addition and scalar multiplication and is complete by performing the same proof as [Proposition 2.17](#). Then, we can define

$$L^p(I, TL_h^p(M, N)) := \cup_{c \in \mathcal{AC}_s^p(I, L_h^p(M, N))} S^p(\pi_{c^*TL_h^p(M, N)}).$$

Furthermore, the mapping

$$\pi_{L^p(I, TL_h^p(M, N))} := (\pi_{TL_h^p(M, N)})_* : L^p(I, TL_h^p(M, N)) \rightarrow \mathcal{AC}_s^p(I, L_h^p(M, N))$$

is surjective and its fibers are given by $\pi_{L^p(I, TL_h^p(M, N))}^{-1}(\{c\}) = S^p(\pi_{c^*TL_h^p(M, N)})$ for all $c \in \mathcal{AC}^p(I, L_h^p(M, N))$. Hence, $\pi_{L^p(I, TL_h^p(M, N))}$ is a Banach bundle. \square

Then, similarly to [Theorem 4.35](#), the mapping that associates its speed to each absolutely continuous curve in a nonlinear Lebesgue space is defined as a section of the Banach bundle given by [Proposition 4.42](#) and the norm of the speed coincide with the metric derivative.

Theorem 4.44 (Speed of absolutely continuous curves in nonlinear Lebesgue spaces for $p > 1$). *Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that (N, b_N) is a connected and complete C^2 Finsler manifold with a spray (or, equivalently, a connection) and that N is modelled on a Banach space satisfying the Radon–Nikodým property. Then, the mapping*

$$\partial_p : \begin{cases} \mathcal{AC}_c^p(I, L_h^p(M, N)) & \longrightarrow & L^p(I, TL_h^p(M, N)) \\ c & \longmapsto & \dot{c} := \partial_p c \end{cases}$$

is a section of $\pi_{L^p(I, TL_h^p(M, N))}$ and satisfies for all $c \in \mathcal{AC}^p(I, L_h^p(M, N))$ that

$$|c'|_p(t) = B_{c(t)}(\dot{c}(t)) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I, \tag{11}$$

Remark 4.45 (On relaxing the connectedness assumption on N). The assumption that N is connected in [Theorem 4.44](#) is solely made to ensure that both d_N and d_{TN} are finite in order to apply [Theorem 3.19](#). However, as remarked in [Remark 3.21](#), the conclusion of [Theorem 3.19](#) holds when d_N is not assumed finite as long as h is assumed constant. Hence, [Theorem 4.44](#) holds without assuming that N is connected, but requires h being constant in that case.

Proof. Let $h \in \mathcal{L}_s^0(M, N)$ and $p \in (1, \infty)$.

Step 1: Construction of the speed. Let $c \in \mathcal{AC}^p(I, L_h^p(M, N))$. Then, by [Theorem 3.19](#), $\tilde{i}_p(c)$ belongs to $L_{\underline{h}}^p(M, \mathcal{AC}^p(I, N))$ and is such that

$$\int_M \int_I |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) < \infty.$$

Also, the measurable set $M_c := \{x \in M : d_\infty(\tilde{i}_p(c)(x), \underline{h}(x)) > 0\}$ has σ -finite μ_M -measure and there is $\hat{c} \in \mathcal{L}_s^0(I \times M, N)$ such that $\hat{c}(t, \cdot) = c(t)$ in $L_h^p(M, N)$ for all $t \in I$ and $\hat{c}(\cdot, x) = \tilde{i}_p(c)(x)$ in $C(I, N)$ for μ_M -a.e. $x \in M$ ([Theorems 3.9](#) and [3.19](#)). Now, define the set

$$\begin{aligned} A &:= \{I \times M : (\partial_N \tilde{i}_p(c)(x))(t) \text{ does not exist}\} \\ &= \{I \times M : \limsup_{\tau, \tau' \rightarrow 0} d_{TN}(g_\tau(t, \tilde{i}_p(c)(x)), g_{\tau'}(t, \tilde{i}_p(c)(x))) > 0\} \end{aligned}$$

which is measurable since $\tilde{i}_p(c)$ is measurable from M to $(\mathcal{AC}^p(I, N), d_\infty)$ and the mappings

$$g_\tau : \begin{cases} I \times \mathcal{AC}^p(I, N) & \longrightarrow TN \\ (t, u) & \longmapsto d\varphi^{-1}(\varphi(u(t))(\varphi(u(t+\tau)) - \varphi(u(t))))/\tau \end{cases}$$

are continuous, hence Borel measurable, for some (thus for any) chart (U, φ) such that $u(t) \in U$ and all $\tau > 0$ such that $u(t+\tau) \in U$. In addition, $\mathcal{L}^1(A^c) = 0$ for μ_M -a.e. $x \in M$ since $\tilde{i}_p(c)(x) \in \mathcal{AC}^p(I, N)$. Now, since $\tilde{i}_p(c)|_{M \setminus M_c} = \underline{h}|_{M \setminus M_c}$, A is a subset of the measurable rectangle $I \times M_c$ and, M_c having σ -finite μ_M -measure and I having finite \mathcal{L}^1 -measure, we get, by the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1], that $\mathcal{L}^1 \otimes \mu_M(A) = 0$ and $\mu_M(A_t) = 0$ for \mathcal{L}^1 -a.e. $t \in I$. Thus, for $\mathcal{L}^1 \otimes \mu_M$ -a.e. $(t, x) \in I \times M$, we can define

$$\eta(t, x) := (\partial_N \tilde{i}_p(c)(x))(t) = (\partial_N \hat{c}(\cdot, x))(t) \in T_{\hat{c}(t, x)},$$

which is measurable and has separable range as the pointwise limit of such mappings. Then, by [Theorem 4.35](#), we get that for $\mathcal{L}^1 \otimes \mu_M$ -a.e. $(t, x) \in I \times M$

$$|\tilde{i}_p(c)(x)'|_N(t) = b_{\hat{c}(t, x)}(\eta(t, x)).$$

Finally, integrating over I and M and using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1] yields that

$$\begin{aligned} \int_{I \times M} b_{\hat{c}(t, x)}(\eta(t, x))^p d\mathcal{L}^1 \otimes \mu_M(t, x) &= \int_{I \times M} |\tilde{i}_p(c)(x)'|_N^p(t) d\mathcal{L}^1 \otimes \mu_M(t, x) \\ &= \int_{I \times M_c} |\tilde{i}_p(c)(x)'|_N^p(t) d\mathcal{L}^1 \otimes \mu_M(t, x) \\ &= \int_{M_c} \int_I |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) \\ &= \int_M \int_I |\tilde{i}_p(c)(x)'|_N^p(t) dt d\mu_M(x) < \infty. \end{aligned}$$

Since N is equipped with a spray (or a connection), this even yields that

$$\int_{I \times M} d_{TN}(\eta(t, x), (s_{0,N} \circ \hat{c})(t, x))^p d\mathcal{L}^1 \otimes \mu_M(t, x) < \infty$$

with $s_{0,N}$ the zero section of π_{TN} . Thus, defining $\hat{c}^* s_{0,N} := s_{0,N} \circ \hat{c}$, we have that $\eta \in L_{\hat{c}^* s_{0,N}}^p(I \times M, TN)$ and we can define $\dot{c} : t \mapsto \eta(t, \cdot)$. Then, $\dot{c}(t)$ belongs to $L^0(M, TN)$ for a \mathcal{L}^1 -a.e. $t \in I$. To prove the measurability of \dot{c} from I to

($L^0(M, TN), D_{p, TN}$), note that, by density of $E_{\hat{c}^* s_{0,N}, \sigma}(I \times M, TN) \cap L_{\hat{c}^* s_{0,N}}^p(I \times M, TN)$ in $L_{\hat{c}^* s_{0,N}}^p(I \times M, TN)$ (see assertion (i) of [Proposition 3.7](#)), there exists a sequence $(\hat{g}_n)_{n \in \mathbb{N}}$ in $E_{\hat{c}^* s_{0,N}, \sigma}(I \times M, TN) \cap L_{\hat{c}^* s_{0,N}}^p(I \times M, TN)$ such that $\hat{g}_n \rightarrow \eta$ in $L_{\hat{c}^* s_{0,N}}^p(I \times M, TN)$ as $n \rightarrow \infty$. Thus, define $g_n : t \mapsto \hat{g}_n(t, \cdot)$, which is measurable as an almost simple mapping from I to $(L^0(M, TN), D_{p, TN})$ with base mapping given by the mapping $t \mapsto \hat{c}^* s_{0,N}(t, \cdot)$, which is Borel measurable from I to $(L^0(M, TN), D_{p, TN})$ since $D_{p, TN}(\hat{c}^* s_{0,N}(t, \cdot), c^* s_{0,p}(t)) = 0$ for all $t \in I$ with $c^* s_{0,p} := s_{0,p} \circ c$ the zero section of $\pi_{c^* TL_h^p(M, N)}$, which is continuous, hence Borel measurable, as the composition of $s_{0,p} := (s_{0,N})_*$, the zero section of $\pi_{TL_h^p(M, N)}$, which is continuous ([Remark 4.40](#)), and of c , which is also continuous. Then, noticing that the union of the sets on which the \hat{g}_n 's and η differ from $\hat{c}^* s_{0,N}$ is contained in a measurable rectangle $I \times \hat{M}$ such that \hat{M} has σ -finite μ_M -measure (see step 2 in the proof of assertion (i) of [Theorem 3.9](#)), we get, using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1], that

$$d_{p,p}(g_n, \dot{c}) = \hat{D}_p(\hat{g}_n, \eta) \longrightarrow 0$$

as $n \rightarrow \infty$, so that, up to the extraction of a subsequence, we can assume for \mathcal{L}^1 -a.e. $t \in I$ that $D_p(g_n(t), \dot{c}(t)) \rightarrow 0$ as $n \rightarrow \infty$. Hence, \dot{c} is measurable as the pointwise limit of measurable mappings. In addition, since $\mu_M(A_t) = 0$ for \mathcal{L}^1 -a.e. $t \in I$, we have for \mathcal{L}^1 -a.e. $t \in I$ that $\dot{c}(t)(x) = \eta(t, x) \in T_{\hat{c}(t,x)}N = T_{c(t)(x)}N$ and $s_{0,N}(\hat{c}(t, x)) = s_{0,N}(c(t)(x))$ for μ_M -a.e. $x \in M$. Thus, $\dot{c}(t) \in T_{c(t)}L_h^p(M, N)$ for \mathcal{L}^1 -a.e. $t \in I$ and, using the Fubini–Tonelli theorem [[Coh13](#), Proposition 5.2.1], we have $\dot{c} \in L_{c^* s_{0,p}}^p(I, TL_h^p(M, N))$. Hence, \dot{c} belongs to $S^p(\pi_{c^* TL_h^p(M, N)})$.

Step 2: Proof of [Equation \(11\)](#). Since $\mu_M(A_t) = 0$ for \mathcal{L}^1 -a.e. $t \in I$, we get that for \mathcal{L}^1 -a.e. $t \in I$ the equality

$$|\tilde{i}_p(c)(x)'|_N(t) = b_{c(t)(x)}(\dot{c}(t)(x))$$

holds for μ_M -a.e. $x \in M$. Hence, integrating over M , we have

$$|c'|_p(t) = \left(\int_M b_{c(t)(x)}(\dot{c}(t)(x))^p d\mu_M(x) \right)^{1/p} = B_{c(t)}(\dot{c}(t)) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I. \quad \square$$

Finally, combining [Proposition 4.4](#) and [Theorem 4.44](#) shows that the metric D_p on nonlinear Lebesgue spaces coincides with the metric induced by the infimum of the energy functional associated with the fibered norm B_p ([Proposition 4.37](#)) over absolutely continuous curves. This is analogous to the situation on standard Finsler manifolds ([Proposition 4.24](#)); the only difference being the class of curves over which the infimum is taken, since nonlinear Lebesgue spaces lack a differential structure.

Corollary 4.46 (\mathcal{L}^p metrics are induced by \mathcal{L}^p fibered norms). *Let $h \in L^0(M, N)$ and $p \in (1, \infty)$. Suppose that (N, b_N) is a connected, complete and separable C^2 Finsler manifold with a spray (or, equivalently, a connection) and that N is modelled on a Banach space satisfying the Radon–Nikodým property. Then, we have for all $(f, f') \in L_h^p(M, N)^2$ that*

$$D_p(f, f') = \inf \left\{ \mathcal{L}^1(I)^{1-1/p} \left(\int_I B_{c(t)}(\dot{c}(t))^p dt \right)^{1/p} : c \in \mathcal{AC}^p(I, L_h^p(M, N)), c(a) = f, c(b) = f' \right\}.$$

Acknowledgements

This is part of the author's PhD thesis. The author thus thanks his advisor Joan Alexis Glaunès (MAP5, Université Paris Cité) for his support, useful advice, and for introducing him to the subject of metamorphoses of manifold-valued images, which led him to dig into the theory of nonlinear Lebesgue spaces. The author also thanks Théo Dumont (LIGM, Université Gustave Eiffel) for suggestions that significantly helped improve the exposition of the present article as well as Alain Trouvé (Centre Borelli, ENS Paris-Saclay) for careful proofreading of early versions of this article and fruitful technical discussions.

Notation and terminology

This section clarifies some recurrent notions and notation used in the present article:

Spaces

- \mathbb{N} denotes the set of nonnegative integers (including 0).
- \mathbb{N}^* denotes the set of positive integers (excluding 0).
- \mathbb{R} denotes the set of real numbers also referred to as *real line*.
- \mathbb{R}_+ denotes the set of nonnegative real numbers (including 0).
- \mathbb{R}^* denotes the set of nonzero real numbers (excluding 0).
- \mathbb{Q} denotes the set of rational numbers.

Topology

- $f(X) := \{f(x) : x \in X\}$ denotes the *range* of a mapping $f : X \rightarrow Y$ between sets X and Y .
- \bar{A} denotes the *closure* of the subset $A \subset X$ in the topological space X , that is, the intersection of all closed subsets of X that include A .
- \bar{A}^{d_X} denotes the closure of the subset $A \subset X$ in the metric space (X, d_X) .
- A set X is called *countable* if there exists a bijective mapping $\varphi : \mathbb{N} \rightarrow X$.
- A subset S of a (semi-)metric space (X, d_X) is called *dense* when for each element x of X and real number $\varepsilon > 0$ there exists an element s of S such that $d_X(x, s) < \varepsilon$.
- A metric space (X, d_X) is called *separable* when it has a countable dense subset.
- A sequence $(x_n)_{n \in \mathbb{N}}$ in a metric space (X, d_X) is called *Cauchy* when for all real $\varepsilon > 0$ there exists an integer N such that for all $n \geq N$ and $m \geq N$, $d_N(x_n, x_m) < \varepsilon$.
- A metric space (X, d_X) is called *complete* when every Cauchy sequence converges to an element of X .
- A^c denotes the *complement* of a subset A of a set X , that is, $A^c := \{x \in X : x \notin A\}$.
- $A \setminus B := A \cap B^c$ denotes the *set difference* between two subsets A and B of a set X .
- $A \Delta A' := (A \setminus A') \cup (A' \setminus A)$ denotes the *symmetric set difference* between two subsets A and A' of a set X .

Measure theory

- \mathcal{L}^d denotes the d -dimensional Lebesgue measure.
- $\#$ denotes the counting measure.
- A measure μ_M on a measurable set (M, Σ_M) is called *purely infinite* if its range is reduced to $\{0, \infty\}$.
- $\sigma(C)$ denotes the smallest σ -algebra that contains a family C of subsets of a set M .
- A subset of measurable space (M, Σ_M) is called *measurable* when it belongs to Σ_M .
- \mathcal{F}_{μ_M} denotes the set of measurable subsets of a measure space (M, Σ_M, μ_M) with finite μ_M -measure.
- $\Sigma_M|_B := \{B \cap A : A \in \Sigma_M\}$ denotes the restriction of a σ -algebra Σ_M on a set M to subsets of $B \in \Sigma_M$.
- $\mu_M|_B$ denotes the restriction of a measure μ_M on a measurable space (M, Σ_M) to $\Sigma_M|_B$ for some $B \in \Sigma_M$.
- $\mathcal{B}(M)$ denotes the *Borel σ -algebra* on a topological space M .
- A measure μ_M on a topological space M is called *Borel* when it is defined on $\mathcal{B}(M)$.
- A subset Z of a measure space (M, Σ_M, μ_M) is called μ_M -*null* if there exists $A \in \Sigma_M$ such that $Z \subset A$ and $\mu_M(A) = 0$.
- \mathcal{Z}_{μ_M} denotes the set of all μ_M -null sets of a measure space (M, Σ_M, μ_M) .
- A property \mathcal{P} that depends on the choice of point x of a measure space (M, Σ_M, μ_M) is said to hold μ_M -*almost everywhere* (μ_M -a.e. for short) if $\{x \in M : \mathcal{P}(x)\}$ is a μ_M -null set.
- In a measure space (M, Σ_M, μ_M) , $\bar{\Sigma}_M$ denotes the *completion of the σ -algebra Σ_M* with respect to μ_M and is defined as $\bar{\Sigma}_M := \sigma(\Sigma_M \cup \mathcal{Z}_{\mu_M}) = \{A \cup Z : (A, Z) \in \Sigma_M \times \mathcal{Z}_{\mu_M}\}$.
- In a measure space (M, Σ_M, μ_M) , $\bar{\mu}_M$ denotes the *completion of the measure μ_M* and is defined as the mapping $\bar{\mu}_M : \bar{\Sigma}_M \rightarrow [0, \infty]$ such that $\bar{\mu}_M(A \cup Z) = \mu_M(A)$ for all $(A, Z) \in \Sigma_M \times \mathcal{Z}_{\mu_M}$. It is itself a measure on $(M, \bar{\Sigma}_M)$.

References

- [AB06] Charalambos D Aliprantis and Kim C Border. *Infinite dimensional analysis: a hitchhiker's guide*. Springer, 2006.
- [AF05] John Ashburner and Karl J. Friston. Unified segmentation. *Neuroimage*, 26(3):839–851, 2005.
- [AGS05] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows: in metric spaces and in the space of probability measures*. Springer Science & Business Media, 2005.
- [ALS24] Ehsan Abedi, Zhenhao Li, and Timo Schultz. Absolutely continuous and bv-curves in 1-wasserstein spaces. *Calculus of Variations and Partial Differential Equations*, 63(1):16, 2024.
- [Bac14] Miroslav Bacák. *Convex analysis and optimization in Hadamard spaces*, volume 22. Walter de Gruyter GmbH & Co KG, 2014.
- [BBI22] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry*, volume 33. American Mathematical Society, 2022.
- [BML94] Peter J. Basser, James Mattiello, and Denis LeBihan. MR diffusion tensor spectroscopy and imaging. *Biophysical journal*, 66(1):259–267, 1994.
- [BMNN25] Martin Bauer, Facundo Mémoli, Tom Needham, and Mao Nishino. The Z-Gromov-Wasserstein Distance. *Journal of Machine Learning Research*, 26(291):1–57, 2025.
- [Bou13] Nicolas Bourbaki. *Topological vector spaces: Chapters 1–5*. Springer Science & Business Media, 2013.
- [Bur15] Annegret Y Burtscher. Length structures on manifolds with continuous riemannian metrics. *New York J. Math.*, 21:273–296, 2015.
- [Cha13] Nicolas Charon. *Analysis of geometric and functional shapes with extensions of currents: applications to registration and atlas estimation*. PhD thesis, École normale supérieure de Cachan-ENS Cachan, 2013. URL <https://theses.hal.science/tel-00942078v1>.
- [Coh13] Donald L Cohn. *Measure theory*, volume 5. Springer, 2013.
- [CVCV77] Charles Castaing, Michel Valadier, Charles Castaing, and Michel Valadier. *Measurable multifunctions*. Springer, 1977.
- [DiB02] Emmanuele DiBenedetto. *Real analysis*, volume 13. Springer, 2002.
- [DS88] Nelson Dunford and Jacob T Schwartz. *Linear operators, part 1: general theory*, volume 10. John Wiley & Sons, 1988.
- [Eng77] Ryszard Engelking. *General topology*, volume 60. PWN Warszawa, 1977.
- [Glo15] Helge Glockner. Measurable regularity properties of infinite-dimensional lie groups. *arXiv preprint arXiv:1601.02568*, 2015.
- [HVN VW16] Tuomas Hytönen, Jan Van Neerven, Mark Veraar, and Lutz Weis. *Analysis in Banach spaces*, volume 12. Springer, 2016.
- [JSSG14] J Jaramillo, M Jimenez-Sevilla, and L Sanchez-Gonzalez. Characterization of a banach-finsler manifold in terms of the algebras of smooth functions. *Proceedings of the American Mathematical Society*, 142(3):1075–1087, 2014.
- [Jos94] Jürgen Jost. Equilibrium maps between metric spaces. *Calculus of Variations and Partial Differential Equations*, 2:173–204, 1994.
- [Jos97a] Jürgen Jost. Generalized dirichlet forms and harmonic maps. *Calculus of Variations and Partial Differential Equations*, 5(1):1–19, 1997.
- [Jos97b] Jürgen Jost. *Nonpositive curvature: geometric and analytic aspects*. Birkhäuser, 1997.
- [JSSG11] Mar Jiménez-Sevilla and Luis Sánchez-González. On some problems on smooth approximation and smooth extension of lipschitz functions on banach-finsler manifolds. *Nonlinear Analysis: Theory, Methods & Applications*, 74(11):3487–3500, 2011.
- [Kli95] Wilhelm Klingenberg. *Riemannian geometry*, volume 1. Walter de Gruyter, 1995.
- [KS93] Nicholas J Korevaar and Richard M Schoen. Sobolev spaces and harmonic maps for metric space targets. *Communications in Analysis and Geometry*, 1(4):561–659, 1993.
- [Lan12] Serge Lang. *Fundamentals of differential geometry*, volume 191. Springer Science & Business Media, 2012.
- [Lis06] Stefano Lisini. *Absolutely continuous curves in Wasserstein spaces with applications to continuity equation and to nonlinear diffusion equations*. PhD thesis, Università degli Studi di Pavia, 2006. URL https://cvgmt.sns.it/media/doc/paper/848/Tesi_Lisini.pdf.
- [Lis07] Stefano Lisini. Characterization of absolutely continuous curves in wasserstein spaces. *Calculus of variations and partial differential equations*, 28(1):85–120, 2007.
- [LP10] Nina Lebedeva and Anton Petrunin. Curvature bounded below: a definition a la berg-nikolaev. *arXiv preprint arXiv:1008.3318*, 2010.
- [MTE⁺95] John C. Mazziotta, Arthur W Toga, Alan Evans, Peter Fox, and Jack Lancaster. A probabilistic atlas of the human brain: theory and rationale for its development. *Neuroimage*, 2(2):89–101, 1995.
- [Pal66] Richard S Palais. Lusternik-schnirelman theory on banach manifolds. *Topology*, 5(2):115–132, 1966.
- [PT24] Thomas Pierron and Alain Trounev. The graded group action framework for sub-riemannian orbit models in shape spaces. *arXiv preprint arXiv:2406.16930*, 2024.
- [Rie10] Friedrich Riesz. Untersuchungen über systeme integrierbarer funktionen. *Mathematische Annalen*, 69(4):449–497, 1910.
- [Sch73] Laurent Schwartz. *Radon measures on arbitrary topological spaces and cylindrical measures*. Oxford University Press, 1973.
- [Sch16] Alexander Schmeding. Manifolds of absolutely continuous curves and the square root velocity framework. *arXiv preprint arXiv:1612.02604*, 2016.
- [SS09] Elias M Stein and Rami Shakarchi. *Real analysis: measure theory, integration, and Hilbert spaces*. Princeton University Press, 2009.
- [ST25] Guillaume Sérieyss and Alain Trounev. Nonlinear lebesgue spaces: Dense subspaces, completeness and separability. *arXiv preprint arXiv:2512.19208*, 2025.
- [Stu99] Karl-Theodor Sturm. Metric spaces of lower bounded curvature. *Expositiones Mathematicae*, 17:035–048, 1999.
- [Stu01] Karl-Theodor Sturm. Nonlinear markov operators associated with symmetric markov kernels and energy minimizing maps between singular spaces. *Calculus of Variations and Partial Differential Equations*, 12(4):317–357, Jun 2001.
- [Stu02] Karl-Theodor Sturm. Nonlinear markov operators, discrete heat flow, and harmonic maps between singular spaces. *Potential Analysis*, 16:305–340, 2002.
- [Stu23] Karl-Theodor Sturm. *The space of spaces: curvature bounds and gradient flows on the space of metric measure spaces*, volume 290. American Mathematical Society, 2023.

- [TCGC04] J-Donald Tournier, Fernando Calamante, David G Gadian, and Alan Connelly. Direct estimation of the fiber orientation density function from diffusion-weighted mri data using spherical deconvolution. *Neuroimage*, 23(3):1176–1185, 2004.
- [Tro95] Alain Trouvé. An infinite dimensional group approach for physics based models in pattern recognition. *preprint*, 1995.
- [Tuc04] David S Tuch. Q-ball imaging. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, 52(6):1358–1372, 2004.

Appendix A. Omitted results

Appendix A.1. Omitted results of Section 3.1

Let us provide a proof of the alternative density result mentioned in Remark 3.8.

Proposition A.1 (Relaxation of the assumptions in (iii) of Proposition 3.7). *Let $h \in E(M, N)$ and $p \in [1, \infty)$. Suppose that (I, Σ_I, μ_I) is a measure space, with no further assumptions. Then, the closure of the target space of $\text{sec}_{M,p}$ is*

$$\overline{E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^0(I, N))}^{D_{p,p}} = L_{\underline{h}}^p(M, L^0(I, N)).$$

Proof. Let $h \in \mathcal{E}(M, N)$ and $p \in [1, \infty)$. Then, we can assume that there exists a finite indexing set J such that $h(M) = \{y_j \in N : j \in J\}$ and define $B_j := h^{-1}(\{y_j\})$, so that $M = \cup_{j \in J} B_j$ and we have $L_{\underline{h}}^p(M, L^0(I, N)) \cong \prod_{j \in J} L_{\underline{h}}^p(B_j, L^0(I, N))$ since J is finite (otherwise the gluing map may produce non integrable mappings). However, on each B_j , we get that $L_{\underline{h}}^p(B_j, L^0(I, N)) = L_{\underline{h}}^p(B_j, L_{h_j}^p(I, N))$ with $h_j \equiv y_j$ and the density of $E_{\underline{h}}(B_j, E(I, N)) \cap L_{\underline{h}}^p(B_j, L_{h_j}^p(I, N))$ follows from the same arguments as the proof of Proposition 3.7. Hence, we get that $\prod_{j \in J} E_{\underline{h}}(B_j, E(I, N)) \cap L_{\underline{h}}^p(B_j, L_{h_j}^p(I, N))$ is dense in $\prod_{j \in J} L_{\underline{h}}^p(B_j, L_{h_j}^p(M_0, N))$ by [Eng77, Corollary 2.3.5]. Finally, by noticing that $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^0(I, N)) \cong \prod_{j \in J} E_{\underline{h}}(B_j, E(I, N)) \cap L_{\underline{h}}^p(B_j, L_{h_j}^p(I, N))$ since J is finite (otherwise the gluing map may produce mappings that are not almost simple), we get that $E_{\underline{h}}(M, E(I, N)) \cap L_{\underline{h}}^p(M, L^0(I, N))$ is a dense subspace of $(L_{\underline{h}}^p(M, L^0(I, N)), D_{p,p})$. \square

Appendix B. Reminders

Appendix B.1. Length spaces

This section introduces the basic notions of metric geometry on length spaces (see [BBI22, Chapter 2] for a complete exposition on that matter).

In a metric space X with metric $d_X : X^2 \rightarrow [0, \infty)$, the *length* of an absolutely continuous curve (Definition 2.28) is defined as

Definition B.1 (Length of a curve). The *length* of a curve $c \in \mathcal{AC}(I, X)$ induced by d is defined as

$$L_X(c) := \int_I |c'|_X(t) dt.$$

Taking the infimum over curves connecting two points happens to induce a metric, potentially infinite-valued, when there is no absolutely continuous curve connecting the two points.

Definition B.2 (Intrinsic metric). The *intrinsic metric*, $\widehat{d}_X : X^2 \rightarrow [0, \infty]$, induced by d_X is defined as

$$\widehat{d}_X(x, x') := \inf\{L_X(c) : c \in \mathcal{AC}(I, X), c(a) = x, c(b) = x'\}.$$

A natural question arising from this construction is whether the intrinsic metric coincides with the original metric. This is exactly the notion of *length space*.

Definition B.3 (Length spaces). X is called a *length space* if and only if $d_X = \widehat{d}_X$.

Remark B.4 (On showing that a metric space is a length spaces). Note that, by the triangle inequality, $d_X \leq \widehat{d}_X$. Hence, showing that a metric space is a length space usually comes down to showing that, for all $(x, x') \in X^2$ and all $\varepsilon > 0$ (resp. $\kappa > 1$), there exists a curve $c \in \mathcal{AC}(I, X)$ connecting x and x' such that $L_X(c) \leq d_X(x, x') + \varepsilon$ (resp. $L_X(c) \leq \kappa d_X(x, x')$).

If the infimum is a minimum, X is called a *geodesic space*.

Definition B.5 (Geodesic spaces). X is called *geodesic space* if and only if for all $(x, x') \in X^2$ there exists $c \in \mathcal{AC}(I, X)$ connecting x and x' such that $d_X(x, x') = L_X(c)$.

Curves minimizing the length between their endpoints are called *minimizing geodesics* of X and can be reparametrized as *constant speed geodesic*, that is, as curves $c : I \rightarrow X$ satisfying for all $(s, t) \in I^2$ that

$$d_X(c(s), c(t)) = \left| \frac{s-t}{b-a} \right| d_X(c(a), c(b)).$$

Note that a constant speed geodesic is necessarily a minimizing geodesic for its endpoints. Now, let us recall the definition of the p -energy of a curve.

Definition B.6 (Energy of a curve). Let $p \in [1, \infty)$. Then, the p -energy of a curve $c \in \mathcal{AC}^p(I, X)$ is defined as

$$E_{p,X}(c) := \int_I |c'|_X^p(t) dt.$$

The 1-energy being the length, that is, $L_X(c) = E_{1,X}(c)$.

Then, the infimum of the length over $\mathcal{AC}(I, X)$ can be shown to be equal to the infimum over $\mathcal{AC}^p(I, X)$ of the p -th root of the p -energy for all $p \geq 1$.

Proposition B.7 (Energy minimizing distance). Let $p \in [1, \infty)$ and $(x, x') \in X^2$. Then,

$$\widehat{d}_X(x, x') = \inf \left\{ \mathcal{L}^1(I)^{1-1/p} E_{p,X}(c)^{1/p} : c \in \mathcal{AC}^p(I, X), c(a) = x, c(b) = x' \right\}.$$

Proof. The proof follows the same approach as the proof of (i) in [Tro95, Theorem 3.7] or [PT24, Proposition 3.8], which consists in finding a constant-speed approximation for each absolutely continuous curve.

Let $(x, x') \in X^2$ be such that there exists $c \in \mathcal{AC}(I, X)$ connecting x and x' , and $\varepsilon > 0$. Then, define $s_\varepsilon : I \rightarrow I$ as

$$s_\varepsilon(t) := \frac{\int_a^t |c'|_X(u) du + (t-a)\varepsilon \mathcal{L}^1(I)^{-1}}{\int_I |c'|_X(u) du + \varepsilon} \mathcal{L}^1(I).$$

Its inverse map t_ε is \mathcal{L}^1 -a.e. differentiable, with derivative given, for \mathcal{L}^1 -a.e. $s \in I$, by

$$\dot{t}_\varepsilon(s) = \frac{\mathcal{L}^1(I)^{-1} (\int_I |c'|_X(u) du + \varepsilon)}{|c'|_X(t_\varepsilon(s)) + \varepsilon \mathcal{L}^1(I)^{-1}} > 0.$$

Now, setting $\tilde{c} := c \circ t_\varepsilon$, we have, for \mathcal{L}^1 -a.e. $s \in I$,

$$\begin{aligned} |\tilde{c}'|_X(s) &= \lim_{t \rightarrow s} \frac{d_X(c \circ t_\varepsilon(s), c \circ t_\varepsilon(t))}{|t_\varepsilon(s) - t_\varepsilon(t)|} \frac{|t_\varepsilon(s) - t_\varepsilon(t)|}{|s - t|} \\ &= |c'|_X(t_\varepsilon(s)) \dot{t}_\varepsilon(s) \leq \mathcal{L}^1(I)^{-1} \left(\int_I |c'|_X(u) du + \varepsilon \right). \end{aligned}$$

Therefore, for all $p \in [1, \infty)$, $\tilde{c} \in \mathcal{AC}^\infty(I, X) \subset \mathcal{AC}^p(I, X)$ and, by Hölder's inequality,

$$L_X(c) = L_X(\tilde{c}) \leq \mathcal{L}^1(I)^{1-1/p} E_{p,X}(\tilde{c})^{1/p} \leq L_X(c) + \varepsilon.$$

Since ε is arbitrary, this yields the conclusion of [Proposition B.7](#). \square

Remark B.8 (On showing that a metric space is length space with the energy). Thus, showing that X is a length space also comes down to showing that, for all $(x, x') \in X^2$ and all $\varepsilon > 0$ (resp. $\kappa > 1$), there exists a curve $c \in \mathcal{AC}^p(I, X)$ connecting x and x' such that $\mathcal{L}^1(I)^{p-1} E_{X,p}(c) \leq d_X(x, x')^p + \varepsilon$ (resp. $\mathcal{L}^1(I)^{p-1} E_{X,p}(c) \leq \kappa d_X(x, x')^p$).

An important consequence of this result is that, when X is a geodesic space, a curve minimizes the p -energy, $p \in [1, \infty)$, if and only if it is a constant speed geodesic.

Appendix B.2. Measurable multifonctions

We recall below the statement of Aumann's measurable selection theorem [CVCV77, III-§4-28-2)] (see [AB06, Corollary 18.27] for a more recent exposition).

Theorem B.9 (Aumann's measurable selection theorem). *Let (O, Σ_O) be a measurable space, (X, d_X) be complete and separable metric space. Let Γ be a multifunction from P to non-empty subsets of X , whose graph G belongs to $\Sigma_O \otimes \mathcal{B}(X)$. Then, there exists a $(\hat{\Sigma}_O, \mathcal{B}(X))$ -measurable mapping S such that $S(o) \in \Gamma(o)$ for all $o \in O$, where $\hat{\Sigma}_O$ denotes the intersection of all completions of Σ_O with respect to each finite measure on (O, Σ_O) . Such mapping S is called $(\hat{\Sigma}_O, \mathcal{B}(X))$ -measurable selection.*

Appendix B.3. Differentiable manifolds

This section briefly recalls the construction of differentiable manifolds modelled on Banach spaces (see [Lan12, Chapter II] for a complete exposition on that matter).

Let us first recall a definition of *atlases* by paraphrasing [Lan12, Chapter II, §1].

Definition B.10 (Atlases). Let $r \in \mathbb{N} \cup \{\infty\}$. An *atlas of class C^r* on a set X is a family of pairs $\{(U_\alpha, \varphi_\alpha)\}_{\alpha \in A}$ indexed by a set A satisfying the following conditions:

- (i) each U_α is a subset of X and the U_α 's cover X .
 - (ii) each φ_α is a bijection from U_α onto an open subset $\varphi_\alpha(U_\alpha)$ of some Banach space \mathbb{B}_α and for all $(\alpha, \alpha') \in A^2$ the set $\varphi_\alpha(U_\alpha \cap U_{\alpha'})$ is open in \mathbb{B}_α .
 - (iii) the mapping $\varphi_\alpha \circ \varphi_{\alpha'}^{-1} : \varphi_{\alpha'}(U_\alpha \cap U_{\alpha'}) \rightarrow \varphi_\alpha(U_\alpha \cap U_{\alpha'})$ is a C^r isomorphism for each pair of indices $(\alpha, \alpha') \in A^2$.
- Two atlases $\{(V_\beta, \psi_\beta)\}_{\beta \in B}$ and $\{(U_\alpha, \varphi_\alpha)\}_{\alpha \in A}$ on X indexed by some sets A and B are said to be *compatible* if each pair (V_β, ψ_β) is such that the mapping $\varphi_\alpha \circ \psi_\beta^{-1} : \psi_\beta(U_\alpha \cap V_\beta) \rightarrow \varphi_\alpha(U_\alpha \cap V_\beta)$ is a C^r isomorphism for all pair $(U_\alpha, \varphi_\alpha)$ and conversely. This relation defines an equivalence relation between atlases on X .

Remark B.11 (On topologies induced by atlases). Note that an atlas uniquely defines a topology in which the charts $\{(U_\alpha, \varphi_\alpha)\}_{\alpha \in A}$ are such that the U_α 's are open sets and the φ_α 's are topological isomorphisms. This induced topology is not required to be Hausdorff.

Using the equivalence relation of compatibility between atlases, *Banach manifolds* are then defined as follows:

Definition B.12 (Banach manifolds). Let $r \in \mathbb{N} \cup \{\infty\}$. A *C^r Banach manifold* is a set X together with an equivalence class of atlases of class C^r (Definition B.10) on this set.