

Low-dimensional geometry learning for turbulence prediction in optimized stellarators

Xishuo Wei^{1*}, Handi Huang¹, Haotian Chen¹, Hongxuan Zhu^{2,3}, Zhe Bai⁴, Samuel Williams⁴, Zhihong Lin^{1*}

1. University of California-Irvine, Irvine, CA, United States of America

2. Princeton University, Princeton, NJ, United States of America

3. Zhejiang University, Zhejiang, China

4. Lawrence Berkeley National Laboratory, Berkeley CA, United States of America

*Email: xishuow@uci.edu, zhihongl@uci.edu

Abstract

The optimized stellarator is an attractive concept for which the averaged particle radial drift is zero, and the single particle loss can be significantly reduced. But for the reactor design, global physics such as turbulent transport also need to be optimized besides the confined single particle orbit, or properties estimated using local estimations and heuristic formulations. The first-principle global transport code is too computationally expensive to integrate into the optimization process. The fast surrogate global transport model based on machine learning is a good alternative choice, but the amount of data required to train the surrogate model is numerous due to the high degree-of-freedom of the stellarator design. The work shows that the stellarator design with quasi-helically(QH) symmetric geometry is approximately distributed in a low dimensional latent space, which can be explicitly found by deep learning. This discovery makes it possible to generate global gyrokinetic simulation data for training surrogate models to directly optimize the stellarator geometry for turbulent transport, energetic particle instability, and MHD modes. Using the low dimensional latent space and data analysis methods, the relation between linear zonal residues and axis-exursion is found, providing a simple guide to optimize low turbulent transport QH stellarators.

1. Introduction

Quasi-symmetric (QS) stellarators are attractive because they approximately recover an approximate symmetry of the magnetic field strength in appropriate magnetic coordinates. The symmetry implies a conserved quantity for guiding center analog to the canonical toroidal momentum, which substantially improves single-particle confinement and reduces neoclassical losses[1], [2]. In practice, QS configurations are typically obtained through numerical optimization, where a predefined objective function is minimized by iteratively adjusting design parameters that determine the equilibrium magnetic flux surfaces. Using this general strategy, previous studies have successfully constructed QS configurations and other optimized concepts such as quasi-isodynamic configurations[2], [3], [4], [5], [6]. Morden stellarator optimization rarely targets symmetry alone. Optimization commonly imposes basic shaping constraints (for example, aspect ratio and elongation), rotational transform (ι) profile, and many additional physical objectives incorporated through computationally inexpensive proxies. Examples include Mercier criterion for MHD interchange stability, effective ripple as a proxy for neo-classical transport in low collisional regime. The recent W7-X experiments [7], [8] have confirmed that the neoclassical transport can be reduced through magnetic-field optimization, and the core energy confinement is determined by turbulence instead. These results motivate the turbulence stellarator optimization aimed at future reactors. Over the years, many results have used different proxies to minimize the linear drive and nonlinear transport of micro-instabilities.[5], [9], [10], [11], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21]. Notably, Kim et al

[22] has directly optimized the local gyrokinetic simulated heat flux, while Landreman et al [23] trained the surrogate model from the local simulations to predict the heat flux, which can also be used for optimization.

A limitation of using the proxies and local simulation for optimization is that the underlying global nonlinear behavior can be qualitatively different from what simplified metrics suggest. For example, the recent nonlinear MHD simulations indicate that the linearly unstable ballooning activity in standard W7-X EIM configuration can saturate benignly, consistent with the “soft” beta limit rather than a hard disruptive limit [24]. Regarding microturbulence, global gyrokinetic simulations in stellarators have reported regimes in which TEM excited by helically trapped electrons can drive substantial transport in W7-X[25], and the different interaction between zonal flows and turbulence can impose a major impact on the steady-state transport level for different stellarator configurations[26]. These findings urge the consideration of first-principle global simulations during optimization. However, these simulations are typically orders of magnitude more expensive than proxy evaluations, making them difficult to embed directly in the iterative optimization loop. Machine learning surrogate models, on the other hand, provide a potential path forward because they can deliver rapid inference and can be integrated with automatic differentiation in modern frameworks, enabling gradient-based design and optimization workflows. The key obstacle is data. Stellarator optimization problems often involve hundreds to thousands of free parameters, which define a very high-dimensional design space. It is not feasible to populate this space with enough first-principle simulations to train a globally reliable surrogate model. And even if a surrogate model is trained on a limited dataset, a fundamental question is why the surrogate can be expected to generalize for new configurations.

A useful observation is that it is not necessary to cover the entire design space. Physically meaningful stellarators occupy a constrained subset because poor single-particle confinement is unacceptable regardless of other performance metrics. Concepts such as QS and QI impose strong geometric constraints, which can restrict the relevant configurations to a small fraction of the nominal parameter space and can dramatically reduce the intrinsic dimensionality of the realizable designs. Related low dimensional structure has been reported in earlier stellarator design studies, for example in global searches over coil current settings [27]. The near-axis-expansion theory itself is an approach to express the configurations approximately with few parameters. The internal kink mode structure is found to have the low-dimensional feature and can be approximately described by 3 parameters [28]. It is also consistent with the broader idea that high-dimensional datasets can concentrate near low-dimensional manifolds, which motivates latent variable representations such as autoencoders[29]. Actually, the low dimensional feature is very common for naturally generated data, such as images and languages[30], [31], [32]

In this paper, we focus on quasi-helical symmetric (QH) configurations and show that their geometric distribution has an intrinsic dimension that is far smaller than the total number of free parameters used to represent the boundary shape. Using a dataset of more than 13,000 QH equilibria, we train an autoencoder to reconstruct the geometry from a low-dimensional latent representation and find that a three-dimensional latent space is sufficient to capture the essential variation of the QH geometries. Building on these latent representations, we construct surrogate models for two turbulence-relevant quantities, namely the zonal-flow residual level in the long wavelength limit [33], and the volume-averaged steady-state heat flux driven by ITG turbulence. We further show that both the zonal-flow residual and the simulated transport exhibit organized structure when projected into the learned three-dimensional latent space, enabling efficient exploration of the QH subspace and the identification and generation of candidate low-transport configurations.

The remainder of this paper is organized as follows. Section 2 describes the data generation, the dimensionality-reduction method, and model architecture. Section 3 reports results for geometry reconstruction, zonal-flow

residual prediction, and turbulent transport prediction. Section 4 summarizes the conclusions and outlines future directions.

2. Methods




Figure 1 Histogram of the QS error of the generated equilibria. The 13949 equilibria with QS error lower than 0.1 are selected to form the dataset.

We used the optimization code DESC[34], [35] to generate a dataset of QH stellarator equilibria with toroidal period NFP=4. The initial parameters were selected to ensure sufficient diversity in geometry distribution. The initial flux surface cross section was set to be concentric circular, with the inverse aspect ratio ϵ randomly sampled from [0.1, 0.25]. The magnetic axis was set to be parameterized as $R_{axis} = 1 + \xi \cos \phi$, $Z_{axis} = \xi \sin \phi$, where $\xi \in [0, 0.2]$ and ϕ is the toroidal angle in the cylindrical coordinates. The ι profile was prescribed by $\iota(\rho) = \iota_0 + 0.1\rho^2$, where ι_0 is randomly selected from [0.35, 1.85]. The pressure profile was prescribed by $p(\rho) = \left((-0.5 \tanh((\rho - p_1)/p_2) - 0.5) \times p_3 + 1 \right) \times p_0$, with p_0 selected from $[10^3, 3 \times 10^4]$, p_1 selected from [0.2, 0.8], p_2 selected from [0.2, 0.5], and p_3 selected from [0.2, 0.95]. The toroidal magnetic flux of last closed flux surface was fixed at $0.04 \times 2\pi$ Wb. Then each randomly generated configuration was optimized by minimizing the ‘two-term’ quasi-symmetry error

$$\hat{f}_C = \langle |(M\iota - N)(\mathbf{B} \times \nabla\psi) \cdot \nabla B - (MG + NI)\mathbf{B} \cdot \nabla B| \rangle / \langle B \rangle^3,$$

where $M = 1$, $N = 4$. ψ is the toroidal flux function. I and G are the poloidal and toroidal components of the covariant form of \mathbf{B} using Boozer coordinates. $I = \mathbf{B} \cdot \partial\mathbf{r}/\partial\theta_B$, $G = \mathbf{B} \cdot \partial\mathbf{r}/\partial\zeta_B$, where θ_B and ζ_B are the poloidal and toroidal Boozer angle. The bracket $\langle \ \rangle$ means volume averaging. During the optimization, ϵ is constrained to [0.1, 1/3], and the maximum elongation was limited to [1, 3]. The boundary toroidal flux, the pressure profile, and the averaged radial position of boundary were fixed. In addition, the force balance of these equilibria was required by minimizing the relative force balance error,

$$|F| = |J \times B - \nabla p| / \langle |\nabla B|^2 / (2\mu_0) \rangle$$

In order to emphasize the geometrical effect on zonal flows and turbulence, two groups of datasets with fixed $\iota_0 = 0.55$ and $\iota_0 = 1.35$ were also generated, with all other settings the same as the random ι_0 dataset. Since our purpose is to find the hidden patten of QS geometries, the obtained configurations with $\hat{f}_C > 0.1$ or $\langle |F| \rangle > 0.1$ were discarded, while the ones with good symmetry were kept in the dataset. The QS error distribution is shown in Fig.1. 2995 equilibria are generated with random ι_0 , 5197 equilibria with $\iota_0 = 0.55$, and 5757 equilibria with $\iota_0 = 1.35$. Each equilibrium is represented by the coefficients (R_{lmn}, Z_{lmn}) that describe the flux surface shape,

$$R(\rho, \theta, \phi) = \sum_{lmn} R_{lmn} Z_l^m(\rho, \theta) \mathcal{F}^n(\phi)$$

$$Z(\rho, \theta, \phi) = \sum_{lmn} Z_{lmn} Z_l^m(\rho, \theta) \mathcal{F}^n(\phi),$$

Where Z_l^m are the Fourier-Zernike basis functions and \mathcal{F}^n are the Fourier basis functions. (l, m, n) stand for the order of basis function in radial, poloidal and toroidal directions. In the current dataset, the basis functions are truncated at maximum order 8 in radial, poloidal, and toroidal directions, and the number of coefficients for each equilibrium is 765. In other words, the design space of these geometries has a dimension of 765.




Figure 2 The Pearson coefficient of the 250 (R,Z) coefficients with largest averaged absolute value.




Figure 3 Reconstruction by PCA. (Left) Dependence of Experienced Variational Ratio of all (R, Z) coefficients on the PCA dimensions. (Right) A example of the reconstructed last closed flux surface with 11-dimensional PCA compared with the ground truth data generated with DESC.

We first evaluate the linear properties of the data distribution. The Pearson correlation coefficients of the top 250 coefficients with the largest mean absolute values were calculated and shown in Fig.2. It is shown that most large coefficients have a strong correlation around ± 0.5 , with the largest magnitudes up to ± 0.8 . Note that the Pearson correlation only shows the linear relation among the parameters, which suggests the coefficients can be subject to even stronger nonlinear constraints, and a significant dimensionality reduction is feasible. Principal Component Analysis (PCA) was applied to evaluate the linear dimensionality reduction. The dependence of the Explained Variance Ratio (EVR) reaches unity in an 11-dimensional subspace, as shown in Fig.3. Using this 11-d space, the (R, Z) coefficients and the associated flux surface shapes can be reconstructed. Although the reconstructed flux surface shapes resemble the original optimized geometries from DESC, the detailed calculations show substantial deviation in $|B|$ and the associated QS error. Furthermore, the 11-d space remains too large for efficient generation of simulation data and surrogate model training.




Figure 4 (a) Autoencoder architecture. (b) Comparison of collocation point positions calculated from DESC coefficients and reconstructed coefficients at different toroidal angles.

To achieve further dimensionality reduction, we constructed a fully connected autoencoder neural network. The input and the output consist of the 765-dimensional (R, Z) coefficient vector (noted by \vec{C}). The encoder and decoder each contain four hidden layers with the ReLU activation function. A low-dimensional latent layer is located at the bottleneck. The primary loss term is the mean-squared reconstruction error of the coefficients,

$$L_{MSE} = \sqrt{\sum_{i=1,765} (C_{i,output} - C_{i,input})^2} / 765. \text{ That is, the autoencoder is trained to ensure the coefficients to}$$

be effectively compressed from the original 765-d space to the low dimensional latent space. To improve the geometric reconstruction accuracy rather than just the coefficients, we incorporated an additional loss term based on collocation points. Specifically, $20 \times 16 \times 17$ uniformly distributed grid points in (ρ, θ, ϕ) space were used to calculate the value of (R, Z, ϕ) in the cylindrical coordinates by substituting the ground truth (R, Z) coefficients and reconstructed coefficients in Eq (1). The average distance was included in the loss function, $L_{points} =$

$$\sum_{k=1,5440} \sqrt{(R_{k,DESC} - R_{k,pred})^2 + (Z_{k,DESC} - Z_{k,pred})^2} / 5440. \text{ The collocation points calculated from DESC}$$

coefficients and from the reconstructed coefficients can be seen from Fig4(b). The final loss function is $L = L_{MSE} + wL_{points}$. In the first 20 epochs during training, only the coefficient loss was used ($w=0$), which makes the training process quickly converges. Subsequently, w was set to be 0.25, and the collocation point loss was introduced to improve the accuracy of the geometric reconstruction.

3. Results




Figure 5 The reconstruction error dependence on latent space dimensions. In the lower two panels, the models trained on different groups of data are tested.

The latent space dimension was varied, and the reconstruction error L_{MSE} on the test dataset was calculated to determine the minimum intrinsic dimensionality of the dataset. Shown in Fig. 5, the reconstruction error increases sharply when the latent dimension is smaller than 3, indicating that the dataset lies approximately on a three-dimensional nonlinear manifold. To be noticed, the models trained on fixed or random t_0 datasets generalize well across different t_0 values, suggesting that the QH geometric distribution is largely independent of rotational transform profile. An example of the reconstructed equilibrium can be seen in Fig. 6. The left panel shows the reconstruction of (R, Z) coefficients, and the right panel shows the reconstruction of the last closed flux surface. Compared with PCA, the autoencoder achieves significantly higher geometric reconstruction accuracy. Another test shows that removing the collocation points loss degrades flux surface reconstruction, although the coefficients reconstruction stays relatively accurate. On the other hand, only using the collocation loss prevents the convergence of the training process of the autoencoder.




Figure 6 An example of (R, Z) coefficients (left panel) and reconstructed last closed flux surface shape (right panel) compared to the DESC data. Latent space dimension is 3.

The results above mean that the flux surface information can be compressed to very few parameters, which are the coordinates of the latent space. Together with the t profile, the whole equilibrium and the consequent physical quantities determined by the geometric parameters can be inferred from the latent space coordinates. The linear zonal flow (ZF) residual level is one important example of these physical quantities. The study of turbulence transport on different stellarator configurations showed that the ZF residual level can play an important role in determining the nonlinear steady-state ZF level and hence the transport level. Generally, ZF residual level can be characterized by its limit when $k_{\perp}\rho_i$ approaches 0, i.e., the Rosenbluth-Hinton (RH) level. In this sense,

finding the geometrical factors that affect the RH level and a way to quickly predict the RH level is very important to find transport-optimized designs. For this purpose, we used the latent space representation to directly predict the volume averaged RH level. The pre-trained encoder obtained in the reconstruction task maps the coefficients into latent space, followed by 3 fully connected layers with the width (64, 64, 32) for the regression of the RH level. The inputs are still the 765-d coefficients vector. The ground-truth RH level for the training data is calculated using the formula in [36],

$$RH(\psi) = \frac{1}{1 + \Lambda_1/\Lambda_0},$$

$$\Lambda_1 = \frac{m_i n_i (G + NI)^2}{B_0^2 (\iota - N)^2} [1.6\epsilon^{3/2} + \mathcal{O}(\epsilon^2)],$$

$$\Lambda_0 = m_i n_i \langle |\nabla\psi|^2 / B^2 \rangle.$$

Where N is the toroidal period, which is 4 for this dataset. ψ is the toroidal flux function. Here ϵ is a function of flux function which can be calculated from $\epsilon = (B_{max} - B_{min}) / (B_{max} + B_{min})$ on the flux surface. Then the RH levels on different flux surfaces are averaged to get a single value as the ground truth value of the output. Since the ι profile will significantly affect the RH level, we only constructed the surrogate model for the two fixed ι datasets to highlight the geometrical effects. As a result, the predicted RH level agrees well with the theoretical values, which can be seen from Fig.7. For a 3-d latent space, the average relative error is 15% for $\iota_0 = 1.35$ test dataset, and 2.5% for $\iota_0 = 0.55$ test dataset. If we increase the latent space to 7-d, the error reduces to 2.3% for $\iota_0 = 1.35$ test dataset, and 2.2% for $\iota_0 = 0.55$ test dataset.




Figure 7 Prediction of RH level compared to the theoretical values. The $y=x$ line means perfect prediction. (Left) $\iota = 1.35$ (Right) $\iota = 0.55$

The distribution of the RH levels of the training data in 3-d latent space is shown in Fig.8. By analyzing the data in different regions, one important geometric factor is the aspect ratio, which directly influences both $\nabla\psi/B$ term and ϵ term in Eq(1). However, even for equilibria with similar aspect ratio and elongation, significant variation in RH level is observed. This variation correlates with the magnetic axis excursion, defined as the amplitude of the magnetic axis displacement in the R-Z plane as ϕ changes. For configurations with similar aspect ratio and identical ι profile, equilibria with small axis excursion exhibit systematically higher RH levels. A comparison of such an equilibrium pair can be seen in Fig.8. This trend can be interpreted qualitatively using the first order near-axis expansion theory. The Λ_1 term in Eq(1) is constant on the flux surface, so the RH level variation on the same flux surface relies on the variation of Λ_0 . It is found that, due to the large toroidal variation of the Jacobian, the flux averaging of Λ_0 is predominantly related to its value at $\zeta = 0$ (bean shape cross section). From [36], The RH level at $\zeta = 0$ can be written as

$$RH(\zeta = 0) = \frac{1}{1 + 1.6 / (\sqrt{\varepsilon} \mathcal{C} (\iota - N)^2)},$$

$$\mathcal{C} = \alpha / (\bar{\eta} R_a)^2, \alpha = \langle |\nabla \psi|^2 / B^2 \rangle,$$

$$\bar{\eta} = \sqrt{2\psi / B} / \varepsilon,$$

Where R_a stands for the total magnetic axis length over one toroidal period. For equilibria with similar aspect ratio and identical ι profile, the factors α and $\bar{\eta}$ factor vary weakly. Consequently, the magnetic axis length R_a becomes the dominant geometric factor controlling \mathcal{C} , and hence the RH level. Smaller axis excursion corresponds to shorter magnetic axis length, which increases \mathcal{C} and therefore increases the RH level. This interpretation provides a clear physical explanation for the latent space structure observed in the data. The autoencoder has effectively identified a low-dimensional representation in which the dominant geometric control parameters for zonal flow residuals are embedded.




Figure 8 (left) The theoretical RH levels of $\iota_0 = 0.55$ training data in the 3-d latent space. (right) A comparison of two equilibria with similar aspect ratio and elongations, but different axis excursions.

To systematically evaluate the turbulent transport properties across the generated QH stellarator database, more than 15,000 global GTC gyrokinetic simulations were performed. All equilibria were resized to isolate the geometric effects. The minor radius was set to $a = 0.8$ m, and the averaged on-axis B strength was fixed at $B_a = 4.9$ T. The ion and electron density profile are set to be uniform with $n_0 = 8.3 \times 10^{13} \text{ cm}^{-3}$. The ion and electron temperature profiles were set to have the largest gradient near $\rho = 0.5$ with the minimum scale length being $a/L_{Ti} = 4.7$. The on-axis temperatures were set to $T_{i0} = T_{e0} = 4$ keV. This set of parameters can destabilize the ITG mode in most cases. The resulting steady-state turbulent transport varied significantly across configurations. Excluding the cases in which ITG mode was linearly stable, the steady-state ion heat conductivity ranged from approximately $10^{-6} D_B$ to $10^{-4} D_B$, where $D_B = T_{e0} / e B_a$ is the Bohm unit. This broad distribution suggests the strong sensitivity of turbulent transport to magnetic geometry within the QH parameter space. A systematic trend is observed that the configurations with smaller ι_0 generally show lower ion heat conductivity, consistent with the previous conclusion in [26], [36]. When the steady-state transport levels are projected into the three-dimensional latent space identified by the autoencoder, recognizable structure emerges. Although the clustering pattern is less clear than in the RH level analysis, regions associated with reduced transport can still be identified. Taking advantage of this property, we constructed a simple generative model in latent space to explore the transport-optimized designs. A Gaussian mixture model was fitted to equilibria satisfying $\chi_i < 10^{-5} D_B$. New latent coordinates were sampled from the fitted distribution in the 3-d space. Then the data were mapped back to the full geometric coefficients using the trained decoder. It is confirmed by the GTC simulations that the new

samples do have a distribution with a lower average transport level compared to the randomly generated original dataset, which is particularly pronounced for the $\iota_0 = 0.55$ dataset.




Figure 9 Distribution of the ion heat conductivity of different dataset.




Figure 10 Prediction of volume averaged ion heat flux driven by ITG compared with first principle GTC simulations.

Beyond qualitative identification of low-transport regions, a quantitative surrogate model was developed to predict the steady-state transport level for QH geometries. The neural network architecture is the same as that used for RH level prediction, with the output replaced by the steady state χ_i/D_B value for each case. If the encoder weights are frozen and only the regression layers are trained, like the RH case, the prediction accuracy of χ_i remains poor. This indicates that the latent space optimized for geometric reconstruction does not necessarily preserve the geometric features most relevant for nonlinear transport. The best performance was obtained through a two-stage training strategy. First, the regression layers are trained for 30 epochs while keeping the encoder fixed. Then the encoder is fine-tuned jointly with the regression layers using a reduced learning rate. This fine-tuning allows the latent representation to adjust slightly so that the transport-relevant geometric features are better resolved. After fine-tuning, the predicted transport levels agree reasonably with the first-principle GTC results, particularly for the $\iota_0 = 0.55$ dataset. The comparison between the predicted and simulated ion heat conductivity is shown in Fig. 10. For $\iota_0 = 0.55$, the coefficient of determination R^2 approaches 0.9, indicating strong predictive capability. In contrast, the $\iota_0 = 1.35$ dataset exhibits lower prediction accuracy and larger scatter. The difference between the two datasets is also reflected in the χ_i of the

sampled data in Fig.9, where it is easier to sample out the low transport cases for $t_0 = 0.55$ dataset. These results highlight two properties of the distribution of transport levels. First, while the geometric manifold is intrinsically low-dimensional, nonlinear turbulence transport depends on more subtle geometric features than those required for accurate flux surface reconstruction. Second, moderate fine-tuning of latent representation significantly improves transport prediction without sacrificing geometric interpretability.

4. Conclusion and future work

This study demonstrates that the QH stellarator geometries exhibit intrinsic dimensionality as low as 3, and hence it is possible to construct a surrogate model to predict the global features of the stellarator with $\sim 10k$ data points due to this feature. The RH zonal flow residual can be predicted accurately from the coordinates of the latent space, which may have an impact on future stellarator designs. A direct relationship between the RH level and the magnetic axis excursion is identified and supported by the near-axis expansion theory. Furthermore, the large-scale first principle GTC global turbulence simulation confirms significant variation in transport properties across different QH geometries. This dataset can be very helpful for future optimized stellarator designs. For example, the latent space together with the simulation data can be used to directly sample new geometries with low transport level while keeping the QH symmetry. Although the original latent space trained from reconstruction is not appropriate for the quantitative prediction of transport level, it can be fine-tuned to improve the prediction accuracy. Although the prediction is not perfect, this work can be a proof of concept using dimensionality reduction and surrogate models to optimize the global features of stellarators.

Future work will focus on improving the prediction accuracy of the transport level through simulation data cleaning, hyperparameter search, and ensemble learning with diverse network architectures. Integration of the transport surrogate model into the optimization framework will allow simultaneous optimization of quasi-symmetry, turbulence, and other properties. Extension of the methodology to QA and QI geometries will also be investigated. Finally, we will broaden the usage of the autoencoder in stellarator optimization, such as geometry generation and PINN from the latent space.

Acknowledgement

This work is supported by the US Department of Energy (DOE) SciDAC project HifiStell. An award of computer time was provided by the U.S. Department of Energy's (DOE) Innovative and Novel Computational Impact on Theory and Experiment (INCITE) Program. This research used supporting resources at the Argonne and the Oak Ridge Leadership Computing Facilities. The Argonne Leadership Computing Facility at Argonne National Laboratory is supported by the Office of Science of the U.S. DOE under Contract No. DE-AC02-06CH11357. The Oak Ridge Leadership Computing Facility at the Oak Ridge National Laboratory is supported by the Office of Science of the U.S. DOE under Contract No. DE-AC05-00OR22725. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a Department of Energy User Facility (project m306-2024). This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Reference

- [1] A. H. Boozer, “Transport and isomorphic equilibria,” *The Physics of Fluids*, 1983, doi: 10.1063/1.864166.
- [2] M. Landreman and E. Paul, “Magnetic Fields with Precise Quasisymmetry for Plasma Confinement,” *Phys. Rev. Lett.*, vol. 128, no. 3, p. 035001, Jan. 2022, doi: 10.1103/PhysRevLett.128.035001.
- [3] J. Nührenberg and R. Zille, “Quasi-helically symmetric toroidal stellarators,” *Physics Letters A*, vol. 129, no. 2, pp. 113–117, May 1988, doi: 10.1016/0375-9601(88)90080-1.
- [4] P. Helander and J. Nührenberg, “Bootstrap current and neoclassical transport in quasi-isodynamic stellarators,” *Plasma Phys. Control. Fusion*, vol. 51, no. 5, p. 055004, Feb. 2009, doi: 10.1088/0741-3335/51/5/055004.
- [5] A. G. Goodman *et al.*, “Quasi-Isodynamic Stellarators with Low Turbulence as Fusion Reactor Candidates,” *PRX Energy*, vol. 3, no. 2, p. 023010, Jun. 2024, doi: 10.1103/PRXEnergy.3.023010.
- [6] D. W. Dudt, A. G. Goodman, R. Conlin, D. Panici, and E. Kolemen, “Magnetic fields with general omnigenity,” *Journal of Plasma Physics*, vol. 90, no. 1, p. 905900120, Feb. 2024, doi: 10.1017/S0022377824000151.
- [7] C. D. Beidler *et al.*, “Demonstration of reduced neoclassical energy transport in Wendelstein 7-X,” *Nature*, vol. 596, no. 7871, pp. 221–226, Aug. 2021, doi: 10.1038/s41586-021-03687-w.
- [8] O. P. Ford *et al.*, “Turbulence-reduced high-performance scenarios in Wendelstein 7-X,” *Nucl. Fusion*, vol. 64, no. 8, p. 086067, Aug. 2024, doi: 10.1088/1741-4326/ad5e99.
- [9] C. C. Hegna, P. W. Terry, and B. J. Faber, “Theory of ITG turbulent saturation in stellarators: Identifying mechanisms to reduce turbulent transport,” *Phys. Plasmas*, vol. 25, no. 2, p. 022511, Feb. 2018, doi: 10.1063/1.5018198.
- [10] M. Nunami, T.-H. Watanabe, and H. Sugama, “A reduced model for ion temperature gradient turbulent transport in helical plasmas,” *Physics of Plasmas*, 2013, doi: 10.1063/1.4822337.
- [11] Mynick, Pomphrey, and Xanthopoulos, “Optimizing Stellarators for Turbulent Transport,” *Physical Review Letters*, 2010, doi: 10.1103/PhysRevLett.105.095004.
- [12] J. H. E. Proll, H. E. Mynick, P. Xanthopoulos, S. A. Lazerson, and B. J. Faber, “TEM turbulence optimisation in stellarators,” *Plasma Phys. Control. Fusion*, vol. 58, no. 1, p. 014006, Jan. 2016, doi: 10.1088/0741-3335/58/1/014006.
- [13] P. Xanthopoulos *et al.*, “Controlling Turbulence in Present and Future Stellarators,” *Phys. Rev. Lett.*, vol. 113, no. 15, p. 155001, Oct. 2014, doi: 10.1103/PhysRevLett.113.155001.
- [14] R. J. J. Mackenbach, J. H. E. Proll, and P. Helander, “Available Energy of Trapped Electrons and Its Relation to Turbulent Transport,” *Phys. Rev. Lett.*, vol. 128, no. 17, p. 175001, Apr. 2022, doi: 10.1103/PhysRevLett.128.175001.
- [15] R. Jorge *et al.*, “Direct microstability optimization of stellarator devices,” *Phys. Rev. E*, vol. 110, no. 3, p. 035201, Sep. 2024, doi: 10.1103/PhysRevE.110.035201.
- [16] P. Helander, J. H. E. Proll, and G. G. Plunk, “Collisionless microinstabilities in stellarators. I. Analytical theory of trapped-particle modes,” *Physics of Plasmas*, vol. 20, no. 12, p. 122505, Dec. 2013, doi: 10.1063/1.4846818.
- [17] J. H. E. Proll, P. Helander, J. W. Connor, and G. G. Plunk, “Resilience of Quasi-Isodynamic Stellarators against Trapped-Particle Instabilities,” *Phys. Rev. Lett.*, vol. 108, no. 24, p. 245002, Jun. 2012, doi: 10.1103/PhysRevLett.108.245002.
- [18] E. Sánchez, J. L. Velasco, I. Calvo, and S. Mulas, “A quasi-isodynamic configuration with good confinement of fast ions at low plasma β ,” *Nucl. Fusion*, vol. 63, no. 6, p. 066037, May 2023, doi: 10.1088/1741-4326/accd82.
- [19] J. H. E. Proll *et al.*, “Turbulence mitigation in maximum-J stellarators with electron-density gradient,” *J. Plasma Phys.*, vol. 88, no. 1, p. 905880112, Feb. 2022, doi: 10.1017/S002237782200006X.
- [20] S. Stroteich, P. Xanthopoulos, G. Plunk, and R. Schneider, “Seeking turbulence optimized configurations for the Wendelstein 7-X stellarator: ion temperature gradient and electron temperature gradient turbulence,” *Journal of Plasma Physics*, vol. 88, no. 5, p. 175880501, Oct. 2022, doi: 10.1017/S0022377822000757.
- [21] G. T. Roberg-Clark, G. G. Plunk, P. Xanthopoulos, C. Nührenberg, S. A. Henneberg, and H. M. Smith, “Critical gradient turbulence optimization toward a compact stellarator reactor concept,” *Phys. Rev. Res.*, vol. 5, no. 3, p. L032030, Sep. 2023, doi: 10.1103/PhysRevResearch.5.L032030.

- [22] P. Kim *et al.*, “Optimization of nonlinear turbulence in stellarators,” *J. Plasma Phys.*, vol. 90, no. 2, p. 905900210, Apr. 2024, doi: 10.1017/S0022377824000369.
- [23] M. Landreman *et al.*, “How does ion temperature gradient turbulence depend on magnetic geometry? Insights from data and machine learning,” *Journal of Plasma Physics*, vol. 91, no. 4, p. E120, Aug. 2025, doi: 10.1017/S0022377825100536.
- [24] Y. Zhou, K. Aleynikova, C. Liu, and N. M. Ferraro, “Benign Saturation of Ideal Ballooning Instability in a High-Performance Stellarator,” *Phys. Rev. Lett.*, vol. 133, no. 13, p. 135102, Sep. 2024, doi: 10.1103/PhysRevLett.133.135102.
- [25] J. H. Nicolau, X. Wei, P. Liu, G. Choi, and Z. Lin, “Trapped electron mode in a quasi-isodynamic stellarator,” *Nucl. Fusion*, vol. 65, no. 8, p. 086049, Jul. 2025, doi: 10.1088/1741-4326/adf1b7.
- [26] H. Chen, X. Wei, H. Zhu, and Z. Lin, “Geometry effects on zonal flow dynamics and turbulent transport in optimized stellarators,” *Nucl. Fusion*, vol. 65, no. 7, p. 074002, Jun. 2025, doi: 10.1088/1741-4326/adde6f.
- [27] H. E. Mynick, N. Pomphrey, and S. Ethier, “Exploration of stellarator configuration space with global search methods,” *Physics of Plasmas*, vol. 9, no. 3, pp. 869–876, Mar. 2002, doi: 10.1063/1.1445756.
- [28] Z. Bai, X. Wei, W. Tang, L. Oliner, Z. Lin, and S. Williams, “Transfer learning nonlinear plasma dynamic transitions in low dimensional embeddings via deep neural networks,” *Mach. Learn.: Sci. Technol.*, vol. 6, no. 2, p. 025015, Apr. 2025, doi: 10.1088/2632-2153/adca83.
- [29] G. E. Hinton and R. R. Salakhutdinov, “Reducing the Dimensionality of Data with Neural Networks,” *Science*, 2006, doi: 10.1126/science.1127647.
- [30] A. Aghajanyan, S. Gupta, and L. Zettlemoyer, “Intrinsic Dimensionality Explains the Effectiveness of Language Model Fine-Tuning,” in *Proceedings of the 59th Annual Meeting of the Association for Computational Linguistics and the 11th International Joint Conference on Natural Language Processing (Volume 1: Long Papers)*, C. Zong, F. Xia, W. Li, and R. Navigli, Eds., Online: Association for Computational Linguistics, Aug. 2021, pp. 7319–7328. doi: 10.18653/v1/2021.acl-long.568.
- [31] L. Basile, S. Acevedo, L. Bortolussi, F. Anselmi, and A. Rodriguez, “Intrinsic Dimension Correlation: uncovering nonlinear connections in multimodal representations,” Mar. 02, 2025, *arXiv*: arXiv:2406.15812. doi: 10.48550/arXiv.2406.15812.
- [32] E. Altinisik, S. Messaoud, H. T. Sencar, H. Sajjad, and S. Chawla, “Explaining the role of Intrinsic Dimensionality in Adversarial Training,” in *Proceedings of the 42nd International Conference on Machine Learning*, PMLR, Oct. 2025, pp. 1298–1313. Accessed: Mar. 05, 2026. [Online]. Available: <https://proceedings.mlr.press/v267/altinisik25a.html>
- [33] M. N. Rosenbluth and F. L. Hinton, “Poloidal Flow Driven by Ion-Temperature-Gradient Turbulence in Tokamaks,” *Phys. Rev. Lett.*, vol. 80, no. 4, pp. 724–727, Jan. 1998, doi: 10.1103/PhysRevLett.80.724.
- [34] D. W. Dudt and E. Kolemen, “DESC: A stellarator equilibrium solver,” *Physics of Plasmas*, 2020, doi: 10.1063/5.0020743.
- [35] D. W. Dudt, R. Conlin, D. Panici, and E. Kolemen, “The DESC stellarator code suite Part 3: Quasisymmetry optimization,” *Journal of Plasma Physics*, vol. 89, no. 2, p. 955890201, Apr. 2023, doi: 10.1017/S0022377823000235.
- [36] H. Zhu, Z. Lin, and A. Bhattarjee, “Collisionless zonal-flow dynamics in quasisymmetric stellarators,” *J. Plasma Phys.*, vol. 91, no. 1, p. E28, Feb. 2025, doi: 10.1017/S0022377824001739.