

Integral Representations for Multiple Apéry-Like Series

Jorge Antonio González Layja*
Mexico

Abstract

By elementary methods, we obtain integral representations for several families of multiple Apéry-like series. These involve multiple harmonic sums in addition to the characteristic central binomial factors. The derivations are based on systematic applications of integration by parts together with two key Fourier expansions. As applications, we recover known evaluations and derive new identities.

Keywords: Apéry-like series; harmonic sums; central binomial coefficients; Fourier expansions; special functions

AMS Subject Classifications (2020): 11M32, 40C10, 11B65, 11M06

1 Introduction and Preliminaries

For $j \in \mathbb{Z}_{\geq 0}$, we consider the multiple harmonic star sums of depth j and weight $2j$ defined by

$$\zeta_n^*(\{2\}_j) := \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \prod_{i=1}^j \frac{1}{n_i^2},$$

and their odd counterparts

$$t_n^*(\{2\}_j) := \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \prod_{i=1}^j \frac{1}{(2n_i - 1)^2},$$

where $\zeta_n^*(\emptyset) = t_n^*(\emptyset) := 1$. These are finite analogues of the multiple zeta star values and multiple t -star values introduced by Hoffman in [7, 8], respectively.

We obtain, for $k \in \mathbb{Z}_{>0}$, integral representations for

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} t_n^*(\{2\}_j),$$

as well as for the related series

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \frac{4^{n_j}}{\binom{2n_j}{n_j}} \prod_{i=1}^j \frac{1}{n_i^2},$$

$$\sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j + 1) 4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i - 1)^2}.$$

These allow us to recover known evaluations and derive new ones. The resulting formulas are expressed in terms of the polylogarithm, the Legendre chi function, and the inverse tangent integrals. Moreover, our approach is entirely elementary. In particular, no generating-function techniques, hypergeometric transformations, or complex-analytic methods are required.

For completeness, we recall the special functions appearing in our results. The polylogarithm of order s is defined by

$$\text{Li}_s(x) := \sum_{n=1}^{\infty} \frac{x^n}{n^s}, \quad \begin{cases} |x| < 1, & s \in \mathbb{C}, \\ x = 1, & \Re(s) > 1, \\ x = -1, & \Re(s) > 0. \end{cases}$$

*Email: jorgelayja16@gmail.com

In addition, $\text{Li}_s(1) = \zeta(s)$ and $\text{Li}_s(-1) = -\eta(s)$, where ζ and η are the Riemann zeta and Dirichlet eta functions (see [6, p. 189]). It satisfies

$$\text{Li}_s(x) = \int_0^x \frac{\text{Li}_{s-1}(t)}{t} dt.$$

The Legendre chi function is defined by

$$\chi_s(x) := \sum_{n=1}^{\infty} \frac{x^{2n-1}}{(2n-1)^s}, \quad \begin{cases} |x| < 1, & s \in \mathbb{C}, \\ x = \pm 1, & \Re(s) > 1, \end{cases}$$

with $\chi_1(x) = \text{arctanh}(x)$ and $\chi_s(1) = \lambda(s)$, where λ denotes the Dirichlet lambda function (see [6, p. 189]). Moreover, it admits the representation $\chi_s(x) = \frac{1}{2}(\text{Li}_s(x) - \text{Li}_s(-x))$.

For $m \in \mathbb{Z}_{>0}$, the inverse tangent integrals are defined by

$$\text{Ti}_m(x) := \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{2n-1}}{(2n-1)^m}, \quad |x| \leq 1,$$

with $\text{Ti}_1(x) = \text{arctan}(x)$ and $\text{Ti}_m(1) = \beta(m)$, where β denotes the Dirichlet beta function (see [6, p. 190]).

Similarly to the polylogarithm, the Legendre chi function and the inverse tangent integrals satisfy

$$\chi_s(x) = \int_0^x \frac{\chi_{s-1}(t)}{t} dt \quad \text{and} \quad \text{Ti}_m(x) = \int_0^x \frac{\text{Ti}_{m-1}(t)}{t} dt.$$

We now collect several lemmas that will be used to prove the main theorems.

2 Lemmas

Lemma 2.1. *The following identities hold:*

(i) For $m \in \mathbb{Z}_{\geq 0}, n \in \mathbb{Z}_{> 0}$,

$$\int_0^{\frac{\pi}{2}} x^{2m} ((-1)^{n-1} + \cos(2nx)) dx = (2m)! \sum_{j=0}^m \frac{(-1)^{j+n-1}}{(2n)^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1}.$$

(ii) For $m \in \mathbb{Z}_{\geq 0}, n \in \mathbb{Z}_{> 0}$,

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} x^{2m-1} ((-1)^{n-1} + \cos(2nx)) dx \\ &= (2m-1)! \left(\sum_{j=0}^{m-1} \frac{(-1)^{j+n-1}}{(2n)^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} + (-1)^m \frac{1 - (-1)^n}{(2n)^{2m}} \right). \end{aligned}$$

(iii) For $m, n \in \mathbb{Z}_{\geq 0}$,

$$\int_0^{\frac{\pi}{2}} x^{2m} \frac{(-1)^{n-1} + \cos(2nx)}{\cos(x)} dx = 2(2m)! \sum_{j=0}^m \frac{(-1)^{j+n-1} O_n^{(2j+1)}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j}.$$

(iv) For $m \in \mathbb{Z}_{> 0}, n \in \mathbb{Z}_{\geq 0}$,

$$\int_0^{\frac{\pi}{2}} x^{2m-1} \frac{1 - \cos(2nx)}{\sin(x)} dx = 2(2m-1)! \sum_{j=1}^m \frac{(-1)^{j-1} \overline{O}_n^{(2j)}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j},$$

where $O_n^{(m)} := \sum_{k=1}^n \frac{1}{(2k-1)^m}$ and $\overline{O}_n^{(m)} := \sum_{k=1}^n \frac{(-1)^{k-1}}{(2k-1)^m}$ denote the odd harmonic numbers of order m and their alternating analogues, respectively.

Proof. Each part of this lemma follows by repeated integration by parts, together with straightforward rearrangements. For parts (iii) and (iv), we also use the identities (see [4, 1.342])

$$2 \sum_{k=1}^n (-1)^{n+k} \cos((2k-1)x) = \frac{(-1)^{n-1} + \cos(2nx)}{\cos(x)},$$

$$2 \sum_{k=1}^n \sin((2k-1)x) = \frac{1 - \cos(2nx)}{\sin(x)}.$$

□

Lemma 2.2. *The following identities hold:*

(i) For $m \in \mathbb{Z}_{\geq 0}, n \in \mathbb{Z}_{> 0}$,

$$\int_0^{\frac{\pi}{2}} x^{2m} \cos^{2n}(x) dx = (2m)! \frac{\binom{2n}{n}}{4^n} \sum_{j=0}^m \frac{(-1)^j \zeta_n^*(\{2\}_j)}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1}.$$

(ii) For $m \in \mathbb{Z}_{\geq 0}, n \in \mathbb{Z}_{> 0}$,

$$\int_0^{\frac{\pi}{2}} x^{2m} \cos^{2n-1}(x) dx = \frac{(2m)!}{2} \frac{4^n}{n \binom{2n}{n}} \sum_{j=0}^m \frac{(-1)^j t_n^*(\{2\}_j)}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j}.$$

(iii) For $m, n \in \mathbb{Z}_{> 0}$,

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} x^{2m-1} \cos^{2n}(x) dx \\ &= (2m-1)! \frac{\binom{2n}{n}}{4^n} \left(\sum_{j=0}^{m-1} \frac{(-1)^j \zeta_n^*(\{2\}_j)}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} + \frac{(-1)^m}{4^m} \sum_{n \geq n_1 \geq \dots \geq n_m \geq 1} \frac{4^{n_m}}{\binom{2n_m}{n_m}} \prod_{i=1}^m \frac{1}{n_i^2} \right). \end{aligned}$$

(iv) For $m, n \in \mathbb{Z}_{> 0}$,

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} x^{2m-1} \sin^{2n-1}(x) dx \\ &= \frac{(2m-1)!}{2} \frac{4^n}{n \binom{2n}{n}} \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j+1)4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i-1)^2}. \end{aligned}$$

Proof. All parts of this lemma are proved in the same manner. We first establish a recurrence relation by a straightforward application of integration by parts. Evaluating the first few cases reveals the general pattern, which is then verified using the recurrence by induction on m . We illustrate this strategy by proving the identity in part (i).

Let

$$I_{m,n_1} = \int_0^{\frac{\pi}{2}} x^{2m} \cos^{2n_1}(x) dx.$$

Applying integration by parts yields

$$\begin{aligned} I_{m,n_1} &= \int_0^{\frac{\pi}{2}} x^{2m} (\sin(x))' \cos^{2n_1-1}(x) dx \\ &= \frac{m}{n_1} \int_0^{\frac{\pi}{2}} x^{2m-1} (\cos^{2n_1}(x))' dx + (2n_1-1) \int_0^{\frac{\pi}{2}} x^{2m} \sin^2(x) \cos^{2n_1-2}(x) dx \\ &= -\frac{m(2m-1)}{n_1} I_{m-1,n_1} + (2n_1-1) I_{m,n_1-1} - (2n_1-1) I_{m,n_1}. \end{aligned}$$

From this we obtain

$$\frac{1}{2} \frac{m(2m-1)}{n_1^2} I_{m-1,n_1} = \frac{2n_1-1}{2n_1} I_{m,n_1-1} - I_{m,n_1}.$$

By multiplying both sides by $\frac{4^{n_1}}{\binom{2n_1}{n_1}}$, using the identity $\binom{2n_1}{n_1} = \frac{2(2n_1-1)\binom{2n_1-2}{n_1-1}}$, and summing from $n_1 = 1$ to n , we further obtain

$$\begin{aligned} \frac{m(2m-1)}{2} \sum_{n_1=1}^n \frac{4^{n_1}}{n_1^2 \binom{2n_1}{n_1}} I_{m-1, n_1} &= \sum_{n_1=1}^n \left(\frac{4^{n_1-1}}{\binom{2n_1-2}{n_1-1}} I_{m, n_1-1} - \frac{4^{n_1}}{\binom{2n_1}{n_1}} I_{m, n_1} \right) \\ &= I_{m,0} - \frac{4^n}{\binom{2n}{n}} I_{m,n}. \end{aligned}$$

Since $I_{m,0} = \frac{1}{2m+1} \left(\frac{\pi}{2}\right)^{2m+1}$, rearranging gives the recurrence

$$I_{m,n} = \frac{\binom{2n}{n}}{4^n} \left(\frac{1}{2m+1} \left(\frac{\pi}{2}\right)^{2m+1} - \frac{(2m)(2m-1)}{4} \sum_{n_1=1}^n \frac{4^{n_1}}{n_1^2 \binom{2n_1}{n_1}} I_{m-1, n_1} \right).$$

Evaluating the recurrence for the first few values of m yields

$$\begin{aligned} I_{0,n} &= \frac{\pi}{2} \frac{\binom{2n}{n}}{4^n}, \\ I_{1,n} &= \frac{\binom{2n}{n}}{4^n} \left(\frac{1}{3} \left(\frac{\pi}{2}\right)^3 - \frac{1}{2} \left(\frac{\pi}{2}\right) \sum_{n_1=1}^n \frac{1}{n_1^2} \right), \\ I_{2,n} &= \frac{\binom{2n}{n}}{4^n} \left(\frac{1}{5} \left(\frac{\pi}{2}\right)^5 - \left(\frac{\pi}{2}\right)^3 \sum_{n_1=1}^n \frac{1}{n_1^2} + \frac{3}{2} \left(\frac{\pi}{2}\right) \sum_{n_1=1}^n \frac{1}{n_1^2} \sum_{n_2=1}^{n_1} \frac{1}{n_2^2} \right). \end{aligned}$$

These initial cases suggest the stated formula in part (i), which is readily verified by induction on m using the recurrence.

The remaining parts follow by the same argument. \square

Lemma 2.3. *Let $k \in \mathbb{Z}_{>0}$, $x \in \mathbb{R}$. Then*

$$\begin{aligned} \text{(i)} \quad \text{Li}_k(\cos^2(x)) &= \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1} \left(\frac{1+t}{2\sqrt{t}} \right) dt \right) \left((-1)^{n-1} + \cos(2nx) \right), \\ \text{(ii)} \quad \text{Li}_k(\sin^2(x)) &= \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} (-1)^{n-1} \left(\int_0^1 t^{n-1} \ln^{k-1} \left(\frac{1+t}{2\sqrt{t}} \right) dt \right) (1 - \cos(2nx)). \end{aligned}$$

Proof. The approach follows Vălean's strategy in [2, Sect. 6.49]. However, we employ the identity (see [4, 1.461.2])

$$\sum_{n=1}^{\infty} \left((-1)^{n-1} + \cos(2nx) \right) e^{-2ny} = -\frac{\cos^2(x) \tanh(y)}{\cos(2x) - \cosh(2y)},$$

which considerably simplifies the calculations.

By substituting $t = e^{-2y}$, we obtain

$$\frac{1}{2} \int_0^1 t^{n-1} \ln^{k-1} \left(\frac{1+t}{2\sqrt{t}} \right) dt = \int_0^{\infty} \ln^{k-1}(\cosh(y)) e^{-2ny} dy.$$

Multiplying both sides by $(-1)^{n-1} + \cos(2nx)$ and summing over $n \geq 1$ gives

$$\begin{aligned} &\frac{1}{2} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1} \left(\frac{1+t}{2\sqrt{t}} \right) dt \right) \left((-1)^{n-1} + \cos(2nx) \right) \\ &= \int_0^{\infty} \ln^{k-1}(\cosh(y)) \left(\sum_{n=1}^{\infty} \left((-1)^{n-1} + \cos(2nx) \right) e^{-2ny} \right) dy \\ &= - \int_0^{\infty} \frac{\cos^2(x) \tanh(y) \ln^{k-1}(\cosh(y))}{\cos(2x) - \cosh(2y)} dy \\ &\stackrel{\cosh(y)=\frac{1}{t}}{=} \frac{(-1)^{k-1}}{2^{k+1}} \int_0^1 \frac{\cos^2(x) \ln^{k-1}(t)}{1 - \cos^2(x)t} dt. \end{aligned}$$

Hence, using the identity (see [1, Sect. 1.6])

$$\int_0^1 \frac{x \ln^n(t)}{1-xt} dt = (-1)^n n! \operatorname{Li}_{n+1}(x),$$

with $x \mapsto \cos^2(x)$ and $n \mapsto k-1$, the result in part (i) follows.

The identity in part (ii) follows from part (i) upon making the substitution $x \mapsto \frac{\pi}{2} - x$. \square

3 Main Results

Theorem 3.1. *Let $j \in \mathbb{Z}_{\geq 0}, k \in \mathbb{Z}_{>0}$. Then*

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) = -\frac{2^k}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) \operatorname{Li}_{2j}(-t)}{t} dt. \quad (3.1)$$

Proof. We first note that

$$\sum_{n=1}^{\infty} \frac{1}{n^k} \int_0^{\frac{\pi}{2}} x^{2m} \cos^{2n}(x) dx = \int_0^{\frac{\pi}{2}} x^{2m} \operatorname{Li}_k(\cos^2(x)) dx.$$

Applying part (i) of Lemmas 2.2 and 2.3 yields

$$\begin{aligned} (2m)! \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \sum_{j=0}^m \frac{(-1)^j \zeta_n^*(\{2\}_j)}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \\ = \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \int_0^{\frac{\pi}{2}} x^{2m} ((-1)^{n-1} + \cos(2nx)) dx. \end{aligned}$$

By part (i) of Lemma 2.1, we obtain

$$\begin{aligned} (2m)! \sum_{j=0}^m \frac{(-1)^j}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) \\ = (2m)! \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \sum_{j=0}^m \frac{(-1)^{j+n-1}}{(2n)^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \\ = (2m)! \sum_{j=0}^m \frac{(-1)^{j-1}}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \frac{2^k}{(k-1)!} \int_0^1 \left(\sum_{n=1}^{\infty} \frac{(-t)^n}{n^{2j}} \right) \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right)}{t} dt. \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{j=0}^m \frac{(-1)^j}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \left(\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) \right) \\ = \sum_{j=0}^m \frac{(-1)^j}{2^{2j} (2m-2j+1)!} \left(\frac{\pi}{2}\right)^{2m-2j+1} \left(-\frac{2^k}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) \operatorname{Li}_{2j}(-t)}{t} dt \right). \end{aligned}$$

Since the system is lower triangular with nonzero diagonal entries, comparing coefficients yields the desired identity. \square

As consequences of Theorem 3.1, we obtain the following corollary and proposition. Corollary 3.2 recovers a conjecture of Genčev and Rucki [9, Eq. 7.2], later proved by Xu [3, Thm. 2.2] by different methods, whereas Proposition 3.3 appears to be new.

Corollary 3.2. *Let $j \in \mathbb{Z}_{\geq 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) = 2 \eta(2j+1).$$

Proof. By setting $k = 1$ in (3.1), we have

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n 4^n} \zeta_n^*(\{2\}_j) = -2 \int_0^1 \frac{\text{Li}_{2j}(-t)}{t} dt = -2 \text{Li}_{2j+1}(-1).$$

Since $\text{Li}_s(-1) = -\eta(s)$, the claimed result follows. \square

Proposition 3.3. *Let $j \in \mathbb{Z}_{\geq 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^2 4^n} \zeta_n^*(\{2\}_j) = 2 \sum_{k=0}^{2j+2} (-1)^k \eta(k) \eta(2j - k + 2),$$

where we use the convention $\eta(0) = \frac{1}{2}$.

Proof. By setting $k = 2$ in (3.1), it follows that

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^2 4^n} \zeta_n^*(\{2\}_j) &= -4 \int_0^1 \frac{\ln\left(\frac{1+t}{2\sqrt{t}}\right) \text{Li}_{2j}(-t)}{t} dt \\ &= -4 \int_0^1 \frac{\ln(1+t) \text{Li}_{2j}(-t)}{t} dt + 2 \int_0^1 \frac{\ln(t) \text{Li}_{2j}(-t)}{t} dt + 4 \ln(2) \int_0^1 \frac{\text{Li}_{2j}(-t)}{t} dt \\ &= -4 \int_0^1 \frac{\ln(1+t) \text{Li}_{2j}(-t)}{t} dt + 2 \eta(2j+2) - 4 \ln(2) \eta(2j+1). \end{aligned} \tag{3.2}$$

Applying integration by parts to the remaining integral, we obtain

$$\int_0^1 \frac{\ln(1+t) \text{Li}_{2j}(-t)}{t} dt = -\eta(2) \eta(2j) + \int_0^1 \frac{\text{Li}_2(-t) \text{Li}_{2j-1}(-t)}{t} dt.$$

Iterating this $2j - 2$ times and using $\text{Li}_1(-t) = -\ln(1+t)$ yields

$$\begin{aligned} \int_0^1 \frac{\ln(1+t) \text{Li}_{2j}(-t)}{t} dt &= \sum_{k=0}^{2j-2} (-1)^{k-1} \eta(k+2) \eta(2j-k) - \int_0^1 \frac{\text{Li}_{2j}(-t) \ln(1+t)}{t} dt \\ &= \frac{1}{2} \sum_{k=2}^{2j} (-1)^{k-1} \eta(k) \eta(2j-k+2) \\ &= \frac{1}{2} \sum_{k=0}^{2j+2} (-1)^{k-1} \eta(k) \eta(2j-k+2) + \frac{1}{2} \eta(2j+2) - \ln(2) \eta(2j+1). \end{aligned} \tag{3.3}$$

Substituting (3.3) into (3.2) completes the proof. \square

Theorem 3.4. *Let $j \in \mathbb{Z}_{\geq 0}, k \in \mathbb{Z}_{> 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} t_n^*(\{2\}_j) = \frac{2^{k+3}}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t^2}{2t}\right) \text{Ti}_{2j+1}(t)}{1+t^2} dt. \tag{3.4}$$

Proof. By applying part (ii) of Lemma 2.2 and part (i) of Lemma 2.3 to

$$\sum_{n=1}^{\infty} \frac{1}{n^k} \int_0^{\frac{\pi}{2}} x^{2m} \cos^{2n-1}(x) dx = \int_0^{\frac{\pi}{2}} x^{2m} \frac{\text{Li}_k(\cos^2(x))}{\cos(x)} dx,$$

we obtain

$$\begin{aligned} &\frac{(2m)!}{2} \sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{j=0}^m \frac{(-1)^j t_n^*(\{2\}_j)}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \\ &= \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \int_0^{\frac{\pi}{2}} x^{2m} \frac{(-1)^{n-1} + \cos(2nx)}{\cos(x)} dx. \end{aligned}$$

Using part (iii) of Lemma 2.1 gives

$$\begin{aligned}
& \frac{(2m)!}{2} \sum_{j=0}^m \frac{(-1)^j}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} t_n^*(\{2\}_j) \\
&= (2m)! \frac{2^{k+1}}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1} \left(\frac{1+t}{2\sqrt{t}} \right) dt \right) \sum_{j=0}^m \frac{(-1)^{j+n-1} O_n^{(2j+1)}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \\
&\stackrel{t \rightarrow t^2}{=} (2m)! \frac{2^{k+2}}{(k-1)!} \sum_{j=0}^m \frac{(-1)^j}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \int_0^1 \left(\sum_{n=1}^{\infty} (-1)^{n-1} O_n^{(2j+1)} t^{2n-1} \right) \ln^{k-1} \left(\frac{1+t^2}{2t} \right) dt.
\end{aligned} \tag{3.5}$$

Furthermore, for $|t| < 1$, it is clear that

$$\begin{aligned}
\sum_{n=1}^{\infty} (-1)^{n-1} O_n^{(2j+1)} t^{2n-1} &= \sum_{n=1}^{\infty} (-1)^{n-1} t^{2n-1} \sum_{k=1}^n \frac{1}{(2k-1)^{2j+1}} \\
&= \sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2j+1}} \sum_{n=k}^{\infty} (-1)^{n-1} t^{2n-1} \\
&= \frac{1}{1+t^2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1} t^{2k-1}}{(2k-1)^{2j+1}} = \frac{\text{Ti}_{2j+1}(t)}{1+t^2}.
\end{aligned} \tag{3.6}$$

Substituting (3.6) into (3.5), we deduce that

$$\begin{aligned}
& \sum_{j=0}^m \frac{(-1)^j}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \left(\sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} t_n^*(\{2\}_j) \right) \\
&= \sum_{j=0}^m \frac{(-1)^j}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \left(\frac{2^{k+3}}{(k-1)!} \int_0^1 \frac{\ln^{k-1} \left(\frac{1+t^2}{2t} \right) \text{Ti}_{2j+1}(t)}{1+t^2} dt \right).
\end{aligned}$$

As in the proof of Theorem 3.1, the identity follows by comparing coefficients in the resulting lower triangular system. \square

The following proposition is the main result of [5], where it first appears and is obtained using less general identities.

Proposition 3.5. *Let $j \in \mathbb{Z}_{\geq 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{4^n}{n^2 \binom{2n}{n}} t_n^*(\{2\}_j) = 8 \sum_{k=0}^{2j} (-1)^k \beta(k+1) \beta(2j-k+1).$$

Proof. Setting $k=1$ in (3.4) yields

$$\sum_{n=1}^{\infty} \frac{4^n}{n^2 \binom{2n}{n}} t_n^*(\{2\}_j) = 16 \int_0^1 \frac{\text{Ti}_{2j+1}(t)}{1+t^2} dt.$$

As in the proof of part (iii) of Lemma 2 in [5], integration by parts gives

$$\int_0^1 \frac{\text{Ti}_{2j+1}(t)}{1+t^2} dt = \beta(1) \beta(2j+1) - \int_0^1 \frac{\arctan(t) \text{Ti}_{2j}(t)}{t} dt.$$

Iterating this $2j$ times and noting that $\text{Ti}_0(t) = \frac{t}{1+t^2}$, we obtain

$$\int_0^1 \frac{\text{Ti}_{2j+1}(t)}{1+t^2} dt = \sum_{k=0}^{2j} (-1)^k \beta(k+1) \beta(2j-k+1) - \int_0^1 \frac{\text{Ti}_{2j+1}(t)}{1+t^2} dt.$$

Since the final integral coincides with the original one, rearranging completes the proof. \square

Theorem 3.6. *Let $j \in \mathbb{Z}_{\geq 0}$, $k \in \mathbb{Z}_{> 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \frac{4^{n_j}}{\binom{2n_j}{n_j}} \prod_{i=1}^j \frac{1}{n_i^2} = \frac{2^{k+1}}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) \chi_{2j}(t)}{t} dt. \quad (3.7)$$

Proof. From

$$\sum_{n=1}^{\infty} \frac{1}{n^k} \int_0^{\frac{\pi}{2}} x^{2m-1} \cos^{2n}(x) dx = \int_0^{\frac{\pi}{2}} x^{2m-1} \text{Li}_k(\cos^2(x)) dx,$$

and parts (iii) and (i) of Lemmas 2.2 and 2.3, respectively, we obtain

$$\begin{aligned} & (2m-1)! \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \left(\sum_{j=0}^{m-1} \frac{(-1)^j \zeta_n^*(\{2\}_j)}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} + \frac{(-1)^m}{4^m} \sum_{n \geq n_1 \geq \dots \geq n_m \geq 1} \frac{4^{n_m}}{\binom{2n_m}{n_m}} \prod_{i=1}^m \frac{1}{n_i^2} \right) \\ &= \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \int_0^{\frac{\pi}{2}} x^{2m-1} ((-1)^{n-1} + \cos(2nx)) dx. \end{aligned}$$

From part (ii) of Lemma 2.1, it follows that

$$\begin{aligned} & (2m-1)! \left(\sum_{j=0}^{m-1} \frac{(-1)^j}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) \right. \\ & \quad \left. + \frac{(-1)^m}{4^m} \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \sum_{n \geq n_1 \geq \dots \geq n_m \geq 1} \frac{4^{n_m}}{\binom{2n_m}{n_m}} \prod_{i=1}^m \frac{1}{n_i^2} \right) \\ &= (2m-1)! \frac{2^k}{(k-1)!} \left(\sum_{j=0}^{m-1} \frac{(-1)^{j-1}}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \int_0^1 \left(\sum_{n=1}^{\infty} \frac{(-t)^n}{n^{2j}} \right) \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right)}{t} dt \right. \\ & \quad \left. + \frac{(-1)^m}{4^m} \int_0^1 \left(\sum_{n=1}^{\infty} \frac{t^n - (-t)^n}{n^{2m}} \right) \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right)}{t} dt \right) \\ &= (2m-1)! \frac{2^k}{(k-1)!} \left(\sum_{j=0}^{m-1} \frac{(-1)^{j-1}}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) \text{Li}_{2j}(-t)}{t} dt \right. \\ & \quad \left. + \frac{(-1)^m}{4^m} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) (\text{Li}_{2m}(t) - \text{Li}_{2m}(-t))}{t} dt \right). \end{aligned} \quad (3.8)$$

By Theorem 3.1 together with $\chi_s(x) = \frac{1}{2} (\text{Li}_s(x) - \text{Li}_s(-x))$, the right-hand side simplifies to

$$\begin{aligned} & (2m-1)! \left(\sum_{j=0}^{m-1} \frac{(-1)^j}{2^{2j} (2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n^k 4^n} \zeta_n^*(\{2\}_j) \right. \\ & \quad \left. + \frac{(-1)^m}{4^m} \frac{2^{k+1}}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) \chi_{2m}(t)}{t} dt \right). \end{aligned}$$

Therefore, cancelling the identical expressions on both sides of (3.8) and relabeling m as j yields the desired identity. \square

The next corollary recovers a second conjecture of Genčev and Rucki [9, Thm. 7.1].

Corollary 3.7. *Let $j \in \mathbb{Z}_{> 0}$. Then*

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n 4^n} \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \frac{4^{n_j}}{\binom{2n_j}{n_j}} \prod_{i=1}^j \frac{1}{n_i^2} = 4 \lambda(2j+1).$$

Proof. Setting $k = 1$ in (3.7), we have

$$\sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n 4^n} \sum_{n \geq n_1 \geq \dots \geq n_j \geq 1} \frac{4^{n_j}}{\binom{2n_j}{n_j}} \prod_{i=1}^j \frac{1}{n_i^2} = 4 \int_0^1 \frac{\chi_{2j}(t)}{t} dt = 4 \chi_{2j+1}(1).$$

The claim now follows from $\chi_s(1) = \lambda(s)$. \square

Theorem 3.8. *Let $j, k \in \mathbb{Z}_{>0}$. Then*

$$\sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j + 1) 4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i - 1)^2} = \frac{2^{k+3}}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t^2}{2t}\right) \chi_{2j}(t)}{1+t^2} dt.$$

Proof. Using

$$\sum_{n=1}^{\infty} \frac{1}{n^k} \int_0^{\frac{\pi}{2}} x^{2m-1} \sin^{2n-1}(x) dx = \int_0^{\frac{\pi}{2}} x^{2m-1} \frac{\text{Li}_k(\sin^2(x))}{\sin(x)} dx,$$

together with part (iv) of Lemma 2.2 and part (ii) of Lemma 2.3, we derive

$$\begin{aligned} & \frac{(2m-1)!}{2} \sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j + 1) 4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i - 1)^2} \\ &= \frac{2^k}{(k-1)!} \sum_{n=1}^{\infty} (-1)^{n-1} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \int_0^{\frac{\pi}{2}} x^{2m-1} \frac{1 - \cos(2nx)}{\sin(x)} dx. \end{aligned}$$

Applying part (iv) of Lemma 2.1, we obtain

$$\begin{aligned} & \frac{(2m-1)!}{2} \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j + 1) 4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i - 1)^2} \\ &= (2m-1)! \frac{2^{k+1}}{(k-1)!} \sum_{n=1}^{\infty} (-1)^{n-1} \left(\int_0^1 t^{n-1} \ln^{k-1}\left(\frac{1+t}{2\sqrt{t}}\right) dt \right) \sum_{j=1}^m \frac{(-1)^{j-1} \overline{O}_n^{(2j)}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \\ &\stackrel{t \rightarrow t^2}{=} (2m-1)! \frac{2^{k+2}}{(k-1)!} \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \int_0^1 \left(\sum_{n=1}^{\infty} (-1)^{n-1} \overline{O}_n^{(2j)} t^{2n-1} \right) \ln^{k-1}\left(\frac{1+t^2}{2t}\right) dt. \end{aligned} \tag{3.9}$$

In addition, for $|t| < 1$, we have that

$$\begin{aligned} \sum_{n=1}^{\infty} (-1)^{n-1} \overline{O}_n^{(2j)} t^{2n-1} &= \sum_{n=1}^{\infty} (-1)^{n-1} t^{2n-1} \sum_{k=1}^n \frac{(-1)^{k-1}}{(2k-1)^{2j}} \\ &= \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{(2k-1)^{2j}} \sum_{n=k}^{\infty} (-1)^{n-1} t^{2n-1} \\ &= \frac{1}{1+t^2} \sum_{k=1}^{\infty} \frac{t^{2k-1}}{(2k-1)^{2j}} = \frac{\chi_{2j}(t)}{1+t^2}. \end{aligned} \tag{3.10}$$

Thus, by substituting (3.10) into (3.9), we arrive at

$$\begin{aligned} & \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \left(\sum_{n=1}^{\infty} \frac{4^n}{n^{k+1} \binom{2n}{n}} \sum_{n \geq n_1 \geq \dots \geq n_{j-1} > n_j \geq 0} \frac{\binom{2n_j}{n_j}}{(2n_j + 1) 4^{n_j}} \prod_{i=1}^{j-1} \frac{1}{(2n_i - 1)^2} \right) \\ &= \sum_{j=1}^m \frac{(-1)^{j-1}}{(2m-2j)!} \left(\frac{\pi}{2}\right)^{2m-2j} \left(\frac{2^{k+3}}{(k-1)!} \int_0^1 \frac{\ln^{k-1}\left(\frac{1+t^2}{2t}\right) \chi_{2j}(t)}{1+t^2} dt \right). \end{aligned}$$

The desired identity follows by comparing coefficients in the resulting lower triangular structure. \square

References

- [1] C. I. Vălean, *(Almost) Impossible Integrals, Sums, and Series*, Springer, New York, 2019.
- [2] C. I. Vălean, *More (Almost) Impossible Integrals, Sums, and Series*, Springer, Cham, 2023.
- [3] C. Xu, On the proof of the Genčev–Rucki conjecture for multiple Apéry-like series, *arXiv preprint* arXiv:2510.09052, 2025.
- [4] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, 8th edition, Academic Press, New York, 2014.
- [5] J. A. G. Layja, On the Evaluation of Apéry-Like Series Involving Multiple t -Harmonic Star Sums, *arXiv preprint* arXiv:2601.20027, 2026.
- [6] L. Lewin, *Polylogarithms and Associated Functions*, North Holland, New York, 1981.
- [7] M. E. Hoffman, Multiple harmonic series, *Pacific Journal of Mathematics*, **152**(2):275–290, 1992.
- [8] M. E. Hoffman, An odd variant of multiple zeta values, *Communications in Number Theory and Physics*, **13**(3):529–567, 2019.
- [9] M. Genčev and P. Rucki, On a class of multiple Apéry-like series and their reduction, *Mediterranean Journal of Mathematics*, **22**:195, 2025.